Bilateral Alignment of Receptive Fields in the Olfactory Cortex

Julien Grimaud[1,2,3,4], William Dorrell[3,5], Cengiz Pehlevan[3,5], and Venkatesh Murthy[2,3]*.

[1] Molecules, Cells, and Organisms Graduate Program, Harvard University, Cambridge, Massachusetts, USA.
[2] Department of Molecular and Cellular Biology and Center for Brain Science, Harvard University, Cambridge, Massachusetts, USA.
[3] Center for Brain Science, Harvard University, Cambridge, Massachusetts, USA.
[4] Sup'Biotech Research Department, Villejuif, France.
[5] John A. Paulson School of Engineering and Applied Sciences, Harvard University, Cambridge, Massachusetts, USA.
*Lead Author. Correspondence: vnmurthy@fas.harvard.edu

Each olfactory cortical hemisphere receives ipsilateral odor
information directly from the olfactory bulb and contralateral
information indirectly from the other cortical hemisphere. Since
neural projections to the olfactory cortex are disordered and non-
topographic, spatial information cannot be used to align projections
from the two sides like in the visual cortex. Therefore, how bilateral
information is integrated in individual cortical neurons is unknown.
We have found, in mice, that the odor responses of individual
neurons to selective stimulation of each of the two nostrils are
highly matched, such that odor identity decoding optimized with
information arriving from one nostril transfers very well to the other
side. Remarkably, these aligned responses are nevertheless
asymmetric enough to allow decoding of stimulus laterality.
Computational analysis shows that such matched odor tuning is
incompatible with purely random connections but is explained
readily by Hebbian plasticity structuring bilateral connectivity. Our
data reveal that despite the distributed and fragmented sensory
representation in the olfactory cortex, odor information across the
two hemispheres is highly coordinated.
20

Introduction

In senses such as vision and audition, information from paired sensors
(eyes or ears) is combined in the brain to perform specific
computations that are behaviorally relevant – for example, estimating
distance to or the direction of the source of the stimuli (Cumming and
DeAngelis, 2001; Grothe and Pecka, 2014). The existence of two
sensors for olfaction has prompted hypotheses about their function
(Catania, 2013; Esquivelzeta Rabell et al., 2017; Mainland et al., 2002;
Porter et al., 2005; Rajan et al., 2006). Information from the two
nostrils can be differentiated, or integrated, depending on the goal of
such computation for the animal – for example, odor source
localization when smooth gradients exist (Álvarez-Salvado et al.,
2018; Baker et al., 2018; Catania, 2013; Esquivelzeta Rabell et al.,
2017; Gire et al., 2016), or identify relevant smells independent of the
odor source location or nostril stimulated (Bojsen-Moller and
Fahrenkrug, 1971; Dalal et al., 2020; Eccles, 2000; Kikuta et al., 2008;
Kucharski and Hall, 1987, 1988; Mainland et al., 2002). These
computations will dictate the nature of the inter-hemispheric
communication and the mechanisms of integration of bilateral
olfactory cues.

Odors are detected by olfactory sensory neurons in the nose, which
project to the ipsilateral olfactory bulb (OB). The OB sends
information to multiple ipsilateral brain regions, including the
anterior olfactory nucleus (AON) and the anterior and posterior
piriform cortices (APC and PPC, respectively) (Giessel and Datta, 2014;
Mori et al., 1999; Nagayama et al., 2014; Wilson and Sullivan, 2011)

(Figure 1A) In many species, including rodents, the two nares are
separated by a septum that minimizes transfer of inhaled odors
between nostrils (Eccles, 2000; Kikuta et al., 2008; Wilson and
Sullivan, 1999). Therefore, information transfer across hemispheres is
largely neural, and originates in the cortex (Bekkers and Suzuki, 2013;
Boyd et al., 2012; Haberly and Price, 1978; Kikuta et al., 2008; Scott et
al., 1980; Wilson and Sullivan, 2011) (Figure 1A).

Bilateral cortical interactions have been investigated in other senses
including vision, where the brain forms ordered representations of
the world using topographic maps, which allow neural matching
through precise point-to-point mapping (Cang and Feldheim, 2013;
Gu and Cang, 2016; Luo and Flanagan, 2007). However, the olfactory
cortex contains no recognizable topography (Bekkers and Suzuki,
2013; Ghosh et al., 2011; Illig and Haberly, 2003; Miyamichi et al.,
2011; Pashkovski et al., 2020; Sosulski et al., 2011; Stettler and Axel,
2009; Wilson and Sullivan, 2011), making point-to-point matching of
bilateral information through continuous maps implausible. It is
possible that odor representations in the two primary olfactory cortex
hemispheres are essentially independent, and downstream areas
align them for coherent perception, as implicitly proposed by Schaffer
et al 2018.

Framing hypotheses about bilateral olfactory processing requires
basic information about how single cortical neurons respond to the
same odor presented to the two nostrils. Such information is
fundamental for uncovering the computations performed, just as
details about binocular matching and disparity in cortical responses
were for vision (Blake and Wilson, 2011). Individual neurons could
respond to ipsi- and contra-lateral odor stimuli in a congruent or
disparate manner. There is some evidence for responses in olfactory
cortical areas to contralateral odor stimulation, but there is little
information on the alignment of ipsi- and contra-lateral responses
(Kikuta et al., 2008, 2010; Wilson, 1997). Recent evidence suggests
that mitral cells in the OB may have matched responses through
organized feedback projections from the anterior olfactory nucleus
(AON) par externa (Grobman et al., 2018; Yan et al., 2008). A cortical
neuron may then be able to inherit the matched response from mitral
cells (Dalal et al., 2020), but a direct evaluation of cortical matching is
lacking

To investigate how the olfactory cortex combines information from
the two nostrils, we recorded odor-evoked spiking activity of
individual neurons in the OC of awake mice. We observed neuron
responses to selective odor stimulation through the contralateral
nostril in all cortical regions recorded. A significant fraction of OC
neurons showed a strong match of ipsi- and contralateral response
profiles. Population activity in OC could be used to accurately decode
odor identity, as well as side identity. A computational model of the

olfactory system suggested that the bilateral overlap we observed in
mice is only possible with non-random bilateral projections. Our
results provide fundamental and novel insight that defines how ipsi-
and contralateral information streams converge and ultimately
enhance the processing capabilities of the brain.

6 Results

8 A New Method for Unilateral Odor Delivery

A major goal of this study was to compare the bilateral odor receptive
fields of OC neurons of awake mice. To do so, we needed a way to
reliably deliver odors to one nostril at a time. Previous studies have
used two different techniques to achieve such unilateral odor
stimulations. Either a piece of tubing was inserted in each nostril
(Wilson, 1997), or the two nostrils were separated by a plastic septum
(Kikuta et al., 2010, 2008). Since mice have very sensitive muzzles,
neither of these approaches is applicable to awake animals.

We developed a new facial mask for unilateral odor delivery in awake
mice (Figure 1B). On each side of the mask, close to the nostrils, we
used a computer-controlled olfactometer to deliver 15
monomolecular odors, one at the time, to the left or the right nostril
(Figures S1A-S1B). The mask was connected to an airflow sensor for
respiration monitoring (Figures S1C-S1F). We aligned all neuronal
odor-evoked activity to the first sniff after odor onset (see Methods).

We tested the side selectivity of the mask through two methods. First,
we simultaneously monitored the glomerular activity at the surface
of the OBs of OMP-GCaMP3 mice (Isogai et al., 2011) through calcium
imaging, while delivering odors to either nostril (Figure 1C) (n=155
glomeruli, 6 OBs, 3 mice). When odors were presented on the
contralateral nostril, glomeruli showed little to no response (Figures
1D-1F).

We also performed extracellular recordings of mitral/tufted cells
(M/T) in the OB of awake mice using chronically implanted tetrodes
(n=42 single units isolated from 4 mice) (Figure 1G, S1G-S1J, and Table
S1). M/T cells hardly responded to contralateral odor presentations:
out of the 42 isolated neurons, 32 responded to ipsilateral
stimulations, while 10 showed significant contralateral responses
(Figure 1H). Furthermore, while the ipsilateral M/T cell responses
were largely positive, the few contralateral responses were always
negative (Figure 1H).

The ratio of contra to ipsilateral response magnitude elicited by odors
presented within the mask was 0.12 in our glomerular imaging data
(ΔF/F magnitude, average ± SEM, ipsi: 7.9% ± 0.025%; contra: 0.93%
± 7.0e-3%) (Figure 1I) and 0.10 in our M/T recordings (response
magnitude, average ± SEM, ipsi: 1.6Hz ± 0.019Hz; contra: 0.16Hz ±
2.1e-3Hz) (Figure 1J). Therefore, on average, odor responses were 8
to 10 times stronger on the ipsilateral side of the mask, compared to
the contralateral side. Previous work has indicated that M/T cell
responses to a given odor at a concentration of 0.1% were quite
similar to the same odor at a concentration of 1% (Bolding and Franks,
2018, 2017). Since the M/T cell responses we observe are
substantially smaller for contralateral presentations, the effective
odor concentration in the ipsilateral nostril, when odors are
presented contralaterally, is likely to be much less than 10% of the
initial concentration from ipsilateral presentations.

Together, these results confirmed that odors, when delivered to one
nostril through our custom-built mask, activate sensory neurons in

Figure 1: A New Method for Unilateral Odor Delivery. (A) *Projection between the different olfactory regions. Only the OC (in particular: the AON and APC) sends direct contralateral projections.* **(B)** *Diagram of the odor mask. From left to right: front, rotated, and side view.* **(C)** *Calcium imaging setup.* **(D)** *Exemplar glomerular activity. Example: all glomeruli from OB1, isopropyl tiglate. Arrows: see below.* **(E)** *Exemplar glomerular activity over time. Grey: odor on (here: isopropyl tiglate). Blue: odor on the left nostril. Red: odor on the right nostril. Average ± SEM. Glomeruli 1 and 2 correspond to arrows 1 and 2 in panel 1D.* **(F)** *Glomerular responses. Each dot is the response of one glomerulus to one odor (glom-od pair). Histogram: contralateral responses. Red: Gaussian fit.* **(G)** *Tetrode recording setup.* **(H)** *Unilateral response profile of M/T cells. NOP: neuron-odor pair.* **(I)** *Magnitude of glomerular response. Red: average for each side. This panel presents the magnitude of the responses displayed in panel 1F.* **(J)** *Magnitude of M/T cell response. Only the ipsilaterally responding NOPs are shown. Red: average for each side.*

1 the other nostril at least an order of magnitude less than ipsilateral
2 activation.

Unilateral Odor Tuning of Different Regions of the Olfactory Cortex

We next used our unilateral odor delivery system to characterize how
the OC responds to unilateral odor presentations. To do so, we
monitored the activity of the AON and APC through extracellular
recordings in awake mice (n=7 mice, 1315 neurons total; AON: 3 mice,
384 neurons; APC: 4 mice, 931 neurons) (Figures 2A, S2, and Table
S1).

Overall, neurons in the OC had low spontaneous
activity (Figure 2B), in accordance with previous
studies (Bolding and Franks, 2017; Litaudon et al.,
2003; Zhan and Luo, 2010). Spontaneous activity
in the AON was significantly higher than in the APC
(average spontaneous activity ± SEM, AON: 3.9Hz
± 0.012Hz; APC: 2.8Hz ± 3.6e-3Hz; rank-sum test,
p<0.0001).

We found a diversity of responses to ipsi- as well
as contralateral odor stimulation in both the AON
and APC, including neurons that responded to
odors presented to either one or the other nostril,
or to both nostrils (Figures 2C-2D). Importantly, in
both OC regions, a sizeable fraction of neurons
responded to contralateral odor presentations
(fraction of neurons significantly responding to at
least one odor presented on the ipsilateral nostril,
AON: 86%; APC: 81%; contralateral side, AON:
87%; APC: 83%), expanding earlier anesthetized
observations to the awake OC (Kikuta et al., 2008,
2010; Wilson, 1997).

OC neurons typically responded to at most a few
odors from our odor panel (Figure 2E), as
previously reported with bilateral odor
stimulations (Bolding and Franks, 2017; Iurilli and
Datta, 2017). We found a wide range of response
magnitudes across the OC, and contralateral
responses were as strong as ipsilateral ones in the
APC, and slightly weaker in the AON (Figure 2F;
Wilcoxon signed-rank test, ipsi versus contra
distribution, AON: p=8.2e-3; APC: p=0.94).

An idiosyncratic feature of olfactory cortical
responses is that a neuron that responds to a
particular odor with an increase in firing rate,
tends to exhibit increases in firing rates for any
other odor it responds to; similarly, a neuron
having inhibitory response to an odor is also
inhibited by any other odor it responds to (Bolding
and Franks, 2017; Hu et al., 2017; Kikuta et al.,

2008; Otazu et al., 2015; Zhan and Luo, 2010). We find that this
feature applies to both cortical regions we examined (Figure 2G).

Overall, we found reliable unilateral odor responses to both the ipsi-
and contralateral nostrils in the OC.

Strong Bilateral Correlations in the Olfactory Cortex

57 Our data indicate that the OC receives robust contralateral odor
58 information, but it is unclear whether this information is similar to
59 that arriving ipsilaterally. To examine this issue, we related the

Figure 2: Odor Receptive Fields of Neurons in the Olfactory Cortex. (A) Experimental setup. (B) Basal activity per region. Circle and value next to it: median. Thick bar: quartiles. Thin bar: range of values. (C) Exemplar odor responses over time. Purple: odor delivery. Responses were aligned to the first sniff following odor onset (dotted line). PSTH: average ± SEM. Examples taken from mouse APC1 (see Table S1 for a detailed list of all mice recorded for this study). (D) Odor responses, illustration. Neurons randomly picked from mice AON1 (left) and APC1 (right). For each neuron, odors were ranked based on the response magnitude to ipsilateral presentations, which are shown in the left half of each plot. The contralateral responses are then shown with the same rank order of odors for each neuron. Each dot gives the average response ± SEM across the 7 trials, and significant responses are colored red. Asterisk: the ipsi- and contralateral tuning curves of these neurons are visually similar. Further analysis shows that their odor responses are bilaterally correlated. See Figure 3A for details. (E) Distribution of the number of odors eliciting a significant response in a given neuron, per region and side. (F) Distribution of significant response magnitudes for all neurons and odors, per region and side. (G) Positive-negative balance, per region and side. 0 = all negative responses; 1 = all positive.

average odor-evoked responses to ipsilateral
against contralateral presentations.

First, we plotted the ipsi- versus contralateral odor
responses for each neuron individually. Given the
prevalent view of OC connectivity as largely
random, we expected OC neurons to show little to
no match between their ipsi- and contralateral
odor response profiles. To our surprise, we found
many neurons with bilaterally-correlated odor
responses (Figure 3A) (F-test, significance of the
linear model, from left to right neuron, top row:
p=0.41; p=0.84; p=0.54; p=0.59; bottom row:
p=5.3e-6; p=5.5e-4; p=6.6e-3; p=2.2e-4) (see also
Figure 2D, neurons with an asterisk). Overall, for
each mouse recorded in the OC, we consistently
found a sizeable fraction of bilaterally-correlated
neurons (Figure 3B).

We used a bootstrap strategy to determine that the
fraction of bilaterally-correlated neurons in each
mouse was significantly higher than what one
would expect from chance. In brief, for each mouse,
each neuron, we randomly shuffled odor identities
on each nostril separately. We repeated this
shuffling for all neurons and calculated the resulting
percentage of bilaterally-correlated neurons. We
reiterated this procedure 10,000 times in order to
build a "chance" distribution. Finally, we compared
this distribution with the actual fraction of
bilaterally-correlated neurons. In all mice recorded
in the OC, the fraction of bilaterally-correlated
neurons was greater than one would expect from chance (Figures 3B,
3C, and S3A-S3D).

While our initial estimates of the fraction of bilaterally-correlated
neurons include all neurons recorded (Figures 3B, S3A), we verified
that our conclusions remained true when selecting only the
significantly responding (Figures S3B-S3C).

We performed additional analysis to determine the extent to which
response variability masks response correlations. We generated
individual trials from a Poisson process with rates determined from
our recordings. We then enquired how correlated responses would
be for two independent sets of 7 trials generated with the same set
of data. A large number of such sampling revealed that the fraction of
(simulated) neurons that show significant response correlations was
on the order of 40-80% even when the ideal response correlations
should be 100% (Figure S3D). This analysis indicates that the fraction
of significantly correlated neurons in the AON observed in our
recordings may underestimate the real fraction of bilaterally-
correlated neurons.

Overall, the bilateral correlations were strong (Figure 3D), with mostly
positive slopes (Figure 3E), and a distribution of intercepts centered
on zero (Figure 3F). In other words, bilaterally-correlated neurons
appear to respond to unilateral odor presentations in a mostly side-
invariant way, with their ipsi- and contralateral odor responses similar
in sign and amplitude.

We wondered if the bilaterally matched responses occurred
preferentially in a particular neuron type. We used a classic method
for identifying putative excitatory and inhibitory neurons – the width
of extracellular action potentials (Frank et al., 2001; Gur et al., 1999;

Figure 3: Side-Invariant Odor Responses in the Olfactory Cortex. *For panels (A) and (G), red line: linear regression (contra = a x ipsi + b). Solid lines show significant regressions (F-test, p < 5%). **(A)** Exemplar ipsi- versus contralateral responses per neuron. Examples picked from AON1. Each dot is a different odor. The neurons in the bottom panels are bilaterally-correlated. Asterisk: same neuron as in Figure 2D, left panel, red asterisk. **(B)** Percentage of bilaterally-correlated neurons (BCNs) per mouse. Each dot is a mouse. Fractions next to each dot: number of BCNs / total number of neurons. Full circle: the fraction is significantly higher than chance. **(C)** Significance of the percentages of BCNs. Here we show the distribution of chance percentages for AON1, obtained with a bootstrapping procedure. **(D)** to **(F)** Coefficients of determination, slopes, and intercepts of the BCNs in the AON and APC (Nb: number). **(G)** Ipsi- versus contralateral responses per region for odor 1 (isopropyl tiglate). Each dot represents one neuron. For the equivalent graphs across all odors, see Figure S4.*

Hassani et al., 2009; Povysheva et al., 2006; Suzuki and Smith, 1985;
Swadlow, 2003; Weir et al., 2015). Most of the bilaterally-correlated
neurons in the AON and APC had broader spikes and lower
spontaneous activity (Figure S3E-S3H), suggesting they are excitatory.

When testing our mask for cross-contamination (Figure 1) we
estimated that at most 10% of odors puffed on one side of the mask
could leak to the other side. Can this cross-contamination alone
explain the side-invariant responses we observed in the OC? To test
this hypothesis, we first determined how much each odor from our
panel needed to be diluted in the solvent to elicit a PID signal 10 times
weaker than our initial dilution (5% volume/volume) (Figure S3I). We
then recorded individual neurons in the AON of awake mice (n = 2
mice, 120 neurons) while delivering odors on either side of the mask,
both at initial (5%) and new dilutions (i.e. dilutions resulting in PID
signals 10 times weaker) (Figure S3J, Table 1). On average, for a given
side of the mask, new dilutions elicited neuronal responses 6 to 7
times weaker than our initial dilutions (neuronal response, mean ±
SEM, for ipsi presentations, initial dilution: 2.1Hz ± 0.023Hz, final
dilution: 0.34Hz ± 2.7e-3Hz; for contra presentations, initial dilution:
2.5Hz ± 0.031Hz, final dilution: 0.37Hz ± 2.5e-3Hz), and the correlation
between the two responses was significant (linear regression, ipsi
presentations: R^2=0.18, p<0.0001; contra presentations: R^2=0.12,
p<0.0001). In other words, contralateral cross-contamination in the
mask, if any, is not sufficient to explain the presence of side-invariant
odor responses.

Finally, we compared the ipsilateral and contralateral population
representation in the two OC regions recorded for each odor (Figures
3G and S4A-S4C). Consistent with our analysis of individual bilaterally-
correlated neurons, we found that unilateral population

representations were significantly and strongly correlated for all
odors tested in the AON and APC (Figures S4A-S4B) (see Table S2 for
detailed statistics). Analyzing population correlations for each odor
individually ensures that the correlations we observe are not driven
by few neurons responding strongly and similarly to all odors, which
would otherwise create an artificially strong positive correlation.
Furthermore, a bootstrap analysis similar to what we described above
confirmed that strong correlation we observed for each odor
individually were unlikely to occur by chance (Figures S4C-S4D). When
considering each mouse separately, our conclusions held true for
most odors (Figure S4E).

Together, these data indicate that the odor information arriving
contralaterally to the OC strongly matches that arriving ipsilaterally.

15 Accurate Decoding of Odor Identity from Contralateral
16 Activity

Odor selectivity in the OC allows a population representation of odor
identity, and odor identity can be decoded from the activity of a
sufficient number of neurons (Bolding and Franks, 2017). If odor
responses are similar for ipsi- and contralateral presentations,
decoding of identity should be transferrable. To formally test this, we
asked a linear classifier to decode odor identity from OC responses to
stimuli in one nostril, after we trained it with OC responses to stimuli
in the opposite nostril.

In brief, at each instance we created a training set by randomly
removing one trial per odor (Bolding and Franks, 2017). If we were
testing on the same side as we trained (e.g. train ipsilaterally, test
ipsilaterally), then we tested the classifier on the trials outside of the
training set. If we were testing on the opposite side (e.g. train
ipsilaterally, test contralaterally), then one trial per odor from the
opposite side was randomly chosen to form the test set (Figure 4A).
We performed all our decoding analyses over an odor response
window of 2s, as it maximized performance of odor decoding on the
same side (Figure S5A).

Figure 4: Decoding Odor Identity and Laterality. (A) Diagram of the odor decoding process. *(B)* and *(C)* Odor identity decoding per region. Mean ± SEM over 200 repetitions of the decoding process. For (B), the decoder was trained with data from ipsilateral presentations. For (C), the decoder was trained on contralateral odor responses. *(D)* Side identity decoding per region. Mean ± SEM over 200 repetitions of the decoding process. For (B) to (D), grey dotted line: chance.

In both the AON and APC, the decoder performed with similarly high
levels of accuracy when tested on data from the same or the opposite
side (Figures 4B-4C). In other words, odor identity decoding optimized
with responses to stimulation from one nostril transfers very well to
stimulation of the other side.

Note that the seeming decrease of decoding performance from AON
to APC is most probably due to variability across mice (Figure S5B).
More precisely, the three mice with overall weaker and fewer odor-
evoked responses (i.e. mice AON2, APC1, and APC2) showed lower
odor decoding performances, no matter the side tested (Figure S6).
In addition, AON2 is the mouse from which we isolated the fewest
neurons (73 neurons, while all other OC recorded mice have over a
hundred neurons each) (Table S1). Predictably, in these same three
mice, the fractions of bilaterally-correlated neurons, while
significantly higher than chance, were the lowest.

Rodents are able to determine the direction of the odor stimulation
at short distances (Esquivelzeta Rabell et al., 2017; Rajan et al., 2006).
We asked whether population activity in a single hemisphere differs
sufficiently for stimulation of ipsi- and contralateral nostril to allow
decoding of intensity differences between the two sides. Side
identity, or laterality, could indeed be decoded with high accuracy
(Figure 4D), which confirms that odor representations from both
sides, despite similarities, remain distinct and separable.

Together, our data indicate that odor decoding is transferrable across
sides. Ipsi- and contralateral odor representations, while distinct,
share a high degree of similarity.

62 A Computational Model of Bilateral Matching in the Olfactory
63 Cortex

64 How is one OC responding so similarly to inputs from two distinct
65 routes/nostrils? So similarly, in fact, that a linear classifier trained on
66 one set of representations can decode the other?

67 An appealingly simple resolution might suggest itself based on recent
theoretical work. It has been shown that if two correlated
representations are projected through the same random matrix the
output representations are also correlated (Babadi and Sompolinsky,
2014; Schaffer et al., 2018). Thus, one could hypothesize that any
interhemispheric projection between two OCs could preserve the
structure inherited from the underlying OB representations, hence
explaining the similarity. However, this argument is insufficient, as
illustrated in Figure 5. The correlations in glomerular neural
representations of the 15 odors used in our study are shown in Figure
5B. These correlations are partially preserved when projected
through two different random matrices (to represent the OB to OC
projections of each hemisphere) (Figure 5C, upper and lower block
diagonals). However, the two OC hemispheres connected with
random projections will not yield correlated responses to the same
odor presented to the left and right side (Figure 5C). In the
supplementary material we show that, even in the limiting case
where the response of ipsi- and contralateral cortices are identical,
unstructured interhemispheric connectivity ensures that the
responses of one cortex to odors presented in different nostrils will
be uncorrelated. Hence, there must exist some structured
connectivity that is aligning the cortical responses (Figure 5D; note
the much higher similarity values in the two off-diagonal blocks).

We then asked what type of structured interhemispheric connectivity
is sufficient to achieve the observed tight alignment of responses
from different nostrils. Through computational modelling we tested a

Figure 5: Structure is necessary for alignment between responses to different nostrils.
(A) Odor tuning of 50 randomly selected glomeruli. (B) The OB odor-odor cosine similarity matrix measured using calcium imaging. Color scale at right applies to panels (B) to (D). (C) We modeled the two OCs and connected them using a random matrix (see methods), the correlations between odors from the same nostril are preserved, but correlations between nostrils are completely lost (quadrants pointed to by red arrows and red dotted squares). (D) Including a structured cross-cortical connectivity produces correlations between ipsi- and contralateral odors.

S7C).

We then linked two simulated AONs using the connectivity matrix G. The computational analogue of the cortical representations of an odor presented contralaterally is the representation after mapping through the interhemispheric matrix, G. Thus, by comparing the simulated and measured alignment of ipsi- and contralaterally presented odors as we vary α, we can ask if this simple Hebbian structure is sufficient to recapture the degree of alignment observed.

We compared the alignment using four metrics: the overall correlation of odor population representation across sides (Figures 6E-6F; similar to Figure 3G), the fraction of bilaterally-correlated neurons (Figure 6G; similar to Figure 3B), the accuracy of odor identity decoding (Figures 6H-6I and S7D; similar to Figures 4B-4C), and the accuracy of side identity decoding (Figure 6J; similar to Figure 4D).

The four computed measurements best matched our actual AON neuronal recordings with a non-zero α (approximately 0.4). This result did not vary when we changed the size of the modeled OC population (Figures S7E-S7H). This suggests that a significantly Hebbian connectivity between cortices is sufficient to explain the observations.

We chose this particular structured connectivity, the 15 odor Hebbian matrix $G_{Structured}$, for its appealing simplicity and biological plausibility. For example, early in life the animal might have smelt these odors in both nostrils and associatively matched the responses between hemispheres. However, there are many other sufficient alignment schemes that do not rely on having experienced specifically these 15 odors; we implemented two such examples in the supplementary material. The first uses a set of Hebbian matching experiences that span the subspace in which the 15 odorants lie. Roughly 6 of these are needed to achieve similar

simple and plausible model for structured matching, a Hebbian
connectivity matrix that links together the same-odor
representations in each hemisphere. We combined this Hebbian
matrix, ($G_{Structured}$), with a random matrix, (G_{Random}), which models
unstructured elements of connectivity and whose elements are
drawn independently from a Gaussian distribution. We vary the
degree of structure using a parameter α. Explicitly:

8
$$G = \alpha G_{Structured} + (1-\alpha)G_{Random}$$

The parameter α took values between 0 and 1 with α = 0
corresponding to purely random connectivity and α = 1 to purely
structured connectivity. The random and Hebbian components were
scaled appropriately such that α represents the proportion of
contralateral input magnitude that is formed by structured
connectivity.

Our modeling strategy began by creating a cortical representation of
odors presented in the ipsilateral nostril, based on the AON. Random
OB-to-OC connectivity indicates that each cortical olfactory neuron
samples a random subset of glomeruli. This implies that the input to
each cortical neuron is statistically independent, and that the
response of one cortical neuron to different odorants can be
modelled as a thresholded multivariate gaussian whose covariance
matrix can be extracted from activities in the OB (Figure 6A, see
Methods). We validated this model of a single AON against neuronal
recordings: our model replicated the population statistics remarkably
well with minimal parameter adjustment (Figures 6B-6D; Figures S7A-

levels of alignment (Figure S8G-S8J) showing that a smaller number of
more general early learning experiences could also explain the
alignment seen in the data. The second technique uses Singular Value
Decomposition to create a low rank approximation of $G_{Structured}$; again,
a rank of about 7 is necessary to match the data (Figure S8C-S8F).
Effectively, any connectivity that links same-odor representations in
each hemisphere together will be able to match the alignment seen
in the data.

68 Discussion

In many animals, odor information collected by the two nostrils
remains separate until it reaches secondary brain areas (Bojsen-
Moller and Fahrenkrug, 1971; Eccles, 2000; Kikuta et al., 2008). How
the two streams of information are integrated, as well as the
functional role of this integration, remain unclear.

Here, we reveal that a sizeable fraction of neurons in the OC are
bilaterally-correlated, meaning that they have matched ipsi- and
contralateral odor tuning. Functional alignment of responses was also
confirmed by the ability to decode odor identity from neural
populations trained using only odor responses from the opposite
nostril. Finally, our computational model suggests that such bilateral
matching requires structured interhemispheric projections. Our
findings shed light on how brains make sense of this bilateral
experience in the olfactory system.

Figure 6: Modeling Olfactory Cross Cortical Connections. (A) *Diagram of the model. (B)* to *(D) Model validation. Contralateral odor responses from the model are overlaid with actual contralateral AON responses (Figures 2E-2G). (E) to (J) Various measurements of cross-cortical alignment from the model. (E) Contra- vs ipsilateral responses for random (alpha = 0) and semi-structured (alpha = 0.42) projections. (F) Contra- vs ipsilateral response correlations R for various alpha values. (G) Percentage of bilaterally-correlated neurons for various alpha values. (H) Odor decoding accuracy for random (alpha = 0) and semi-structured (alpha = 0.42) projections. Red: same as Figure 3B, AON, "test contra". (I) Odor decoding accuracy for various alpha values. (J) Side decoding accuracy for various alpha values. For (E) to (H), measurements from the model were calculated by sampling 385 neurons 200 times. The best fit alpha (0.42) was found by comparing to AON data (n = 385 neurons from 3 mice).*

nostril-independent recognition of odors: rats trained to associate odors with reward using only one naris can generalize to stimuli delivered through the other nostril (Kucharski and Hall, 1988, 1987; Yan et al., 2008). This behavior requires interhemispheric communication, suggesting a convergence of ipsilateral and contralateral representations in higher brain areas (Dalal et al., 2020).

Robust Contralateral Odor Responses in the Olfactory Cortex

Here, we developed and tested a new method to deliver odor stimuli selectively to individual nostrils in awake, freely breathing mice (Figure 1). We performed several experiments to confirm selective stimulation. First, PID recordings indicated that odors were delivered at the same concentration on either side and are not detectable on the side contralateral to the stimulus. Second, direct measurements of sensory input to the OB indicated that glomerular responses to contralateral stimulation were below detection threshold. It is possible that some low-level cross contamination went undetected by the PID and glomerular recordings, but it is unlikely that such weak inputs could contribute to contralateral responses that are as strong as those to ipsilateral

stimulation, as shown by our recordings in AON to 10-fold dilute
odors (Figure S3J).

Third, recordings from putative M/T cells in the OB using tetrodes
revealed few detectable responses to contralateral nares stimulation.
These results seem at odds with a recent report suggesting that
topographic projections from contralateral AONpE to M/T cells
support bilaterally matched odor responses (Grobman et al., 2018). A
possible source of the discrepancy is the much lower concentration
of odor used in our mask. However our observations confirm former
work from various teams regarding the near absence of contralateral
M/T response in the OB (Kikuta et al., 2008, 2010; Wilson, 1997). Our
results are also consistent with the known projections from the AON
pars externa to the GABAergic granule cells of the contralateral OB
(Schoenfeld and Macrides, 1984; Yan et al., 2008), since the
infrequent significant responses we observed in M/T cells to
contralateral nostril stimulation were inhibitory. The minimal
responses in glomeruli and M/T cells also argues strongly against
bilateral return of odors from the respiratory system during
exhalation.

80 Bilateral Integration in the Olfactory Cortex is not Random

Our first key finding is that responses to the stimulation of the
contralateral nostril could readily be detected in the AON and APC
(Figure 2). This is not entirely surprising, since crossed projections
already occur at the level of the AON and APC. Previous studies
indicated that contralateral stimulation evoked fewer and weaker
responses in the AON (Kikuta et al., 2008). However, these studies

1 Importance of the Integration of Bilateral Information

In many senses, information is collected through a pair of sensors and
is integrated within the brain. Bilateral perception allows us to
perform vital behavioral computations such as sound localization or
depth perception. Since the two nostrils may sample independent
pockets of air (Eccles, 2000; Kikuta et al., 2008; Wilson and Sullivan,
1999), can bilateral information be used to discover features of the
odor stimulus not readily available to single nostrils? Animals may
compare of odor concentration across both nostrils to locate odor
sources (Catania, 2013; Esquivelzeta Rabell et al., 2017; Gire et al.,
2016; Parthasarathy and Bhalla, 2013), which requires bilateral
integration and comparison of different signals. Two brain regions
that seem to fulfil this criterion are the AON and APC (Kikuta et al.,
2008, 2010; Wilson, 1997). Recent studies have indicated that the
AON may be sensitive to the differential activation of left and right
nostrils, therefore computing the direction of the stimulus in front of
the animal (Esquivelzeta Rabell et al., 2017). Our own data shows that
side identity can be decoded from odor responses, which provides
further evidence for differential processing (Figure 4D).

In contrast, many behaviors require the animal to make inferences
independent of the nostril stimulated. One simple example can be
found in our everyday life: one nostril is usually more blocked than
the other, and which nostril gets clogged changes several times a day
(Eccles, 2000). Nevertheless, we can identify smells and have a unified
olfactory experience. In the same fashion, animals need to be able to
identify relevant smells like food or a predator, no matter where the
odor source is relative to their nose. Indeed, there is evidence for

recorded from a small number of neurons in anesthetized animals. While we confirmed the presence of weaker and fewer responses to contralateral stimulations in the AON, the differences were much smaller than previously reported. Another recent study implicated AON in interhemispheric communication using a variety of methods, but the responses of neurons in this area were not measured (Yan et al., 2008). Finally, one previous study reported the presence of neurons responsive to contralateral naris stimulations in the APC of anesthetized rats, but did not test the correlation of the responses between the two nares (Wilson, 1997).

We found that the odor information arriving ipsilaterally to the OC strongly matches that arriving contralaterally (Figure 3). Our control experiments indicated that less than 10% of an odor delivered to one nostril may leak to the other nostril, and we found that this potential 10% leak is not sufficient to explain the presence of bilateral neurons. Furthermore, our data offer strong hints that principal neurons are a major part of the ensemble of neurons that have matched responses to stimuli from both nostrils (Figure S3). This feature would allow downstream areas receiving olfactory cortical inputs to readily have access to side-stable odor responses. Indeed, we showed explicitly that odor decoding is transferrable across sides (Figure 4), as expected from the presence of matched responses in the OC.

Finally, we considered how the prevalent view of OC connectivity could explain our data (Figure 5). Recent studies have shown that two representations projected through the same random matrix remain correlated (Babadi and Sompolinsky, 2014; Schaffer et al., 2018). In addition, another study reported that population response correlations between odors is shaped in the cortex, presumably by experience, such that some response similarities in the OB are enhanced and others are suppressed (Pashkovski et al., 2020). However, these studies do not address alignment due to projections between cortices. Here we implemented a computational model of the olfactory system to test matching of ipsilateral and contralateral odor tuning in the OC as a function of structure in bilateral projections. We showed that, in our model, purely random projections cannot explain our OC results. That is, even if the two hemispheres have the same ordered representations for similar odors, unstructured interhemispheric cortical projections will disrupt these correlations.

Reconciling Our Results with Known Connectivity Schemes

How can the responses of cortical neurons to selective stimulation of each of the two nostrils be matched? First, there may be topographically matched crossed projection somewhere in the olfactory system allowing bilateral response matching. The only known candidate for such matching is the projection from AON pars externa to OB (Schoenfeld and Macrides, 1984; Yan et al., 2008). This projection, however, should result in net inhibition of M/T cells since its major target are the GABAergic granule cells (Schoenfeld and Macrides, 1984; Yan et al., 2008) (but see Grobman et al., 2018). Furthermore, previous studies (Kikuta et al., 2010, 2008; Wilson, 1997) and our own data (Figure 1) suggest that this projection is too weak to support the widespread bilateral matching we observed in the OC. However, M/T cells that respond to contralateral odor presentation in a correlated way (Grobman et al 2018) may report to ipsilateral OC neurons the same information for ipsi- and contralateral stimulation under some conditions, for example, high odor concentrations. Even so, unstructured interhemispheric cortical projections will disrupt this potential alignment in the cortex. Therefore, additional mechanisms, such as Hebbian plasticity, are needed to explain our observations.

A second way in which matched responses may arise is through fine-scale order in anatomical connectivity that somehow allows the same glomerular output channels from both OBs to converge on individual neurons. Such structure seems implausible and has no precedent but remains formally possible.

A third explanation that seems likely because of its simplicity and local nature is that synaptic connectivity in cortical regions is refined by Hebbian plasticity (Best and Wilson, 2003; Franks and Isaacson, 2005; Johenning et al., 2009) to allow matched responses from both sides. During their normal experience, mice will frequently inhale similar odors through both nostrils. This can allow associative plasticity in cortical areas to select for matched inputs from both hemispheres, similar to, for example, the well-studied map development in the visual system (Bednar and Wilson, 2016; Gu and Cang, 2016; Hubel and Wiesel, 1962). This hypothesis of experience-dependent matching of bilateral inputs to individual cortical neurons is open to experimental testing.

Methods

Contact for Reagent and Resource Sharing

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Venkatesh N. Murthy (vnmurthy@fas.harvard.edu).

Experimental Model and Subject Details

Male C57Bl/6J and OMP-GCaMP3 (Isogai et al., 2011) mice were used in this study. All the mice were 3 to 8 months old at the time of the experiments. Mice were singly housed after chronic tetrode implantation. A summary of the mice used for tetrode recording can be found in Table S1. All experiments were performed in accordance with the guidelines set by the National Institutes of Health and approved by the Institutional Animal Care and Use Committee at Harvard University.

Bilateral Odor Stimulation

The odor panel was composed of 15 monomolecular odors, purchased from Millipore Sigma: isopropyl tiglate, ethyl tiglate, propyl acetate, isoamyl acetate, ethyl valerate, hexanal, heptanal, allyl butyrate, citronellal, hexyl tiglate, 4-allyl anisole, isobutyl propionate, 2-heptanone, ethyl propionate, and eucalyptol (Figure S1A). The odors were diluted in diethyl phthalate, also purchased from Millipore Sigma (final odor dilution: 5% volume/volume). In addition to these 15 odors, we also tested the response to diethyl phthalate alone, referred as the "blank" trials (Methods, "Neuronal Recording" section). The olfactometer was controlled by a custom LabVIEW script (National Instruments).

Odors were delivered through a custom-built facial mask (Figure 1B). The mask was made of a large piece of tubing surrounding the mouse's muzzle. On each side of the mask, close to each nostril, a piece of tubing delivered odors at a small flow rate (50 standard cubic centimeters per minute). Also on the mask, right below each odor delivery tube, a vacuum ensured proper air cleaning. We confirmed that the mask could deliver similar amounts of odor on each side of the mask with a photoionization detector (miniPID 200B, Aurora Scientific, Canada) (Figure S1B). We confirmed the absence of

contralateral stimulations through calcium imaging at the surface of the OB and M/T cell recordings (Figure 1).

Odors were delivered for 2s for each trial. To avoid odor habituation, a period of 15s separated consecutive trials (i.e., 15s from the end of a trial to the beginning of the next trial). Each odor was delivered in two configurations: ipsilateral and contralateral to the implantation. Each odor delivery was repeated 7 times. Therefore, each recording session contained $(15\ odors + 1\ blank\ trial) \times 2\ sides \times 7\ repetitions = 224$ odor presentations. The order of odor presentations was randomly shuffled for each recording session.

Breathing Monitoring

The face mask was connected to an airflow sensor (Honeywell AWM3300V, Morris Plains, NJ) to monitor the animal's respiration during the experiments (Figure 1B) and later align neuronal activity with the sniffs (Figure 2C) (Bolding and Franks, 2017). The accuracy of the face mask in monitoring respiration was tested against intranasal pressure transients (Figures S1C-S1F) as described previously (Reisert et al., 2014).

Chronic Tetrode Implantation

Mice were anesthetized with an intraperitoneal injection of ketamine and xylazine (100 mg/kg and 10 mg/kg, respectively), then placed in a stereotaxic apparatus. After the skull was cleaned and gently scratched, a custom-made titanium head bar was glued to it. A small craniotomy was performed above the implantation site, before 8 custom-built tetrodes (Chang et al., 2013) were lowered together into the brain (coordinates for the AON pars principalis: antero-posterior 3mm, medio-lateral 1.5mm, dorso-ventral 2.3mm - APC: antero-posterior 1.6mm, medio-lateral -2.8mm, dorso-ventral 3.4mm - PPC: antero-posterior 0.3mm, medio-lateral 3.9mm, dorso-ventral 3.5mm - OB: antero-posterior 1.2 mm, medio-lateral 1.1 mm, dorso-ventral 0.3mm; all antero-posterior and medio-lateral coordinates are given relative to Bregma, except for the OB, for which they are given relative to the junction of inferior cerebral vein and superior sagittal sinus; all dorso-ventral coordinates are given relative to the brain surface). A reference electrode was implanted on the occipital crest. All the mice were implanted in the right hemisphere, except mouse APC2, which was implanted in the left hemisphere. We did not find any difference between APC2 and the other mice. The whole system was stabilized with dental cement. Mice were given a week of recovery before any new manipulation.

Neuronal Recording

A week after chronic tetrode implantation, mice were progressively habituated to stay calm while head-fixed on the recording setup. The habituation process typically took a week. Brain activity was then recorded once a day, always at the same period of the day, while delivering odors to the ipsilateral or contralateral nostril. The mice were awake, head-restrained, and freely-breathing during all the recordings.

Since the tetrode headstage allowed for the tetrodes to be finely adjusted up or down, the tetrodes were slightly lowered in the brain after each recording session (around 40µm deep), ensuring that different neurons were recorded each day.

Electrical activity was amplified, filtered (0.3-6kHz), and digitized at 30kHz (Intan Technologies, RHD2132, connected to an Open Ephys

board). Single units were sorted offline manually using MClust script (Schmitzer-Torbert et al., 2005) written for MATLAB (MathWorks). Units with more than 1% of their inter spike intervals below 2ms refractory period were discarded. Units displaying large changes of amplitude or waveform during the recording were also discarded (Rey et al., 2015) (Figures S1G-S1I). The position of the tetrodes in the brain was confirmed post-mortem through electrolesion (200µA for 4s per channel) (Figure S2).

The average response R to an odor o (in Hz) over the 7 trials t was calculated as followed:

$$R = \sum_{t=1}^{7}(Dur_o^t - Bef_o^t)/7 - \sum_{t=1}^{7}(Dur_b^t - Bef_b^t)/7$$

where, for each trial t, Dur_o^t (resp. Dur_b^t) is the firing rate during the odor (resp. "blank") delivery, while Bef_o^t (resp. Bef_b^t) is the firing rate before the odor (resp. "blank") delivery.

Dur_o^t (resp. Dur_b^t) was calculated for the first 2s following the first sniff after odor (resp. "blank") onset. Bef_o^t (resp. Bef_b^t) was calculated for the 4s preceding odor (resp. "blank") onset.

To determine the significance of a response to an odor o over the 7 trials t, we compared the activity evoked by the odor ($Dur_o^t - Bef_o^t$) to the "blank" activity ($Dur_b^t - Bef_b^t$) (Wilcoxon signed-rank test, critical value set to 5%) (Bolding and Franks, 2017).

Finally, we investigated whether a neuron that responds to a particular odor with an increase in firing rate, tends to exhibit increases in firing rates for any other odor it responds to; and similarly, whether a neuron having inhibitory response to an odor is also inhibited by any other odor it responds to. To do so, we calculated a "positive-negative balance" metric. For each neuron significantly responding to at least one odor, the positive-negative balance corresponds to the fraction of significant responses that are positive. For example, if a neuron has a positive-negative balance of 0.5, then we know that, among the odors eliciting a significant response, half of them elicit an increase of activity, while the other half elicit a decrease. Similarly, a balance of 0 means that all the significant odor responses are negative, and a balance of 1 indicates that all the significant responses are positive.

Acute Calcium Imaging

Mice were anesthetized with an intraperitoneal injection of ketamine/xylazine mixture (100 mg/kg and 10 mg/kg, respectively). After the skull was cleaned and gently scratched, a custom-made titanium head bar was glued to it. A large craniotomy was performed above the two OBs and covered with 1% low melting point agarose (Thermo Fisher). While the mouse was still anesthetized, calcium activity was imaged above the craniotomies with a custom-made epifluorescence microscope (10X objective, Olympus, NA 0.3; blue excitation LED, Thorlabs; emission filter, 500–550nm, Thorlabs; CMOS camera, The Imaging Source) at 4 frames/s, while presenting odors with the custom-built facial mask (Figures 1A and 1B).

The acquired sequences of images were converted to ΔF/F images (change in fluorescence over resting fluorescence). Then the maximal response at each pixel was calculated for each of the 15 odors, for each stimulus configuration (right or left nostrils), and these maximum response maps were used to manually identify glomeruli for further analysis. Finally, for each odor, for each stimulus configuration, the response at each glomerulus was quantified by

1 integrating the ΔF/F signal 4s after odor onset (which corresponds to
2 the 2s of odor presentation, and the 2s immediately following).

Decoding Task

Neuronal Decoding

For each neuron we created a feature vector by calculating the
summed spike count in a 2s window after the mouse first sniffs the
odor and subtracting from it the same sum from the 'blank' odor
averaged across seven trials.

Odor identity decoding ability was assessed using linear classifiers and
leave-out cross validation on a pseudopopulation made up of all
neurons recorded in a particular cortical area – AON or APC (Bolding
and Franks, 2017). At each instance we chose a particular dataset
(either ipsilateral or contralateral odor presentation) and formed
from it a training set by randomly removing one trial per odor. We
calculated mean responses for each odor by averaging across the
training set. We then chose our test set in two related ways. If we
were testing on the same side as we trained (e.g., train ipsilaterally,
test ipsilaterally), then the 15 trials (one per odor) outside the training
set were used as the test set. If we were testing on the opposite side
(e.g., train ipsilaterally, test contralaterally), then one trial per odor
from the opposite side was randomly chosen to form the test set.
Classification accuracy was measured by assigning each test trial to
the closest Euclidean mean odor response and calculating the
percentage of trials correctly assigned (Bolding and Franks, 2017).
This process was repeated 200 times using different random test and
train trial choices.

In order to examine the effect of changing number of neurons on
classification accuracy, we randomly selected neurons from the
pseudopopulation and performed classification using only the
selected cells. This process is repeated 200 times. We then report the
average accuracy and standard deviation over those 200 selections of
neurons and, for each set of neurons, 200 sets of train/test trials in
Figures 4 and S5.

A very similar process is employed for side decoding. One trial per side
was removed, the remaining trials are averaged to form a mean side
response for both ipsilateral and contralateral presentations. The
removed trials are then classified to the nearer mean side response
and accuracy is the percentage correctly assigned. This is repeated
200 times for different random removed trial choices.

41
Comparisons to the Theoretical Model

Our model predicts the trial-averaged neural responses. In order to
make comparisons to the neuronal data, we designed a new
classification task. For simulations and trial averaged neuronal data
we performed odor decoding as follows: we assigned each of the 15
odor centroids from one side (contralateral or ipsilateral) to their
closest neighbor on the other side. This classification was correct if an
odor from one side was classified to the same odor on the other side.
The accuracy is then the proportion of odors correctly assigned.
Again, average accuracy and standard deviation over 200 randomly
chosen neurons are reported in Figures 4 and S5. One exception to
the procedure above is Figure S7D, right panel, where we classify
centroids to their nearest angular neighbor.

For side classification comparisons, the two centroids corresponding
to a given odor presented on both sides are removed. From the
remaining 28 centroids, two points representing ipsilateral and
contralateral cortex are made by averaging over all remaining
centroids from a particular side. The two points we removed are then
classified as ipsilateral or contralateral depending on which mean
they are nearest to. This procedure is repeated 15 times, removing a
different odor each time. We report the average accuracy and
standard deviation over 200 randomly chosen pseudopopulations in
Figures 4 and S5.

65
Computational Modeling

We built a feed-forward neural network model to explore the
implications of a random or structured connection between the
olfactory cortex in each hemisphere. We effectively model one
cortical region in each hemisphere, based on the AON.

To begin, we considered the cortical response to odors presented
ipsilaterally, later we will link these to explore interhemispheric
effects. Each cortical neuron's activity to an ipsilateral odor o is a
weighted sum of glomerular activities passed through an activation
function:

$$y_{o,i} = f\left(\Sigma_{j=1}^{N_x} J_{ij} x_{o,j}\right) = f(h_{o,i}),$$

where **J** is the glomerular to cortical connectivity matrix, $x_{o,j}$ and $y_{o,i}$
are the firing rates of glomerulus j and cortical neuron i respectively
to odor o, $h_{o,i}$ is the input to cortical neuron i, j runs from 1 to N_x (the
total number of glomeruli), and f is the activation function.

The zero point of these neurons is set to be basal firing rate plus
response to a blank odor. Therefore, we modeled only odor related
perturbations from the blank and base to match all comparisons to
the blank and base subtracted data.

The nonlinearity, f, was chosen to recreate features of the
electrophysiological recordings. First, it was found that the odor
related perturbations are equally likely to be positive or negative (Of
all the significant neuron-odor pairs, 46% are positive in the AON, 48%
in the APC, and 48% in the PPC) therefore we chose a nonlinearity
that was symmetric around zero. Negative activity does not mean the
neurons have a negative firing rate, but that their firing rate is
inhibited relative to their response to base plus blank. Second,
consistent with previous findings (Bolding and Franks, 2017; Iurilli and
Datta, 2017; Poo and Isaacson, 2009), the cortical response to odors
was relatively sparse. Therefore, we chose the shrink function:

$$f(h_{o,i}) = \begin{cases} h_{o,i} - \varphi & if \quad h_{o,i} \geq \varphi \\ 0 & if \quad |h_{o,i}| < \varphi, \\ h_{o,i} + \varphi & if \quad h_{o,i} \leq \varphi \end{cases}$$

where φ is a threshold that controls sparseness.

Based on previous work we then modelled the connectivity matrix
between the bulb and the cortex to be completely random (Schaffer
et al., 2018) with elements either equal to 0, with probability $1 - \xi$,
or drawn independently from a zero mean distribution with finite
variance $\frac{\sigma^2}{N_x}$, with probability ξ. ξ, therefore, controls the sparseness of
the matrix[1]. Now by Central Limit Theorem $h_{o,i}$ is distributed in a
Gaussian manner and with zero mean.

---

[1] Note that negative weights do not mean that OB inputs are directly inhibitory; they rather model OB inputs projecting to inhibitory OC neurons, which in turn project to our OC neurons of interest.

Since input to each cortical neuron is statistically independent, to fully
specify its distribution, we must simply set the odor-odor covariance
matrix of the Gaussian. Under the assumption that the size of the OB
output response is identical for each odor ($|x_o|^2 = |x_{o'}|^2$ $\forall$ odors o
and o'), which can be justified by the divisive normalization that
appears to happen in the OB (Banerjee et al., 2015; Zhu et al., 2013),
the elements of the covariance matrix can be found as follows:

8
$$\mathbb{E}_i[h_{o,i}^2] = \xi\sigma^2 \frac{1}{N_x}|x_o|^2 = \gamma^2$$

9
$$\mathbb{E}_i[h_{o,i}h_{o',i}] = \xi\sigma^2 \frac{1}{N_x}x_o \cdot x_{o'} = \gamma^2\cos(\theta_{o,o'}),$$

where $\mathbb{E}_i$ refers to an expectation over the cortical population and we
defined a new variable $\gamma^2 := \xi\sigma^2 \frac{1}{N_x}|x_o|^2$ that controls the variance,
and $\theta_{o,o'}$ is the angle between two glomerular representations. For
the 15 odors under consideration, we used calcium imaging to extract
the cosine similarity which serves as the Gaussian covariance. To
complete our ipsilateral specification we must, therefore, only choose
γ and a threshold ϕ – we shall rationalize these terms in two ways.

The threshold controls sparseness of cortical response, this we match
to the AON recordings. The sparseness of the recordings is measured
as the proportion of neuron-odor pairs for which the Wilcoxon signed
rank test p value comparing the set of seven trials to mean blank odor
response is below 5% (22.4% Ipsilateral, 25.2% Contralateral).
Labeling the sparseness as S, because of the assumed Gaussianity of
cortical inputs, we get:

24
$$S = \frac{1}{\sqrt{2\pi\gamma^2}}\left(\int_\phi^\infty e^{-\frac{h^2}{2\gamma^2}}\,dh + \int_{-\infty}^{-\phi} e^{-\frac{h^2}{2\gamma^2}}\,dh\right) = \text{erfc}\left[\frac{\phi}{\gamma\sqrt{2}}\right],$$

25 where erfc is the cumulative error function. Therefore, the sparseness
26 sets the ratio of threshold to variance:

27
$$\frac{\phi}{\gamma} = \sqrt{2}\,\text{erfc}^{-1}[S].$$

Finally, we use the ipsilateral distribution of significant response
magnitudes in the AON (Figure 2E, top panel) to fit the conversion of
numbers to firing rate (Hz), γ. Our model predicts the significant
response magnitude distribution to be the tail of a Gaussian, by
changing γ we are simply rescaling the x axis of this distribution, we
choose the γ that permits the best model-data fit. This fully specified
the ipsilateral simulations and fit the data remarkably well (see
Figures 6 and S7).

To compare ipsilateral to contralateral responses we create two
ipsilateral responses according to the scheme above. One of these is
then mapped through a cross cortical connectivity matrix to create a
contralateral response. We use this contralateral response and the
second ipsilateral creation for our comparisons to mice.

Our final task, therefore, is to design a cross-cortical connectivity
matrix. In the body of the paper, we use a simple Hebbian structured
matching of each odor between hemispheres. Here we include two
additional connectivity schemes that are low dimensional
approximations to G_{Struct}. The first uses SVD to create a low rank
matrix closest to G_{Struct} as measured by the Frobenius norm. The
second uses the principal components to create G_{Struct} using a
smaller number of Hebbian matching experiences. These show that
the particular form of the structured connectivity is not important. In
each case we combine the structured matrix with a random matrix
and examine how much structure is required to fit the observed data.

The random component has elements drawn from a zero mean
normal distribution with variance 1 (this choice of variance is
unimportant as the matrix will be rescaled in future):

55
$$G_{ij}^{Rand} \sim \mathcal{N}(0,1)$$

56 with probability ξ, else 0.

57 And the structured Hebbian component maps from contralateral to
58 ipsilateral odors (Dayan and Abbott, 2001):

59
$$G_{Struct}^{Unscaled} = \sum_{o=1}^{P}(\mathbf{Y_o^{Ipsi}} - \mathbf{C^{Ipsi}})(\mathbf{Y_o^{Opp}} - \mathbf{C^{Opp}})^{\mathbf{T}},$$

where $\mathbf{Y_o^{Ipsi}}$ is the response vector to odor o presented ipsilaterally in
the ipsilateral cortex, $\mathbf{C^{Ipsi}}$ is response in the ipsilateral cortex
averaged over odors, P=15 is the total number of odors, and $\mathbf{Y_o^{Opp}}$ is
the response in the opposite cortex to an odor presented in the
opposite nostril.

These two matrices currently have outputs that are of arbitrary
magnitudes. To ensure each contribution has a roughly similar sized
output, the structured section is rescaled by a factor:

68
$$G_{Struct} = \chi\,G_{Struct}^{Unscaled} \text{ where } \chi = \frac{\Sigma_{o=1}^{P}\left|G_{Rand}Y_o^{Opp}\right|}{\Sigma_{o=1}^{P}\left|G_{Struct}^{Unscaled}Y_o^{Opp}\right|},$$

69 before being combined using the parameter α to create a cross
70 cortical link with varying levels of randomness/structure:

71
$$G = \alpha G_{Struct} + (1-\alpha)G_{Rand}.$$

Since the size of the outputs of each matrix are equal, we can now
interpret alpha as the proportion of contralateral input magnitude
that is formed by structured connectivity.

The magnitude of the elements of this matrix are then scaled again
such that the resulting contralateral representation has the desired
sparseness when the same piriform threshold is applied as was used
to create the ipsilateral sparseness. Finally, the conversion to Hz, γ, is
used to compare to data.

We varied alpha and looked at how four measures of cross-cortical
alignment changed (Figures 6E-6J; see main text for details) and how
simulations compared to our actual AON data. We chose the optimal
alpha to be the point with the minimum z^2-score across the four – the
smallest squared sum of sigmas (error bars) between the point and
the AON data line in all four plots.

We modeled our AON with 50,000 neurons, this is a rough
underestimate based on other paleocortical neuronal densities
(Srinivasan and Stevens, 2017) and measurements of AON size from
the Allen Brain Atlas. Our model, however, was robust to changes in
number of neurons, for example we tried a 25,000-neuron version
(Figures S7E-S7H) and the optimal alpha did not change.

Our first low rank approximation uses Singular Value Decomposition
to factorise G_{Struct}:

94
$$G_{Struct} = U\Sigma V^T,$$

95 where U and V are orthogonal matrices and Σ is a diagonal matrix.
96 We vary the dimensionality of our approximation through setting all
97 but the largest Q singular values in Σ to 0, creating Σ^* which we then
98 use to recreate G_{Struct}:

99
$$G_{Struct\,2}^{Unscaled} = U\Sigma^*V^T.$$

The last type of structured connectivity is a simple extension, comprising a Hebbian matching of vectors between the two hemispheres. However rather than matching the odors explicitly we try to align the odorant subspace in which the odorants sit. To do this we calculate the principal components of the contralateral representation and design a connectivity scheme that aligns the first few of the contralateral principal components between cortices:

$$G_{Struct\,3}^{Unscaled} = \sum_{q=1}^{Q} \mathbf{Y}_q^{\mathbf{Ipsi}} \mathbf{Y}_q^{\mathbf{Opp^T}},$$

Where Q is the number of principal components used, $\mathbf{Y}_q^{\mathbf{Opp}}$ is the q-th principal component of the opposite cortical representation, and $\mathbf{Y}_q^{\mathbf{Ipsi}}$ is the ipsilateral representation of the same mixture of odors that comprises $\mathbf{Y}_q^{\mathbf{Opp}}$. More explicitly, the q-th principal component in the opposite cortex is a mixture of normalized odorant vectors:

$$\mathbf{Y}_q^{\mathbf{Opp}} = \sum_{o=1}^{P} \rho_o^q \mathbf{Y}_o^{\mathbf{Opp}},$$

Where ρ^q is the vector of coefficients for the q-th principal components. We then use the same set of coefficients to create $\mathbf{Y}_q^{\mathbf{Ipsi}}$:

$$\mathbf{Y}_q^{\mathbf{Ipsi}} = \sum_{o=1}^{P} \rho_o^q \mathbf{Y}_o^{\mathbf{Ipsi}}.$$

The dimensionality of each approximation, Q, must be chosen such that most of the response variation is captured; however, the low dimensionality of the responses means high alignment can be achieved despite a small Q. For example, in Figures S8A and S8B we show how cross-cortical decoding accuracy varies with the dimensionality of each approximate structured matrix. Only 7 singular values (Figure S8C-S8F), or 6 principal components (explaining 73% of the variance, Figures S8G-S8J), are needed for the alignment metrics to reach the values seen in neuronal recordings.

Quantification and Statistical Analysis

All quantifications and statistical analyses were performed using custom scripts in MATLAB (MathWorks), except (1) the circular statistical analyses, which were performed with the CircStat toolbox written by Philipp Berens (Berens, 2009), and (2) the Hartigan's dip tests, which were performed with the code written by Ferenc Mechler (Mechler and Ringach, 2002). The linear regressions were obtained with the least square method. The data is given as median ± standard deviation (SD), unless specified otherwise. For all statistical tests, the critical value was set to 5%. All statistical tests were two-sided, with the exception of the bootstrap strategies (where we asked whether the measurements were significantly higher than chance).

Additionally, the significance of the linear regressions was tested using either an F-test (Figures 3, S3, and S4). The significance of the percentage of bilaterally correlated neurons was tested using a bootstrap strategy (see figure caption for details). Table S2 summarizes all the statistical tests performed in this study

Data and Code Availability

All data and code used to compute quantities presented in this study is available on the Murthy lab GitHub page:
https://github.com/VNMurthyLab/IpsiContra_v2.

Author Contributions

V.N.M and J.G designed the experimental setup. C.P and W.D. designed the computational modeling strategy with inputs from all authors. J.G. acquired the electrophysiological data. J.G. and W.D. analyzed the data. J.G. and W.D. wrote the manuscript with inputs form all authors.

Acknowledgements

Research reported in this paper was supported, in part, by grants from the NIH (R01 DC016289, R01DC011291), the Harvard Merit Fellowship, and the Herchel Smith Scholarship. We thank Vikrant Kapoor, as well as all members of the Murthy and Pehlevan labs for helpful feedback. We also thank Massimo Vergassola and Gautam Reddy for critical discussion. Finally, we thank Naoshige Uchida and Kenneth Blum for their important feedback on a previous version of this manuscript.

Competing Interests

The authors declare no competing interests.

Reference

Álvarez-Salvado E, Licata AM, Connor EG, McHugh MK, King BM, Stavropoulos N, Victor JD, Crimaldi JP, Nagel KI. 2018. Elementary sensory-motor transformations underlying olfactory navigation in walking fruit-flies. *Elife* **7**. doi:10.7554/eLife.37815

Babadi B, Sompolinsky H. 2014. Sparseness and Expansion in Sensory Representations. *Neuron* **83**:1213–26. doi:10.1016/j.neuron.2014.07.035

Baker KL, Dickinson M, Findley TM, Gire DH, Louis M, Suver MP, Verhagen J V, Nagel KI, Smear MC. 2018. Algorithms for Olfactory Search across Species. *J Neurosci* **38**:9383–9389. doi:10.1523/JNEUROSCI.1668-18.2018

Banerjee A, Marbach F, Anselmi F, Koh MS, Davis MB, Garcia da Silva P, Delevich K, Oyibo HK, Gupta P, Li B, Albeanu DF. 2015. An Interglomerular Circuit Gates Glomerular Output and Implements Gain Control in the Mouse Olfactory Bulb. *Neuron* **87**:193–207. doi:10.1016/j.neuron.2015.06.019

Bednar JA, Wilson SP. 2016. Cortical Maps. *Neuroscientist* **22**:604–617. doi:10.1177/1073858415597645

Bekkers JM, Suzuki N. 2013. Neurons and circuits for odor processing in the piriform cortex. *Trends Neurosci* **36**:429–38. doi:10.1016/j.tins.2013.04.005

Berens P. 2009. CircStat : A MATLAB Toolbox for Circular Statistics. *J Stat Softw* **31**:1–21. doi:10.18637/jss.v031.i10

Best AR, Wilson DA. 2003. A postnatal sensitive period for plasticity of cortical afferents but not cortical association fibers in rat piriform cortex. *Brain Res* **961**:81–87. doi:10.1016/S0006-8993(02)03847-7

Blake R, Wilson H. 2011. Binocular vision. *Vision Res* **51**:754–70. doi:10.1016/j.visres.2010.10.009

Bojsen-Moller F, Fahrenkrug J. 1971. Nasal swell-bodies and cyclic changes in the air passage of the rat and rabbit nose. *J Anat* **110**:25–37.

Bolding KA, Franks KM. 2018. Recurrent cortical circuits implement concentration-invariant odor coding. *Science* **361**:eaat6904. doi:10.1126/science.aat6904

Bolding KA, Franks KM. 2017. Complementary codes for odor identity and intensity in olfactory cortex. *Elife* **6**. doi:10.7554/eLife.22630

Boyd AM, Sturgill JF, Poo C, Isaacson JS. 2012. Cortical Feedback Control of
Olfactory Bulb Circuits. *Neuron* **76**:1161–1174.
doi:10.1016/j.neuron.2012.10.020

Cang J, Feldheim DA. 2013. Developmental mechanisms of topographic map
formation and alignment. *Annu Rev Neurosci* **36**:51–77.
doi:10.1146/annurev-neuro-062012-170341

Catania KC. 2013. Stereo and serial sniffing guide navigation to an odour
source in a mammal. *Nat Commun* **4**:1441. doi:10.1038/ncomms2444

Chang EH, Frattini SA, Robbiati S, Huerta PT. 2013. Construction of
Microdrive Arrays for Chronic Neural Recordings in Awake Behaving
Mice. *J Vis Exp* **77**:e50470. doi:10.3791/50470

Cumming BG, DeAngelis GC. 2001. The Physiology of Stereopsis. *Annu Rev
Neurosci* **24**:203–238. doi:10.1146/annurev.neuro.24.1.203

Dalal T, Gupta N, Haddad R. 2020. Bilateral and unilateral odor processing
and odor perception. *Commun Biol* **3**:150. doi:10.1038/s42003-020-
0876-6

Dayan P, Abbott LF. 2001. Theoretical Neuroscience: Computational and
Mathematical Modeling of Neural Systems. Cambridge, MA: The MIT
Press.

Eccles R. 2000. Nasal airflow in health and disease. *Acta Otolaryngol*
120:580–95. doi:10.1080/000164800750000388

Esquivelzeta Rabell J, Mutlu K, Noutel J, Martin del Olmo P, Haesler S. 2017.
Spontaneous Rapid Odor Source Localization Behavior Requires
Interhemispheric Communication. *Curr Biol* **27**:1542-1548.e4.
doi:10.1016/j.cub.2017.04.027

Frank LM, Brown EN, Wilson MA. 2001. A comparison of the firing properties
of putative excitatory and inhibitory neurons from CA1 and the
entorhinal cortex. *J Neurophysiol* **86**:2029–40.
doi:10.1152/jn.2001.86.4.2029

Franks KM, Isaacson JS. 2005. Synapse-specific downregulation of NMDA
receptors by early experience: A critical period for plasticity of
sensory input to olfactory cortex. *Neuron* **47**:101–114.
doi:10.1016/j.neuron.2005.05.024

Ghosh S, Larson SD, Hefzi H, Marnoy Z, Cutforth T, Dokka K, Baldwin KK.
2011. Sensory maps in the olfactory cortex defined by long-range viral
tracing of single neurons. *Nature* **472**:217–20.
doi:10.1038/nature09945

Giessel AJ, Datta SR. 2014. Olfactory maps, circuits and computations. *Curr
Opin Neurobiol* **24**:120–32. doi:10.1016/j.conb.2013.09.010

Gire DH, Kapoor V, Arrighi-Allisan A, Seminara A, Murthy VN. 2016. Mice
develop efficient strategies for foraging and navigation using complex
natural stimuli. *Curr Biol* **26**:1261–1273.
doi:10.1016/j.cub.2016.03.040

Grobman M, Dalal T, Lavian H, Shmuel R, Belelovsky K, Xu F, Korngreen A,
Haddad R. 2018. A Mirror-Symmetric Excitatory Link Coordinates
Odor Maps across Olfactory Bulbs and Enables Odor Perceptual Unity.
Neuron **99**:800–813. doi:10.1016/j.neuron.2018.07.012

Grothe B, Pecka M. 2014. The natural history of sound localization in
mammals--a story of neuronal inhibition. *Front Neural Circuits* **8**:116.
doi:10.3389/fncir.2014.00116

Gu Y, Cang J. 2016. Binocular matching of thalamocortical and intracortical
circuits in the mouse visual cortex. *Elife* **5**:e22032.
doi:10.7554/eLife.22032

Gur M, Beylin A, Snodderly DM. 1999. Physiological properties of macaque
V1 neurons are correlated with extracellular spike amplitude,
duration, and polarity. *J Neurophysiol* **82**:1451–64.
doi:10.1152/jn.1999.82.3.1451

Haberly LB, Price JL. 1978. Association and commissural fiber systems of the

olfactory cortex of the rat. I. Systems originating in the piriform cortex
and adjacent areas. *J Comp Neurol* **178**:711–740.
doi:10.1002/cne.901780408

Hassani OK, Lee MG, Henny P, Jones BE. 2009. Discharge Profiles of Identified
GABAergic in Comparison to Cholinergic and Putative Glutamatergic
Basal Forebrain Neurons across the Sleep-Wake Cycle. *J Neurosci*
29:11828–11840. doi:10.1523/JNEUROSCI.1259-09.2009

Hu R, Zhang J, Luo M, Hu J. 2017. Response Patterns of GABAergic Neurons in
the Anterior Piriform Cortex of Awake Mice. *Cereb Cortex* **27**:3110–
3124. doi:10.1093/cercor/bhw175

Hubel DH, Wiesel TN. 1962. Receptive fields, binocular interaction and
functional architecture in the cat's visual cortex. *J Physiol* **160**:106–54.
doi:10.1113/jphysiol.1962.sp006837

Illig KR, Haberly LB. 2003. Odor-evoked activity is spatially distributed in
piriform cortex. *J Comp Neurol* **457**:361–373. doi:10.1002/cne.10557

Isogai Y, Si S, Pont-Lezica L, Tan T, Kapoor V, Murthy VN, Dulac C. 2011.
Molecular organization of vomeronasal chemoreception. *Nature*
478:241–245. doi:10.1038/nature10437

Iurilli G, Datta SR. 2017. Population Coding in an Innately Relevant Olfactory
Area. *Neuron* **93**:1180–1197. doi:10.1016/j.neuron.2017.02.010

Johenning FW, Beed PS, Trimbuch T, Bendels MHK, Winterer J, Schmitz D.
2009. Dendritic compartment and neuronal output mode determine
pathway-specific long-term potentiation in the piriform cortex. *J
Neurosci* **29**:13649–61. doi:10.1523/JNEUROSCI.2672-09.2009

Kikuta S, Kashiwadani H, Mori K. 2008. Compensatory Rapid Switching of
Binasal Inputs in the Olfactory Cortex. *J Neurosci* **28**:11989–11997.
doi:10.1523/JNEUROSCI.3106-08.2008

Kikuta S, Sato K, Kashiwadani H, Tsunoda K, Yamasoba T, Mori K. 2010.
Neurons in the anterior olfactory nucleus pars externa detect right or
left localization of odor sources. *Proc Natl Acad Sci* **107**:12363–12368.
doi:10.1073/pnas.1003999107

Kucharski D, Hall WG. 1988. Developmental Change in the Access to
Olfactory Memories. *Behav Neurosci* **102**:340–8. doi:10.1037/0735-
7044.102.3.340

Kucharski D, Hall WG. 1987. New routes to early memories. *Science* **238**:786–
8. doi:10.1126/science.3672125

Litaudon P, Amat C, Bertrand B, Vigouroux M, Buonviso N. 2003. Piriform
cortex functional heterogeneity revealed by cellular responses to
odours. *Eur J Neurosci* **17**:2457–61. doi:10.1046/j.1460-
9568.2003.02654.x

Luo L, Flanagan JG. 2007. Development of continuous and discrete neural
maps. *Neuron* **56**:284–300. doi:10.1016/j.neuron.2007.10.014

Mainland JD, Bremner E a, Young N, Johnson BN, Khan RM, Bensafi M, Sobel
N. 2002. Olfactory plasticity: one nostril knows what the other learns.
Nature **419**:802. doi:10.1038/419802a

Mechler F, Ringach DL. 2002. On the classification of simple and complex
cells. *Vision Res* **42**:1017–33. doi:10.1016/S0042-6989(02)00025-1

Miyamichi K, Amat F, Moussavi F, Wang C, Wickersham I, Wall NR, Taniguchi
H, Tasic B, Huang ZJ, He Z, Callaway EM, Horowitz MA, Luo L. 2011.
Cortical representations of olfactory input by trans-synaptic tracing.
Nature **472**:191–196. doi:10.1038/nature09714

Mori K, Nagao H, Yoshihara Y. 1999. The olfactory bulb: coding and
processing of odor molecule information. *Science* **286**:711–5.
doi:10.1126/science.286.5440.711

Nagayama S, Homma R, Imamura F. 2014. Neuronal organization of olfactory
bulb circuits. *Front Neural Circuits* **8**:98. doi:10.3389/fncir.2014.00098

Otazu GH, Chae H, Davis MB, Albeanu DF. 2015. Cortical Feedback
Decorrelates Olfactory Bulb Output in Awake Mice. *Neuron* **86**:1461–

1477. doi:10.1016/j.neuron.2015.05.023

Parthasarathy K, Bhalla US. 2013. Laterality and symmetry in rat olfactory behavior and in physiology of olfactory input. *J Neurosci* **33**:5750–60. doi:10.1523/JNEUROSCI.1781-12.2013

Pashkovski SL, Iurilli G, Brann D, Chicharro D, Drummey K, Franks KM, Panzeri S, Datta SR. 2020. Structure and flexibility in cortical representations of odour space. *Nature* **583**:253–258. doi:10.1038/s41586-020-2451-1

Poo C, Isaacson JS. 2009. Odor Representations in Olfactory Cortex: "Sparse" Coding, Global Inhibition, and Oscillations. *Neuron* **62**:850–61. doi:10.1016/j.neuron.2009.05.022

Porter J, Anand T, Johnson B, Khan RM, Sobel N. 2005. Brain mechanisms for extracting spatial information from smell. *Neuron* **47**:581–92. doi:10.1016/j.neuron.2005.06.028

Povysheva N V., Gonzalez-Burgos G, Zaitsev A V., Kröner S, Barrionuevo G, Lewis DA, Krimer LS. 2006. Properties of excitatory synaptic responses in fast-spiking interneurons and pyramidal cells from monkey and rat prefrontal cortex. *Cereb Cortex* **16**:541–552. doi:10.1093/cercor/bhj002

Rajan R, Clement JP, Bhalla US. 2006. Rats smell in stereo. *Science* **311**:666–70. doi:10.1126/science.1122096

Reisert J, Golden GJ, Matsumura K, Smear M, Rinberg D, Gelperin A. 2014. Comparing thoracic and intra-nasal pressure transients to monitor active odor sampling during odor-guided decision making in the mouse. *J Neurosci Methods* **221**:8–14. doi:10.1016/j.neumeth.2013.09.006

Rey HG, Pedreira C, Quian Quiroga R. 2015. Past, present and future of spike sorting techniques. *Brain Res Bull* **119**:106–17. doi:10.1016/j.brainresbull.2015.04.007

Schaffer ES, Stettler DD, Kato D, Choi GB, Axel R, Abbott LF. 2018. Odor Perception on the Two Sides of the Brain: Consistency Despite Randomness. *Neuron* **98**:736–742. doi:10.1016/j.neuron.2018.04.004

Schmitzer-Torbert N, Jackson J, Henze D, Harris K, Redish AD. 2005. Quantitative measures of cluster quality for use in extracellular recordings. *Neuroscience* **131**:1–11. doi:10.1016/j.neuroscience.2004.09.066

Schoenfeld TA, Macrides F. 1984. Topographic organization of connections between the main olfactory bulb and pars externa of the anterior olfactory nucleus in the hamster. *J Comp Neurol* **227**:121–135.
doi:10.1002/cne.902270113

Scott JW, McBride RL, Schneider SP. 1980. The organization of projections from the olfactory bulb to the piriform cortex and olfactory tubercle in the rat. *J Comp Neurol* **194**:519–534. doi:10.1002/cne.901940304

Sosulski DL, Lissitsyna Bloom M, Cutforth T, Axel R, Datta SR. 2011. Distinct representations of olfactory information in different cortical centres. *Nature* **472**:213–216. doi:10.1038/nature09868

Srinivasan S, Stevens CF. 2017. A quantitative description of the mouse piriform cortex. *bioRxiv* 99002.

Stettler DD, Axel R. 2009. Representations of Odor in the Piriform Cortex. *Neuron* **63**:854–864. doi:10.1016/j.neuron.2009.09.005

Suzuki SS, Smith GK. 1985. Burst characteristics of hippocampal complex spike cells in the awake rat. *Exp Neurol* **89**:90–95. doi:10.1016/0014-4886(85)90267-5

Swadlow HA. 2003. Fast-spike Interneurons and Feedforward Inhibition in Awake Sensory Neocortex. *Cereb Cortex* **13**:25–32. doi:10.1093/cercor/13.1.25

Weir K, Blanquie O, Kilb W, Luhmann HJ, Sinning A. 2015. Comparison of spike parameters from optically identified GABAergic and glutamatergic neurons in sparse cortical cultures. *Front Cell Neurosci* **8**:460. doi:10.3389/fncel.2014.00460

Wilson DA. 1997. Binaral interactions in the rat piriform cortex. *J Neurophysiol* **78**:160–9. doi:10.1152/jn.1997.78.1.160

Wilson DA, Sullivan RM. 2011. Cortical processing of odor objects. *Neuron* **72**:506–19. doi:10.1016/j.neuron.2011.10.027

Wilson DA, Sullivan RM. 1999. Respiratory airflow pattern at the rat's snout and an hypothesis regarding its role in olfaction. *Physiol Behav* **66**:41–44. doi:10.1016/S0031-9384(98)00269-8

Yan Z, Tan J, Qin C, Lu Y, Ding C, Luo M. 2008. Precise Circuitry Links Bilaterally Symmetric Olfactory Maps. *Neuron* **58**:613–624. doi:10.1016/j.neuron.2008.03.012

Zhan C, Luo M. 2010. Diverse patterns of odor representation by neurons in the anterior piriform cortex of awake mice. *J Neurosci* **30**:16662–16672. doi:10.1523/JNEUROSCI.4400-10.2010

Zhu P, Frank T, Friedrich RW. 2013. Equalization of odor representations by a network of electrically coupled inhibitory interneurons. *Nat Neurosci* **16**:1678–86. doi:10.1038/nn.3528

Title

Bilateral Alignment of Receptive Fields in the Olfactory Cortex

Authors

Julien Grimaud[1,2,3,4], William Dorrell[3,5], Cengiz Pehlevan[3,5], and Venkatesh Murthy[2,3*].

[1] Molecules, Cells, and Organisms Graduate Program, Harvard University, Cambridge, Massachusetts, USA.
[2] Department of Molecular and Cellular Biology and Center for Brain Science, Harvard University, Cambridge, Massachusetts, USA.
[3] Center for Brain Science, Harvard University, Cambridge, Massachusetts, USA.
[4] Sup'Biotech Research Department, Villejuif, France.
[5] John A. Paulson School of Engineering and Applied Sciences, Harvard University, Cambridge, Massachusetts, USA.
*Lead Author. Correspondence: vnmurthy@fas.harvard.edu

Supplementary Information

Note S1: Random Connections Produce Zero Correlations – a Derivation.

Figure S1: A New Method for Unilateral Odor Delivery.

Figure S2: Post-Mortem Confirmation of Tetrode Placement in the Olfactory Cortex.

Figure S3: Bilaterally-Correlated Neurons.

Figure S4: Correlation of Population Odor Representations.

Figure S5: Ipsi- and Contralateral Odor Decoding.

Figure S6: Variability Across Mice.

Figure S7: Modeling Olfactory Cross Cortical Connections.

Figure S8: Low-dimensional structured connectivity is sufficient to produce alignment.

Table S1: Mice Recorded in this Study.

Table S2: Statistical Tests Performed in this Study.

Note S1: Random Connections Produce Zero Correlations – a Derivation.

We will prove that if the same response is mapped through two different random matrices the resulting representations will be uncorrelated. Let $\mathbf{x}$ be our N-dimensional input pattern, for example the bulb response to an odor. Project it through two different random matrices to the same set of cortical neurons. The two matrices, $\mathbf{J}$ and $\mathbf{G}$, each have independently identically distributed elements drawn from a mean zero distribution with finite variance. This creates two further representations which we shall later pass through an activation function. First, however, we shall consider the pre-thresholded representations:

$$h_i = \sum_{j=1}^{N} J_{ij}\, x_j \text{ and } \quad g_i = \sum_{j=1}^{N} G_{ij}\, x_j.$$

We can use the Multivariate Central Limit Theorem to argue that these two representations are jointly normally distributed with zero mean:

$$\begin{pmatrix} h_i \\ g_i \end{pmatrix} = \sum_{j=1}^{N} \begin{pmatrix} J_{ij} x_j \\ G_{ij} x_j \end{pmatrix} \sim \mathcal{N}(0, \Sigma)$$

The correlation matrix can be derived and shown to be diagonal:

$$\Sigma_{hh} = \mathbb{E}_i \left[\sum_j J_{ij}\, x_j \sum_k J_{ik} x_k \right] \neq 0$$

$$\Sigma_{hg} = \mathbb{E}_i \left[\sum_j J_{ij}\, x_j \sum_k G_{ik}\, x_k \right] = 0,$$

where the last equality follows from the independent, zero mean, nature of elements of $\mathbf{G}$ and $\mathbf{J}$.

Now, since h_i and g_i are jointly normally distributed random variables with a diagonal covariance matrix, they are also independent. We then pass these representations through a nonlinearity. Using the fact that functions of independent variables are independent, the resulting output representations are also independent and hence uncorrelated.

This demonstration can be extended to many cases of direct interest. If we let h and g depend on different odors they are still independent and uncorrelated by a simple generalization of the previous argument:

$$h_i = \sum_j J_{ij}\, x_{1,j} \text{ and } \quad g_i = \sum_j G_{ij}\, x_{2,j}.$$

The experiments consider another similar case. Interpret h (g) as the unthresholded representation in the ipsilateral (contralateral) AON to an odor presented ipsilaterally (contralaterally). If the cross cortical matrix is random then thresholding these responses and mapping them cross cortically will not change their independence from one and other.

Previous work has shown that, if two correlated representations are projected through a random matrix, the resulting representations remain correlated (Babadi and Sompolinsky, 2014; Schaffer et al., 2018). The key difference in our work is that we are considering two different random matrices, one in each hemisphere. Hence, this derivation has shown that projecting a pair of correlated representations through two different random matrices eliminates the correlations between resulting outputs.

Therefore, since observed correlations between odors presented ipsilaterally and contralaterally are not zero, the cross cortical connectivity must be structured in some way.

Figure S1: A New Method for Unilateral Odor Delivery.

Related to Figure 1. **(A)** Odorants used in this study. **(B)** Control of the symmetry of the olfactometer. A PID measured the changes of odor concentration over time at the end of the right or the left end of the facial mask. Each measurement was repeated three times; therefore, each graph shows six over-imposed traces, three with the PID on the right, three on the left. Note the similarity between the traces obtained on the left and right of the mask, and the absence of odor detected on the contralateral side. Grey: odor presentation (here: isopropyl tiglate). All 15 odors were tested and showed similar results. **(C)** Diagram of the setup used for assessing the reliability of the face mask for respiration monitoring. The breathing of awake mice was monitored simultaneously with the mask and a pressure sensor connected to a chronic intranasal cannula (n = 3 animals, 15 min of continuous recording per mouse). **(D)** Exemplar breathing traces, simultaneously recorded from the intranasal cannula (green) and the face mask (purple). Note the synchrony of both signals and the apparent absence of phase shift. **(E)** Histogram of instantaneous breathing rates, calculated with the exhalation (top panel) or inhalation (bottom panel) onsets. The data shown here comes from the full monitoring session of one exemplar mouse. The distributions obtained from the intranasal cannula (light bars) or the face mask (dark bars) are almost identical. **(F)** Delay from cannula to mask, calculated with the exhalation (blue) or inhalation (red) onsets. The data from the three mice were combined here. Values: median ± SD. The phase shift between the intranasal cannula and the face mask is small and reliable. **(G)** Example of tetrode recording. The four electrodes belong to the same tetrode. Grey: odor delivery. Green: unsorted electrode traces. Blue and red: two single units. Purple: sniffing acquired simultaneously (up: inhalation). These exemplar traces were recorded in

mouse APC1. **(H)** Example of single unit clustering. The red and blue units are the same as in (G). **(I)** Example of clustered single units. The red and blue units are the same as in (G). Lighter traces: all the peaks from the recording session were over-imposed. Darker traces: average peaks. **(J)** Spike count versus inter-spike interval (ISI) for the two units shown in (G) (blue and red). Note the presence of a refractory period.

Figure S2: Post-Mortem Confirmation of Tetrode Placement in the Olfactory Cortex.
Related to Figure 2. Sagittal sections of two exemplar animals, implanted in the AON (left) and APC (right). The round shape on the ventral end of the tetrode scar in the left panel was caused by post-mortem electrolesion. For animals recorded in the APC, the tetrodes went all the way through the brain, until they reached the ventral part of the skull. As a consequence, it was not possible to save the most anterior part of the section in the right panel. A: anterior. P: posterior. D: dorsal. V: ventral. CC: corpus callosum. LV: lateral ventricle. LOT: lateral olfactory tract. Scale bar: 0.25mm (left) or 1mm (center, right). DAPI staining. We did not trace the border between striatum and amygdala as we could not confidently determine it.

Figure S3: Bilaterally-Correlated Neurons.

Related to Figure 3. **(A) to (D)** Percentage of bilaterally-correlated neurons per mouse. **(A)** Identical to Figure 3B. **(B)** Same as Figure 3B, except we kept only the neurons responding to at least one odor on one side for this analysis. **(C)** Same as Figure 3B, except we kept only the neurons responding to at least one odor on each side for this analysis (note that the odors eliciting responses can be different on each side). **(D)** Correlation across artificially-generated trials. For each mouse, each side, each neuron, each odor, two sets of 7 artificial trials were generated using a Poisson process. Then for each side, an analysis similar to Figure 3B was performed, except we looked for correlations between the first and second sets of 7 artificial trials, instead of correlations between ipsi and contralateral trials. A full circle means the percentage is higher than chance (significance is determined with a bootstrapping method similar to Figure 3C). Panel (D) allows us to estimate the apparent percentage of bilaterally-correlated neurons one would expect if all neurons were bilaterally-correlated. **(E)** Spike traces from mouse PPC1. The average spike of all the neurons recorded from APC1 are shown. The spike amplitudes were normalized for display. Note the apparent bimodal distribution of spike widths. **(F)** Histograms of the spike widths at half-maximum amplitude, per OC region. For each histogram, a Hartigan's

dip test was performed to confirm that the distribution of spike widths was not unimodal (AON p = 0.020, APC p = 0.016). The corresponding p-values are reported next to each graph. We also used a maximum likelihood estimate approach to fit a mixture of two Gaussian distributions to each distributions (AON: weight of the 1st Gaussian relative to the mixture w = 0.07, μ1 = 85.6, σ1 = 4.5, μ2 = 130.1, σ2 = 22.9; APC: w = 0.11, μ1 = 94.2, σ1 = 8.1, μ2 = 146.3, σ2 = 26.2). For each distribution, the vertical bar indicates spike width at which the Gaussian mixture reaches its minimum between the two peaks: we chose this value as the definition for "narrow" and "broad" spikes (AON 95μs; APC 110μs). Using this criterion, we found 85% of broad spikes in the AON and 89% in the APC. **(G)** Basal firing rate of the neurons with narrow versus broad spikes. Based on the histograms in (F), spikes were categorized as narrow or broad. White circle and value next to it: median. Thick bar: quartiles. Thin bar: range of values. The neurons with narrow spikes showed a significantly higher basal activity than the broad ones. This finding, along with the presence of a bimodal spike width distribution, suggests that the neurons with narrow spikes are mostly inhibitory, while the neurons with larger spikes are mostly excitatory. The purple bar shows the same analysis performed on the bilaterally-correlated (noted "cor") neurons of the AON and APC (Kruskal-Wallis test across the "narrow", "broad', and "cor" categories: p = 3.2e-6; post-hoc Wilcoxon rank-sum test, "narrow" versus "broad" p < 0.0001, "narrow" versus "cor" p = 0.0008, "broad" versus "cor" p = 0.60). **(H)** Comparison of the spike width distribution of all AON and APC neurons versus bilaterally-correlated neurons. Unlike the overall distribution in the AON and APC, the spike width distribution for the bilaterally-correlated neurons is not significantly different from a unimodal distribution (Hartigan's dip test, p = 0.06) Almost all the bilaterally-correlated neurons have a broad spike (97% neurons have a broad spike), suggesting they are putative excitatory neurons. **(I)** PID signal amplitude at different odor dilutions. All amplitudes are reported relative to the amplitude of the dilution used for the rest of this study (5%). Each grey dot is one PID measurement (10 repeats per condition), the black dots are the average amplitudes. For each odor, the red dot indicates the dilutions at which the PID signal is about 10 times weaker than the initial 5% dilution. **(J)** Response magnitude of the initial versus weaker dilutions. Each graph is one side of the mask. Each dot is one neuron-odor pair (n = 2 mice recorded in the AON, 120 neurons). We only tested the 6 odors shown in panel I (initial dilution: 5%; new dilution: see red dot in I). Only the neuron-odor pairs significantly responding to the initial 5% concentration are shown. Red: average for each side. When we stimulated the same side of the mask with 10 times less odorant, we obtained neuronal responses about 6 to 7 times weaker than with our initial 5% concentration. Therefore, cross-contamination in the mask, if any, cannot explain the presence of side-invariant odor responses.

Figure S4: Correlation of Population Odor Representations.
Related to Figure 3. **(A)** and **(B)** Ipsi- versus contralateral responses per odor in both OC regions tested. (A) AON. (B) APC. Red line: linear regression (contra = a x ipsi + b). Solid lines show significant regressions (F-test, p < 5%). Odor # is the same as Figure S1A. Each dot is one neuron. The top left graph for each region is also shown in Figure 3G. **(C)** Same as panel (A), odor 1, except neuron identity has been randomly shuffled on each side. Note the absence of significant correlation. **(D)** Significance of the regression slope for panel (A), odor 1. We built a chance distribution by repeating the shuffling shown in panel (C) 1000 times. We then compared the actual value of the slope to this distribution to determine its significance. Here we show the distribution of chance slopes for the AON, odor 1. The results of our bootstrap strategy matched the results of the F-test mentioned above. **(E)** We performed the same analysis as panels (A) and (B), but we looked at each mouse separately. Here we report the R^2 value of each regression, as well as whether the correlation was significant (F-test). N.s. means that the correlation was not statistically significant ($p \geq 5\%$).

Figure S5: Ipsi- and Contralateral Odor Decoding.

Related to Figure 4. **(A)** Decoding accuracy with varying odor response time windows. For each data point, response vectors were built by summing spikes from 0s to the time indicated on the x-axis (responses aligned to the first sniff after odor onset, see Methods for details). This example shows the odor decoding for all neurons in the AON, train ipsi, test ipsi. Odor decoding accuracy is maximal at 2s, therefore we decided to perform all our decoding analysis over an odor response window of 2s. **(B)** Same as Figures 4B-4C, except the decoding process was applied to each mouse separately. Accuracy at n = 50 neurons is significantly higher than chance for all mice. The mice with poorer decoding accuracy were usually the ones in which neurons showed fewer / weaker responses on average, as well as fewer bilaterally-correlated neurons (see Figure S6).

Figure S6: Variability Across Mice.

Related to Figures 3 and 4. In this figure we point at factors that may explain the variability we observed across mice in terms of percentage of bilaterally-correlated neurons, as well as odor decoding accuracy. **(A)** Mean number of odors eliciting response (related to Figure 2E). **(B)** Mean response magnitude (related to Figure 2F). **(C)** Percentage of bilaterally-correlated neurons (related to Figure 3B). Solid dot: significant percentage. **(D)** Mean accuracy of the odor decoding process with model trained on responses to ipsilateral presentations (related to Figure S5B). **(E)** Similar to (D) with training on contralateral presentations.

Figure S7: Modeling Olfactory Cross Cortical Connections.

Related to Figure 6. **(A) to (C)** Same as Figures 6B-6D, except for we are comparing the distributions of mouse ipsilateral data and our simulations of ipsilateral responses with optimal alpha (0.42). Mouse data is identical to the contralateral lines from Figures 2E-2F, top panels. **(D)** Same as Figure 6I, except we decoded ipsilateral odors using the contralateral presentations. (Left) Using closest Euclidean neighbor decoding is markedly worse in simulations than mice, due mainly to the difference in magnitude of responses that results as you increase the structure parameter. (Right) Using closest angular neighbors instead recaptures better performance, and the best alpha overlaps with the AON performance. This suggests that, in the left panel, the decoder limited performance rather than lack of information. **(E) to (H)** Same as Figures 6F, 6G, 6I, and 6J respectively, except each OC layer contains 25,000 neurons (instead of 50,000 neurons in Figure 6, see Methods). These show that the optimal alpha level (0.42) is robust to changes in the number of cortical neurons simulated.

Figure S8: Low-dimensional structured connectivity is sufficient to produce alignment.
(A) and **(B)** Odor decoding accuracy using 385 neurons as Q is varied, averaged over 200 iterations of decoding and 10 different creations of G_{Struct}, **(A)** approximates using SVD and **(B)** using PCA. **(C)** – **(F)** are the same as Figure

6F, G, I, and J but using an approximate G_{Struct} created with 7 singular values. **(G) – (J)** are the same but using 6 PCA components.

Brain region	Mouse name	Age at the first recording	Number of sessions	Number of neurons
AON	AON1	3 m.o.	11	193
	AON2	3 m.o.	17	73
	AON3	3 m.o.	14	118
APC	APC1	3 m.o.	15	375
	APC2	4 m.o.	12	272
	APC3	3 m.o.	11	141
	APC4	3 m.o.	11	143
PPC	PPC1	5 m.o.	12	184
	PPC2	5 m.o.	11	155
	PPC3	4 m.o.	10	166
OB	OB1	7 m.o.	1	7
	OB2	7 m.o.	1	9
	OB3	8 m.o.	1	11
	OB4	8 m.o.	2	15

Table S1: Mice Recorded in this Study.

All tetrode recordings were performed in awake, head-restrained mice.

Figure	Test	P-value	Post-hoc test	Post-hoc p-values
Figure 1F, histogram	Z-test for the null hypothesis that the contralateral responses come from a Gaussian distribution with μ = 0 and σ = 0.56.	p = 0.84	N/A	N/A
Figure 2B	Wilcoxon rank-sum test, AON versus APC	p < 1.0e-4	N/A	N/A
Figure 2E	Wikcoxon signed-rank test, ipsi versus contra, per region	AON, p = 9.4e-3 APC, p = 0.51	N/A	N/A
Figure 2E	Wilcoxon rank-sum test, AON versus APC	"ipsi" distributions, p = 0.55 "contra" distributions, p = 0.68	N/A N/A	N/A N/A
Figure 2F	Wilcoxon signed-rank test, ipsi versus contra, per region	AON, p = 8.2e-3 APC, p = 0.94	N/A	N/A
Figure 2F	Wilcoxon rank-sum test, AON versus APC	"ipsi" distributions, p < 1.0e-4 "contra" distributions, p < 1.0e-4	N/A N/A	N/A N/A
Figure 2G	Wilcoxon rank-sum test, ipsi versus contra, per region	AON, p = 0.99 APC, p = 0.41	N/A	N/A
Figure 2G	Wilcoxon rank-sum test, AON versus APC	"ipsi" distributions, p = 0.10 "contra" distributions, p = 0.22	N/A	N/A
Figure 3A	F-test, significance of the linear model	Top row, from left to right: p = 0.41; p = 0.84; p = 0.54; p = 0.59 Bottom row, from left to right: p = 5.3e-6; p = 5.5e-4; p = 6.6e-3; p = 2.2e-4	N/A	N/A
Figure 3G	F-test, significance of the linear model	AON, p < 1.0e-4 APC, p < 1.0e-4	N/A	N/A
Figures 4B and 4C	Wikcoxon signed-rank test, distribution of accuracies obtained at n = 350 neurons versus chance accucacy (1/15).	p < 1.0e-4 for all regions, all conditions	N/A	N/A
Figure 4D	Wikcoxon signed-rank test, distribution of accuracies obtained at n = 350 neurons versus chance accucacy (1/2).	p < 1.0e-4 for all regions	N/A	N/A
Figure S1E top panel	Wilcoxon rank-sum test between the two distributions	p = 0.25	N/A	N/A
Figure S1E bottom panel	Wilcoxon rank-sum test between the two distributions	p = 0.62	N/A	N/A
Figures S3F and S3H	Hartigan dip test of unimodality	AON, p = 0.020 APC, p = 0.016 AON bilat-corr neurons, p = 0.23	N/A	N/A
Figure S3G	Kruskal-Wallis test across neuron types (narrow, broad, bilatetally-correlated)	p < 1.0e-4	Wilcoxon rank-sum test	narrow vs broad, p < 1.0e-4 narrow vs bilat-corr, p < 1.0e-4 broad vs bilat-corr, p < 1.0e-4
Figures S4A and S4B	F-test, significance of the linear model	p < 1.0e-4 for regions, all odors	N/A	N/A
Figure S4C	F-test, significance of the linear model	p = 0.42	N/A	N/A
Figure S5B	Wikcoxon signed-rank test, distribution of accuracies obtained at n = 50 neurons versus chance accucacy (1/15).	AON, p < 1.0e-4 for all mice, all conditions APC, p < 1.0e-4 for all mice, all conditions	N/A	N/A

Table S2: Statistical Tests Performed in this Study.

The statistical tests showing significant differences are in red (critical value: 5%). For post-hoc tests, the p-values reported in the table have been corrected for multiple testing problem (Bonferroni method).