A CRISPR/Cas9-BASED APPROACH FOR EDITING IMMORTALISED HUMAN MYOBLASTS TO MODEL DUCHENNE MUSCULAR DYSTROPHY IN VITRO

P. Soblechero-Martín[1], E. Albiasu-Arteta[1], A. Anton-Martinez[1], I. Garcia-Jimenez[1], G. González-Iglesias[1], I. Larrañaga-Aiestaran[1], A. López-Martinez[1], J. Poyatos-García[2], E. Ruiz-Del-Yerro[1], F. Gonzalez[3], V. Arechavala-Gomeza[1,4]

[1]Neuromuscular Disorders, Biocruces Bizkaia Health Research Institute, Barakaldo, Spain
[2]La Fe Health Research Institute, Hospital La Fe, Valencia, Spain
[3]Pluripotent Stem Cells and Activation of Endogenous Tissue Programs for Organ Regeneration (PR Lab), Institute for Bioengineering of Catalonia (IBEC), Barcelona, Spain
[4]Ikerbasque, Basque Foundation for Science, Bilbao, Spain

Address for correspondence:

Prof. Virginia Arechavala-Gomeza

Biocruces Bizkaia Health Research Institute

Barakaldo, Bizkaia, 48903, Spain

Phone: +34 946007967

E-mail: virginia.arechavalagomeza@osakidetza.eus

ORCID: 0000-0001-7703-3255

Summary

We report two novel immortalised myoblast culture models for studying Duchenne

muscular dystrophy (DMD), generated through CRISPR/Cas9 gene editing: one

recapitulates a common *DYSTROPHIN* (*DMD*) deletion and the other a regulatory

mutation leading to *UTROPHIN (UTRN)* ectopic upregulation.

Abstract

CRISPR/Cas9-mediated gene editing may allow treating and studying rare genetic

disorders by respectively, correcting disease mutations in patients, or introducing

them in cell cultures. Both applications are highly dependent on Cas9 and sgRNA

delivery efficiency. While gene editing methods are usually efficiently applied to cell

lines such as HEK293 or hiPSCs, CRISPR/Cas9 editing *in vivo* or in cultured myoblasts

prove to be much less efficient, limiting its use. After a careful optimisation of

different steps of the editing protocol, we established a consistent approach to

generate human immortalised myoblasts disease models through CRISPR/Cas9

editing. Using this protocol we successfully created a coding deletion of exon 52 of

the *DYSTROPHIN (DMD)* gene in wild type immortalised myoblasts modelling

Duchenne muscular dystrophy (DMD), and a microRNA binding sites deletion in the

regulatory region of the *UTROPHIN (UTRN)* gene leading to utrophin upregulation in

in Duchenne muscular dystrophy patient immortalised cultures. Sanger sequencing

confirmed the presence of the corresponding genomic alterations and protein

expression was characterised using myoblots. To show the utility of these cultures

as platforms for assessing the efficiency of DMD treatments, we used them to

evaluate the impact of exon skipping therapy and ezutromid treatment. Our editing

protocol may be useful to others interested in genetically manipulating myoblasts

and the resulting edited cultures for studying DMD disease mechanisms and

assessing therapeutic approaches.

KEYWORDS: Duchenne muscular dystrophy, Dystrophin, Utrophin, Gene editing

INTRODUCTION

Duchenne muscular dystrophy (DMD) is a fatal X-linked recessive disease affecting one out of 3.500-5.000 newborn males. It is commonly caused by deletions disrupting the open reading frame of the *DYSTROPHIN (DMD)* gene causing a lack of dystrophin protein (Hoffman et al., 1987). Patients carrying out of frame mutations present a severe phenotype, while those carrying in-frame mutations may result in hypomorphic alleles, a partially functional dystrophin and milder phenotypes, such as in Becker muscular dystrophy (BMD)(Anthony et al., 2011). Dystrophin plays a major role in membrane stabilization during muscle contraction, linking the actin cytoskeleton to the sarcolemma (Muntoni et al., 2003) and also contributes to extracellular signalling (Lai et al., 2009). Lack of dystrophin in DMD patients' muscles, leads to progressive muscle wasting and degeneration. DMD children suffer from loss of ambulation in the first or second decade of life and premature death by cardiac and respiratory complications (Eagle et al., 2002).

Although no definitive cure for DMD is available, three drugs have been recently approved by different regulatory agencies. Ataluren, approved by the European Medicines Agency (EMA), induces readthrough of premature stop codons during mRNA translation, generating a full length dystrophin protein (Finkel, 2010). Eteplirsen and golodirsen, approved by the US Food and Drug Administration (FDA), are antisense oligonucleotides. Eteplirsen targets *DMD* exon 51 and golodirsen, exon 53. Both antisense oligonucleotides modulate splicing by exon skipping, restoring *DMD* reading frame and leading to a shorter but functional protein (Kinali et al., 2009; Muntoni et al., 2018). Exon skipping therapies aim to attenuate the phenotype and phenocopy milder BMD-like genotypes, to potentially improve disease outcome. Ataluren, eteplirsen and golodirsen are designed for rescuing specific patient mutations only present respectively in 13%, 13% and 8% of patients (Aartsma-Rus et al., 2006). It is therefore important to test and assess alternative exon-skipping strategies targeting other *DMD* exons in different phases of clinical assays (Arechavala-Gomeza et al., 2012). Alternatively, a number of compounds applicable to all DMD patients are also under evaluation, targeting secondary DMD pathologies or trying to compensate for the lack of dystrophin.

UTROPHIN *(UTRN)* is an autosomal paralog of dystrophin, expressed in skeletal muscle
cells during embryonic development, but restricted to neuromuscular and
myotendinous junctions in the mature muscle fibre (Tinsley et al., 1992).
Overexpression of utrophin in skeletal muscle in DMD animal models can partially
compensate the lack of dystrophin and improve DMD phenotype (Cerletti et al., 2003;
Tinsley et al., 1998; Tinsley et al., 1996). Importantly, ectopic and high levels of
utrophin in myoblasts are not associated with toxicity, making utrophin upregulation
an interesting therapeutic strategy applicable to all patients, no matter their particular
mutation (Fairclough et al., 2013; Fisher et al., 2001). Ezutromid/SMT-C1100 was the
first utrophin modulator evaluated in clinical assays, but was recently abandoned due
to lack of evidence of utrophin restoration, nor clinical improvement for patients
(Ricotti et al., 2016; Tinsley et al., 2015). Alternatively, other studies proposed new
strategies to upregulate *UTRN* by removing the binding sites of microRNAs repressing
UTRN expression through gene editing (Amenta et al., 2011; Goyenvalle et al., 2010;
Morgoulis et al., 2019; Pisani et al., 2018). Finally, utrophin upregulation has been
efficiently achieved in multiple animal models with non-immunogenic response using
gene therapy (Song et al., 2019).

CRISPR/Cas9 currently represents the most efficient and versatile genome-engineering
tool, introducing small and large DNA modifications, including large genomic deletions
in different cell types and organisms (Wright et al., 2016). Hence, in the presence of
two single guide (sg) RNAs targeting two different loci on the same chromosome, Cas9
can induce two DNA double strand breaks (DDSBs), leading in some cases to deletion
of the excised DNA segment through repair by the non-homologous end joining (NHEJ)
pathway (He et al., 2015). Similar to antisense oligo-mediated exon skipping therapies
at RNA level, CRISPR/Cas9 can therefore be used to remove mutations by deleting
mutated exons and restore the open reading frame of the *DMD* gene (Amoasii et al.,
2018; Ousterout et al., 2015; Young et al., 2016). The advantage of this approach is
that the genetic modification, once introduced, is stable over cell cycles. However, it´s
efficiency is currently too low to provide a real therapeutic alternative *in vivo*, not even
mentioning immunogenicity and off-target problems linked with the use of Cas9
(Charlesworth et al., 2019) . These difficulties have been recently reported in the

golden retriever muscular dystrophy dog (GRMD), were no dystrophin restoration at protein level was obtained after gene editing using this technology (Mata Lopez et al., 2020).

In order to easily and rapidly assess the efficiency of current and novel therapies to treat DMD, *in vitro* cellular models are particularly useful. However, only a few human immortalized muscle cell lines derived from DMD patients are currently available (Mamchaoui et al., 2011). Due to the wide spectrum of DMD mutations and the difficulties to obtain DMD patient muscle biopsies, an efficient approach to edit DMD mutations in immortalised myoblasts would be extremely valuable to generate a more exhaustive panel of DMD-myoblast models. Such lines would provide a powerful resource for *in vitro* drug screening and study disease rescue mechanisms.

Here, we report an optimised CRISPR/Cas9 approach to edit myoblasts to create DMD disease models. We successfully used this protocol to created two new cell lines: in objective 1, control myoblasts were edited to remove *DMD* exon 52 (a common mutation in DMD patients); in objective 2, DMD patient's myoblasts were edited to generate a utrophin ectopic expression rescue model by deleting miRNA binding sites in *UTRN* regulatory region.

RESULTS

Optimisation of gene editing in myoblasts and generation of two new cell culture models.

- Guide selection for CRISPR/Cas9 system.

We had two different gene editing objectives: **objective 1** aimed to delete exon 52 of the *DMD* gene to generate a disease model in control immortalised human myoblasts; **objective 2** was to delete in the *UTRN* gene a binding site for *UTRN*-repressing microRNAs in DMD immortalized human myoblasts.

Our strategy to perform CRISPR/Cas9 editing in myoblasts was to design two sgRNAs flanking the region to be deleted in order to generate two DSBs leading to removal of that region (Figure 1).

- <u>Testing sgRNAs in HEK293 cultures.</u>

As transfection of myoblast is very inefficient, all the different combinations of the sgRNAs cutting before and after the target region (5x5), were tested in HEK293 cultures first (Figure 2). The combination of sgRNAs that was most efficient in HEK293 cells for each objective was selected to be used in the transfection of human immortalized myoblasts.

- <u>Myoblast transfection and single cell sorting workflow</u>

Myoblasts were transfected with the two GFP-plasmids encoding each of the selected sgRNA selected for the edition of the target region. After comparing four different transfection reagents and an electroporation method, transfection of myoblasts with ViaFect^{TM} (Promega) was selected (See Supplementary figure 1. After transfection, GFP-positive cells were sorted by FACS (See Supplementary figure 1), seeded as single cells in 96 well plates (five plates per condition) and expanded in culture (see schematic workflow in Figure 3).

- <u>Selection of edited clones:</u>

A limited number of clones derived from GFP-positive single cells grew enough for further analysis: 20/314 for objective 1 and 35/480 for objective 2. To confirm the presence of the desired deletions, a genomic PCR was performed with specific primers for each targeted gene (Figure 4 and Supplementary Table 1). Amplicons corresponding in size with the expected deletions were analysed by Sanger sequencing and the expected deletions were confirmed in all the positive clones, which corresponds to editing efficiency between 5-6%. To evaluate any potential off-target effects, each selected sgRNA was analysed *in silico* using the bioinformatics web-tool CRISPOR (Haeussler et al., 2016) . We selected the six more likely off-target sites for each sgRNA and analysed each one of them through PCR, followed by Sanger sequencing in edited clones (Table 1). We found no off-target effects in any of the 12 sites studied for each clone sites (Supplementary figure 2).

Analysis of dystrophin and utrophin expression in edited clones.

We compared dystrophin expression in myotubes of the DMDΔ52-Model to controls and DMD cultures and confirmed that it was abolished by immunocytochemistry

(Figure 5A), western blot analysis (Figure 5B) and myoblots (Figure 5C). Dystrophin

levels in this model, where exon 52 had been removed by CRISPR/Cas9 editing, were

statistically no different than those seen in a culture from a DMD patient.

It was difficult to corroborate by immunocytochemistry the increase of utrophin

expression between unedited DMD and DMD-UTRN-Model myotubes (Figure 5D) but

this increase was evident by western blot (a 175% increase, Figure 5E) and myoblot

analysis (close to 50% increase, Figure 5F). We also evaluated myoblast differentiation

in all the cultures by myoblot, and we could observe a decrease in the MF20

differentiation marker in all the edited clones, no matter the deletion, compared with

their corresponding controls (Supplementary Figure 3).

Evaluation of therapies in newly generated model cell lines.

To assess if the DMDΔ52-Model cell culture could be useful to test potential mutation

specific therapies for DMD, we evaluated the exon skipping efficiency of an antisense

oligonucleotide in this culture. We treated the DMDΔ52-Model cultures with an

antisense oligonucleotide drug that can skip exon 51 (van Deutekom et al., 2007) and

restore *DMD* open reading frame. After treatment with this drug, we confirmed that

exon skipping had taken place at RNA level (Figure 6A), and the restoration of

dystrophin expression by myoblot analysis (Figure 6B).

To test our DMD-UTRN-Model as a positive utrophin overexpression control, we

cultured it alongside the original unedited DMD cultures, which we treated with

several concentrations of ezutromid and we evaluated the expression of utrophin in all

cultures. We observed that utrophin was hardly modified in DMD cultures treated with

ezutromid while a robust overexpression was confirmed in the DMD-UTRN-Model

compared to the unedited DMD cultures. (Figure 6C)

DISCUSSION

CRISPR/Cas9 as a possible therapeutic approach for DMD has been explored in the

past in *in vitro* models of the disease. In most of these cases, a "permanent" exon

skipping approach was selected where a shorter protein would be produced

(Ousterout et al., 2015; Young et al., 2016), while in some others full length dystrophin

was the result of the edition (Lattanzi et al., 2017; Li et al., 2014; Wojtal et al., 2015).

Several studies have shown efficacy in mice models (Long et al., 2016; Nelson et al.,

2016; Tabebordbar et al., 2016), an most recently in dogs, which is currently the most

advanced example of its application to DMD (Amoasii et al., 2018) but there are also

examples of the challenges associated to gene editing (Mata Lopez et al., 2020). Before

this can be a credible therapeutic option, several delivery and manufacturing problems

will need to be overcome. In the meantime, this methodology is very useful for

researchers looking for disease models: muscle biopsies are not routinely collected

during diagnosis of this disorder and seldom cultured. This means there are few good

culture models of the disease. Another tool created to facilitate research, was the

immortalisation of some of the available cultures (Mamchaoui et al., 2011), which

increases the possibility of performing more experiments with a given culture. We

have used immortalised cultures to further increase this possibility.

Our preliminary experiments to perform gene editing in the *DMD* gene of HEK293 cells

were successful. However, a big roadblock was encountered in the form of very poor

transfection efficiency of our chosen gene editing reagents in myoblasts (Ousterout et

al., 2015; Wojtal et al., 2015). This is a problem shared by many research laboratories,

and we consider that our protocol, although still not very efficient, may help those

facing the same difficulties. We have successfully applied this protocol to edit 2

different regions in two different cell backgrounds (Control and DMD), and we

consider that those described and fully characterised in this manuscript could be

relevant research models that we would be happy to share. The first of these models is

an immortalised DMD disease cell culture model, (DMDΔ52-Model) that lacks exon 52

of the *DMD* gene, which disrupts the ORF and dystrophin expression. This model could

be useful to evaluate mutation-independent drug treatments, and also exon skipping

drugs that aim to skip exons 51 or 53 (Arechavala-Gomeza et al., 2007; Popplewell et

al., 2010) as skipping either exon in this case would restore the ORF and dystrophin

expression. We have demonstrated that DMDΔ52-Model lacks dystrophin expression

and that this can be reverted through treatment with an exon 51 skipping drug.

An immortalised cell culture model constitutively expressing utrophin, DMD-UTRN-

Model, is both a proof of principle of a possible therapeutic option to overexpress

utrophin as a substitute for dystrophin, and a valuable research tool. The search of
drugs that could overexpress utrophin is ongoing (Guiraud and Davies, 2017) after the
recent failure of ezutromid to show relevant results in clinical trials. Many research
projects, including drug re-purposing screening, are ongoing to find new candidates to
test in clinical trials. However, there are no reliable positive controls that could be used
to compare such treatments. We propose that our cell model could serve that
purpose, offering researchers useful custom controls for their studies. We have tested
this hypothesis and compared the stable utrophin overexpression quantified in the
DMD-UTRN-Model with the one that is obtained after treatment with ezutromid, the
lead market candidate in this field until very recently, with positive results. Previous
studies in muscle sections show that DMD patients already overexpress utrophin, in
many cases 4 to 5-fold the levels seen in control muscle sections (Arechavala-Gomeza
et al., 2010). Our choice to target this particular *UTR* region, increases basal
overexpression in DMD cultures, and the amount of overexpression varies significantly
when evaluated by western blot (more than 2.5 times) or our preferred method,
myoblots (close to 1.5 times). We like to consider that myoblot evaluation reflects
more closely the actual protein expression, as it is not subjected to many of the
inherent problems of western blotting when evaluating very large proteins (Ruiz-Del-
Yerro et al., 2018). This is why we cannot comment yet on the differences in
expression between our study and other published studies that also aimed to
overexpress utrophin by gene editing, but which 1) targeted different promoter
regions (*UTRN* A or *UTRN* B) of utrophin and 2) evaluated their results by western blot
analysis (Wojtal et al., 2015). We would be interested on studying this matter further
to analyse the differences in utrophin expression when targeting different regions.

As a conclusion, we have optimised a gene editing method to be applied to myoblasts,
a rather difficult target, and expect our experience to be useful to other muscle
researchers. The two new cell culture models described are already being used as
efficient tools in our search for new therapies for DMD and we are looking forward
expanding those experiments.

 MATERIALS AND METHODS

CRISPR/Cas9 tools

Each edition requires 2 sgRNAs. Specific sgRNA guides were designed using the online bioinformatics tool http://crispr.mit.edu (Ran et al., 2013). Ten different guides (five before and five after the target region) were designed targeting exon 52 flanking regions in *DMD* gene and another ten targeting a repressor binding site in the UTR 3' region of *UTRN* gene and selected according to their score number (Figure 1). They were cloned into a plasmid containing *Cas9* from *S. pyogenes* with 2A-*EGFP* pSpCas9(BB)-2A-GFP (PX458) (Addgene plasmid # 48138, deposited by Feng Zhang. All sgRNAs were cloned using BbsI sites.

Cell cultures

Immortalized myoblasts derived from muscle biopsies from healthy controls and DMD patients were provided by the CNMD Biobank, London, UK and the Institut de Myologie Paris, France, and cultured using skeletal muscle medium (SMM) (Promocell, Germany) and differentiation media as previously described (Ruiz-Del-Yerro et al., 2018).

Human embryonic kidney 293 (HEK 293) cells, used in the preliminary selection of the best sgRNA combinations for our experiments, were purchased from the European Collection of Authenticated Cell cultures (ECCAC) via Sigma-Aldrich, Spain, and maintained following the manufacturer's protocols.

Cell culture transfection

All different sgRNAs combinations were transfected into HEK 293 cells using lipofectamine 2000® (Thermo Fisher Scientific), according to the manufacturer's protocol. Myoblasts seeded in 6 well plates at 70-80% confluence were transfected with 1.5ug of each plasmid with the most efficient guide RNA combination using ViaFect™ (Promega) transfection reagent (1:5 ratio).

FACS Sorting of GFP positive myoblasts:

48 hours after transfection myoblasts were trypsinised and collected for FACS sorting (fluorescence activated cell sorting), at the Cell Analytics Facility (BD FACS Jazz)

Achucarro Basque Center for Neuroscience - (Leioa, Spain). GFP-positive cells were

seeded individually in 96 well plates for clonal selection. The first colonies were visible

around 7 days post-sorting. Clones were expanded from single cell to near-confluence

and expanded into larger well plates to be harvested 15-30 days post-sorting.

Myoblasts often developed elongated and stressed shapes during this clonal expansion

after single cell sorting. Harvested cultures were aliquoted: some aliquots were frozen

for archival; others were pelleted for DNA analysis, while replicates were cultured

further for characterization by immunocytochemistry, western blot and myoblots.

Analysis of gene editing

DNA was extracted from cell pellets using a QIAamp® DNA Mini Kit, Qiagen. PCR

amplification targeting the edited regions was carried out using Taq DNA Polymerase

(Recombinant), Invitrogen, under the following conditions: preheating 3' 94ºC, 25

cycles of 94º for 3', 94º 20'', 63º 20'', 72º 1' and 72º 5' and DMD-Seq-D52-DOWN-F2

and DMD-Seq-D52-DOWN-R2 primers (see Table 1). PCR products were resolved in 2%

TAE-agarose gels and purified with QIAquick® Gel extraction Kit, Qiagen. PCR

amplicons corresponding to the expected length were analysed by Sanger sequencing

at the sequencing platform of Biocruces Bizkaia Health Research Institute using DMD-

Seq-D52-DOWN-F2 and DMD-Seq-D52-DOWN-R2 primers (see Table 1).

Off-target analysis of mutations in clonal lines

Potential off-target region loci of each sgRNA used were predicted using CRISPOR

bioinformatics tool http://crispor.tefor.net/. The six most probable off-target

sequences per guide were analysed in the edited clones using genomic PCR and Sanger

sequencing. Primer sets flanking off-target sites and the corresponding internal

primers used for Sanger sequencing are listed in table 2.

Primary Antibodies

Anti-dystrophin: Dys1 (Leica Biosystems), Mandys1, Mandys106 (The MDA Monoclonal

Antibody Resource)

Anti-utrophin: Mancho7 (The MDA Monoclonal Antibody Resource)

Immunostaining assays

Immunostaining assays

Original and edited clones for objectives 1 (DMDΔ52-Model) and 2 (DMD-UTRN-
Model) were cultured and immunostained for dystrophin or utrophin expression.
Edited clones were seeded into chamber slides and treated with a MyoD virus,
(Applied Biological Materials Inc, Canada) to facilitate differentiation into myotubes
(Roest et al., 1996). After seven days differentiating, samples were fixed with 4% PFA.
Cultures were permeabilised with Triton 0.5% and then blocked for half an hour with
BSA 2%. Afterwards, immunostaining was performed overnight at 4ºC with the
required antibodies. Primary antibodies used for dystrophin staining were a mix of
Dys1, Mandys1 and Mandys106 at 1:100 dilution and for utrophin staining was
Mancho 7 diluted at 1:50. The following day, after being washed with PBS Tween 0.1%,
cells were stained with Alexa Fluor 488 goat anti-mouse (Invitrogen) for 1 hour at
room temperature. Hoechst 1/2000 was used for nuclei staining and chamber slides
were mounted with PermaFluor™ Aqueous Mounting Medium (Thermoscientific).
Images were captured with a LEICA DMI 6000B microscope at the Microscopy Platform
at Biocruces Bizkaia Health Research Institute.

In-cell western assay (myoblots)

Myoblots were performed as described before (Ruiz-Del-Yerro et al., 2018). In short,
clones were seeded in 96-well plates and incubated for 24 h in SMMC, after which they
were treated with MyoD virus and incubated in differentiation media for 7 days. Then,
plates were fixed with ice-cold methanol, permeabilised and blocked before incubation
with the required primary antibodies overnight: anti-dystrophin mix (Dys1, Mandys1
and Mandys106 at 1:100), anti-utrophin (Mancho 7 antibody at 1:50), and anti-myosin
heavy chain antibody (MF20 at 1:100) that was used to evaluate differentiation. Next
day, plates were incubated with the secondary antibodies. Biotin-mediated
amplification (Abcam 6788 goat antimouse IgG biotin 1:2000) was used to increase
dystrophin signal. Secondary antibodies, IRDye 800cw streptavidin 1:2000 and IRDye
800CW goat anti-mouse 1:500, were prepared together with CellTag 700 Stain (LI-
COR® Biosciences) at 1:1000 dilution and incubated for 1 hour at RT and protected

from light. After incubation, plates were analysed using the Odyssey® CLx Imager (LI-
COR® Biosciences).

Treatment with antisense exon skipping drugs

Cultures in 96 wells and P6 wells were treated with a 2'MOE-phosphorotioate
antisense oligonucleotide (AO) aiming to skip *DMD* exon 51 (5⍰-[
T*C*A*A*G*G*A*A*G*A*T*G*G*C*A*T*T*T*C*T]-3⍰, Eurogentec, Belgium) by
transfection with Lipofectamine as described in (Arechavala-Gomeza et al., 2007;
Popplewell et al., 2010) and analysed by either myoblot (96 well plates) or RT-PCR
(pellets extracted from 6 well plates).

RT-PCR

RNA was extracted from cell pellets (RNeasy mini kit, Qiagen) according to the
manufacturer's protocol. Reverse transcription of the samples was performed using
(SuperScrip^{TM} IV Reverse Transcriptase, Invitrogen) according to the manufacturer's
protocol. cDNA samples were amplified by nested PCR using specific primers sets
(supplementary table 1) and Taq DNA Polymerase (Recombinant), Invitrogen, as
described in (Arechavala-Gomeza et al., 2007). Samples were resolved in TAE-agarose
and PCR amplicons of interest were first analysed with Gel Doc TM EZ Imager, BIORAD
and then purified with (QIAquick® Gel extraction Kit, QIAGEN) for sequencing analysis.
Before DNA extractions bands were semi-quantified using Image J.

Treatment with utrophin overexpression drugs

Ezutromid was diluted first in DMSO and finally in differentiation medium to different
concentrations and added to myoblasts in 96 well plates 7 days after differentiation.
Twenty four hours after treatment, medium was removed and plates were fixed with
ice-cold methanol for myoblot analysis.

Western blot

Cell cultures were seeded into P6 plates and trypsinized after 7 days of differentiation.
Then, cell pellets were solubilized in lysis/loading buffer and denatured at 95°C for 5
min. The samples were loaded onto a NuPAGE® Novex® 3–8% Tris-Acetate Gel3–8%
(Thermo Fisher Scientific) and run in Novex Tris-Acetate SDS Running Buffer (Thermo

Fisher Scientific) for 60 min at 70 V + 120 min at 150 V at 4ºC. Protein wet transfer was

performed overnight at 4°C using an Immobilon®-FL PVDF membrane (Merck™). Next

day, membranes were stained with Revert ™ 700 Total Protein Stain (Li-Cor) for total

protein measurement, blocked with Intercept (PBS) Blocking Buffer (Li-Cor) for 2 hours

and incubated overnight at 4°C with the primary antibodies (1/200 anti-dystrophin

antibody Abcam15277, 1/50 anti-utrophin antibody Mancho 7 or 1/500 anti a-actinin

antibody, Sigma-Aldrich A7732). After washing steps with PBS-Tween 0.1%,

membranes were incubated with secondary antibodies for 1 hour (1/5000 IRDye

800CW goat anti-rabbit 926–32211 or IRDye 680RD goat anti-mouse 926-68070, Li-

Cor) at room temperature, washed again and scanned using an Odyssey Clx imaging

system. Bands quantification was performed using Image Studio ™ software.

Statistical analysis

Mann-Whitney U test was used throughout this study to calculate P values for

determination of statistical significance (**P values<0.05, ****p-value<0.0001).

Statistical analysis was performed using GraphPad Prism software.

ACKNOWLEDGEMENTS

We acknowledge the use of cell cultures provided by the Queen Square Centre for

Neuromuscular Disorders BioBank (CNMD Biobank, London, UK) and the Institut de

Myologie (Paris, France, immortalised cultures).

We gratefully acknowledge the use of the antibodies provided by Professor Glenn

Morris from the Muscular Dystrophy Association (MDA) Monoclonal Antibody

Resource, which distributes free antibodies for research in neuromuscular diseases

worldwide from Oswestry, United Kingdom.

The MF20 antibody developed by D.A. Fischman, Weill Cornell Medical College, was

obtained from the Developmental Studies Hybridoma Bank, created by the NICHD of

the NIH and maintained at The University of Iowa, Department of Biology, Iowa City, IA

52242.

Cell sorting experiments were performed at the Cell Analytics Facility at Achucarro-
Basque Centre of Neuroscience (Leioa, Spain).

We would like to thank Ms. Karmele Alapont-Celaya for her assistance in the
completion of some laboratory tasks during this project. Drs. Gustavo Pérez-Nanclares
and Ana Belén de la Hoz at the Genetics-Genomics Facility, Biocruces Bizkaia Health
Research Institute, are also gratefully acknowledged for their help with sequencing
reactions. Special thanks are due to Drs. Maria Dolores-García Vázquez at the Cell
Culture Facility and Dr Javier Díez-García at the Microscopy Facility, both at Biocruces
Bizkaia Health Research Institute, for their assistance at their corresponding platforms.

COMPETING INTERESTS

The authors declare no competing interests

FUNDING
This work was supported by funding from Health Institute Carlos III (ISCIII, Spain) and
the European Regional Development Fund, (ERDF/FEDER), 'A way of making Europe':
Grant PI15/00333; Basque Government (grants 2016111029 and 2016111091) and
Duchenne Parent Project Spain (grant 05/2016). P. S-M holds a Rio Hortega Fellowship
from ISCIII (CM19/00104). V.A-G holds a Miguel Servet Fellowship from the ISCIII
(CPII17/00004), part-funded by ERDF/FEDER. V.A-G also acknowledges funding from
Ikerbasque (Basque Foundation for Science). F. G is supported by a Ramon y Cajal
Grant in Biomedicine (RYC-2014-16751) from the Ministry of Economy and
Competitivity (MINECO), Spain.

DATA AVAILABILITY

Under request

AUTHOR CONTRIBUTIONS STATEMENT
P S-M, formal analysis, investigation, data curation, writing or original draft, review and
editing, visualization, supervision. E. A-A, investigation, data curation, writing -review
and editing. A. A-M, investigation, writing - review and editing. I. G-J, methodology,
investigation, review and editing, supervision. G. G-I, investigation, writing - review
and editing. I. L-A, investigation, writing - review and editing. A. L-M, investigation,
writing - review and editing. J. P-G, investigation, writing - review and editing. E. R-D-
Y, methodology, investigation, writing-review and editing, supervision. F. G

conceptualization, formal analysis, writing-review and editing. V. A-G:

conceptualization, methodology, formal analysis, investigation, resources, data

curation, writing or original draft, review and editing, visualization, supervision, project

administration and funding acquisition.

REFERENCES

Aartsma-Rus, A., Van Deutekom, J. C., Fokkema, I. F., Van Ommen, G. J. and Den
Dunnen, J. T. (2006). Entries in the Leiden Duchenne muscular dystrophy mutation database:
an overview of mutation types and paradoxical cases that confirm the reading-frame rule.
Muscle Nerve **34**, 135-44.

Amenta, A. R., Yilmaz, A., Bogdanovich, S., McKechnie, B. A., Abedi, M., Khurana, T.
S. and Fallon, J. R. (2011). Biglycan recruits utrophin to the sarcolemma and counters
dystrophic pathology in mdx mice. *Proc Natl Acad Sci U S A* **108**, 762-7.

Amoasii, L., Hildyard, J. C. W., Li, H., Sanchez-Ortiz, E., Mireault, A., Caballero, D.,
Harron, R., Stathopoulou, T. R., Massey, C., Shelton, J. M. et al. (2018). Gene editing restores
dystrophin expression in a canine model of Duchenne muscular dystrophy. *Science*.

Anthony, K., Cirak, S., Torelli, S., Tasca, G., Feng, L., Arechavala-Gomeza, V., Armaroli,
A., Guglieri, M., Straathof, C. S., Verschuuren, J. J. et al. (2011). Dystrophin quantification and
clinical correlations in Becker muscular dystrophy: implications for clinical trials. *Brain* **134**,
3547-59.

Arechavala-Gomeza, V., Anthony, K., Morgan, J. and Muntoni, F. (2012). Antisense
oligonucleotide-mediated exon skipping for Duchenne muscular dystrophy: progress and
challenges. *Curr Gene Ther* **12**, 152-60.

Arechavala-Gomeza, V., Graham, I. R., Popplewell, L. J., Adams, A. M., Aartsma-Rus,
A., Kinali, M., Morgan, J. E., van Deutekom, J. C., Wilton, S. D., Dickson, G. et al. (2007).
Comparative analysis of antisense oligonucleotide sequences for targeted skipping of exon 51
during dystrophin pre-mRNA splicing in human muscle. *Hum Gene Ther* **18**, 798-810.

Arechavala-Gomeza, V., Kinali, M., Feng, L., Brown, S. C., Sewry, C., Morgan, J. E. and
Muntoni, F. (2010). Immunohistological intensity measurements as a tool to assess
sarcolemma-associated protein expression. *Neuropathol Appl Neurobiol* **36**, 265-74.

Cerletti, M., Negri, T., Cozzi, F., Colpo, R., Andreetta, F., Croci, D., Davies, K. E.,
Cornelio, F., Pozza, O., Karpati, G. et al. (2003). Dystrophic phenotype of canine X-linked
muscular dystrophy is mitigated by adenovirus-mediated utrophin gene transfer. *Gene Ther*
10, 750-7.

Charlesworth, C. T., Deshpande, P. S., Dever, D. P., Camarena, J., Lemgart, V. T.,
Cromer, M. K., Vakulskas, C. A., Collingwood, M. A., Zhang, L., Bode, N. M. et al. (2019).
Identification of preexisting adaptive immunity to Cas9 proteins in humans. *Nat Med* **25**, 249-
254.

Eagle, M., Baudouin, S. V., Chandler, C., Giddings, D. R., Bullock, R. and Bushby, K.
(2002). Survival in Duchenne muscular dystrophy: improvements in life expectancy since 1967
and the impact of home nocturnal ventilation. *Neuromuscul Disord* **12**, 926-9.

Fairclough, R. J., Wood, M. J. and Davies, K. E. (2013). Therapy for Duchenne muscular
dystrophy: renewed optimism from genetic approaches. *Nat Rev Genet* **14**, 373-8.

Finkel, R. S. (2010). Read-through strategies for suppression of nonsense mutations in
Duchenne/ Becker muscular dystrophy: aminoglycosides and ataluren (PTC124). *J Child Neurol*
25, 1158-64.

Fisher, R., Tinsley, J. M., Phelps, S. R., Squire, S. E., Townsend, E. R., Martin, J. E. and
Davies, K. E. (2001). Non-toxic ubiquitous over-expression of utrophin in the mdx mouse.
Neuromuscul Disord **11**, 713-21.

Goyenvalle, A., Babbs, A., Powell, D., Kole, R., Fletcher, S., Wilton, S. D. and Davies,
K. E. (2010). Prevention of dystrophic pathology in severely affected dystrophin/utrophin-
deficient mice by morpholino-oligomer-mediated exon-skipping. *Mol Ther* **18**, 198-205.

Guiraud, S. and Davies, K. E. (2017). Pharmacological advances for treatment in
Duchenne muscular dystrophy. *Curr Opin Pharmacol* **34**, 36-48.

Haeussler, M., Schönig, K., Eckert, H., Eschstruth, A., Mianné, J., Renaud, J.-B.,
Schneider-Maunoury, S., Shkumatava, A., Teboul, L., Kent, J. et al. (2016). Evaluation of off-

target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. *Genome Biology* **17**, 148-148.

He, Z., Proudfoot, C., Mileham, A. J., McLaren, D. G., Whitelaw, C. B. and Lillico, S. G. (2015). Highly efficient targeted chromosome deletions using CRISPR/Cas9. *Biotechnol Bioeng* **112**, 1060-4.

Hoffman, E. P., Brown, R. H., Jr. and Kunkel, L. M. (1987). Dystrophin: the protein product of the Duchenne muscular dystrophy locus. *Cell* **51**, 919-28.

Kinali, M., Arechavala-Gomeza, V., Feng, L., Cirak, S., Hunt, D., Adkin, C., Guglieri, M., Ashton, E., Abbs, S., Nihoyannopoulos, P. et al. (2009). Local restoration of dystrophin expression with the morpholino oligomer AVI-4658 in Duchenne muscular dystrophy: a single-blind, placebo-controlled, dose-escalation, proof-of-concept study. *Lancet Neurol* **8**, 918-28.

Lai, Y., Thomas, G. D., Yue, Y., Yang, H. T., Li, D., Long, C., Judge, L., Bostick, B., Chamberlain, J. S., Terjung, R. L. et al. (2009). Dystrophins carrying spectrin-like repeats 16 and 17 anchor nNOS to the sarcolemma and enhance exercise performance in a mouse model of muscular dystrophy. *J Clin Invest* **119**, 624-35.

Lattanzi, A., Duguez, S., Moiani, A., Izmiryan, A., Barbon, E., Martin, S., Mamchaoui, K., Mouly, V., Bernardi, F., Mavilio, F. et al. (2017). Correction of the Exon 2 Duplication in DMD Myoblasts by a Single CRISPR/Cas9 System. *Mol Ther Nucleic Acids* **7**, 11-19.

Li, H. L., Fujimoto, N., Sasakawa, N., Shirai, S., Ohkame, T., Sakuma, T., Tanaka, M., Amano, N., Watanabe, A., Sakurai, H. et al. (2014). Precise Correction of the Dystrophin Gene in Duchenne Muscular Dystrophy Patient Induced Pluripotent Stem Cells by TALEN and CRISPR-Cas9. *Stem Cell Reports*.

Long, C., Amoasii, L., Mireault, A. A., McAnally, J. R., Li, H., Sanchez-Ortiz, E., Bhattacharyya, S., Shelton, J. M., Bassel-Duby, R. and Olson, E. N. (2016). Postnatal genome editing partially restores dystrophin expression in a mouse model of muscular dystrophy. *Science* **351**, 400-3.

Mamchaoui, K., Trollet, C., Bigot, A., Negroni, E., Chaouch, S., Wolff, A., Kandalla, P. K., Marie, S., Di Santo, J., St Guily, J. L. et al. (2011). Immortalized pathological human myoblasts: towards a universal tool for the study of neuromuscular disorders. *Skelet Muscle* **1**, 34.

Mata Lopez, S., Balog-Alvarez, C., Vitha, S., Bettis, A. K., Canessa, E. H., Kornegay, J. N. and Nghiem, P. P. (2020). Challenges associated with homologous directed repair using CRISPR-Cas9 and TALEN to edit the DMD genetic mutation in canine Duchenne muscular dystrophy. *PLoS ONE* **15**, e0228072.

Morgoulis, D., Berenstein, P., Cazacu, S., Kazimirsky, G., Dori, A., Barnea, E. R. and Brodie, C. (2019). sPIF promotes myoblast differentiation and utrophin expression while inhibiting fibrosis in Duchenne muscular dystrophy via the H19/miR-675/let-7 and miR-21 pathways. *Cell Death Dis* **10**, 82.

Muntoni, F., Frank, D., Sardone, V., Morgan, J., Schnell, F., Charleston, J., Desjardins, C., Phadke, R., Sewry, C., Popplewell, L. et al. (2018). Golodirsen Induces Exon Skipping Leading to Sarcolemmal Dystrophin Expression in Duchenne Muscular Dystrophy Patients With Mutations Amenable to Exon 53 Skipping (S22.001). *Neurology* **90**.

Muntoni, F., Torelli, S. and Ferlini, A. (2003). Dystrophin and mutations: one gene, several proteins, multiple phenotypes. *Lancet Neurol* **2**, 731-40.

Nelson, C. E., Hakim, C. H., Ousterout, D. G., Thakore, P. I., Moreb, E. A., Castellanos Rivera, R. M., Madhavan, S., Pan, X., Ran, F. A., Yan, W. X. et al. (2016). In vivo genome editing improves muscle function in a mouse model of Duchenne muscular dystrophy. *Science* **351**, 403-7.

Ousterout, D. G., Kabadi, A. M., Thakore, P. I., Majoros, W. H., Reddy, T. E. and Gersbach, C. A. (2015). Multiplex CRISPR/Cas9-based genome editing for correction of dystrophin mutations that cause Duchenne muscular dystrophy. *Nat Commun* **6**, 6244.

Pisani, C., Strimpakos, G., Gabanella, F., Di Certo, M. G., Onori, A., Severini, C.,
Luvisetto, S., Farioli-Vecchioli, S., Carrozzo, I., Esposito, A. et al. (2018). Utrophin up-
regulation by artificial transcription factors induces muscle rescue and impacts the
neuromuscular junction in mdx mice. *Biochim Biophys Acta*.
Popplewell, L. J., Adkin, C., Arechavala-Gomeza, V., Aartsma-Rus, A., de Winter, C. L.,
Wilton, S. D., Morgan, J. E., Muntoni, F., Graham, I. R. and Dickson, G. (2010). Comparative
analysis of antisense oligonucleotide sequences targeting exon 53 of the human DMD gene:
Implications for future clinical trials. *Neuromuscul Disord* **20**, 102-10.
Ran, F. A., Hsu, P. D., Wright, J., Agarwala, V., Scott, D. A. and Zhang, F. (2013).
Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* **8**, 2281-2308.
Ricotti, V., Spinty, S., Roper, H., Hughes, I., Tejura, B., Robinson, N., Layton, G.,
Davies, K., Muntoni, F. and Tinsley, J. (2016). Safety, Tolerability, and Pharmacokinetics of
SMT C1100, a 2-Arylbenzoxazole Utrophin Modulator, following Single- and Multiple-Dose
Administration to Pediatric Patients with Duchenne Muscular Dystrophy. *PLoS ONE* **11**,
e0152840.
Roest, P. A., van der Tuijn, A. C., Ginjaar, H. B., Hoeben, R. C., Hoger-Vorst, F. B.,
Bakker, E., den Dunnen, J. T. and van Ommen, G. J. (1996). Application of in vitro Myo-
differentiation of non-muscle cells to enhance gene expression and facilitate analysis of muscle
proteins. *Neuromuscul Disord* **6**, 195-202.
Ruiz-Del-Yerro, E., Garcia-Jimenez, I., Mamchaoui, K. and Arechavala-Gomeza, V.
(2018). Myoblots: dystrophin quantification by in-cell western assay for a streamlined
development of Duchenne muscular dystrophy (DMD) treatments. *Neuropathol Appl Neurobiol*
44, 463-473.
Song, Y., Morales, L., Malik, A. S., Mead, A. F., Greer, C. D., Mitchell, M. A., Petrov, M.
T., Su, L. T., Choi, M. E., Rosenblum, S. T. et al. (2019). Non-immunogenic utrophin gene
therapy for the treatment of muscular dystrophy animal models. *Nat Med* **25**, 1505-1511.
Tabebordbar, M., Zhu, K., Cheng, J. K. W., Chew, W. L., Widrick, J. J., Yan, W. X.,
Maesner, C., Wu, E. Y., Xiao, R., Ran, F. A. et al. (2016). In vivo gene editing in dystrophic
mouse muscle and muscle stem cells. *Science* **351**, 407-411.
Tinsley, J., Deconinck, N., Fisher, R., Kahn, D., Phelps, S., Gillis, J. M. and Davies, K.
(1998). Expression of full-length utrophin prevents muscular dystrophy in mdx mice. *Nat Med*
4, 1441-4.
Tinsley, J., Robinson, N. and Davies, K. E. (2015). Safety, tolerability, and
pharmacokinetics of SMT C1100, a 2-arylbenzoxazole utrophin modulator, following single-
and multiple-dose administration to healthy male adult volunteers. *J Clin Pharmacol* **55**, 698-
707.
Tinsley, J. M., Blake, D. J., Roche, A., Fairbrother, U., Riss, J., Byth, B. C., Knight, A. E.,
Kendrick-Jones, J., Suthers, G. K., Love, D. R. et al. (1992). Primary structure of dystrophin-
related protein. *Nature* **360**, 591-3.
Tinsley, J. M., Potter, A. C., Phelps, S. R., Fisher, R., Trickett, J. I. and Davies, K. E.
(1996). Amelioration of the dystrophic phenotype of mdx mice using a truncated utrophin
transgene. *Nature* **384**, 349-53.
van Deutekom, J. C., Janson, A. A., Ginjaar, I. B., Frankhuizen, W. S., Aartsma-Rus, A.,
Bremmer-Bout, M., den Dunnen, J. T., Koop, K., van der Kooi, A. J., Goemans, N. M. et al.
(2007). Local dystrophin restoration with antisense oligonucleotide PRO051. *N Engl J Med* **357**,
2677-86.
Wojtal, D., Kemaladewi, D. U., Malam, Z., Abdullah, S., Wong, T. W., Hyatt, E.,
Baghestani, Z., Pereira, S., Stavropoulos, J., Mouly, V. et al. (2015). Spell Checking Nature:
Versatility of CRISPR/Cas9 for Developing Treatments for Inherited Disorders. *Am J Hum Genet*.
Wright, A. V., Nunez, J. K. and Doudna, J. A. (2016). Biology and Applications of
CRISPR Systems: Harnessing Nature's Toolbox for Genome Engineering. *Cell* **164**, 29-44.

Young, C. S., Hicks, M. R., Ermolova, N. V., Nakano, H., Jan, M., Younesi, S.,
Karumbayaram, S., Kumagai-Cresse, C., Wang, D., Zack, J. A. et al. (2016). A Single CRISPR-
Cas9 Deletion Strategy that Targets the Majority of DMD Patients Restores Dystrophin
Function in hiPSC-Derived Muscle Cells. *Cell Stem Cell* **18**, 533-40.

Figure legends

Table 1. Off-target sites analyses.

Top 6 off-target sequences of Ob1sgRNA2, Ob1sgRNA6, Ob2sgRNA22 and

Ob2_sgRNA26 identified with CRISPOR webtool, including the mismatches between

sgRNAs, the off-target sequence, the chromosomes and loci targeted. All of them were

analysed by PCR and Sanger sequencing, and no off-targets were detected.

Figure 1. Editing approach and sgRNA design.

(A and B). Schematic representation of our strategy for editing the *DMD* (A) and the

UTRN loci (B). A pair of flanking sgRNAs are co-transfected in order to delete *DMD*

exon 52 (A) or the microRNA cluster contained in the 5'UTR of *UTRN* (B). (C and D). List

of sgRNAs for editing the *DMD* (C) and the *UTRN* loci (D), showing sgRNAs sequences,

PAM sequences and scores of all sgRNA tested. (E and F) Genomic location of the

sgRNAs targeting the *DMD* (E) and *UTRN* loci (F).

Figure 2. sgRNAs pairs test in HEK293 cells.

(A and B). Representation of all the different sgRNAs combinations tested for editing

the *DMD* (A) and the *UTRN* loci (B). (C and D) Representative PCR analysis of HEK293

cells transfected with some of the sgRNAs combinations tested. Upper bands

correspond to wild type or non-edited cells, while the lower bands correspond to the

edited ones. Selected combinations are highlighted: Obj1sgRNA2 (C); Obj2

sgRNA22+sgRNA26 (D)

Figure 3. Cloning and editing workflow, and efficiency diagram.

(A) Scheme of the workflow followed to obtain the edited myoblast cell lines. 48 hours

after plasmid transfection GFP positive myoblasts were single cell sorted using FACS.

Clones were expanded until confluence for DNA extraction. (B) Efficiency of the

different steps during the workflow process.

Figure 4. Genotyping *DMD* and *UTRN* deletion breakpoints in edited myoblast clones.

(A and B) PCR genotyping (A) and Sanger sequencing (B) of *DMD* edited clones (A). (C
and D) PCR genotyping (C) and Sanger sequencing (D) of *UTRN* edited clones (C).
Larger products in agarose gels (A and C) indicate non-edited clones, and shorter ones
correspond with the expected deletion. (B and D) Sequences of the smaller bands
confirmed the expected gene editing for objective 1: DMDΔ52-Model and objective 2:
DMD-UTRN-Model (B).

**643 Figure 5. Functional consequences of gene editing: dystrophin and utrophin
expression.**

Dystrophin expression in control myotubes compared to DMDΔ52-Model cultures and
DMD myotubes studied by immunocytochemistry (A), western blotting (B) and
myoblots (C) where n=24 wells per cell type were compared (****p-value<0.0001).
Utrophin expression in DMD myotubes compared to DMD-UTRN-Model studied by
immunocytochemistry (D), western blotting (E) and myoblots (F), where n=48 replicate
wells were included per cell type. (****p-value<0.0001) (P values were determined
with Mann-Whitney U test).

652 Figure 6. Evaluation of potential therapies in the generated cell culture models.

A) DMDΔ52-Model cell line was treated with an antisense to skip exon 51 and the
effect was evaluated by RT-PCR and nested PCR analysis. Gel picture shows a pattern
corresponding with the correct skipping, which was confirmed by Sanger sequencing.
The same experiment evaluated by myoblot using n=20 wells per condition (B) showed
the restoration of dystrophin expression in the treated cultures. (**P values<0.05. P
values were determined with Mann-Whitney U test).
C) The DMD-UTRN-Model was used as a positive control in an experiment in which
unedited DMD cultures were treated with different ezutromid concentrations to up-
regulate utrophin expression. Myoblot analysis using n=8 wells per condition of the
treated cultures shows that ezutromid had no significant effect in DMD cultures while
utrophin expression is significantly increased in DMD-UTRN-Model compared to
unedited DMD cultures. (**P values<0.05. P values were determined with Mann-
Whitney U test).

Suplementary figure 1. Myoblasts transfection optimization and selection of positive cells by FACS.

A) GFP expression 48h after transfection in control myoblasts using four different transfection reagents and following manufacturers conditions. B) Control myoblasts expressing GFP 48h post transfection with ViaFect™ reagent and dot plot showing GFP positive cells (3,22%) selected using FACS.

Supplementary figure 2. PCR Analysis of Off-target Effects. Representative gel and sequence of PCR analysis performed for all targets.

A) Agarose gel showing the amplification of the six predicted off-targets regions for sgRNA2 and the six for sgRNA6 (the combination used for our DMD edition model) amplified in control myoblasts and DMDΔ52-Model. B) All the amplicons were sequenced and no differences between control myoblasts and DMDΔ52-Model were found.

Supplementary figure 3. MF20 differentiation marker depletion in edited cells.

MF20 expression determined by myoblot in DMDΔ52-Model compared to control myoblasts (A) and in DMD-UTRN-Model compared to DMD myoblasts (B). Myoblot analysis was performed using n=12 (A) or n=18 (B) wells for MF20 staining. (**P values<0.05; **** P values<0.0001 P values were determined with Mann-Whitney U test).

Supplementary table 1. Gene edition and antisense experiments primer sets.

Supplementary table 2. Off-target primer sets.

Off Target name	Guide sequence	Off target sequence	Mis	Chrom	Locus	PCR analysis	Sequencing
Ob1_2_Off1	GCTGAAGAACCCTGATACTAAGG (Ob1_sgRNA2	GCTGGAGAACCCTGATACTGTGG	2	chr1	intergenic:RP4-781L3.1-RP4-706G24.1	No off-target editing	Confirmed
Ob1_2_Off2		TCTGGAGAACCCTAATACTAAGG	3	chr8	intergenic:RP11-24P4.1-AC009695.1	No off-target editing	Confirmed
Ob1_2_Off3		ACTGAAGAATCCAGAAACTAGGG	4	chr7	intergenic:NOBOX-RP4-545C24.5	No off-target editing	Confirmed
Ob1_2_Off4		GCTAGAGAAACCTGAAACTAAGG	4	chr8	intergenic:RP11-536K17.1-EIF3H	No off-target editing	Confirmed
Ob1_2_Off5		TCTGGAGAACCCTAATACTGTGG	4	chr3	intron:TMEM45A	No off-target editing	Confirmed
Ob1_2_Off6		TCTGAAGAATCCTGATATTTTGG	4	chr2	intron:AC019100.3	No off-target editing	Confirmed
Ob1_6_Off1	GACCAACAGCCAAGGATATGAGG (Obj1 sgRNA6)	CACCATCAGCCAAGAATATGCGG	3	chr11	intergenic:RP11-430H10.3-RP11-958J22.1	No off-target editing	Confirmed
Ob1_6_Off2		TAACAACAGCCAAAGACATGAGG	4	chr14	exon:RP11-1012A1.4/RDH11	No off-target editing	Confirmed
Ob1_6_Off3		GTAAAAGAGCCAAGGATATGAGG	4	chr10	intron:RP11-556E13.1	No off-target editing	Confirmed
Ob1_6_Off4		TACTAGCAGCCAAGGATATCTGG	4	chr2	intergenic:AC007377.1-SLC8A1	No off-target editing	Confirmed
Ob1_6_Off5		GAGCGACAGCCAAGAATATTCGG	4	chr3	intron:CD96	No off-target editing	Confirmed
Ob1_6_Off6		AATCAACAGCCAAGAATGTGGGG	4	chr5	intergenic:CTD-2201E9.4-SEMA5A	No off-target editing	Confirmed
Ob2_22_Off1	GGTTCTCTTTAGCTGGGATCTGG (Obj2 sgRNA22)	TGTTCTCTCTAACTGGGATCTGG	3	chr18	intergenic:RP11-411B10.6-RP11-411B10.5	No off-target editing	Confirmed
Ob2_22_Off2		TGTTCTCTAGAGCTGGGATCTGG	3	chr21	intron:LCA5L	No off-target editing	Confirmed
Ob2_22_Off3		TGTTCTCTCCAACTGGGATCTGG	4	chr22	intron:PPP6R2	No off-target editing	Confirmed

Ob2_22_Off4		GAATCCTTTTAGCTGGGATCAGG	4	chr19	intron:ZNF536	No off-target editing	Confirmed
Ob2_22_Off5		GGTTCATCTTAGCTGGGATATGG	4	chr13	intron:FLT1	No off-target editing	Confirmed
Ob2_22_Off6		TGTTCTCTCTAACTGGGGTCTGG	4	chr21	intergenic:PPP6R2P1-AP001347.6	No off-target editing	Confirmed
Ob2_26_Off1		AAGCTTTCCTGGATATGACAAGG	4	chr4	intron:RNF150	No off-target editing	Confirmed
Ob2_26_Off2		GTGCTTACTTGGGTAAGACGTGG	3	chr17	intergenic:RP11-212E8.1-RP11-642M2.1	No off-target editing	Confirmed
Ob2_26_Off3	GTGCTTTCTTGGGTATGACATGG (Obj2 sgRNA26)	GAGTTAACTTGGGTATGACAGGG	4	chr4	intron:RGS12	No off-target editing	Confirmed
Ob2_26_Off4		GTGCTCTCATGAGAATGACAGGG	4	chr4	intergenic:GABRG1-RP11-320H14.1	No off-target editing	Confirmed
Ob2_26_Off5		GAGCTTTCCTGGGAATGACAGGG	3	chr1	intergenic:FOXO6-RNA5SP45	No off-target editing	Confirmed
Ob2_26_Off6		GTGCTTTATAGGATATAACATGG	4	chr6	intron:GSTA3	No off-target editing	Confirmed

691   **Table 1. Potential off-target sites.**

A)

sgRNAs targeting intron 51 *DMD* gene	sgRNAs targeting intron 52 *DMD* gene
sgRNA 1	sgRNA 6
sgRNA 2	sgRNA 7
sgRNA 3	sgRNA 8
sgRNA 4	sgRNA 9
sgRNA 5	sgRNA 10

B)

sgRNAs targeting 3'UTR region *UTRN* gene	sgRNAs targeting 3'UTR region *UTRN* gene
sgRNA 21	sgRNA 26
sgRNA 22	sgRNA 27
sgRNA 23	sgRNA 28
sgRNA 24	sgRNA 29
sgRNA 25	sgRNA 30

C)

Heks control
sgRNAs 2+6
sgRNAs 1+6
sgRNAs 5+6

800pb
300pb

D)

Heks control
sgRNAs 21+26
sgRNAs 21+27
sgRNAs 22+26

700pb
400pb

A) Lanes: C-, C+, Clone 1, Clone 2, Clone 3, Clone 4, Clone 5, Clone 6, **Clone 7**, Clone 8, Clone 9, H₂O. Markers: 700bp, 300bp. Non-edited clones, Edited clones. Clone 7= DMDΔ52- Model

B) sgRNA 2, complete sequence, sgRNA 6. ccaacgctgaagaaccctgatactaagggatattt / cagtctctctggaactgttggtcttgttggtccaagc — Non-edited control. EXON 52 deletion (476bp). Intron 51, Intron 52 — Edited clone: DMDΔ52-Model

C) Lanes: C-, C+, Clone 1, Clone 2, Clone 3, Clone 4, Clone 5, Clone 6, Clone 7, **Clone 8**, Clone 9, H₂O. Markers: 700bp, 400bp. Non-edited clones, Edited clones. Clone 8= DMD-UTRN -Model

D) sgRNA 22, complete sequence, sgRNA 26. ACTTTGGGTTCTCTTAGCTGGGATCTGGCCAAGAA / TATCAGGGCCATGTCATACCCAAGAAAGGCATCTATTT — Non-edited control. microRNAs binding site deletion (237 bp) — Edited clone: DMD-UTRN-Model

A)

500pb

200pb

91%

| EXON49 | EXON50 | EXON51 | EXON53 |

| EXON49 | EXON50 | EXON53 |

B)

Dystrophin expression by myoblot

Dystrophin antibody/Cell number signal

2.0

1.5

1.0

0.5

0.0

**

DMD.S2-Model non treated

DMD.S2-Model AO 100nM

C)

Utrophin expression by myoblot

Utrophin antibody/cell number signal

2.0

1.5

1.0

0.5

0.0

**

DMDdS8a DMSO

DMDdS8a exo1uM

DMDdS8a exo3uM

DMDdS8a exo5uM

DMDdS8a exo7uM

DMD-UTRN-Model