Duchenne Muscular Dystrophy Cell Culture Models Created By CRISPR/Cas 9 Gene Editing And Their Application To Drug Screening

P. Soblechero-Martín[1,2], E. Albiasu-Arteta[1], A. Anton-Martinez[1], L. de la Puente-Ovejero[1], I. Garcia-Jimenez[1], G. González-Iglesias[1], I. Larrañaga-Aiestaran[1], A. López-Martinez[1], J. Poyatos-García[3], E. Ruiz-Del-Yerro[1], F. Gonzalez[4], V. Arechavala-Gomeza[1,5]

[1]Neuromuscular Disorders, Biocruces Bizkaia Health Research Institute, Barakaldo, Spain
[2]Osakidetza Basque Health Service, Bilbao-Basurto Integrated Health Organisation, Basurto University Hospital, Clinical Laboratory Service, Bilbao, Spain
[3]La Fe Health Research Institute, Hospital La Fe, Valencia, Spain
[4]Pluripotent Stem Cells and Activation of Endogenous Tissue Programs for Organ Regeneration (PR Lab), Institute for Bioengineering of Catalonia (IBEC), Barcelona, Spain
[5]Ikerbasque, Basque Foundation for Science, Bilbao, Spain

Address for correspondence:

Prof. Virginia Arechavala-Gomeza

Biocruces Bizkaia Health Research Institute

Barakaldo, Bizkaia, 48903, Spain

Phone: +34 946007967

E-mail: virginia.arechavalagomeza@osakidetza.eus

ORCID: 0000-0001-7703-3255

Abstract

Gene edition methods are an attractive putative therapeutic option for Duchenne
muscular dystrophy and they have an immediate application in the generation of
research models. To generate two new edited myoblast cultures that could be
useful *in vitro* drug screening, we have optimised a CRISPR/Cas9 gene edition
protocol. We have successfully used it in wild type immortalised myoblasts to delete
exon 52 of the dystrophin gene: DMDΔ52-Model, modelling a common Duchenne
muscular dystrophy mutation; and in patient's immortalised cultures we have
deleted an inhibitory microRNA target region of the utrophin UTR, leading to
utrophin upregulation. We have characterised these cultures and, to show their use
in the assessment of DMD treatments, we have performed exon skipping in the
DMDΔ52-Model and have used the unedited cultures/ DMD-UTRN-Model combo to
assess utrophin overexpression after drug treatment. While the practical use of
DMDΔ52-Model is limited to the validation to our gene edition protocol, DMD-
UTRN-Model offers a possible therapeutic gene edition target as well as a useful
positive control in the screening of utrophin overexpression drugs.

KEYWORDS: Duchenne muscular dystrophy, CRISPR/Cas9 gene editing, inhibitory
microRNA target region, dystrophin, utrophin.

INTRODUCTION

Duchenne muscular dystrophy (DMD) is a fatal X-linked recessive disease affecting

one out of 5.000 newborn males. It is commonly caused by deletions disrupting the

open reading frame of the *DMD* gene causing a lack of dystrophin protein[1]. Patients

carrying out of frame mutations present a severe phenotype, while those carrying in-

frame mutations may result in hypomorphic alleles, a partially functional dystrophin

and milder phenotypes, such as in Becker muscular dystrophy (BMD)[2]. Dystrophin

plays a major role in membrane stabilization during muscle contraction, linking the

actin cytoskeleton to the sarcolemma[3] and also contributes to extracellular signalling[4].

Lack of dystrophin in DMD patients' muscles leads to progressive muscle wasting and

degeneration. DMD children suffer from loss of ambulation in the first or second

decade of life and premature death by cardiac and respiratory complications[5].

Although no definitive cure for DMD is available, four drugs have been recently

approved by different regulatory agencies: ataluren, approved by the European

Medicines Agency (EMA), induces readthrough of premature stop codons during

mRNA translation, generating a full length dystrophin protein[6]; and the US Food and

Drug Administration (FDA), has recently approved three antisense oligonucleotide

drugs, which modulate splicing by exon skipping, restoring *DMD* reading frame and

leading to a shorter but functional protein[7,8]: eteplirsen, golodirsen and viltolarsen.

Exon skipping therapies aim to attenuate the phenotype and phenocopy milder BMD-

like genotypes, to potentially improve disease outcomes. Eteplirsen targets *DMD* exon

51 and golodirsen and viltolarsen, exon 53. All approved drugs are mutation specific

and designed to rescue specific patient mutations only present, respectively, in 13%

(ataluren), 13% (eteplirsen) and 8% (golodirsen and viltolarsen) of patients[9]. It is

therefore important to assess exon-skipping strategies targeting other *DMD* exons, and

many such drugs are in different phases of clinical assays[10].

Other than these approved therapies, gene transfer has emerged as a promising

approach applicable to all DMD patients. SRP-9001, a human micro-dystrophin gene

transfer therapy, driven by the muscle-specific promoter MHCK7 and delivered by the

recombinant adeno-associated virus rAAVrh74 has recently got the Fast Track

designation by the FDA after showing a robust posttreatment increase of micro-
dystrophin expression in patients muscles and positive safety and tolerability data in
clinical trials[11] suggesting the potential of this therapy to provide clinically meaningful
functional improvement in DMD patients.

Alternatively to these therapies aiming to restore dystrophin expression, many
compounds targeting secondary effects of dystrophin deficiency or looking for
alternatives to dystrophin are also under evaluation.

In vitro cellular models are particularly useful to assess the efficiency of novel
therapies for DMD. However, only a few human immortalized muscle cell lines derived
from DMD patients are currently available[12]. Due to the wide spectrum of DMD
mutations and the difficulties to obtain DMD patient muscle biopsies, DMD-myoblasts
models would provide a powerful resource for *in vitro* drug screening and study
disease rescue mechanisms.

UTROPHIN *(UTRN)* is an autosomal paralog of dystrophin, expressed in skeletal muscle
cells during embryonic development, but restricted to neuromuscular and
myotendinous junctions in the mature muscle fibre[13]. Overexpression of utrophin in
skeletal muscle in DMD animal models can partially compensate the lack of dystrophin
and improve DMD phenotype[14-16]. Importantly, ectopic and high levels of utrophin in
myoblasts are not associated with toxicity, making utrophin upregulation an
interesting therapeutic strategy applicable to all patients, regardless of their specific
mutation[17,18]. Ezutromid/SMT-C1100 was the first utrophin modulator evaluated in
clinical assays, but was recently abandoned due to lack of evidence of utrophin
restoration, nor clinical improvement demonstrated in patients [19,20]. Alternatively,
other studies proposed new strategies to upregulate *UTRN* by blocking the inhibitory
target region of microRNAs repressing *UTRN* expression[21]. Recently, utrophin
upregulation has been efficiently achieved using gene therapy in multiple animal
models with non-immunogenic side effects[22]. Several new compounds that aim to
overexpress utrophin are currently being developed[23,24], and this preclinical
development could benefit from a gold standard or an adequate positive control to
use in these assays.

CRISPR/Cas9 currently represents a very efficient and versatile genome-engineering tool, introducing small and large DNA modifications, including large genomic deletions in different cell types and organisms[25]. In the presence of two single guide (sg) RNAs targeting two different loci on the same chromosome, Cas9 can induce two DNA double strand breaks (DDSBs), leading in some cases to deletion of the excised DNA segment through repair by the non-homologous end joining (NHEJ) pathway[26]. Similar to antisense oligo-mediated exon skipping therapies at RNA level, CRISPR/Cas9 can therefore be used to remove mutations by deleting mutated exons and restore the open reading frame of the *DMD* gene[27-29]. The advantage of this approach is that the genetic modification, once introduced, is stable over cell cycles. CRISPR/Cas9 has been successfully employed to correct mutations and/or restore the open reading frame recovering dystrophin expression both *in vitro* and *in vivo*[30,31]. However, some hurdles have been reported, such as difficulties transfecting myoblast or a recent study in the golden retriever muscular dystrophy dog (GRMD), where no dystrophin restoration at protein level was evident after gene editing using this technology[32]. More importantly, as well as these preclinical problems others such as possible off-target problems or immunogenicity linked with the use of Cas9[33] may delay their clinical application. However, while gene editing as therapeutic option still needs further development, CRISPR/Cas9 methodology has been applied to provide a large number of new animal models to further understand DMD pathology and perform preclinical studies[34]. In our quest to optimise the preclinical development of new therapies for DMD, we have developed a genome editing strategy applicable in control and patient myoblasts. Here, we report two new cell cultured models that can be successfully used for preclinical assessment of new DMD therapies: a culture that replicates a patient's deletion **(DMDΔ52-Model)** and another that overexpresses utrophin **(DMD-UTRN-Model)**.

RESULTS

Generation of cell culture models by gene editing

We completed two different gene editing projects: **objective 1** aimed to delete exon 52 of the *DMD* gene (a common mutation in DMD patients) in control immortalised

human myoblasts to generate a disease model **(DMDΔ52-Model)**; **objective 2 (DMD-**
UTRN-Model) was to delete in the *UTRN* gene of DMD immortalized human myoblasts
an inhibitory microRNA target region to generate a utrophin ectopic expression rescue
model.

Each edition required to design two sgRNAs flanking the region to be deleted to
generate two DDSBs leading to the removal of that region (Figure 1A and D). All the 25
different combinations of the sgRNAs designed, cutting before (x5) and after (x5) the
target region, were tested in HEK293 cultures first (Supplementary figure 1A and B).
The most efficient combination of sgRNAs in HEK293 cells for each objective was
selected to be used in the transfection of human immortalized myoblasts
(Supplementary figure 1C and D).
To accomplish objective 1, two GFP-plasmids, each encoding Cas9 nuclease and either
sgRNA 2 or sgRNA 6, were transfected into human immortalized control myoblasts. For
objective 2, the selected GFP-plasmids, encoded Cas9 nuclease and either sgRNA 22 or
sgRNA 26 and were transfected into human immortalized DMD myoblasts. After FACS
sorting of individual GFP-positive cells, clones were expanded for DNA extraction
(Supplementary figure 2). Clones were analysed to confirm the presence of the desired
deletions by genomic PCR performed with specific primers for each targeted gene
(Figure 1B and 1E), amplicons corresponding in size with the expected deletions were
analysed by Sanger sequencing, and the expected deletions were confirmed in all the
positive clones (Figure 1C and 1F).
For **objective 1** two positive clones were obtained, clone number 2 and 7 but only
clone 7 was used for further analysis and called **DMDΔ52-Model** (Figure 1B and 1C).
For **objective 2,** two clones were edited in only one allele, corresponding to numbers 3
and 4 and other two were completely edited, numbers 2 and 8. In this case, clone
number 8 was selected to be used for further analysis and was called **DMD-UTRN-**
Model (Figure 1E and 1F).
To evaluate any potential off-target effects, each selected sgRNA was analysed *in silico*
using the bioinformatics web-tool CRISPOR[35]. We selected the six more likely off-target
sites for each sgRNA and analysed each one of them through PCR, followed by Sanger

sequencing in edited clones (Table 2 and 3). We found no off-target effects in any of
the 12 sites studied for each clone sites (Supplementary figure 3).

Analysis of dystrophin and utrophin expression in edited clones.

We compared dystrophin expression in myotubes of the DMDΔ52-Model to controls
and DMD cultures and confirmed that it was abolished by immunocytochemistry
(Figure 2A), western blot analysis (Figure 2B), myoblots (Figure 2C) and droplet digital
PCR (ddPCR) (Figure 2D). Dystrophin levels in this model, where exon 52 had been
removed by CRISPR/Cas9 editing, were statistically no different than those seen in a
culture from a DMD patient.

Immunocytochemistry showed the increase of utrophin expression between unedited
DMD and DMD-UTRN-Model myotubes (Figure 3A) and this increase was corroborated
by western blot (a 175% increase, Figure 3B), myoblot analysis (close to 50% increase,
Figure 3C) and ddPCR (Figure 3D).

We also quantified dystrophin and utrophin expression in the edited cultures by
myoblot during their differentiation process and compared these with control and
DMD myotubes (Supplementary figure 4, bar chart).

Analysis of differentiation markers expression in edited clones.

As we suspected that the editing and cloning process could have affected the
differentiation of the edited models, the fusion index (%) of edited and non-edited
myotubes after MF20 and Hoechst inmunocytochemistry was calculated (Figure 2E and
3E) and the expression of different myogenic regulatory factors like the myogenic
factor 5 (Myf5) and the myosin heavy chain isoform 3 (MyH3) were analysed at
different time points during myotube formation in DMDΔ52-Model and DMD-UTRN-
Model as well as in their corresponding unedited cultures by ddPCR (Figure 2F and 3F).
Fusion index was clearly lower in DMDΔ52-Model compared to control myoblasts
(Figure 2E) while no significant differences were found between DMD-UTRN-Model
and DMD cultures (Figure 3E). As expected, Myf5 expression decreased while MyH3
increased during the differentiation process in both models and followed the same
pattern in their origin cells, however we see that MyH3 marker is significantly lower in
the edited models at day 5 and day 7 after starting fusion (Figure 2F and 3F).

The MF20 differentiation marker was also analysed by myoblot in edited cultures and
we could observe a decrease in both edited clones, no matter the deletion, compared
with their corresponding controls (Supplementary figure 4, red lines).

Evaluation of therapies in newly generated model cell lines.
To assess if the DMDΔ52-Model cell culture could be useful to test potential mutation
specific therapies for DMD, we evaluated the exon skipping efficiency of an antisense
oligonucleotide in this culture. We treated the DMDΔ52-Model cultures with an
antisense oligonucleotide drug that can skip exon 51[36] and restore *DMD* open reading
frame. After treatment with this drug, we confirmed that exon skipping had taken
place at RNA level (Figure 4A), and a limited restoration of dystrophin expression by
myoblot analysis (Figure 4B).

To test our DMD-UTRN-Model as a positive utrophin overexpression control, we
cultured it alongside the original unedited DMD cultures, which we treated with
several concentrations of ezutromid and we evaluated the expression of utrophin in all
cultures. We observed that utrophin was hardly modified in DMD cultures treated with
ezutromid while a robust overexpression was confirmed in the DMD-UTRN-Model
compared to the unedited DMD cultures (Figure 4C). We also confirmed that this
overexpression was stable during a time course experiment (Supplementary figure 4D).

DISCUSSION

CRISPR/Cas9 has been explored in the past as a possible therapeutic approach for
DMD in *in vitro* models of the disease. In most of these cases, a "permanent" exon
skipping approach was selected where a shorter protein would be produced[27,28], while
in some others full length dystrophin was the result of the edition[37-39]. Several studies
have shown efficacy in mice models[40-42], an most recently in dogs, which is currently
the most advanced example of its application to DMD[29] but there are also examples of
the challenges associated to gene editing[32]. Before this can be a credible therapeutic
option, several delivery and manufacturing problems will need to be overcome. In the
meantime, this methodology is very useful for researchers looking for cell culture
models: muscle biopsies are not routinely collected during diagnosis of this disorder
and seldom cultured. This means there are few good culture models of the disease.

Another tool created to facilitate research was the immortalisation of some of the
available cultures[12], which increases the possibility of performing more experiments
with a given culture. We have used immortalised cultures to further increase this
possibility.

Although we are concerned about the efficiency limitations of our gene editing
protocol, specially due to transfection difficulties in myoblasts reported also by other
laboratories[27,38]; we have successfully applied it to edit 2 different regions in two
different cell backgrounds (Control and DMD), and we consider that those described
and fully characterised in this manuscript could be relevant research models that we
would be happy to share. The first of these models is an immortalised DMD disease
cell culture model, (DMDΔ52-Model) that lacks exon 52 of the *DMD* gene, which
disrupts the ORF and dystrophin expression. This model could be useful to evaluate
mutation-independent drug treatments, and also exon skipping drugs that aim to skip
exons 51 or 53[43,44] as skipping either exon in this case would restore the ORF and
dystrophin expression. We have demonstrated that DMDΔ52-Model lacks dystrophin
expression and that this can be reverted through treatment with an exon 51 skipping
drug. However, the main interest of this model was demonstrating the efficacy of the
technique, as immortalised cultures from patients with the same deletion are currently
available. We have replicated this protocol to generate other DMD-like cultures
(details will be included in a manuscript being currently drafted).

An immortalised cell culture model constitutively expressing utrophin, DMD-UTRN-
Model, is both a proof of principle of a possible therapeutic option to overexpress
utrophin as a substitute for dystrophin, and a valuable research tool. The search of
drugs that could overexpress utrophin is ongoing[45] after the recent failure of
ezutromid to show relevant results in clinical trials. Many research projects, including
drug re-purposing screening, are ongoing to find new candidates to test in clinical
trials[46]. However, there are no reliable positive controls that could be used to compare
such treatments. We propose that our cell model could serve that purpose, offering
researchers useful custom controls for their studies. We have tested this hypothesis
and compared the stable utrophin overexpression quantified in the DMD-UTRN-Model
with the utrophin that is expressed after treatment of the unedited DMD patient

cultures with ezutromid, the lead market candidate in this field until very recently,

with positive results. Previous studies in muscle sections show that DMD patients

already overexpress utrophin, in many cases 4 to 5-fold the levels seen in control

muscle sections[47]. Our choice to target this particular *UTR* region, increases basal

overexpression in DMD cultures, and the amount of overexpression varies significantly

when evaluated by western blot (more than 2.5 times) or our preferred method,

myoblots (close to 1.5 times). We like to consider that myoblot evaluation reflects

more closely the actual protein expression, as it is not subjected to many of the

inherent problems of western blotting when evaluating very large proteins[48]. This is

why we cannot comment yet on the differences in expression between our study and

other published studies that also aimed to overexpress utrophin by gene editing, but

which 1) targeted different promoter regions (*UTRN* A or *UTRN* B) of utrophin and 2)

evaluated their results by western blot analysis[38]. We would be interested on studying

this matter further to analyse the differences in utrophin expression when targeting

different regions.

As a conclusion, we have created two new cell culture models that we have already

used as efficient tools in our search for new therapies for DMD, after optimization of a

gene editing method to be applied to myoblasts, a rather difficult target. We expect

our experience to be useful to other muscle researchers and we are looking forward

expanding those experiments.

METHODS

CRISPR/Cas9 tools

Specific sgRNA guides were designed using the online bioinformatics tool http://crispr.mit.edu[49]. Ten different guides (five before and five after the target region) were designed targeting exon 52 flanking regions in *DMD* gene and another ten targeting a repressor binding site in the UTR 3' region of *UTRN* gene and selected according to their score number (Table 1). They were cloned into a plasmid containing *Cas9* from *S. pyogenes* with 2A-*EGFP* pSpCas9 (BB)-2A-GFP (PX458) (Addgene plasmid # 48138, deposited by Feng Zhang). All sgRNAs were cloned using BbsI sites.

Cell cultures

Human embryonic kidney 293 (HEK 293) cells, used in the preliminary selection of the best sgRNA combinations for our experiments, were purchased from the European Collection of Authenticated Cell cultures (ECCAC) via Sigma-Aldrich, Spain, and maintained following the manufacturer's protocols.

Immortalized myoblasts derived from muscle biopsies from healthy controls and DMD patients were provided by the CNMD Biobank, London, UK and the Institut de Myologie Paris, France, and cultured using skeletal muscle medium (SMM) (Promocell, Germany) and differentiation media as previously described[48].

Cell culture transfection

All different sgRNAs combinations were transfected into HEK 293 cells using lipofectamine 2000® (Thermo Fisher Scientific), according to the manufacturer's protocol. Myoblasts seeded in 6 well plates at 70-80% confluence were transfected with 1.5ug of each plasmid with the most efficient guide RNA combination using ViaFect™ (Promega) transfection reagent (1:5 ratio).

FACS workflow of GFP positive myoblasts

48 hours after transfection, myoblasts were trypsinised and collected for fluorescence activated cell sorting (FACS) at the Cell Analytics Facility (BD FACS Jazz) Achucarro Basque Center for Neuroscience - (Leioa, Spain). GFP-positive cells were seeded

individually in 96 well plates for clonal selection. The first colonies were visible around
7 days post-sorting. Clones were expanded from single cell to near-confluence and
expanded into larger well plates to be harvested 15-30 days post-sorting. Myoblasts
often developed elongated and stressed shapes during this clonal expansion after
single cell sorting. Harvested cultures were aliquoted: some aliquots were frozen for
archival; others were pelleted for DNA analysis, while replicates were cultured further
for characterization by immunocytochemistry, western blot and myoblots (see
schematic workflow in Supplementary figure 2).

Analysis of gene editing
DNA was extracted from cell pellets using a QIAamp® DNA Mini Kit, Qiagen. PCR
amplification targeting the edited regions was carried out using Taq DNA Polymerase
(Recombinant), Invitrogen, under the following conditions: preheating 3' 94ºC, 25
cycles of 94º for 3', 94º 20'', 63º 20'', 72º 1' and 72º 5' and DMD-Seq-D52-DOWN-F2
and DMD-Seq-D52-DOWN-R2 primers (see Suplementary Table 1). PCR products were
resolved in 2% TAE-agarose gels and purified with QIAquick® Gel extraction Kit,
Qiagen. PCR amplicons corresponding to the expected length were analysed by Sanger
sequencing at the sequencing platform of Biocruces Bizkaia Health Research Institute
using   DMD-Seq-D52-DOWN-F2   and   DMD-Seq-D52-DOWN-R2   primers   (see
Supplementary Table 1).

Off-target analysis of mutations in clonal lines
Potential off-target region loci of each sgRNA used were predicted using CRISPOR
bioinformatics tool http://crispor.tefor.net/. The six most probable off-target
sequences per guide (Table 2 and 3) were analysed in the edited clones using genomic
PCR and Sanger sequencing. Primer sets flanking off-target sites and the corresponding
internal primers used for Sanger sequencing are listed in Supplementary Table 2.

Primary Antibodies

Anti-dystrophin: Dys1 (Leica Biosystems), Mandys1, Mandys106 (The MDA Monoclonal

Antibody Resource) and anti-dystrophin antibody (Abcam 15277).

Anti-utrophin: Mancho7 (The MDA Monoclonal Antibody Resource).

Anti-myosin heavy chain (MF20: Developmental Studies Hybridoma Bank).

Anti a-actinin (Sigma-Aldrich A7732).

Immunostaining assays

Original and edited clones for objectives 1 (DMDΔ52-Model) and 2 (DMD-UTRN-

Model) were cultured and immunostained for dystrophin or utrophin expression.

Edited clones were seeded into chamber slides and treated with a MyoD virus,

(Applied Biological Materials Inc, Canada) to facilitate differentiation into myotubes[50].

After seven days differentiating, samples were fixed with 4% PFA. Cultures were

permeabilised with Triton 0.5% and then blocked for half an hour with BSA 2%.

Afterwards, immunostaining was performed overnight at 4ºC with the required

antibodies. Primary antibodies used for dystrophin staining were a mix of Dys1,

Mandys1 and Mandys106 at 1:100 dilution and for utrophin staining was Mancho 7

diluted at 1:50. The following day, after being washed with PBS Tween 0.1%, cells were

stained with Alexa Fluor 488 goat anti-mouse (Invitrogen) for 1 hour at room

temperature. Hoechst 1/2000 was used for nuclei staining and chamber slides were

mounted with PermaFluor™ Aqueous Mounting Medium (Thermoscientific). Images

were captured with a LEICA DMI 6000B microscope at the Microscopy Platform at

Biocruces Bizkaia Health Research Institute.

In-cell western assay (myoblots)

Myoblots were performed as described before[48]. In short, clones were seeded in 96-

well plates and incubated for 24 h in SMMC, after which they were treated with MyoD

virus and incubated in differentiation media for 7 days. Then, plates were fixed with

ice-cold methanol, permeabilised and blocked before incubation with the required

primary antibodies overnight: anti-dystrophin mix (Dys1, Mandys1 and Mandys106 at

1:100), anti-utrophin (Mancho 7 antibody at 1:50), and anti-myosin heavy chain

antibody (MF20 at 1:100) that was used to evaluate differentiation. Next day, plates

were incubated with the secondary antibodies. Biotin-mediated amplification (Abcam

6788 goat antimouse IgG biotin 1:2000) was used to increase dystrophin signal.

Secondary antibodies, IRDye 800cw streptavidin 1:2000 and IRDye 800CW goat anti-

mouse 1:500, were prepared together with CellTag 700 Stain (LI-COR® Biosciences) at

1:1000 dilution and incubated for 1 hour at RT and protected from light. After

incubation, plates were analysed using the Odyssey® CLx Imager (LI-COR® Biosciences).

Treatment with antisense exon skipping drugs

Cultures in 96 wells and P6 wells were treated with a 2'MOE-phosphorotioate

antisense oligonucleotide (AO) aiming to skip *DMD* exon 51 (5☐-[

T*C*A*A*G*G*A*A*G*A*T*G*G*C*A*T*T*T*C*T]-3☐, Eurogentec, Belgium) by

transfection with Lipofectamine as described in[43,44] and analysed by either myoblot (96

well plates) or RT-PCR (pellets extracted from 6 well plates).

RT-PCR

RNA was extracted from cell pellets (RNeasy mini kit, Qiagen) according to the

manufacturer's protocol. Reverse transcription of the samples was performed using

(SuperScrip[TM] IV Reverse Transcriptase, Invitrogen) according to the manufacturer's

protocol. cDNA samples were either used for digital droplet PCR analysis or amplified

by nested PCR using specific primers sets (Supplementary Table 1) and Taq DNA

Polymerase (Recombinant), Invitrogen, as described in[44] for exon skipping analysis.

Amplified samples were resolved in TAE-agarose and PCR amplicons of interest were

first analysed with Gel Doc TM EZ Imager, BIORAD and then purified with (QIAquick®

Gel extraction Kit, QIAGEN) for sequencing analysis. Before DNA extractions bands

were semi-quantified using Image J.

Treatment with utrophin overexpression drugs

Ezutromid was diluted first in DMSO and finally in differentiation medium to different
concentrations and added to myoblasts in 96 well plates 7 days after differentiation.
Twenty-four hours after treatment, medium was removed, and plates were fixed with
ice-cold methanol for myoblot analysis.

Western blot

Cell cultures were seeded into P6 plates and trypsinized after 7 days of differentiation.
Then, cell pellets were solubilized in lysis/loading buffer and denatured at 95°C for 5
min. Samples were loaded onto a NuPAGE® Novex® 3–8% Tris-Acetate Gel3–8%
(Thermo Fisher Scientific) and run in Novex Tris-Acetate SDS Running Buffer (Thermo
Fisher Scientific) for 60 min at 70 V + 120 min at 150 V at 4ºC. Protein wet transfer was
performed overnight at 4°C using an Immobilon®-FL PVDF membrane (Merck™). Next
day, membranes were stained with Revert ™ 700 Total Protein Stain (Li-Cor) for total
protein measurement, blocked with Intercept (PBS) Blocking Buffer (Li-Cor) for 2 hours
and incubated overnight at 4°C with the primary antibodies (1/200 anti-dystrophin
antibody Abcam15277, 1/50 anti-utrophin antibody Mancho 7 or 1/500 anti a-actinin
antibody, Sigma-Aldrich A7732). After washing steps with PBS-Tween 0.1%,
membranes were incubated with secondary antibodies for 1 hour (1/5000 IRDye
800CW goat anti-rabbit 926–32211 or IRDye 680RD goat anti-mouse 926-68070, Li-
Cor) at room temperature, washed again and scanned using an Odyssey Clx imaging
system. Bands quantification was performed using Image Studio ™ software.

Droplet digital PCR (ddPCR)

Gene expression was detected and quantified using a QX200™ Droplet Digital™ PCR
system (Bio-Rad). The reaction was performed using 2 µl of cDNA in a 20 µl reaction
volume containing: 1 µl of Utrophin Taqman probe (ID: Hs01125975, FAM labelled),
1µl of MYF5 ddPCR™ GEX Assay probe (ID: dHsaCPE5026295, HEX labelled) or MYH2
ddPCR™ GEX Assay probe (ID: dHsaCPE5050991, HEX labelled), 10 µl of ddPCR™
Supermix for Probes (no dUTP) (Bio-Rad) and 6 µl of DNase/RNase-free H2O.
To generate the droplets, 20 µL of the previous ddPCR reaction and 70 µL of Droplet
Generation Oil for Probes (Bio-Rad) were added to the 8 channel droplet generation
cartridge according to manufacturer instructions and this cartridge was placed in the

QX200 droplet generator (Bio-Rad). Then, 40 μL of the resulting droplet emulsion was

transferred to a semi-skirted 96 well PCR plate (Eppendorf), sealed with foil and

amplified on a thermal cycler using the following amplification conditions: enzyme

activation 10', 40 cycles of 94°C for 30'' and 55°C for 1', and heat deactivation 10' 98°C.

Plates containing the amplified droplets were loaded into the QX200 droplet reader

and results were analysed using QuantaSoft software^{TM} (Bio-Rad).

Statistical analysis

Mann-Whitney U test was used throughout this study to calculate P values for

determination of statistical significance (*p-value<0.05, ** p-value<0.01, ****p-

value<0.0001). Statistical analysis was performed using GraphPad Prism software.

ACKNOWLEDGEMENTS

We acknowledge the use of cell cultures provided by the Queen Square Centre for

Neuromuscular Disorders BioBank (CNMD Biobank, London, UK) and the Institut de

Myologie (Paris, France, immortalised cultures).

We gratefully acknowledge the use of the antibodies provided by Professor Glenn

Morris from the Muscular Dystrophy Association (MDA) Monoclonal Antibody

Resource, which distributes free antibodies for research in neuromuscular diseases

worldwide from Oswestry, United Kingdom.

The MF20 antibody developed by D.A. Fischman, Weill Cornell Medical College, was

obtained from the Developmental Studies Hybridoma Bank, created by the NICHD of

the NIH and maintained at The University of Iowa, Department of Biology, Iowa City, IA

52242.

Cell sorting experiments were performed at the Cell Analytics Facility at Achucarro-

Basque Centre of Neuroscience (Leioa, Spain).

We would like to thank Ms. Karmele Alapont-Celaya for her assistance in the

completion of some laboratory tasks during this project. Drs. Gustavo Pérez-Nanclares

and Ana Belén de la Hoz at the Genetics-Genomics Facility, Biocruces Bizkaia Health

Research Institute, are also gratefully acknowledged for their help with sequencing
reactions. Special thanks are due to Drs. Maria Dolores-García Vázquez at the Cell
Culture Facility and Dr Javier Díez-García at the Microscopy Facility, both at Biocruces
Bizkaia Health Research Institute, for their assistance at their corresponding platforms.

COMPETING INTERESTS

The authors declare no competing interests

FUNDING

This work was supported by funding from Health Institute Carlos III (ISCIII, Spain) and
the European Regional Development Fund, (ERDF/FEDER), 'A way of making Europe':
Grant PI15/00333; Basque Government (grants 2016111029, 2018222035 and
2020333012) and Duchenne Parent Project Spain (grant 05/2016). P. S-M holds a Rio
Hortega Fellowship from ISCIII (CM19/00104). V.A-G holds a Miguel Servet Fellowship
from the ISCIII (CPII17/00004), part-funded by ERDF/FEDER. J P-G acknowledges
funding from Fundación Isabel Gemio. A. L-M acknowledges funding by Biocruces
Bizkaia Health Research Institute (BC/I/DIV/19/001). F. G is supported by a Ramon y
Cajal Grant in Biomedicine (RYC-2014-16751) from the Ministry of Economy and
Competitivity (MINECO), Spain. V.A-G also acknowledges funding from Ikerbasque
(Basque Foundation for Science).

DATA AVAILABILITY

Under request

AUTHOR CONTRIBUTIONS STATEMENT

P S-M, formal analysis, investigation, data curation, writing or original draft, review and
editing, visualization, supervision. E. A-A, investigation, data curation, writing -review
and editing. A. A-M, investigation, writing - review and editing. L. d-l-P-O,
investigation, writing - review and editing. I. G-J, methodology, investigation, review
and editing, supervision. G. G-I, investigation, writing - review and editing. I. L-A,
investigation, writing - review and editing. A. L-M, investigation, writing - review and
editing. J. P-G, investigation, writing - review and editing. E. R-D-Y, methodology,
investigation, writing-review and editing, supervision. F. G conceptualization, formal
analysis, writing-review and editing. V. A-G: conceptualization, methodology, formal
analysis, investigation, resources, data curation, writing or original draft, review and
editing, visualization, supervision, project administration and funding acquisition.

REFERENCES

1   Hoffman, E. P., Brown, R. H., Jr. & Kunkel, L. M. Dystrophin: the protein product of the
494   Duchenne muscular dystrophy locus. *Cell* **51**, 919-928, doi:10.1016/0092-
495   8674(87)90579-4 (1987).
496   2   Anthony, K. *et al.* Dystrophin quantification and clinical correlations in Becker muscular
497   dystrophy: implications   for   clinical   trials. *Brain* **134**, 3547-3559,
498   doi:10.1093/brain/awr291 (2011).
499   3   Muntoni, F., Torelli, S. & Ferlini, A. Dystrophin and mutations: one gene, several
500   proteins, multiple phenotypes. *Lancet Neurol* **2**, 731-740, doi:10.1016/s1474-
501   4422(03)00585-4 (2003).
502   4   Lai, Y. *et al.* Dystrophins carrying spectrin-like repeats 16 and 17 anchor nNOS to the
503   sarcolemma and enhance exercise performance in a mouse model of muscular
504   dystrophy. *J Clin Invest* **119**, 624-635, doi:10.1172/JCI36612 (2009).
505   5   Eagle, M. *et al.* Survival in Duchenne muscular dystrophy: improvements in life
506   expectancy since 1967 and the impact of home nocturnal ventilation. *Neuromuscul*
507   *Disord* **12**, 926-929, doi:10.1016/s0960-8966(02)00140-2 (2002).
508   6   Finkel, R. S. Read-through strategies for suppression of nonsense mutations in
509   Duchenne/ Becker muscular dystrophy: aminoglycosides and ataluren (PTC124). *J Child*
510   *Neurol* **25**, 1158-1164, doi:10.1177/0883073810371129 (2010).
511   7   Kinali, M. *et al.* Local restoration of dystrophin expression with the morpholino
512   oligomer AVI-4658 in Duchenne muscular dystrophy: a single-blind, placebo-
513   controlled, dose-escalation, proof-of-concept study. *Lancet Neurol* **8**, 918-928,
514   doi:10.1016/S1474-4422(09)70211-X (2009).
515   8   Muntoni, F. *et al.* Golodirsen Induces Exon Skipping Leading to Sarcolemmal
516   Dystrophin Expression in Duchenne Muscular Dystrophy Patients With Mutations
517   Amenable to Exon 53 Skipping (S22.001). *Neurology* **90** (2018).
518   9   Aartsma-Rus, A., Van Deutekom, J. C., Fokkema, I. F., Van Ommen, G. J. & Den Dunnen,
519   J. T. Entries in the Leiden Duchenne muscular dystrophy mutation database: an
520   overview of mutation types and paradoxical cases that confirm the reading-frame rule.
521   *Muscle Nerve* **34**, 135-144, doi:10.1002/mus.20586 (2006).
522   10   Arechavala-Gomeza, V., Anthony, K., Morgan, J. & Muntoni, F. Antisense
523   oligonucleotide-mediated exon skipping for Duchenne muscular dystrophy: progress
524   and challenges. *Curr Gene Ther* **12**, 152-160, doi:10.2174/156652312800840621
525   (2012).
526   11   Mendell, J. R. *et al.* Assessment of Systemic Delivery of rAAVrh74.MHCK7.micro-
527   dystrophin in Children With Duchenne Muscular Dystrophy: A Nonrandomized
528   Controlled Trial. *JAMA neurology*, doi:10.1001/jamaneurol.2020.1484 (2020).
529   12   Mamchaoui, K. *et al.* Immortalized pathological human myoblasts: towards a universal
530   tool for the study of neuromuscular disorders. *Skelet Muscle* **1**, 34, doi:10.1186/2044-
531   5040-1-34 (2011).
532   13   Tinsley, J. M. *et al.* Primary structure of dystrophin-related protein. *Nature* **360**, 591-
533   593 (1992).
534   14   Tinsley, J. *et al.* Expression of full-length utrophin prevents muscular dystrophy in mdx
535   mice. *Nat Med* **4**, 1441-1444 (1998).
536   15   Cerletti, M. *et al.* Dystrophic phenotype of canine X-linked muscular dystrophy is
537   mitigated by adenovirus-mediated utrophin gene transfer. *Gene Ther* **10**, 750-757,
538   doi:10.1038/sj.gt.3301941 (2003).
539   16   Tinsley, J. M. *et al.* Amelioration of the dystrophic phenotype of mdx mice using a
540   truncated utrophin transgene. *Nature* **384**, 349-353, doi:10.1038/384349a0 (1996).
541   17   Fisher, R. *et al.* Non-toxic ubiquitous over-expression of utrophin in the mdx mouse.
542   *Neuromuscul Disord* **11**, 713-721 (2001).

18 Fairclough, R. J., Wood, M. J. & Davies, K. E. Therapy for Duchenne muscular
dystrophy: renewed optimism from genetic approaches. *Nat Rev Genet* **14**, 373-378,
doi:10.1038/nrg3460 (2013).
19 Tinsley, J., Robinson, N. & Davies, K. E. Safety, tolerability, and pharmacokinetics of
SMT C1100, a 2-arylbenzoxazole utrophin modulator, following single- and multiple-
dose administration to healthy male adult volunteers. *J Clin Pharmacol* **55**, 698-707,
doi:10.1002/jcph.468 (2015).
20 Ricotti, V. *et al.* Safety, Tolerability, and Pharmacokinetics of SMT C1100, a 2-
Arylbenzoxazole Utrophin Modulator, following Single- and Multiple-Dose
Administration to Pediatric Patients with Duchenne Muscular Dystrophy. *PLoS One* **11**,
e0152840, doi:10.1371/journal.pone.0152840 (2016).
21 Mishra, M. K., Loro, E., Sengupta, K., Wilton, S. D. & Khurana, T. S. Functional
improvement of dystrophic muscle by repression of utrophin: let-7c interaction. *PLoS*
One **12**, e0182676, doi:10.1371/journal.pone.0182676 (2017).
22 Song, Y. *et al.* Non-immunogenic utrophin gene therapy for the treatment of muscular
dystrophy animal models. *Nat Med* **25**, 1505-1511, doi:10.1038/s41591-019-0594-0
(2019).
23 Ito, M., Ehara, Y., Li, J., Inada, K. & Ohno, K. Protein-Anchoring Therapy of Biglycan for
Mdx Mouse Model of Duchenne Muscular Dystrophy. *Hum Gene Ther* **28**, 428-436,
doi:10.1089/hum.2015.088 (2017).
24 Pisani, C. *et al.* Utrophin up-regulation by artificial transcription factors induces muscle
rescue and impacts the neuromuscular junction in mdx mice. *Biochim Biophys Acta*
Mol Basis Dis **1864**, 1172-1182, doi:10.1016/j.bbadis.2018.01.030 (2018).
25 Wright, A. V., Nunez, J. K. & Doudna, J. A. Biology and Applications of CRISPR Systems:
Harnessing Nature's Toolbox for Genome Engineering. *Cell* **164**, 29-44,
doi:10.1016/j.cell.2015.12.035 (2016).
26 He, Z. *et al.* Highly efficient targeted chromosome deletions using CRISPR/Cas9.
Biotechnology and bioengineering **112**, 1060-1064, doi:10.1002/bit.25490 (2015).
27 Ousterout, D. G. *et al.* Multiplex CRISPR/Cas9-based genome editing for correction of
dystrophin mutations that cause Duchenne muscular dystrophy. *Nat Commun* **6**, 6244,
doi:10.1038/ncomms7244 (2015).
28 Young, C. S. *et al.* A Single CRISPR-Cas9 Deletion Strategy that Targets the Majority of
DMD Patients Restores Dystrophin Function in hiPSC-Derived Muscle Cells. *Cell Stem*
Cell **18**, 533-540, doi:10.1016/j.stem.2016.01.021 (2016).
29 Amoasii, L. *et al.* Gene editing restores dystrophin expression in a canine model of
Duchenne muscular dystrophy. *Science*, doi:10.1126/science.aau1549 (2018).
30 Min, Y. L. *et al.* CRISPR-Cas9 corrects Duchenne muscular dystrophy exon 44 deletion
mutations in mice and human cells. *Sci Adv* **5**, eaav4324, doi:10.1126/sciadv.aav4324
(2019).
31 Duchêne, B. L. *et al.* CRISPR-Induced Deletion with SaCas9 Restores Dystrophin
Expression in Dystrophic Models In Vitro and In Vivo. *Mol Ther* **26**, 2604-2616,
doi:10.1016/j.ymthe.2018.08.010 (2018).
32 Mata Lopez, S. *et al.* Challenges associated with homologous directed repair using
CRISPR-Cas9 and TALEN to edit the DMD genetic mutation in canine Duchenne
muscular dystrophy. *PLoS One* **15**, e0228072, doi:10.1371/journal.pone.0228072
(2020).
33 Charlesworth, C. T. *et al.* Identification of preexisting adaptive immunity to Cas9
proteins in humans. *Nat Med* **25**, 249-254, doi:10.1038/s41591-018-0326-x (2019).
34 Lim, K. R. Q., Nguyen, Q., Dzierlega, K., Huang, Y. & Yokota, T. CRISPR-Generated
Animal Models of Duchenne Muscular Dystrophy. *Genes* **11**,
doi:10.3390/genes11030342 (2020).

35   Haeussler, M. *et al.* Evaluation of off-target and on-target scoring algorithms and
integration into the guide RNA selection tool CRISPOR. *Genome biology* **17**, 148-148,
doi:10.1186/s13059-016-1012-2 (2016).
36   van Deutekom, J. C. *et al.* Local dystrophin restoration with antisense oligonucleotide
PRO051. *N Engl J Med* **357**, 2677-2686, doi:10.1056/NEJMoa073108 (2007).
37   Li, H. L. *et al.* Precise Correction of the Dystrophin Gene in Duchenne Muscular
Dystrophy Patient Induced Pluripotent Stem Cells by TALEN and CRISPR-Cas9. *Stem cell*
reports, doi:10.1016/j.stemcr.2014.10.013 (2014).
38   Wojtal, D. *et al.* Spell Checking Nature: Versatility of CRISPR/Cas9 for Developing
Treatments for Inherited Disorders. *Am J Hum Genet*, doi:10.1016/j.ajhg.2015.11.012
(2015).
39   Lattanzi, A. *et al.* Correction of the Exon 2 Duplication in DMD Myoblasts by a Single
CRISPR/Cas9 System. *Mol Ther Nucleic Acids* **7**, 11-19, doi:10.1016/j.omtn.2017.02.004
(2017).
40   Nelson, C. E. *et al.* In vivo genome editing improves muscle function in a mouse model
of Duchenne muscular dystrophy. *Science* **351**, 403-407, doi:10.1126/science.aad5143
(2016).
41   Tabebordbar, M. *et al.* In vivo gene editing in dystrophic mouse muscle and muscle
stem cells. *Science* **351**, 407-411, doi:10.1126/science.aad5177 (2016).
42   Long, C. *et al.* Postnatal genome editing partially restores dystrophin expression in a
mouse   model   of   muscular   dystrophy.   *Science*   **351**,   400-403,
doi:10.1126/science.aad5725 (2016).
43   Popplewell, L. J. *et al.* Comparative analysis of antisense oligonucleotide sequences
targeting exon 53 of the human DMD gene: Implications for future clinical trials.
Neuromuscul Disord **20**, 102-110, doi:10.1016/j.nmd.2009.10.013 (2010).
44   Arechavala-Gomeza, V. *et al.* Comparative analysis of antisense oligonucleotide
sequences for targeted skipping of exon 51 during dystrophin pre-mRNA splicing in
human muscle. *Hum Gene Ther* **18**, 798-810, doi:10.1089/hum.2006.061 (2007).
45   Guiraud, S. & Davies, K. E. Pharmacological advances for treatment in Duchenne
muscular dystrophy. *Curr Opin Pharmacol* **34**, 36-48, doi:10.1016/j.coph.2017.04.002
(2017).
46   Loro, E. *et al.* High-throughput identification of post-transcriptional utrophin up-
regulators for Duchenne muscle dystrophy (DMD) therapy. *Sci Rep* **10**, 2132,
doi:10.1038/s41598-020-58737-6 (2020).
47   Arechavala-Gomeza, V. *et al.* Immunohistological intensity measurements as a tool to
assess sarcolemma-associated protein expression. *Neuropathol Appl Neurobiol* **36**,
265-274, doi:10.1111/j.1365-2990.2009.01056.x (2010).
48   Ruiz-Del-Yerro, E., Garcia-Jimenez, I., Mamchaoui, K. & Arechavala-Gomeza, V.
Myoblots: dystrophin quantification by in-cell western assay for a streamlined
development of Duchenne muscular dystrophy (DMD) treatments. *Neuropathol Appl*
Neurobiol **44**, 463-473, doi:10.1111/nan.12448 (2018).
49   Ran, F. A. *et al.* Genome engineering using the CRISPR-Cas9 system. *Nature protocols*
8, 2281-2308, doi:10.1038/nprot.2013.143 (2013).
50   Roest, P. A. *et al.* Application of in vitro Myo-differentiation of non-muscle cells to
enhance gene expression and facilitate analysis of muscle proteins. *Neuromuscul*
Disord **6**, 195-202, doi:10.1016/0960-8966(96)00006-5 (1996).

 Figure legends

Table 1. List of sgRNAs

List of sgRNAs for editing the *DMD* and the *UTRN* loci, showing sgRNAs sequences,

PAM sequences and scores of all sgRNA tested.

Table 2. Off-target sites analyses.

Top 6 off-target sequences of Ob1sgRNA2 and Ob1sgRNA6 identified with CRISPOR

webtool, including the mismatches between sgRNAs, the off-target sequence, the

chromosomes and loci targeted. All of them were analysed by PCR and Sanger

sequencing, and no off-targets were detected.

Table 3. Off-target sites analyses.

Top 6 off-target sequences of Ob2sgRNA22 and Ob2_sgRNA26 identified with CRISPOR

webtool, including the mismatches between sgRNAs, the off-target sequence, the

chromosomes and loci targeted. All of them were analysed by PCR and Sanger

sequencing, and no off-targets were detected.

Figure 1. Editing approach and genotyping *DMD* and *UTRN* deletion breakpoints in

edited myoblast clones.

(A and D). Schematic representation of our strategy for editing the *DMD* (A) and the

UTRN loci (D). A pair of flanking sgRNAs were co-transfected in order to delete *DMD*

exon 52 (A) or the inhibitory microRNA target region contained in the 5'UTR of *UTRN*

(D). (B and C) PCR genotyping of *DMD* edited clones (B) and Sanger sequencing of

clone 7 or DMDΔ52-Model (C) . (E and F) PCR genotyping of *UTRN* edited clones (E) and

Sanger sequencing of clone 8 or DMD-UTRN-Model (F). Larger products in agarose gels

(B and E) indicate non-edited clones, and shorter ones correspond with the expected

deletion. (C and F) Sequences of the smaller bands confirmed the expected gene

editing for objective 1: DMDΔ52-Model (C) and objective 2: DMD-UTRN-Model (F).

Figure 2. Characterization of DMDΔ52-Model cultures.

Dystrophin expression in DMDΔ52-Model cultures compared to control myotubes and

DMD myotubes were studied by immunocytochemistry (A), western blotting (B) and

myoblots (C). Myoblot experiments, where n=24 wells per cell type were compared,

were performed twice. (D) Dystrophin expression in DMDΔ52-Model cultures was

compared to control myotubes by ddPCR. For ddPCR experiments three technical

replicates per sample and condition were run in parallel and a no template control

(NTC) was included as negative control. (E) Differentiated myotubes of DMDΔ52-

Model and control cultures were inmunostained with MF20 and Hoechst antibodies.

Fusion index was calculated as the ratio between the number of nuclei in

differentiated myotubes (defined as >2 nuclei and MF20-positive cells) compared to

the total number of nuclei. For quantification, five fields per cell line were randomly

chosen and more than 200 nuclei were counted. Analysis was performed using ImageJ

software. (F) Differentiation markers, Myf5 and MyH3, were studied by ddPCR at

different fusion times in DMDΔ52-Model cultures compared to control myotubes. For

ddPCR experiments three technical replicates per sample and condition were run in

parallel and a no template control (NTC) was included as negative control. (*p-value

<0.05, ** p-value <0.01, ****p-value<0.0001). (P values were determined with Mann-

Whitney U test).

Figure 3. Characterization of DMD-UTRN-Model cultures.

Utrophin expression in DMD myotubes compared to DMD-UTRN-Model studied by

immunocytochemistry (A), western blotting (B) and myoblots (C). Myoblot

experiments, where n=48 wells per cell type were compared, were performed twice.

(D) Utrophin expression in DMD-UTRN-Model cultures was compared to DMD

myotubes by ddPCR. For ddPCR experiments three technical replicates per sample and

condition were run in parallel and a no template control (NTC) was included as

negative control. (E) Differentiated myotubes of DMD-UTRN-Model and DMD cultures

were inmunostained for MF20 and Hoechst. Fusion index was calculated as the ratio

between the number of nuclei in differentiated myotubes (defined as >2 nuclei and

MF20-positive cells) compared to the total number of nuclei. For quantification, five

fields per cell line were randomly chosen and more than 200 nuclei were counted.
Analysis was performed using ImageJ software. (F) Differentiation markers, Myf5 and
MyH3, were studied by ddPCR at different fusion times in DMD-UTRN-Model cultures
compared to DMD myotubes. For ddPCR experiments three technical replicates per
sample and condition were run in parallel and a no template control (NTC) was
included as negative control. (*p-value <0.05, ** p-value <0.01, ****p-value<0.0001).
(P values were determined with Mann-Whitney U test).

Figure 4. Evaluation of potential therapies in the generated cell culture models.

(A) DMDΔ52-Model cell line was treated with an antisense to skip exon 51 and the
effect was evaluated by RT-PCR and nested PCR analysis. Gel picture shows a pattern
corresponding with the correct skipping, which was confirmed by Sanger sequencing.
The same experiment evaluated by myoblot using n=20 replicate wells per condition
(B) showed the restoration of dystrophin expression in the treated cultures compared
to non-treated (*P values<0.05. P values were determined with Mann-Whitney U test).
(C) The DMD-UTRN-Model was used as a positive control in an experiment in which
unedited DMD cultures were treated with different ezutromid concentrations to up-
regulate utrophin expression. Myoblot analysis using n=8 replicate wells per condition
of the treated cultures shows that ezutromid had no significant effect in DMD cultures
while utrophin expression is significantly increased in DMD-UTRN-Model compared to
unedited DMD cultures. (**P values <0.01. P values were determined with Mann-
Whitney U test).
720

CODE	SCORE	SEQUENCE	PAM
sgRNA1	79	CTGAAGAACCCTGATACTAA	GGG
sgRNA2	78	GCTGAAGAACCCTGATACTA	AGG
sgRNA3	69	AACAAATATCCCTTAGTATC	AGG
sgRNA4	65	ACAAATATCCCTTAGTATCA	GGG
sgRNA5	60	TAAGGGATATTTGTTCTTAC	AGG
sgRNA6	48	ATTTCTAAAAGTGTTTTGGC	TGG
sgRNA7	44	AAAAAAGATGTTACTGTATA	AGG
sgRNA8	44	AAAAAGATGTTACTGTATAA	GGG
gsRNA9	29	TTTACTTTGTATTATGTAAA	AGG
sgRNA10	24	TTTTATTTCTAAAAGTGTTT	TGG
sgRNA21	71	AACTTTGGGTTCTCTTTAGC	TGG
sgRNA22	66	GGTTCTCTTTAGCTGGGATC	TGG
sgRNA23	63	TATTTTAGAATAGGTTGGGT	GGG
sgRNA24	62	ACTTTGGGTTCTCTTTAGCT	GGG
sgRNA25	62	TCTAACTTTAAGCCTCCTTC	TGG
sgRNA26	76	GTGCTTTCTTGGGTATGACA	TGG
sgRNA27	68	CAAAGTCTAGAGCTTTTATC	AGG
sgRNA28	66	CAACTTGGAGTTGAGAGCTC	AGG
sgRNA29	64	TCAACTCCAAGTTGTAGATT	TGG
sgRNA30	63	TCCATCTTCATCCATTGCAT	TGG

Table 1. List of sgRNAs

Off Target name	Guide sequence	Off target sequence	Mis	Chrom	Locus	PCR analysis	Sequencing
Ob1_2_Off1		GCTGGAGAACCCTGATACTGTGG	2	chr1	intergenic:RP4-781L3.1-RP4-706G24.1	No off-target editing	Confirmed
Ob1_2_Off2		TCTGGAGAACCCTAATACTAAGG	3	chr8	intergenic:RP11-24P4.1-AC009695.1	No off-target editing	Confirmed
Ob1_2_Off3	GCTGAAGAACCCTGATACTAAGG (Ob1_sgRNA2)	ACTGAAGAATCCAGAAACTAGGG	4	chr7	intergenic:NOBOX-RP4-545C24.5	No off-target editing	Confirmed
Ob1_2_Off4		GCTAGAGAAACCTGAAACTAAGG	4	chr8	intergenic:RP11-536K17.1-EIF3H	No off-target editing	Confirmed
Ob1_2_Off5		TCTGGAGAACCCTAATACTGTGG	4	chr3	intron:TMEM45A	No off-target editing	Confirmed
Ob1_2_Off6		TCTGAAGAATCCTGATATTTTGG	4	chr2	intron:AC019100.3	No off-target editing	Confirmed
Ob1_6_Off1		CACCATCAGCCAAGAATATGCGG	3	chr11	intergenic:RP11-430H10.3-RP11-958J22.1	No off-target editing	Confirmed
Ob1_6_Off2		TAACAACAGCCAAAGACATGAGG	4	chr14	exon:RP11-1012A1.4/RDH11	No off-target editing	Confirmed
Ob1_6_Off3	GACCAACAGCCAAGGATATGAGG (Obj1 sgRNA6)	GTAAAAGAGCCAAGGATATGAGG	4	chr10	intron:RP11-556E13.1	No off-target editing	Confirmed
Ob1_6_Off4		TACTAGCAGCCAAGGATATCTGG	4	chr2	intergenic:AC007377.1-SLC8A1	No off-target editing	Confirmed
Ob1_6_Off5		GAGCGACAGCCAAGAATATTCGG	4	chr3	intron:CD96	No off-target editing	Confirmed
Ob1_6_Off6		AATCAACAGCCAAGAATGTGGGG	4	chr5	intergenic:CTD-2201E9.4-SEMA5A	No off-target editing	Confirmed

742   **Table 2. Objective 1 potential off-target sites.**

Off Target name	Guide sequence	Off target sequence	Mis	Chrom	Locus	PCR analysis	Sequencing
Ob2_22_Off1	GGTTCTCTTTAGCTGGGATCTGG (Obj2 sgRNA22)	TGTTCTCTCTAACTGGGATCTGG	3	chr18	intergenic:RP11-411B10.6-RP11-411B10.5	No off-target editing	Confirmed
Ob2_22_Off2		TGTTCTCTAGAGCTGGGATCTGG	3	chr21	intron:LCA5L	No off-target editing	Confirmed
Ob2_22_Off3		TGTTCTCTCCAACTGGGATCTGG	4	chr22	intron:PPP6R2	No off-target editing	Confirmed
Ob2_22_Off4		GAATCCTTTTAGCTGGGATCAGG	4	chr19	intron:ZNF536	No off-target editing	Confirmed
Ob2_22_Off5		GGTTCATCTTAGCTGGGATATGG	4	chr13	intron:FLT1	No off-target editing	Confirmed
Ob2_22_Off6		TGTTCTCTCTAACTGGGGTCTGG	4	chr21	intergenic:PPP6R2P1-AP001347.6	No off-target editing	Confirmed
Ob2_26_Off1	GTGCTTTCTTGGGTATGACATGG (Obj2 sgRNA26)	AAGCTTTCCTGGATATGACAAGG	4	chr4	intron:RNF150	No off-target editing	Confirmed
Ob2_26_Off2		GTGCTTACTTGGGTAAGACGTGG	3	chr17	intergenic:RP11-212E8.1-RP11-642M2.1	No off-target editing	Confirmed
Ob2_26_Off3		GAGTTAACTTGGGTATGACAGGG	4	chr4	intron:RGS12	No off-target editing	Confirmed
Ob2_26_Off4		GTGCTCTCATGAGAATGACAGGG	4	chr4	intergenic:GABRG1-RP11-320H14.1	No off-target editing	Confirmed
Ob2_26_Off5		GAGCTTTCCTGGGAATGACAGGG	3	chr1	intergenic:FOXO6-RNA5SP45	No off-target editing	Confirmed
Ob2_26_Off6		GTGCTTTATAGGATATAACATGG	4	chr6	intron:GSTA3	No off-target editing	Confirmed

Table 3. Objective 2 potential off-target sites.

A)

| EXON49 | EXON50 | EXON51 | EXON53 |

| EXON49 | EXON50 | EXON53 |

B)

Dystrophin expression by myoblots

C)

Utrophin expression by myoblot