Toxic stress-specific cytoprotective responses regulate learned behavioral decisions in *C. elegans*

Gábor Hajdú, Eszter Gecse, István Taisz[#], István Móra & Csaba Sőti*

Department of Medical Chemistry, Semmelweis University, Budapest, Hungary

[#]*Current address: MRC Laboratory of Molecular Biology, Neurobiology Division, Cambridge, UK*

***Corresponding author.** Department of Medical Chemistry, Semmelweis University, P.O.Box 2, Budapest, H-1428, Hungary. Tel.: + 36 1 4591500 extn. 60130; Fax: + 36 1 4591500 extn. 60141; E-mail: soti.csaba@med.semmelweis-univ.hu

Keywords: cytoprotection, behavioral decision, aversion, associative memory, stress and detoxification responses

Abstract

Protection of organismal integrity involves both molecular cytoprotective responses and behavioral defenses. Here we show that exposure of *Caenorhabditis elegans* to high concentrations of naturally attractive food-derived odors, benzaldehyde and diacetyl, induces toxicity and aversive behavior. Benzaldehyde preconditioning activates systemic cytoprotective responses involving DAF-16/FOXO, SKN-1/Nrf and Hsp90 in somatic cells, which confer behavioral tolerance to benzaldehyde and cross-tolerance to methyl-salicylate, but not to diacetyl. In contrast, diacetyl preconditioning augments diacetyl avoidance and does not induce apparent molecular defenses. Reinforcement of the experiences by massed training forms relevant associative memories. Memory retrieval by the odor olfactory cues leads to avoidance of food contaminated by diacetyl and context-dependent behavioral decision to avoid benzaldehyde only if there is an alternative, food-indicative odor. Our findings reveal a regulatory link between cytoprotective mechanisms and behavior which facilitates self-protection in real and anticipated stresses and might have relevance in maladaptive avoidant human behaviors.

Introduction

Adequate, coordinated responses of multicellular organisms are key to adapt to and overcome fundamental alterations of the environment (Bijlsma and Loeschcke, 2005; Kültz, 2005; Selye, 1973). These responses originate from intracellular molecular defenses, such as the oxidative, xenobiotic, metabolic and proteotoxic stress responses, which guard homeostasis and confer cytoprotection against the respective stresses, promoting physiological adaptation, fitness and longevity at the organismal level (Kenyon, 2010). Adaptation also involve complex behavioral responses orchestrated by the neuroendocrine system (Badyaev, 2005; Baldwin, 2013; McCarty, 2016). For instance, sensory cues representing danger evoke aversive behavior as a result of perception of multiple sensory stimuli, neuronal processing and decision making both in humans and in other species (Ardiel and Rankin, 2010; Barrett and Broesch, 2012; Puzzo et al., 2016). In some cases, the neural impulse of perceived danger is so intense that the organism decides to avoid co-occurring cues representing life-sustaining qualities such as food (Baldwin, 2013; Garcia, 2017). Besides external sensory cues, decision making is modulated by neural context like arousal, motivation, and reward (Botvinick and Braver, 2015; Robbins and Everitt, 1996). Importantly, behavioral decisions are also influenced by sensory cues that evoke associative memories of past events (Cosmides and Tooby, 2013). Moreover, exaggerated, inadequate avoidant behavior is characteristic to human anxiety disorders such as phobias (Garcia, 2017), where sometimes intense physical symptoms of toxicity and disgust are evoked by olfactory cues. Although the neuroendocrine mechanisms of stress are extensively studied, the contribution of somatic, especially intracellular defenses to behavioral regulation is largely unexplored.

The soil nematode *Caenorhabditis elegans* with its 959 cells is a versatile model system to study the link between cytoprotective stress responses and behavior. Worms, using a well-defined network of 302 neurons are capable of complex behavioral decisions (Bargmann, 2006; Bargmann et al., 1993; Hobert, 2003). Flavors and volatiles have great impact on decision making of nematodes, informing about possible nutrition and danger *via* neuronal processing of olfactory and gustatory cues (Bargmann, 2006). Besides well characterized escape responses, tissue damaging insults, such as toxins and pathogens, induce a network of evolutionary conserved cytoprotective defenses in each somatic cell and in specialized tissues (Kenyon, 2010). Fixing the actual damage as well as eliminating damaging agents are key mechanisms of cellular protection (Blackwell et al., 2015). Nematodes and mammals share diverse molecular processes to recognize and overcome toxic, stressor agents. One of *C. elegans'* key

stress-responsive molecule is the forkhead type, FOXO ortholog DAF-16 transcription factor
regulated by the INS/IGF-1 signaling pathway (Ogg et al., 1997). DAF-16 is ubiquitously
expressed, localized in the cytosol, and is activated by nuclear translocation in response to
oxidative and genotoxic agents, starvation, desiccation and heat stress (Zečić and Braeckman,
2020). Loss-of-function mutation in the *daf-16* locus results in compromised stress resistance,
dauer development and shorter lifespan (Hsu et al., 2003).

The Nrf-2 orthologue SKN-1 transcription factor is the major xenobiotic and oxidative
stress regulator in nematodes (Blackwell et al., 2015; Park et al., 2009). Its nuclear translocation
is induced by dietary restriction, pathogen attack, the INS/IGF-1 and TIR-1/PMK-1 pathways
to modulate cellular respiration, enhance oxidative stress resistance, immunity and systemic
detoxification defenses (Inoue et al., 2005; Papp et al., 2012; Park et al., 2009). SKN-1
cooperates with numerous stress related pathways and regulators including DAF-16 and the *C.*
elegans heat-shock transcription factor orthologue HSF-1 to fine-tune cytoprotective gene
expression patterns (Blackwell et al., 2015). Upregulation of specific and overlapping
molecular stress responses explains an adaptive process called physiological conditioning
hormesis in stress biology (Calabrese et al., 2007). In course of hormesis, a conditioning stress
exposure results in increased survival under a subsequent, lethal stress of the same or different
kind, a phenomenon called stress tolerance or cross tolerance, respectively. To clearly
discriminate physiological and behavioral terms, we use the term preconditioning for
physiological conditioning and introduce a new term, behavioral tolerance for diminished
avoidance.

In this study, we set out to investigate how cytoprotective molecular defenses influence
stress-induced behaviors and learned behavioral decisions. To this end, we employed two food-
derived volatile odors, benzaldehyde (BA) and diacetyl (DA), which are attractive at low, but
aversive at high concentrations (Nuttley et al., 2001; Yoshida et al., 2012). The advantage of
these odors is that they contain both the chemosensory cue as well as a dual, attractive or
aversive property. Our results indicate a critical role of the ability to mount specific systemic
cytoprotective responses in shaping adaptive behavioral decisions based on associative
learning.

4

Results

Concentrated benzaldehyde and diacetyl induce toxicity and food avoidance

Low concentrations of food odors are attractive to *C. elegans*, whereas high concentrations induce an aversive response (Yoshida et al., 2012). Specifically, worms exhibit a biphasic chemotaxis curve towards concentrated 100% benzaldehyde (ccBA) called benzotaxis (Nuttley et al., 2001 and Figure 1-figure supplement 1A). We hypothesized that the second, aversive phase is a defensive behavioral response to ccBA toxicity. Indeed, we found that longer ccBA exposures using the aversive concentration ranges induced extensive paralysis in a dose- and time-dependent manner (Figure 1A). To investigate whether another concentrated food odor may induce toxicity at aversive concentrations, we tested the chemically unrelated diacetyl. Undiluted diacetyl (ccDA) also triggered biphasic chemotaxis behavior (Figure 1-figure supplement 1B) and dose-dependent paralysis at approximately four-fold higher doses compared to ccBA (Figure 1B). Furthermore, aversive, but lower doses of ccBA and ccDA both impaired thermotolerance (Figure 1-figure supplement 1C), demonstrating compromised organismal integrity and stress resistance in response to odor toxicity. Therefore, we used non-paralyzing doses of odors as a source of toxic stress throughout this study.

Figure 1. Concentrated benzaldehyde (ccBA) and diacetyl (ccDA) induce toxicity and food avoidance in the aversive concentration range.

Time-dependence curves of paralysis to various doses of ccBA (A) or ccDA (B) using a hanging drop assay. (C) Representative images of food leaving behavior in response to various concentrations of BA. BA was placed in ethanol vehicle in a total volume of 1µl in the middle of bacterial lawn. (D) Dose dependence of BA and DA induced food avoidance. BA or DA was

placed in a total volume of 1 µl or 4 µl in the middle of bacterial lawn. (E) Time dependence of ccBA and ccDA induced food avoidance. Data are expressed as mean ± SEM, N = number of independent experiments. *p < 0.05; **p < 0.01; ***p < 0.001. Mean durations of odor exposure that induced 50% paralysis by log rank (Mantel-Cox) test were as follows: ccBA – 2 µl: 5.27 h ± 0.17 hours, 4 µl: 2.94 ± 0.21 hours, 8 µl: 0.94 ± 0.14 hours. ccDA? – 8 µl: 5.68 ± 0.20 hours, 16 µl: 3.46 ± 0.17 hours. P-values compared to 1 µl BA or 4 µl DA treatments are <0.001 in all conditions.

Figure 1 supplement 1. Concentrated benzaldehyde (ccBA) and diacetyl (ccDA) induce behavioral and physiological alterations characteristic to toxicity.

Although food is not necessary for adult worms' survival, worms are continuously feeding and seldom leave the bacterial lawn under laboratory conditions (Avery and You, 2012). Therefore, to establish a more stringent test for behavioral adaptation, we placed a ccBA drop in the middle of an *E. coli* OP50 lawn and monitored food avoidance. We observed that while unexposed worms remained on the lawn after 50 minutes, the majority of ccBA-exposed worms left the food (Figure 1C). We observed similar, concentration- and time-dependent food aversion phenotypes with both ccBA and ccDA (Figure 1D, E). These findings indicate that the perception of toxic stress initiates a decision to leave the lawn, giving up the benefit of nutrients for the protection of physical integrity.

Opposing behavioral patterns elicited by toxic benzaldehyde and diacetyl exposure

We observed that transient ccBA and ccDA exposure increased motility (Figure 2-figure supplement 1A), indicating that perception of toxic stress increases locomotor activity which may help instantly avoid the threat. Interestingly, the increased motility returned to baseline after removing ccBA, but showed a sustained elevation after the removal of ccDA (Figure 2-figure supplement 1A). Moreover, we found that after an extended, 2-hour exposure to ccBA, animals started to return to the bacterial lawn, whereas the same exposure to ccDA further increased aversion (Figure 2-figure supplement 1B). Thus, the adverse physiological effects of ccBA might be eliminated faster than those of ccDA. We reasoned that a preconditioning exposure might differentially affect behavior. To test this, after exposure to the same sublethal doses of odors we investigated the lawn avoidance behavior of naive and preconditioned worms (Figure 2A). Indeed, we found that preconditioning with ccBA diminished ccBA-induced

aversion, while that with ccDA further increased avoidance of ccDA (Figure 2A, B). For the increased capacity of worms to remain in the presence of toxic ccBA we coined the term "behavioral tolerance", to the analogy of physiological stress tolerance. Thus, ccBA preconditioning induces behavioral tolerance, while ccDA preconditioning induces sensitization. The effect developed by a 2-hour preconditioning and was only moderately altered by further increasing the duration of the preconditioning exposure (Figure 2-figure supplement 1C and D).

Figure 2. ccBA preconditioning induces behavioral tolerance, while ccDA preconditioning induces sensitization.

(A) Experimental setup for preconditioning and food aversion test. Animals were exposed to a hanging drop of concentrated odor (preconditioned, PC) or vehicle (naive), washed and assayed for food aversion. (B) ccBA induced food aversion of naive and ccBA PC animals at different time points. (C) ccDA induced food aversion of naive and ccDA PC animals at different time points. Data are expressed as mean ± SEM, N = number of independent experiments. *p < 0.05; **p < 0.01; ***p < 0.001.

Figure 2 supplement 1. Opposing effects of odor pre-exposure on odor induced behaviors.

Concentrated benzaldehyde, but not concentrated diacetyl, activates specific systemic cytoprotective responses

In agreement with our findings on the toxicity of ccBA, previous studies demonstrated that BA induced oxidative stress (Andersen, 2006; Tabatabaie and Floyd, 1996). Therefore, we tested various oxidative stress response pathways that might be involved in the organismal adaptation to ccBA. Using the TJ356 strain expressing GFP-tagged DAF-16, we observed that ccBA exposure induced a strong nuclear translocation of DAF-16, comparable to that induced by heat stress. However, DAF-16 remained cytosolic in response to ccDA (Figure 3A and Figure 3-figure supplement 1A). The shift in DAF-16 localization exhibited a clear BA dose-dependence (Figure 3-figure supplement 1B). These congruent changes in DAF-16 translocation and food avoidance (cf. Figure 1D) indicate a potential link between cytoprotective responses and behavioral tolerance.

Next, we tested several other stress and detoxification pathways using GFP-tagged marker strains. Translocation of the oxidative-xenobiotic stress master regulator SKN-1::GFP in the LD001 strain was induced by ccBA, but not by ccDA, comparable to that seen upon the oxidative agent paraquat (PQ) treatment (Figure 3B). Further, ccBA, but not ccDA induced the expression of the phase I marker cytochrome P450 enzyme *cyp-35b* and the phase II enzyme *gst-4* (Figure 3C-F). The induction of *cyp-35b* was abolished by *daf-16* RNAi, while that of *gst-4* was abolished by *skn-1* RNAi (Figure 3D, F). Importantly, neither the HSF-1 and DAF-16 target *hsp-16.2*, and the HSF-1 target and endoplasmic reticulum unfolded protein response (UPR) reporter *hsp-4*, nor the SKN-1 dependent *gcs-1* and the DAF-16 dependent *sod-3* reporter was induced by ccBA (Figure 3-figure supplement 1C). These findings demonstrate that a specific stress and detoxification response involving a subset of DAF-16 and SKN-1 activated genes is required for the molecular defense against ccBA toxicity.

198 **Figure 3. ccBA, but not ccDA, activates specific systemic cytoprotective responses.**

199 Representative epifluorescent microscopic images of DAF-16::GFP nuclear translocation in

200 response to heat stress, ccBA or ccDA (A), as well as *SKN-1::GFP* nuclear translocation in

201 response to paraquat, ccBA or ccDA (B, arrowheads). Representative epifluorescent

202 microscopic images (C) and quantification (D) of *cyp-35b::GFP* expression in response to

ccBA or ccDA in worms fed by control EV and *daf-16* RNAi. Representative epifluorescent microscopic images (E) and quantification (F) of *gst-4::GFP* expression in response to ccBA or ccDA in nematodes fed by control EV and *skn-1* RNAi. Data are expressed as mean ± SEM, N = number of independent experiment replicates. n.s.: not significant, *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

Figure 3 supplement 1. Characteristics of DAF-16 nuclear translocation and specificity of ccBA-induced cytoprotective responses.

ccBA-induced cytoprotective responses confer behavioral tolerance to ccBA, but not to ccDA

We asked whether the cytoprotective responses induced by ccBA which are known to induce physiological stress tolerance might play a role in the generation of behavioral decisions. To this end, we preconditioned N2 and *daf-16* null mutant nematodes with ccBA and studied their food avoidance to ccBA. We found that naive *daf-16* mutants showed avoidant behavior comparable to wildtype, however, they failed to decrease their aversion in response to preconditioning (Figure 4A). Similar phenotype was obtained by silencing the evolutionarily conserved molecular chaperone Hsp90, which was shown to regulate DAF-16 activity (Somogyvári et al., 2018) (Figure 4B). Likewise, *skn-1* silencing similarly prevented the development of behavioral tolerance, whereas the activation of SKN-1 by knocking down the WDR-23 protein responsible for its degradation (Choe et al., 2009) augmented behavioral tolerance towards ccBA (Figure 4C). In sharp contrast, after ccDA preconditioning, neither *skn-1*, nor *wdr-23* RNAi altered the behavioral sensitization towards ccDA exposure (Figure 4D). These results indicate that the specific cytoprotective responses induced by toxic ccBA exposure actively participate in the development of behavioral tolerance.

Figure 4. ccBA-induced cytoprotective responses confer behavioral tolerance to ccBA, but not to ccDA.

(A) ccBA induced food aversion of naive and ccBA preconditioned N2 wildtype and *daf-16(mu86)* mutant animals, as well as (B) N2 fed by EV and *hsp-90* RNAi bacteria. (C) ccBA induced food aversion of naive and ccBA preconditioned nematodes fed by control EV, *skn-1* and *wdr-23* RNAi, respectively. (D) ccDA induced food aversion of naive and ccDA preconditioned nematodes fed by control EV, *skn-1* and *wdr-23* RNAi, respectively. Data are expressed as mean ± SEM. N = number of independent experiments. *p < 0.05; **p < 0.01; ***p < 0.001.

Behavioral cross-tolerance is mediated by chemical structure-specific cytoprotective responses

Xenobiotic-induced stress and detoxification responses are related to the chemical structure of the toxin as well as the nature of damage they induce (Galal et al., 2015). We reasoned that the observed BA-dependent cytoprotective machinery might also be induced in response to a chemically similar toxic compound. BA is both spontaneously as well as enzymatically oxidized to benzoic acid during its detoxification (Andersen, 2006; Jorissen and van der Beek, 2010; Sankar et al., 2014) (Figure 5A). The chemical structure of the volatile plant stress hormone methyl-salicylate (MS) (Park et al., 2007), harboring an aromatic benzene ring and an

esterified carboxyl group is closely related to that of benzaldehyde and benzoic acid (Figure 5A). We found that similarly to ccBA, concentrated MS (ccMS) was also toxic and induced food avoidance behavior (Figure 5-figure supplement 1A and B). Moreover, ccMS and ccBA shared identical molecular defense responses, including DAF-16 translocation, *cyp-35b::GFP* and *gst-4P::gfp* expression (Figure 5B). Importantly, preconditioning with either ccMS or ccBA reduced food aversion in response to a subsequent ccMS exposure (Figure 5C). However, ccBA preconditioning did not affect food aversion in the presence of ccDA, indicating that DAF-16 and SKN-1 dependent processes are unable to reduce ccDA toxicity (Figure 5D). We conclude that the BA-specific cytoprotective responses confer behavioral cross-tolerance towards a toxin harboring a similar chemical structure, but not towards another compound, DA, which is unrelated chemically and probably by mechanism of action.

Figure 5. Behavioral cross-tolerance is mediated by chemical structure-specific cytoprotective responses.

(A) Benzaldehyde, its metabolite benzoic acid and methyl-salicylate (MS) share similar chemical structures. (B) Representative epifluorescent microscopic images showing the effect of concentrated MS (ccMS) exposure on DAF-16::GFP nuclear translocation, *cyp-35b*::GFP and *gst-4P::gfp* reporters expression. (C) ccMS (MS PC) as well as ccBA (BA PC) preconditioning significantly and comparably decrease ccMS-induced lawn avoidance, while ccBA preconditioning does not reduce ccDA-induced lawn avoidance (D). Error bars represent mean ± SEM compared to the respective naive values, N = number of independent experiments. n.s.: not significant, **p<0.01, ***p<0.001.

Figure 5 supplement 1. Concentrated methyl-salicylate (ccMS) exposure impairs survival and triggers food avoidance behavior.

Deficient or efficient cellular protection generate relevant learned behaviors to stress-associated olfactory cues

The lack of behavioral tolerance in case of ccDA preconditioning indicates inefficient cellular protection, in agreement with our findings (see Figure 3). Moreover, the phenomenon of behavioral sensitization, a significantly faster and more pronounced aversive response towards the odor suggests a role for avoidant associative learning. To test this, we investigated alterations in behaviors towards attractive (1%) doses of DA and BA after pre-exposure of toxic, concentrated doses of the respective odors (Figure 6A). Indeed, worms preconditioned with ccDA significantly reduced their chemotaxis towards naturally attractive 1% DA (Figure 6B) and even chose to leave the food in the presence of 1% DA (Figure 6C). We also investigated decision making by providing both DA and BA naturally associated with food in an odor choice assay. The aversive change of the DA olfactory cue was underscored by an almost complete shift in odor preference to BA (Figure 6D). In contrast, worms preconditioned with ccBA did not leave the bacterial lawn in the presence of 1% BA (Figure 6F). Moreover, they maintained their chemotaxis towards, 1% BA (Figure 6E), when the olfactory cue of BA was the only option. However, they displayed reduced preference to BA in the simultaneous presence of attractive DA (Figure 6G). These results are consistent with the formation of distinct, avoidant or tolerant learned behaviors associated to the sensory cues of DA and BA, respectively, after a previous encounter with their toxic doses, which appear to stem from the prior internal experience resulting from a deficient or efficient cytoprotection.

Figure 6. Learned stress-associated behaviors are shaped by prior cellular defenses against stress.

(A) Experimental design for toxic odor preconditioning induced associative learning. Animals were exposed to a hanging drop of concentrated odor (ccBA or ccDA, preconditioned, PC) or vehicle (naive), washed and assayed for chemotaxis, food aversion and odor preference using 1% of the odors. ccDA preconditioning significantly reduces chemotaxis to (B), and stimulates lawn avoidance from (C) 1% DA, and entirely shifts odor preference to 1% BA (D). ccBA preconditioning does not affect chemotaxis to (E), and lawn avoidance from (F) 1% BA, but

induces a modest reduction of preference to 1% BA in the presence of 1% DA (G). Odor choice was quantified by scoring worms on BA and DA spot. Error bars represent mean ± SEM. N = number of independent experiments. n.s.: not significant, *p < 0.05; **p < 0.01; ***p < 0.001.

A stress-associated memory of somatic resilience enables context-dependent decision making

The association of naturally attractive odors with learned stress-reactive behaviors raises the possibility that the learned experiences may give rise to associative memories to cope with a similar anticipated future insult. On the other hand, forgetting irrelevant experiences is also important as both the organism and the environment is changing. Indeed, a 2-hour recovery period after a single ccBA preconditioning of two or four hours significantly attenuates behavioral tolerance in the food leaving assay (Figure 7B). We reasoned that repeated exposure of the same conditions might reinforce the co-occurring experience resulting in a lasting neural representation. To test this, we employed a protocol of massed training known to induce short and intermediate term associative memories (STAM and ITAM) (Amano and Maruyama, 2011; Kauffman et al., 2011) (Figure 7A). By definition, STAM decays within, whereas ITAM persists over, one hour. Massed training of one-hour exposure to ccBA four times resulted in a potent behavioral tolerance that was retained after a 2-hr recovery (Figure 7C). A single 4-hour preconditioning, a physiological stress of the same duration induced comparable food aversion immediately after training (cf. Figs. 6F and 7C), which indicates similar levels of cytoprotection and makes it unlikely that the behavioral tolerance is a result of higher physiological stress tolerance. Likewise, ccDA massed training induced a robust food avoidance in the presence of 1% DA which persisted after 2-hr recovery (Figure 7D). Again, massed-trained worms exhibited comparable aversive behaviors against 1% DA as if they encountered ccDA (cf. Figures 2C, 6C and 7D). These results are consistent with the reinforcement of toxic stress-associated neurosensory integration into different associative memories of active coping or passive avoidance. The stability of memory after two hours indicates the formation of ITAM. Finally, we asked how the coping memories affect the choice between the stress-associated and a natural attractive odor olfactory cue. Massed training with ccBA almost entirely shifted the preference towards DA (Figure 7E), potentiating the change already observed by a single preconditioning (Figure 6G). This phenomenon also shows an apparent similarity to the complete disappearance of DA preference after a single preconditioning with ccDA (cf. Figures

7E and 6D). Nonetheless, in contrast to the compelling avoidant behavior to the memory of uncompensated harm, the memory of cellular protection not only provides the ability to cope with anticipated toxicity for food, but allows a context-dependent decision to spare resources when the organism also perceives the cue of a potentially toxin-free food. This result also suggests that the memory of a stressful insult contains the representation of the original valence of the olfactory cue, the internal experience of stress-induced harm and that of the induced cytoprotection.

Figure 7. **Reinforcement of stress-associated internal experiences form distinctive intermediate term associative memories.**

(A) Experimental design for toxic odor preconditioning induced associative memory. Animals were exposed to a hanging drop of concentrated odor (1 µl ccBA or 4 µl ccDA, preconditioned, PC) or vehicle (naive), using a single preconditioning (B) or a 4x massed training protocol (C-E), then assayed for food aversion or odor preference immediately or after the indicated

recovery periods. (B) A 2-hour recovery period decreases the food aversion elicited by a single 2-hour or 4-hour ccBA preconditioning. (C) Nematodes exposed to a 4x1-hour ccBA massed training retain the behavioral tolerance to ccBA after a 2-hour recovery period. (D) Nematodes exposed to a 4x1-hour ccDA massed training maintain their avoidant behavior to 1% DA after a 2-hour recovery period. (E) A 4x1-hour ccBA massed training induces a robust shift in odor preference towards 1% DA. Odor choice was quantified by scoring worms on BA, DA and on the empty agar surface (0). Error bars represent mean ± SEM, N = number of independent experiments. n.s.: not significant, *p < 0.05, **p<0.01, ***p<0.001.

Discussion

In this study we have set up a paradigm in *C. elegans* to assess the impact of cytoprotective responses on behavioral decisions. We have shown that the innately attractive odors benzaldehyde and diacetyl, when employed at high concentrations, induce toxicity and behavioral aversion of the odor-contaminated bacterial lawn. ccBA-induced systemic cytoprotective responses involving DAF-16, SKN-1 and Hsp90 conferred behavioral tolerance to ccBA and cross-tolerance to concentrated methyl-salicylate (ccMS), while neither behavioral tolerance nor apparent molecular defenses were observed upon exposure to ccDA. Massed training generated an associative memory that made diluted DA aversive but enabled animals to decide whether to approach or to avoid diluted BA depending on alternative choice. Our study suggests that the (in)ability of *C. elegans* to counteract toxic stress with cytoprotective mechanisms regulates behavior during stress and determines learned behavioral decisions upon re-encounter with stress-associated olfactory cues (Figure 8).

Previous studies reported either decreased attraction or aversion to high concentrations of food-derived odors (Bargmann et al., 1993; Nuttley et al., 2001; Troemel et al., 1995; Yoshida et al., 2012). One study proposed that the change in odor preference towards concentrated benzaldehyde was due to olfactory adaptation (Nuttley et al., 2001). Although this mechanism might hold true, several findings of our study argue for an alternative explanation. The *(i)* strong dose-dependent food aversion (Figure 1C-E) *(ii)* increased motility (Figure 2-figure supplement 1A) *(iii)* progressive dose-dependent paralysis (Figure 1A, B) *(iv)* compromised thermotolerance (Figure 1-figure supplement 1C) *(v)* induction of robust systemic cytoprotective responses by ccBA (Figure 3) *(vi)* the comparable ccBA concentration dependence of aversion and DAF-16 nuclear translocation (Figure 1D and Figure 3-figure

19

supplement 1B) (*vii*) the manipulation of these responses modulates the development of
behavioral tolerance to ccBA (Figure 4) (*viii*) RNAi is unable to penetrate neurons (Winston et
al., 2002) all indicate somatic toxicity as an underlying mechanism. This interpretation gains
support from the facts that low concentrations of benzaldehyde and isoamyl alcohol mediate
attraction via activating the AWC chemosensory neuron, whilst high concentrations activate
the polymodal nociceptive ASH neuron, which in turn drives repulsion (Aoki et al., 2011;
Bargmann et al., 1993; Troemel et al., 1995; Yoshida et al., 2012). Consistent with our study,
various studies in mammals describe the toxic effects of benzaldehyde (Andersen, 2006; Kluwe
et al., 1983; McKim et al., 1987) and diacetyl (Brass and Palmer, 2017; Clark and Winter, 2015;
Shibamoto, 2014) such as inhalation toxicity and long-term impairment in lung function.
Together with the evidence presented by the above reports, our study suggests that tissue
damage caused by odor toxicity stimulates aversion (Figure 8). Perhaps neural (Nuttley et al.,
2001) and somatic (our study) inputs are integrated to increase the robustness of behavior. Our
findings also draw attention to nematode associative learning experiments where different
conditions are paired with concentrated odors (Amano and Maruyama, 2011; Nuttley et al.,
2002; Sasakura and Mori, 2013), because odor toxicity might stimulate repulsion independently
of, or synergistically with, the unconditioned stimulus. Therefore, behavioral experiments using
diluted odors as the conditioned stimulus are recommended.

Our observations on the toxic odor-induced food aversion (Figure 1) indicates this
neurobehavioral response is a first line of defense against dangerous insults, which preserves
physical integrity and spares resources. ccBA exposed worms, however, started to return to
food during the second hour and a preconditioning exposure also diminished ccBA avoidance
(Figure 2-figure supplement 1B and Figure 2B). Reduced avoidance coincided with DAF-16
and SKN-1 activation and induction of phase 1 and 2 xenobiotic detoxification reporters (Figure
3), consistent with the aromatic structure and toxic profile of ccBA. Such cytoprotective stress
and detoxification responses co-operate to ensure survival, stress tolerance, immunity and
longevity (Blackwell et al., 2015; Kenyon, 2010; Singh and Aballay, 2006; Tullet et al., 2008),
forming a cellular defense. Our findings that *daf-16* knockout, *hsp-90*, *skn-1* and *wdr-23* RNAi
in somatic cells specifically modulate ccBA avoidance (Figure 4) show that specific stress-
responsive regulators control aversion, revealing a novel regulatory role of somatic
cytoprotective responses on behavioral decisions.

We found ccBA-induced behavioral cross-tolerance to concentrated methyl-salicylate
(ccMS) (Figure 5C). Our results on identical cytoprotective responses shared by ccBA and

ccMS (Figure 5) suggest that the responses stimulated by ccBA preconditioning also eliminate the toxic agent and repair damage during ccMS exposure. Indeed, high doses of methyl-salicylate cause heavy toxicity in mammals (Greene et al., 2017). Thus, the preservation/restoration of tissue integrity by toxin-specific cytoprotective responses suppresses aversion. Consistent with this idea, ccDA treatment resulted in sustained hypermotility after ccDA removal and increased ccDA aversion (Figure 2-figure supplement 1A and Figure 2C). Furthermore, ccDA did not appear to activate considerable molecular defenses (Figure 3) and neither SKN-1 modulation (Figure 4D) nor the systematic induction of defenses by ccBA preconditioning (Figure 5D) affected ccDA aversion. These results also exclude the unlikely possibility that induction of systemic cytoprotective responses *per* se inhibits aversive behavior. Hence, in the absence of adequate molecular defenses, the disruption of cellular integrity may represent a danger signal which induces aversion.

The mechanisms by which toxin-induced disruption of cellular integrity elicits aversion is unclear. The stress-activated JNK and p38 MAP kinases are conserved signal transducers of cell stress including abiotic, xenobiotic, oxidative, proteo-, genotoxic and pathogen stresses (Kyriakis and Avruch, 2012). Indeed, both the JNK-1 ortholog KGB-1 (Melo and Ruvkun, 2012) and the p38 ortholog PMK-1 (Kim et al., 2002; Lee and Mylonakis, 2017) were shown to monitor homeostasis and transmit signals to induce aversion in response to toxicity and infection. Further, yet unidentified signals emerge from the bloated intestine in response to infection (Singh and Aballay, 2019a), the major inner barrier and site of immunity and detoxification. Both DAF-16 and SKN-1 positive nuclei appear to be located around the gut lumen in our experiments (Figure 3). The apparent impermeability of the cuticle to chemicals (Xiong et al., 2017) suggests that signals evoked by toxic odors are also likely to emanate primarily from the intestine. Toxic stress-induced neuronal processing and integratory mechanisms are also to be identified. Potential candidates are the NPR neuropeptide receptor and serotonin signaling pathways, which were shown to be required for aversive behavior and aversive olfactory learning against pathogens, the repellent 2-nonanone, and for methyl-salicylate (Harris et al., 2019; Horspool and Chang, 2018; Luo et al., 2015; Melo and Ruvkun, 2012; Singh and Aballay, 2019a, 2019b; Zhang et al., 2005). However, other, yet, unidentified pathways might also be involved. We note that worms exposed to ccBA stayed at the edge of the food lawn but deserted at higher concentrations (see e.g. the 8% ccBA and above in Figure 1C), similarly to the observations made using the repellent 2-nonanone (Harris et al., 2019), suggesting the multisensory integration of attractive and aversive impulses for decision making.

Taken together, the elucidation of the mechanisms of cell stress-elicited behavioral decisions is an intriguing subject of future research.

Our findings that after a single stress exposure, characteristic stress-related behavioral responses are retrieved by the associated olfactory cues (Figure 6) are indicative of associative learning. Similar phenomena, where somatic/cellular stress regulates learned aversion are the compromise in vital functions by toxins or RNAi targeting essential cellular processes and the intestinal bloating caused by pathogen bacteria (Melo and Ruvkun, 2012; Singh and Aballay, 2019b). Although the prevalence of new behaviors decreases with time, repeated stress exposures not only maintain (food leaving on diluted DA after ccDA massed training) but also enhance (odor choice after ccBA massed training) the respective behavioral choices (Figure 7D, E). These findings are consistent with forgetting and intermediate term associative memory formation by reinforcement of the experience (Amano and Maruyama, 2011; Stein and Murphy, 2014). The elicitation of opposing behaviors by DA and BA olfactory cues after preconditioning demonstrates that the absence or presence of adequate cytoprotective responses at the time of stresses is a critical regulator of future behavioral decisions to anticipated stress (Figure 8). The internal experience of disrupted tissue homeostasis by ccDA in the absence of cytoprotection is integrated into an associative memory of danger, which reverses the naturally attractive valence of DA and upon retrieval gives rise to avoidant behavior. Efficient cytoprotection in response to ccBA restores homeostasis and forms a memory of protection, which upon retrieval elicits a tolerant behavior. Besides, the increased preference of DA over BA after ccBA preconditioning (Figure 6G and 7E) suggests that the natural valence of BA is integrated with the cost to maintain it, probably through the associated experience of toxic stress. Such representation allows individuals to consider whether investment of resources for self-protection are needed or not to obtain food. Thus, although the BA-associated memory of somatic resilience creates a flexible choice depending on the local context of alternative food odor, whereas the DA-associated memory of danger gives rise to a stereotypic aversion, both behavioral decisions depend on the respective neural contexts of prior experiences (Figure 8).

An evolutionarily conserved adaptive response to stress is the "fight-or-flight" response, originally coined for the neuroendocrine system (Cannon, 1939; Selye, 1973). Together with recent studies we here show the co-occurrence of behavioral "flight" responses with molecular stress and immune "fight" responses combating stress (Gecse et al., 2019; Melo and Ruvkun, 2012; Singh and Aballay, 2019a). Moreover, our studies reveal a regulatory link between intracellular cytoprotective responses and behavior suggesting a coordinated action of the fight

and the flight responses to preserve organismal integrity. Beyond ensuring physical survival,
the maintenance of cellular homeostasis by stress responses equips nematodes with behavioral
tolerance to stay in or to approach real and anticipated stressful locations. Further, temporary
avoidance in stresses that overwhelm molecular defenses allows the restoration of physiological
integrity and the strengthening of cytoprotective mechanisms. Moreover, genetically weakened
or absent molecular defenses narrows fitness by reinforcing avoidant behavior. An important
implication of our work is that even a memory of past stresses accompanied by insufficient
cellular defenses may condition to avoidance. Avoidant behaviors are characteristic to various
human mental disorders, such as phobias or panic attacks and eating disorders. These diseases
are accompanied by intense physical sensations and emotions to jettison or to avoid perceived
danger or toxic threat, which happen in response to specific or unidentified sensory cues
(Garcia, 2017; Pittig et al., 2018). The foundations of stress and detoxification responses and
learning are conserved between nematodes and humans. Thus, it might be conceivable that
unconscious memories of prior stressful somatic experiences govern emotions and behavioral
decisions in response to sensory cues.

Figure 8. Model for the regulation of learned behavioral decisions by cytoprotective responses.

(A) Toxic stress-induced disruption of cellular homeostasis (red symbols) prevails in the absence of adequate cytoprotection (left) and emits danger signals (red arrow) towards the nervous system (dark grey nerve ring in the head). Stress-specific cytoprotective responses (right) restore cellular homeostasis (green symbols) which probably suppresses/changes danger signals (green arrow).

(B) Simultaneously, signal processing in the nervous system generate an internal experience of danger (damaged shield) or stress-and-protection (intact shield). Integration with the co-occurring olfactory cues forms either a danger-based (red nerve ring) or stress-resilient (red-green nerve ring) associative memory.

(C) Memory retrieval by the respective olfactory cues evokes either a stereotypical avoidant behavior or a flexible behavioral decision depending on the external context, such as the absence or presence of another food-indicative odor beside the stress-associated olfactory cue.

Materials and Methods

Materials

The reagents benzaldehyde, diacetyl and methyl-salicylate were obtained from Sigma Aldrich. ccBA and ccDA abbreviate undiluted (concentrated) benzaldehyde and diacetyl, respectively. All other chemicals were obtained from Sigma or Fluka, if not otherwise mentioned.

C. elegans strains and maintenance

All strains used were provided by the Caenorhabditis Genetics Center: N2 (Bristol) wild type; TJ356 [daf-16p::daf-16a/b::GFP + rol-6(su1006)]; LD001 {Is007 [skn-1::gfp]};TJ375 [hsp-16.2p::GFP]; CF1038 [daf-16(mgDf50)]; CY573 [bvIs5(cyp-35B1p::GFP + gcy-7p::GFP)]; MJCU017 {kIs17[gst-4::gfp, pDP#MM016B]X}; LD1171 {Is003 [gcs-1::gfp]}; SJ4005 [hsp-4::GFP]; CF1553 {muIs84[pAD76(sod-3::GFP)]};. Strains were grown and maintained as previously described (Brenner, 1974). Animals were synchronized by allowing adults to lay eggs for 4 hours. All experiments were performed using day 1 adults, if not otherwise indicated.

Odor preconditioning and massed training

Preconditioning treatments were performed using the hanging drop method to prevent direct contact of concentrated volatiles with worms in the presence of bacterial food source to prevent the associated experience of starvation. More precisely, 1µl and 4µl drop of concentrated benzaldehyde (ccBA) or diacetyl (ccDA), respectively, was placed on the lid of 6 cm NGM plates seeded with OP50, containing a synchronous population of 200-300 young adults. The plate was sealed with parafilm to maintain a relatively constant dose of volatile. If not otherwise stated, preconditioning time was three hours. Massed training protocol was designed as described (Amano and Maruyama, 2011; Kauffman et al., 2011) employing four sequential one-hour exposures to hanging drops of 2 µl ccBA, 4µl ccDA or vehicle with intermittent washes in M9 buffer.

Acute toxicity and thermotolerance measurements

Toxicity and thermotolerance assays were carried out at 20°C, and at 35°C, respectively, by using approximately 25-40 worms per plate in three replicates in 3cm NGM plates in case of toxicity, and 6cm NGM plates in case of thermotolerance. Both toxicity and thermotolerance were measured by counting paralyzed worms using „head lifting" behavior of moveless animals (CROLL, 2009). If an apparently paralyzed worm was not able to display at least "head lifting" movements following gentle fall of assay plate into experimental surface, it was counted as „paralyzed". Paralysis index was calculated as the average of $N_{paralyzed}/N_{total}$ at each time point. Animals that crawled off the agar surface were censored.

Chemotaxis assays

Chemotaxis experiments were carried out as previously described (Bargmann et al., 1993) and carried out earlier by our lab (Gyurkó et al., 2015), with modifications. Briefly, synchronous population of young adults were washed twice in M9 buffer, then 80-100 worms were placed in the middle of a 10 cm CTX assay plate containing the odors without anesthetics in order to monitor the actual decisions at indicated time points. In kinetic chemotaxis, plates were streaked at each centimetre to measure the weighted distribution of worms at indicated time points. The Weighted Chemotaxis Index (WCI) were calculated as previously described (Nuttley et al., 2001).

Food avoidance assay

Bacterial lawn-avoidance experiments were performed as previously conceived (Milward et al., 2011), with modifications. Briefly, 50-80 synchronous day-1 adults were washed twice with

M9 buffer and dropped onto the OP50 lawn in the middle of 6 cm NGM plates. Worms were allowed to settle for 30 minutes, unless otherwise indicated. A drop of given odor were placed on a piece of parafilm in the middle of the OP50 lawn. Animals on or off the lawn were counted at each indicated time point. Worms incapable to move or crawled off the agar surface were censored. Food-leaving index was calculated as the average of N_{off} /N_{total} taken from three technical replicates.

Motility assay

Motility was characterized as described (Sawin et al., 2000) and performed earlier (Gyurkó et al., 2015) by counting body bends for 1 minute using 10-15 animals in each condition. After measuring baseline motility on an OP50-seeded NGM agar plate, a toxic dose of odor hanging drop was placed on the lid and motility was measured at the indicated time points.

RNA interference

RNAi strains were obtained from Source Bioscience (Notthingam, UK). RNAi treatments were performed as previously described (Timmons, 2006). RNAi feeding clones were grown overnight in LB medium containing 100 μg/ml ampicillin. Worms were grown on plates containing 1 mM IPTG, 50 μg/ml ampicillin and 6.25 μg/ml tetracyclin and seeded with *E. coli* HT115 strains harboring the L4440 empty vector (EV) control and specific RNAi vectors, respectively, from hatching.

Fluorescence microscopy

Analysis and quantification of fluorescence was carried out as previously described (Papp et al., 2012), with modifications. After treatments, at least 20 worms per condition were picked individually and immobilized by 20 mM NaN_3 washed in M9 buffer onto a 2% agarose pad. Microscopic examination was carried out on a NIKON Eclipse E400 type fluorescence microscope linked to a Diagnostic Instruments SPOT 500 camera in case of TJ356, TJ375, CY573, MJCU017, LD1171, SJ4005, CF1553 strains; and OLYMPUS CKX53 Fluorescence microscope, OLYMPUS DP74 Cooled color camera in case of LD001 strain, using green fluorescent filters. Images are representatives of at least three independent experiments. Fluorescence intensity measurements were quantified with ImageJ. Visualization of skn-1::GFP nuclear punctae were carried out by OLYMPUS CellSens v2.3 Imaging software.

Odor preference assay

Odor preference was carried out in standard CTX plates. 80-100 naive and preconditioned
young adults were washed off twice in M9 buffer and dropped into the middle of the assay
plate. Odors were placed into two sides of the assay plate and worms were allowed to migrate
for 50 minutes. Data are expressed as % of animals migrated into a 1 cm drawn circle around
the respective odors as well as % of animals remained out of circles.

Statistical analysis

Kaplan–Meier log-rank tests using the program IBM SPSS Statistics were carried out to
evaluate toxicity assays. Food avoidance and chemotaxis assays were examined by one-way
ANOVA with Fisher's LSD post-hoc test. Odor preference assays were analyzed by two-way
ANOVA with Fisher's LSD post-hoc test after evaluation of normal distribution significance
by Shapiro-Wilk test. Significance in fluorescence intensity were calculated by unpaired
Student's t-test following evaluation of normal distribution significance by Kolmogorov-
Smirnov test and Shapiro-Wilk test. One-way ANOVA with Fisher's LSD post-hoc tests,
Shapiro-Wilk and Kolmogorov-Smirnov tests, and unpaired Student's t-test were carried out
using IBM SPSS Statistics, while two-way ANOVA with Fisher's LSD post-hoc tests were
performed with STATISTICA. Data were expressed as mean ± standard error of the mean
(SEM) Statistical levels of significance are shown in each figure as follows: $*p < 0.05$; $**p < $
0.01; $***p < 0.001$.

References

Amano H, Maruyama IN. 2011. Aversive olfactory learning and associative long-term memory in

Caenorhabditis elegans. *Learn Mem* **18**:654–665. doi:10.1101/lm.2224411

Andersen A. 2006. Final report on the safety assessment of benzaldehyde. *Int J Toxicol* **25 Suppl**

1:11–27. doi:10.1080/10915810600716612

Aoki R, Yagami T, Sasakura H, Ogura K -i., Kajihara Y, Ibi M, Miyamae T, Nakamura F, Asakura T, Kanai

Y, Misu Y, Iino Y, Ezcurra M, Schafer WR, Mori I, Goshima Y. 2011. A Seven-Transmembrane

Receptor That Mediates Avoidance Response to Dihydrocaffeic Acid, a Water-Soluble Repellent

in Caenorhabditis elegans. *J Neurosci* **31**:16603–16610. doi:10.1523/JNEUROSCI.4018-11.2011

Ardiel EL, Rankin CH. 2010. An elegant mind: learning and memory in Caenorhabditis elegans. *Learn*

Mem **17**:191–201. doi:10.1101/lm.960510

Avery L, You Y-J. 2012. C. elegans feeding. *WormBook* 1–23. doi:10.1895/wormbook.1.150.1

Badyaev A V. 2005. Stress-induced variation in evolution: from behavioural plasticity to genetic

assimilation. *Proc R Soc B Biol Sci* **272**:877–886. doi:10.1098/rspb.2004.3045

Baldwin D V. 2013. Primitive mechanisms of trauma response: An evolutionary perspective on

trauma-related disorders. *Neurosci Biobehav Rev* **37**:1549–1566.

doi:10.1016/J.NEUBIOREV.2013.06.004

Bargmann C. 2006. Chemosensation in C. elegans. *WormBook* 1–29. doi:10.1895/wormbook.1.123.1

Bargmann CI, Hartwieg E, Horvitz HR. 1993. Odorant-selective genes and neurons mediate olfaction

in C. elegans. *Cell* **74**:515–527. doi:10.1016/0092-8674(93)80053-H

Barrett HC, Broesch J. 2012. Prepared social learning about dangerous animals in children. *Evol Hum*

Behav **33**:499–508. doi:10.1016/J.EVOLHUMBEHAV.2012.01.003

Bijlsma R, Loeschcke V. 2005. Environmental stress, adaptation and evolution: An overviewJournal of

Evolutionary Biology. pp. 744–749. doi:10.1111/j.1420-9101.2005.00962.x

Blackwell TK, Steinbaugh MJ, Hourihan JM, Ewald CY, Isik M. 2015. SKN-1/Nrf, stress responses, and

aging in Caenorhabditis elegans. *Free Radic Biol Med* **88**:290–301.

doi:10.1016/J.FREERADBIOMED.2015.06.008

Botvinick M, Braver T. 2015. Motivation and Cognitive Control: From Behavior to Neural Mechanism.

Annu Rev Psychol **66**:83–113. doi:10.1146/annurev-psych-010814-015044

Brass DM, Palmer SM. 2017. Models of toxicity of diacetyl and alternative diones. *Toxicology* **388**:15–20. doi:10.1016/j.tox.2017.02.011

Brenner S. 1974. The genetics of Caenorhabditis elegans. *Genetics* **77**:71–94. doi:10.1002/cbic.200300625

Calabrese EJ, Bachmann KA, Bailer AJ, Bolger PM, Borak J, Cai L, Cedergreen N, Cherian MG, Chiueh CC, Clarkson TW, Cook RR, Diamond DM, Doolittle DJ, Dorato MA, Duke SO, Feinendegen L, Gardner DE, Hart RW, Hastings KL, Hayes AW, Hoffmann GR, Ives JA, Jaworowski Z, Johnson TE, Jonas WB, Kaminski NE, Keller JG, Klaunig JE, Knudsen TB, Kozumbo WJ, Lettieri T, Liu S-Z, Maisseu A, Maynard KI, Masoro EJ, McClellan RO, Mehendale HM, Mothersill C, Newlin DB, Nigg HN, Oehme FW, Phalen RF, Philbert MA, Rattan SIS, Riviere JE, Rodricks J, Sapolsky RM, Scott BR, Seymour C, Sinclair DA, Smith-Sonneborn J, Snow ET, Spear L, Stevenson DE, Thomas Y, Tubiana M, Williams GM, Mattson MP. 2007. Biological stress response terminology: Integrating the concepts of adaptive response and preconditioning stress within a hormetic dose–response framework. *Toxicol Appl Pharmacol* **222**:122–128. doi:10.1016/J.TAAP.2007.02.015

Cannon WB. 1939. The wisdom of the body, 2nd ed., The wisdom of the body, 2nd ed.

Choe KP, Przybysz AJ, Strange K. 2009. The WD40 repeat protein WDR-23 functions with the CUL4/DDB1 ubiquitin ligase to regulate nuclear abundance and activity of SKN-1 in Caenorhabditis elegans. *Mol Cell Biol* **29**:2704–2715. doi:10.1128/MCB.01811-08

Clark S, Winter CK. 2015. Diacetyl in Foods: A Review of Safety and Sensory Characteristics. *Compr Rev Food Sci Food Saf* **14**:634–643. doi:10.1111/1541-4337.12150

Cosmides L, Tooby J. 2013. Evolutionary Psychology: New Perspectives on Cognition and Motivation. *Annu Rev Psychol* **64**:201–229. doi:10.1146/annurev.psych.121208.131628

CROLL NA. 2009. Components and patterns in the behaviour of the nematode Caenorhabditis elegans. *J Zool* **176**:159–176. doi:10.1111/j.1469-7998.1975.tb03191.x

Galal A, Walker L, Khan I. 2015. Induction of GST and Related Events by Dietary Phytochemicals: Sources, Chemistry, and Possible Contribution to Chemoprevention. *Curr Top Med Chem* **14**:2802–2821. doi:10.2174/1568026615666141208110721

Garcia R. 2017. Neurobiology of fear and specific phobias. *Learn Mem* **24**:462–471.

doi:10.1101/lm.044115.116

Gecse E, Gilányi B, Csaba M, Hajdú G, Sőti C. 2019. A cellular defense memory imprinted by early life
toxic stress. *Sci Rep* **9**:18935. doi:10.1038/s41598-019-55198-4

Greene T, Rogers S, Franzen A, Gentry R. 2017. A critical review of the literature to conduct a toxicity
assessment for oral exposure to methyl salicylate. *Crit Rev Toxicol* **47**:98–120.
doi:10.1080/10408444.2016.1236071

Gyurkó MD, Csermely P, Sőti C, Steták A. 2015. Distinct roles of the RasGAP family proteins in C.
elegans associative learning and memory. *Sci Rep* **5**:15084. doi:10.1038/srep15084

Harris G, Wu T, Linfield G, Choi M-K, Liu H, Zhang Y. 2019. Molecular and cellular modulators for
multisensory integration in C. elegans. *PLOS Genet* **15**:e1007706.
doi:10.1371/journal.pgen.1007706

Hobert O. 2003. Behavioral plasticity inC. elegans: Paradigms, circuits, genes. *J Neurobiol* **54**:203–
223. doi:10.1002/neu.10168

Horspool AM, Chang HC. 2018. Neuron-specific regulation of superoxide dismutase amid pathogen-
induced gut dysbiosis. *Redox Biol* **17**:377–385. doi:10.1016/j.redox.2018.05.007

Hsu A-L, Murphy CT, Kenyon C. 2003. Regulation of aging and age-related disease by DAF-16 and
heat-shock factor. *Science* **300**:1142–1145. doi:10.1126/science.1083701

Inoue H, Hisamoto N, An JH, Oliveira RP, Nishida E, Blackwell TK, Matsumoto K. 2005. The C. elegans
p38 MAPK pathway regulates nuclear localization of the transcription factor SKN-1 in oxidative
stress response. *Genes Dev* **19**:2278–2283. doi:10.1101/gad.1324805

Jorissen WP, van der Beek PAA. 2010. The oxidation of benzaldehyde. *Recl des Trav Chim des Pays-*
Bas **49**:138–141. doi:10.1002/recl.19300490207

Kauffman A, Parsons L, Stein G, Wills A, Kaletsky R, Murphy C. 2011. C. elegans positive butanone
learning, short-term, and long-term associative memory assays. *J Vis Exp* **49**. 1–10.
doi:10.3791/2490

Kenyon CJ. 2010. The genetics of ageing. *Nature* **464**:504–512. doi:10.1038/nature08980

Kim DH, Feinbaum R, Alloing G, Emerson FE, Garsin DA, Inoue H, Tanaka-Hino M, Hisamoto N,
Matsumoto K, Tan M-W, Ausubel FM. 2002. A conserved p38 MAP kinase pathway in
Caenorhabditis elegans innate immunity. *Science* **297**:623–626. doi:10.1126/science.1073759

Kluwe WM, Montgomery CA, Giles HD, Prejean JD. 1983. Encephalopathy in rats and nephropathy in

rats and mice after subchronic oral exposure to benzaldehyde. *Food Chem Toxicol* **21**:245–250.

doi:10.1016/0278-6915(83)90055-8

Kültz D. 2005. Molecular and evolutionary basis of the cellular stress response. *Annu Rev Physiol*

67:225–257. doi:10.1146/annurev.physiol.67.040403.103635

Kyriakis JM, Avruch J. 2012. Mammalian MAPK Signal Transduction Pathways Activated by Stress and

Inflammation: A 10-Year Update. *Physiol Rev* **92**:689–737. doi:10.1152/physrev.00028.2011

Lee K, Mylonakis E. 2017. An Intestine-Derived Neuropeptide Controls Avoidance Behavior in

Caenorhabditis elegans. *Cell Rep* **20**:2501–2512. doi:10.1016/j.celrep.2017.08.053

Luo J, Xu Z, Tan Z, Zhang Z, Ma L. 2015. Neuropeptide receptors NPR-1 and NPR-2 regulate

Caenorhabditis elegans avoidance response to the plant stress hormone methyl salicylate.

Genetics **199**:523–531. doi:10.1534/genetics.114.172239

McCarty R. 2016. Learning about stress: neural, endocrine and behavioral adaptations. *Stress*

19:449–475. doi:10.1080/10253890.2016.1192120

McKim JM, Schmieder PK, Niemi GJ, Carlson RW, Henry TR. 1987. Use of respiratory-cardiovascular

responses of rainbow trout (*Salmo gairdneri*) in identifying acute toxicity syndromes in fish:

Part 2. malathion, carbaryl, acrolein and benzaldehyde. *Environ Toxicol Chem* **6**:313–328.

doi:10.1002/etc.5620060408

Melo JA, Ruvkun G. 2012. Inactivation of conserved C. elegans genes engages pathogen- and

xenobiotic-associated defenses. *Cell* **149**:452–466. doi:10.1016/j.cell.2012.02.050

Milward K, Busch KE, Murphy RJ, de Bono M, Olofsson B. 2011. Neuronal and molecular substrates

for optimal foraging in Caenorhabditis elegans. *Proc Natl Acad Sci U S A* **108**:20672–20677.

doi:10.1073/pnas.1106134109

Nuttley WM, Atkinson-Leadbeater KP, Van Der Kooy D. 2002. Serotonin mediates food-odor

associative learning in the nematode Caenorhabditiselegans. *Proc Natl Acad Sci U S A*

99:12449–12454. doi:10.1073/pnas.192101699

Nuttley WM, Harbinder S, van der Kooy D. 2001. Regulation of distinct attractive and aversive

mechanisms mediating benzaldehyde chemotaxis in Caenorhabditis elegans. *Learn Mem* **8**:170–

181. doi:10.1101/lm.36501

Ogg S, Paradis S, Gottlieb S, Patterson GI, Lee L, Tissenbaum HA, Ruvkun G. 1997. The Fork head

transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in C. elegans.

Nature **389**:994–999. doi:10.1038/40194

Papp D, Csermely P, Sőti C. 2012. A role for SKN-1/Nrf in pathogen resistance and

immunosenescence in Caenorhabditis elegans. *PLoS Pathog* **8**:e1002673.

doi:10.1371/journal.ppat.1002673

Park S-K, Tedesco PM, Johnson TE. 2009. Oxidative stress and longevity in Caenorhabditis elegans as

mediated by SKN-1. *Aging Cell* **8**:258–269. doi:10.1111/j.1474-9726.2009.00473.x

Park S-W, Kaimoyo E, Kumar D, Mosher S, Klessig DF. 2007. Methyl salicylate is a critical mobile signal

for plant systemic acquired resistance. *Science* **318**:113–116. doi:10.1126/science.1147113

Pittig A, Treanor M, LeBeau RT, Craske MG. 2018. The role of associative fear and avoidance learning

in anxiety disorders: Gaps and directions for future research. *Neurosci Biobehav Rev* **88**:117–

140. doi:10.1016/j.neubiorev.2018.03.015

Puzzo D, Fiorito J, Purgatorio R, Gulisano W, Palmeri A, Arancio O, Nicholls R. 2016. Molecular

Mechanisms of Learning and Memory. *Genes, Environ Alzheimer's Dis* pp. 1–27.

doi:10.1016/B978-0-12-802851-3.00001-2

Robbins TW, Everitt BJ. 1996. Neurobehavioural mechanisms of reward and motivation. *Curr Opin*

Neurobiol **6**:228–236. doi:10.1016/S0959-4388(96)80077-8

Sankar M, Nowicka E, Carter E, Murphy DM, Knight DW, Bethell D, Hutchings GJ. 2014. The

benzaldehyde oxidation paradox explained by the interception of peroxy radical by benzyl

alcohol. *Nat Commun* **5**:3332. doi:10.1038/ncomms4332

Sasakura H, Mori I. 2013. Behavioral plasticity, learning, and memory in C. elegans. *Curr Opin*

Neurobiol **23**:92–99. doi:10.1016/j.conb.2012.09.005

Sawin ER, Ranganathan R, Horvitz HR. 2000. C. elegans Locomotory Rate Is Modulated by the

Environment through a Dopaminergic Pathway and by Experience through a Serotonergic

Pathway. *Neuron* **26**:619–631. doi:10.1016/S0896-6273(00)81199-X

Selye H. 1973. The Evolution of the Stress Concept: The originator of the concept traces its

development from the discovery in 1936 of the alarm reaction to modern therapeutic

applications of syntoxic and catatoxic hormones. *Am Sci* **61**:692–699.

Shibamoto T. 2014. Diacetyl: Occurrence, Analysis, and Toxicity. *J Agric Food Chem* **62**:4048–4053.

doi:10.1021/jf500615u

Singh J, Aballay A. 2019a. Microbial Colonization Activates an Immune Fight-and-Flight Response via

Neuroendocrine Signaling. *Dev Cell* **49**:89–99.e4. doi:10.1016/j.devcel.2019.02.001

Singh J, Aballay A. 2019b. Intestinal infection regulates behavior and learning via neuroendocrine

signaling. *Elife* **8** e50033. doi:10.7554/eLife.50033

Singh V, Aballay A. 2006. Heat-shock transcription factor (HSF)-1 pathway required for

Caenorhabditis elegans immunity. *Proc Natl Acad Sci U S A* **103**:13092–13097.

doi:10.1073/pnas.0604050103

Somogyvári M, Gecse E, Sőti C. 2018. DAF-21/Hsp90 is required for C. elegans longevity by ensuring

DAF-16/FOXO isoform A function. *Sci Rep* **8**:12048. doi:10.1038/s41598-018-30592-6

Stein GM, Murphy CT. 2014. C. elegans positive olfactory associative memory is a molecularly

conserved behavioral paradigm. *Neurobiol Learn Mem* **115**:86–94.

doi:10.1016/j.nlm.2014.07.011

Tabatabaie T, Floyd RA. 1996. Inactivation of Glutathione Peroxidase by Benzaldehyde. *Toxicol Appl*

Pharmacol **141**:389–393. doi:10.1006/TAAP.1996.0304

Timmons L. 2006. Delivery Methods for RNA Interference in <i>C. elegans</i>C. Elegans.

New Jersey: Humana Press. pp. 119–126. doi:10.1385/1-59745-151-7:119

Troemel ER, Chou JH, Dwyer ND, Colbert HA, Bargmann CI. 1995. Divergent seven transmembrane

receptors are candidate chemosensory receptors in C. elegans. *Cell* **83**:207–218.

doi:10.1016/0092-8674(95)90162-0

Tullet JMA, Hertweck M, An JH, Baker J, Hwang JY, Liu S, Oliveira RP, Baumeister R, Blackwell TK.

2008. Direct Inhibition of the Longevity-Promoting Factor SKN-1 by Insulin-like Signaling in C.

elegans. *Cell* **132**:1025–1038. doi:10.1016/j.cell.2008.01.030

Winston WM, Molodowitch C, Hunter CP. 2002. Systemic RNAi in C. elegans requires the putative

transmembrane protein SID-1. *Science* **295**:2456–2459. doi:10.1126/science.1068836

Xiong H, Pears C, Woollard A. 2017. An enhanced C. elegans based platform for toxicity assessment.

Sci Rep **7**:9839. doi:10.1038/s41598-017-10454-3

Yoshida K, Hirotsu T, Tagawa T, Oda S, Wakabayashi T, Iino Y, Ishihara T. 2012. Odour concentration-

dependent olfactory preference change in C. elegans. *Nat Commun* **3**:739.

doi:10.1038/ncomms1750

Zečić A, Braeckman BP. 2020. DAF-16/FoxO in Caenorhabditis elegans and Its Role in Metabolic

Remodeling. *Cells* **9**:109. doi:10.3390/cells9010109

Zhang Y, Lu H, Bargmann CI. 2005. Pathogenic bacteria induce aversive olfactory learning in

Caenorhabditis elegans. *Nature* **438**:179–184. doi:10.1038/nature04216

Acknowledgments

We thank the *Caenorhabditis* Genetics Center for *C. elegans* strains, Wormbase for collecting and providing data on *C. elegans*. We are grateful to Somogyvári Milán for technical and methodological help, and other members of the Stress Group for discussions. C.S. is thankful for the Merit Prize of the Semmelweis University.

Additional Information

Funding

This work was funded by grants from the Hungarian Science Foundation (OTKA K 116525), from the Semmelweis University (STIA_18_M/6800313263) and by the European Commission (GENiE, COST BM1408) to C.S.

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author Contributions

Gábor Hajdú: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Writing—original draft, Writing—review and editing

Eszter Gecse: Data curation, Investigation, Methodology

István Taisz: Data curation, Investigation, Writing—review and editing

István Móra: Data curation, Investigation

Csaba Sőti: Conceptualization, Data curation, Formal analysis, Methodology, Funding acquisition, Project administration, Writing—original draft, Writing—review and editing

Author ORCIDs

Csaba Sőti https://orcid.org/0000-0002-5975-3352

Competing Interests:

The authors declare no competing interests.

 Supplementary figures

Figure 1 – figure supplement 1. Concentrated benzaldehyde (ccBA) and diacetyl (ccDA) induce behavioral and physiological alterations characteristic to toxicity.

Undiluted, 100% BA and DA trigger initial attraction that diminishes over time as opposed to sustained attraction to diluted 1% odors (A, B). Odor sources were placed in the positive side in three drops (see Methods) opposite to three drops of ethanol vehicle. (C) Continuous exposure to ccBA and ccDA impairs thermotolerance. Error bars represent mean +/− SEM. N = number of independent experiments. *p < 0.05; **p < 0.01; ***p < 0.001. Mean durations of heat shock that induced 50% paralysis by log rank (Mantel-Cox) test were as follows: 14.46 ±

823 0.23 hours for vehicle treated control, 10.74 ± 0.42 hours for ccBA exposed (p=0.0001
824 compared to control), 12.45 ± 0.43 hours for ccDA exposed (p=0.011 compared to control).

Figure 2 - figure supplement 1. Opposing effects of odor pre-exposure on odor induced behaviors.

(A) Motility assays show a reversible vs. a sustained elevation in locomotion in response to a transient exposure to ccBA *vs.* ccDA. (B) Food aversion data showing that extended odor exposure to ccBA decreases, whereas that to ccDA further increases aversive behavior. (C) ccBA induced food avoidance as a function of duration of preconditioning exposure. (D) ccDA induced food avoidance as a function of duration of preconditioning exposure. Data are expressed as mean ± SEM. N = number of independent experiments. n.s.: not significant, *p < 0.05; **p < 0.01; ***p < 0.001.

Figure 3 - figure supplement 1. Characteristics of DAF-16 nuclear translocation and specificity of ccBA-induced cytoprotective responses.

(A) ccBA induces nuclear translocation of DAF-16::GFP comparable to that upon heat shock (HS) (quantification of fluorescence intensities of the experiment from Figure 3A). (B) Concentration dependence of ccBA-induced DAF-16::GFP nuclear translocation. (C) Representative epifluorescent microscopic images of *hsp-16.2::GFP*, *hsp-4::GFP*, *gcs-1::GFP and sod-3::GFP* stress reporters upon ccBA toxicity and heat shock (HS) as a control. Data are expressed as mean of three independent experiments by 40-60 animals evaluated per condition. n.s.: not significant, *p < 0.05; **p < 0.01; ***p < 0.001.

Figure 5 - figure supplement 1. Concentrated methyl-salicylate (ccMS) exposure impairs survival and triggers food avoidance behavior.

(A) Survival curves of nematodes exposed to various doses of ccMS using a hanging drop assay. (B) Time-dependence of ccMS induced food avoidance behavior. Error bars represent mean $\pm$ SEM, N = number of independent experiments. *$p < 0.05$, **$p<0.01$, ***$p<0.001$. Mean paralysis values by log rank (Mantel-Cox) test were as follows: 6.65 ± 0.32 hours for 4 µl ccMS, 4.72 ± 0.42 hours for 6 µl ccMS, $p=0.001$.