Inflammatory state of lymphatic vessels and miRNA profiles associated with relapse in ovarian cancer patients.

Sarah C Johnson[1], Sanjukta Chakraborty[2], Anastasios Drosou[3], Paula Cunnea[4], Dimitrios Tzovaras[3], Katherine Nixon [4], David C Zawieja [2], Mariappan Muthuchamy [2], Christina Fotopoulou[4][¶] and James E Moore Jr. [1][¶][*].

[1] Department of Bioengineering, Imperial College London

[2] College of Medicine, Texas A&M University

[3] Information Technologies Institute Centre for Research & Technology Hellas, Greece

[4] Department of Surgery and Cancer, Imperial College London.

* Corresponding author

Email. james.moore.jr@imperial.ac.uk (JM)

¶ *Christina Fotopoulou and Prof. James E Moore Jr. should be considered as joint senior authors.*

Abstract

Lymphogenic spread is associated with poor prognosis in epithelial ovarian cancer (EOC), yet little is known regarding roles of non-peri-tumoural lymphatic vessels (LVs) outside the tumour microenvironment that may impact relapse. The aim of this feasibility study was to assess whether inflammatory status of the LVs and/or changes in the miRNA profile of the LVs have potential prognostic and predictive value for overall outcome and risk of relapse. Samples of macroscopically normal human lymph LVs (n=10) were isolated from the external iliac vessels draining the pelvic region of patients undergoing debulking surgery. This was followed by quantification of the inflammatory state (low, medium and high) and presence of cancer-infiltration of each LV using immunohistochemistry. LV miRNA expression profiling was also performed, and analysed in the context of high versus low inflammation, and cancer-infiltrated versus non-cancer-infiltrated. Results were correlated with clinical outcome data including relapse with an average follow-up time of 13.3 months. The presence of a high degree of inflammation correlated significantly with patient relapse (p=0.033). Cancer-infiltrated LVs showed a moderate but non-significant association with relapse (p=0.07). Differential miRNA profiles were identified in cancer-infiltrated LVs and those with high versus low inflammation. In particular, several members of the let-7 family were consistently down-regulated in highly inflamed LVs (>1.8-fold, p<0.05) compared to the less inflamed ones. Down-regulation of the let-7 family appears to be associated with inflammation, but whether inflammation contributes to or is an effect of cancer-infiltration requires further investigation.

Introduction

Epithelial Ovarian cancer (EOC) accounts for over 4000 deaths in the UK every year [1]. Lymph node (LN) metastases are very common in EOC-patients and associated with a more dismal overall prognosis [2-3]. Total macroscopic tumour clearance with removal of bulky LN is one of the most important prognostic factors for survival post-surgery in EOC [3-4]. There is no therapeutic value in the systematic removal of clinically normal appearing LN in the advanced disease setting, as shown in prospective randomised trials [5]. The involvement and changes in the connecting lymphatic vessels (LVs) and how they are related to the future risk of metastasis and relapse have never been investigated, to our knowledge.

LVs are specialised in the uptake and transport of macromolecules from tissue fluid, consequently permitting lymphogenic spread of cancer cells. Within the tumour microenvironment, intra-tumoural LVs are present, however functional tests in mice have indicated that intra-tumoural lymphatics are non-functional [6,7]. The collapse of intra-tumoural LVs is due to some combination of the high interstitial pressure observed in multiple types of tumours, a lack of lymphatic valves, or induced mechanical pressure of growing tumour cells [6]. However, in both mice and humans, functional peri-tumoural lymphatics remain [6]. LVs can facilitate metastases both passively and through active changes in lymphatic endothelial cell (LEC) expression that aid cancer-cell infiltration, allowing subsequent drainage to LNs [8]. Tumour cells can also arrest within the draining LVs and form in-transit metastases, and in some cases escape the LVs, contributing to loco-regional metastasis, potentially aided by altered LV barrier integrity [8].

Lymphatic endothelial cells (LECs) that line the LVs can also modulate immune cell

function, in both anti-tumour immune response and cancer-induced immunosuppression [9].

For example, LEC expression of Protein-Death-Ligand 1 (PD-L1), the ligand for T cell

inhibitory receptor, PD-1, can contribute to peripheral tolerance in LNs and tumours [10].

Tumour associated LECs can scavenge and cross-present tumour-antigen to cognate CD8[+] T

cells, displaying the ability to endogenise allergen ovalbumin, then present the exogenous

antigen on Major Histocompatibility Complex (MHC) class I molecules, and contribute to

inducing tolerance of tumour-specific CD8[+] T cells [11]. LECs of the LN can acquire peptide

MHCII-from dendritic cells and induce tolerance of CD4[+] T cells via subsequent

presentation, in addition to up-regulating MHCII in response to viral-induced inflammation

or Interferon-gamma (IFN-γ) [12]. LVs are also sensitive to inflammation-induced signalling

molecules which can promote lymphangiogenesis, and induce further secretion of

inflammatory molecules that may drain to the LNs [13]. Chronic inflammation in collecting

LVs induce changes including LV dilation and inhibition of contractile ability but the effects

on metastasis are unknown [14].

MicroRNAs provide vital post-transcriptional regulation of gene expression, a function that is

often disrupted by cancer [15]. The ability of a single miRNA to inhibit multiple target genes

has led to several studies of altered tumour miRNA expression in ovarian cancer and

response to chemotherapy. This includes comparison of miRNA expression in different

tumour subsets, in healthy versus diseased ovaries, in patient versus healthy serum, benign

versus malignant tumours, expression associated with reduced patient survival, and

expression associated with drug-resistance [16]. For example, a study using both EOC cell

lines and EOC tumour tissue showed highly elevated miR-221, decreased miR-21 expression,

and down-regulation of several members of the let-7 family [17].

Expression of miRNAs in LVs and LECs is altered during inflammation and by nearby

tumours. In inflamed LECs, upregulated miR-1236 reduced expression of vascular-

endothelial growth factor-3 receptor, inhibiting lymphangiogenesis [18]. A separate study

showed that inflamed dermal LECs increased miR-155 expression, a miRNA with known

involvement in PD-L1 expression which may contribute to peripheral tolerance changes [19].

Furthermore, a panel of differentially expressed miRNAs was identified in inflamed rat LECs

and human LEC culture, indicating effects on inflammation, angiogenesis, endothelial

mesenchymal transition, cell proliferation and senescence pathways [13]. Tumour expression

profiling has identified changes in lymphangiogenesis-inducing miRNAs, such as miR-126 in

oral cancer and miR-128 in non-small cell lung cancer, that could influence metastasis

[20,21]. A study using human LEC co-culture models of gastric cancer, representing

tumoural or draining LVs, identified changes in differential miRNA expression associated

with lymphangiogenesis and with positive lymphatic metastasis [22]. Further miRNA

expression studies of draining LVs are lacking but many gene dysregulations were identified

in LECs isolated from afferent LVs that drained a metastatic gastric tumour in rats, including

up-regulation of chemokine CXCL1 [23].

The aim of this feasibility study was to provide insight into, and assess the prognostic value

of the changes in the inflammatory state and miRNA expression of the transporting LVs in

EOC. It is expected that these changes will be indicative of the immune response in the LVs

and the downstream LN, and therefore could be predictive of patient prognosis.

Methods

LV sample collection

LVs were collected from patients undergoing debulking surgery for advanced stage (FIGO IIC, III or IV) epithelial ovarian cancer at Hammersmith Hospital, Imperial College Healthcare NHS Trust, London, United Kingdom (Tables 1 and S1). Informed consent was provided from each patient. Ethics committee approval was obtained from Hammersmith and Queen Charlotte's and Chelsea Research Ethics Committee (REC reference: 05/Q0406/178), and tissue samples were provided by the Imperial College Healthcare NHS Trust Tissue Bank (ICHTB). The following LV processing steps are summarised in S1 Fig.

Macroscopically normal appearing LVs along the external iliac vessels with surrounding fibro-fatty tissue were removed with the nearby LN as part of the debulking procedure. Vessels were ligated with VIcyl03 or PDS-4-0, avoiding bipolar coagulation and other thermic energy to limit vessel damage. All patients had no macroscopic residual disease after surgery.

Table 1. Summary of the LV microscopic state and clinical data.

Vessel Sample	Type of marker			Stage	Histology	LN histology	Relapse (mos)*
	MPO	WT1	Pax-8				
1	+++	+++	++	IVB	HGS	N0	y(11)
2	+++	+++	+++	IVA	HGS	Nx	y(12)
3	+++	-	-	IIC	HGS	Nx	y(13)
4	+	++	+	IV	HGS	Nx	y(9)
5	-	-	-	IIIB	CC	Nx	n
6	-	-	-	IIIC	HGS	Nx	n
7	-	-	-	IIIC	HGS	N1	n
8	-	-	-	IIIC	HGS	Nx	n
9	+	-	-	IVB	HGS	Nx	n
10	+	+	++	IIC	HGS	N0	n

Key	MPO	WT1/Pax-8	Histology	LN histology
	+++ high	+++=15 cells	HGS=high grade serous	N0= Negative status
		++ 15> cells	CC=clear cell	
	+ medium	>=5	carcinoma	N1=Positive status
	- low	+ < 5 cells		Nx= Unassessed status

143
144 Sections of the LVs were stained with antibodies for the inflammatory marker

145 myeloperoxidase (MPO), ovarian cancer markers Wilms-Tumour-1 (WT1) and Paired Box

146 Gene 8 (Pax-8), and the lymphatic endothelial marker, Podoplanin (PDPN).

147 *Mean time since surgery = 13.2 ±3.6 months

Vessel isolation and storage

Tissue specimens were transported directly from surgical theatre to the laboratory in ice-cold

sterile Phosphate Buffered Saline (VWR, UK). LVs were identified and isolated along with a

small amount of surrounding tissue under a dissection microscope within a sterile laminar

flow cabinet (Fig 1 A-C). Each vessel was divided into two sections. One half of each vessel

was further cleaned of surrounding fatty tissue, placed in RNA stabilising solution

(RNAlater, ThermoFisher, UK), frozen on dry ice and stored at -20°C. The other half was

fixed in 4% paraformaldehyde (VWR, UK) for 45 minutes, followed by submersion in 15%

sucrose for 3 hours, then 30% sucrose for 6-12 hours. This was followed by embedding in

OCT (Scigen, UK) using dry ice and isopropyl alcohol (ThermoFisher, UK) and stored at -

80°C.

Fig 1. Lymphatic vessel isolation and classification. A. LV isolation technique. B. Tissue

sample containing LV. C. Isolated and cleaned vessel for RNA extraction. D&E,

Quantification of LVs inflammation was determined using the ratio of MPO staining to cell

nuclei (DAPI), in this example 1.0%/6.3%= 0.16, to divide samples into low, medium and

high inflammatory states with thresholds of <3%, 3-10% and >10% respectively.

MiRNA extraction and analysis

Samples in RNAlater were individually defrosted and homogenized in 700μL of QIAzol

(Qiagen, UK), using a PT 1300 D homogenisor with a 3mm dispersing aggregate head

(Kinematica, Switzerland) at 20,000 RPM in a sterile 1.5 ml tube. RNA was extracted using a

miRNeasy Mini Kit (Qiagen, UK) following the standard protocol and quality was verified

using a NanoDropTM 2000 at A260/280nm and A260/230nm absorbance (Thermofisher,

UK). Each sample was reverse-transcribed to cDNA using a miScript II RT Kit (Qiagen),

then qPCR was carried out using a miScriptSybrGreen real-time PCR kit (Qiagen) and pre-

defined miScript miRNA PCR Array (Qiagen, UK). The array contained primer pairs for 8

housekeeping genes and controls plus 88 human miRNAs that have previously been

identified or predicted to target genes involved in the regulation of inflammatory responses

and autoimmunity. One well was populated with a negative control sample. The array was

run using a StepOnePlus Real-Time PCR System (Applied Biosystems), according to the

manufacturer's instructions.

Immunofluorescence staining and imaging

Samples embedded in OCT were sectioned at 10-16μm and stored at -80°C. Both standard

Haematoxylin and Eosin (H&E) and immunofluorescent staining were subsequently

performed. For immunofluorescent staining, slides were washed with Tris Buffered Saline

(TBS) (Sigma-Aldrich, UK) and 0.05% Triton-X (Sigma) and blocked for 1hour with 10%

goat serum/TBS (Sigma-Aldrich). Slides were incubated overnight at 4°C with primary

antibodies diluted in 5% goat serum/TBS at ratios of 1:200 for inflammatory marker

Myeloperoxidase (MPO) (ab9535), 1:200 for ovarian carcinoma marker Wilms Tumour-1

(WT1) (ab89901), 1:50 for ovarian carcinoma marker Paired-Box-Gene 8 (Pax-8)

(ab189249), and 1:200 for LEC marker Podoplanin (PDPN) (ab10288). Alexa-488

conjugated secondary antibody, anti-Mouse (ab150113) or anti-Rabbit (ab150077) were

applied at a dilution of 1:250 in 5% goat serum/TBS at room temperature for 1 hour.

Samples were stained with DAPI (4',6-diamidino-2-phenylindole) solution (Sigma-Aldrich)

and mounted with Fluoroshield (Sigma-Aldrich). Confocal Z-stacks and multi-channel

images were processed and analysed using FIJI imaging software [24].

Assessment of H&E staining confirmed cryo-sections of LV samples stored at -80°C

maintained sufficient morphology to quantify inflammation and cancer cell presence. The

identification of LVs was confirmed by the positive staining with Podoplanin. The LV were

subsequently classified into a low, medium or high inflammatory group with thresholds of

<3%, 3-10% and >10% of MPO staining to nuclei (DAPI) staining ratio (Fig 1 D and E). The

level of cancer-cell infiltration was determined by the number of cancer cells detected in a

LV cross-section, as shown by positive staining with WT1 and Pax-8.

Statistical analysis

Clinical correlation between different combinations of LV state (inflammation, high-

inflammation, cancer-infiltration), and patient relapse, and cancer-stage were assessed by

applying Fishers exact test, to identify significant non-random associations, and calculation

of Cohen's Kappa coefficient of inner agreement (κ), interpreted according to the methods of

Vierra and Garett 2005 [25]. A P value below 0.05 was considered statistically significant.

Kaplan Meier estimates of probability of non-relapse were also calculated for cancer-

infiltration, inflammation and age. Prior to further analysis, the miRNA expression of each

LV (ΔCT) was normalized to the expression of a stably expressed gene in each array,

confirmed with NormFinder (http://moma.dk/normfinder-software) and GeNorm

(http://www.biogazelle.com/qbaseplus) software. SNORD96 was selected as a reference

"housekeeping" gene. The fold change between imaging-informed groups of LVs was

calculated as ΔΔCT Group2/ΔΔCT Group1 where $\Delta\Delta CT = 2^{\Delta}CT$. Fold regulation was

normalised to values greater than 1, noting up- or down-regulation. Principle component

analysis (PCA) was used to transform multivariate data linearly into a set of uncorrelated

variables to identify patterns in the variance amongst the data. We applied PCA to the

miRNA expression recorded across all our samples to calculate two principle components

(PC) that account for the greatest sources of miRNA variation within samples, using online

software ClustVis (https://biit.cs.ut.ee/clustvis/) [26]. In cases where more than 1 miRNA

was identified as significantly up-regulated or significantly down-regulated, pathway analysis

was carried out using Diana-MiRPath v3 [27]. After removing miRNA that did not show a

>1.8 fold-regulation change, a student-t-test, assuming equal variance, was applied to further

confirm variation between groups. To visualise the distributions of the individual miRNA for

which statistical significant difference was identified between groups, the probability density

functions for miRNA expression in each group were estimated using Kernel Density

Estimated distributions.

A number of widely used automatic classification methods were then used to predict the

binary outcomes of the individuals (relapse vs. non-relapse, cancer infiltration vs. no cancer

infiltration, medium or inflammation versus low inflammation), based on the LV miRNA

expression of the individuals as predictor (input) variables. We used the supervised

classification methods Logistic Regression, K-Nearest Neighbours, Support Vector Machine,

Random Forest, and Gaussian Naïve Bayes. Evaluation of each method was performed using

a leave-one-out cross validation procedure: each patient was considered in turn as a new

patient (i.e. the test patient), while the remainder were considered as the training set. The

predicted and actual outcome for each test patient was then compared with method accuracy

measured as the percentage of correct predictions. Two other measures of accuracy were also

used: the Cohen's Kappa coefficient (κ) of inner agreement between predicted and actual

values, and the McNemar's test measuring the significance of equality of predicted

probability (inner accuracy) between groups for each outcome. In each method, we also split

the cases according to the binary outcome variable and measured the percentage of correct

predictions in each group. A set of only the miRNA that were significantly differentially

expressed between groups was then used to train new classifiers. We then assessed whether

use of the new classifiers could improve predictions when compared to the performance of

classifiers built using the entire miRNA expression set.

Data availability

Further analytical results are available in the supplementary material. All raw and normalised

expression data files have been deposited within Imperial College London's Research Data

Repository at http://doi.org/10.14469/hpc/7192.

Results

Post-operative assessment of patients

Ten patients were included in the present analysis. Nine of those patients had a high grade

serous histology, and one patient had a clear cell carcinoma (Table 1). 60% of the lymph

vessels were harvested at primary debulking surgery and 40% at interval debulking surgery

(S1 Table). Only one positive para-aortic LN was detected at final histopathology. In the

13.2 ± 3.6 months follow up period, 4 of the 10 patients had relapsed. Of the four patients that

relapsed, three vessels demonstrated high inflammation and one vessel showed medium

inflammation. Vessels from patients that did not relapse showed low inflammation in 4 cases

and medium inflammation in the remaining 2 cases. Of the four relapse cases, cancer-cell

infiltration was detected in three vessels (two with high inflammation and one showing

medium inflammation). The overall relative range of inflammation in the lymphatic vessels

was small with high inflammation defined as an average of >10% MPO:DAPI ratio and a

narrow range of an average of 3-10% MPO:DAPI ratio for medium inflammation. All vessels

showed less inflammation than similarly stained ovarian tumour tissue (Fig 2I).

Consequently, to compare LVs with a notable difference in inflammation we focused initially

on comparison of the LVs with high inflammation versus low inflammation.

Fig 2. Lymphatic vessel imaging. A. H & E stained sections of the LVs, show that the

nuclei were sufficiently preserved for subsequent Paired-Box-Gene 8 (Pax-8) and WT1

(Wilms-Tumour-1) staining. B-D. LVs with high, medium and low/no inflammatory state

determined with myeloperoxidase (MPO) staining. E. Confirmation of LV status was carried

out using LEC marker Podoplanin (PDPN) that stained around the LV lumen. F-H. Imaging

with WT1, (F) negative LV, (G) positive LV (H) positive control. I. Tumour MPO positive

control. J-L. Imaging with Pax-8, (J) negative LV, (K) positive LV and (L) positive control.

An Ovarian tumour sample was used as a positive control and stained for both (M) WT1 and

(Q) Pax-8. Scale=140μm.

Clinical correlation of LV inflammatory state or cancer-cell

infiltration and patient relapse

Of the four cancer-infiltrated LVs identified, 2 presented with high inflammation and the

remaining 2 samples presented with medium inflammation. Cancer-free LVs presented with

generally lower degrees of inflammation (Table 1). Tumour cells were detected in 50% of

LVs with medium inflammation and two-thirds of LVs displaying high inflammation. The

LVs presented with high inflammation in 3 of 10 samples, two of which were also positive

for cancer cells**.** Three LV samples presented with medium inflammation, two of which

contained cancer cells, and the remaining 4 LV samples presented with low inflammation and

no detectable cancer cell infiltration (Table 1). The samples were macroscopically normal in

appearance, however positive staining for the ovarian tumour markers WT1 and Pax8 was

observed in four of the 10 samples collected (Fig 2 G,K). The LVs with cancer cells showed

a moderate agreement (κ=0.583) with patient relapse but this was not significant (Table 2)

(p=0.114). A moderate but statistically non-significant agreement was found between cancer-

infiltrated LVs and Stage IV and inflammation (κ =0.583, 0.61 respectively).

Table 2: Cohen's Kappa Coefficient for inner-agreement between different sample

states (1=perfect agreement) and Fisher's exact test for identification of significant non-

random association (p<0.05) (n=10).

Cohen's Kappa Co-efficient					
	Relapse*	Inflam.	High Inflam.	Cancer-cell inf.	Stage IV
Relapse	x	0.615	**0.783**	0.583	0.583
Inflam.	0.615	x	x	0.615	0.615
High Inflam.	**0.783**	x	x	0.348	0.348
Cancer-cell inf.	0.583	0.615	0.348	x	0.583
Stage IV	0.583	0.615	0.348	0.583	x
Fishers Exact Test (p-value)					
	Relapse*	Inflam.	High Inflam.	Cancer-cell inf.	Stage IV
Relapse	x	0.071	**0.033**	0.114	0.114
Inflam.	0.071	x	x	0.071	0.071
High Inflam.	**0.033**	x	x	0.3	0.3
Cancer-cell inf.	0.114	0.071	0.3	x	0.114
Stage IV	0.114	0.071	0.3	0.114	x

Inflam = comparison of LVs with high or medium inflammation to low inflammation. High

Inflam. = comparison of LVs with high inflammation to low or medium inflammation.

Cancer-cell inf. = cancer-cell infiltration.

Mean time since surgery= 13.2 months ±3.6

As the degree of inflammation in the LVs increased, so did correlation with patient relapse.

The presence of LVs showing high inflammation significantly correlated with patient relapse

(κ= 0.783, p=0.03) (Table 3). The presence of medium or high inflammation within LVs

showed a moderate agreement with patient relapse (κ= 0.615, p=0.07) and patients with Stage

IV, relapse and cancer-cell infiltration (κ=0.615, p=0.07) (Table 2). The presence of highly

inflamed LVs compared to the presence of low inflammation only, resulted in a stronger

correlation involving high inflammation status (Table 3). LVs exhibiting high degrees of

inflammation showed near perfect agreement with patient relapse compared to those with low

inflammatory state (n=7, κ=1.0, p = 0.029). -Additionally, there were indications of a

relationship between highly inflamed vessels with cancer-infiltration and Stage IV cancer

(κ=0.695), but this lacked statistical significance (p=0.143). These results therefore support

the earlier identified correlation between high inflammation and relapse. Kaplan Meier

estimates of non-relapse according to inflammatory state, cancer-infiltration and age

thresholds (>55, >60 , >70, >75 years) indicate that probability of non-relapse is decreased in

patients with LVs displaying medium or low inflammation and patients with cancer-

infiltrated LVs (S2 Fig). Probability of non-relapse showed that patients >55 years age were

more likely to relapse (S3 Fig).

Table 3: Comparison of LVs with high inflammation vs those with low inflammation

only (n=7) using Cohen's Kappa Coefficient and Fisher's exact test, as described in

Table 2.

	Relapse*	Cancer	Stage IV
Cohen's Kappa Co-efficient	**1**	0.695	0.695
Fishers Exact Test	**0.029**	0.143	0.143

331 *Mean time since surgery= 13.2 months ±3.6*

Lymphatic vessel miRNA expression during EOC

Expression analysis of a panel of miRNA involved in auto-immunity and inflammation was

carried out. This showed that let-7 was differentially expressed in the highly-inflamed LVs

(Table 4, S2 Table). Significant differential expression of 8 miRNA was identified when

comparing miRNA expression of LVs presenting with high inflammation versus low

inflammation (p<0.05, n=7). Of the 8 miRNA, 6 belonged to the let-7 family with miR-23a-

3p and miR-23b-3p completing the set. Differential expression can also be clearly visualised

when the estimated probability density functions, calculated using kernel density estimation

for inner groups, for the differentially expressed individual miRNA are compared (S4 Fig).

Correction for multiple testing by applying a post-hoc Bonferroni correction showed that

differential expression of three miRNA, let-7b, let-7c and let-7d remained significant

(p<0.0017) . Comparison of miRNA expression in LVs displaying low inflammation versus

medium or high inflammation showed a similar trend towards down-regulation of the let-7

family (Table 4, S3 Table). Six miRNA were down-regulated >1.8 fold with significant

down-regulation of let-7g-5p and let-7i-5p (p<0.05).

Table 4: Differential expression of miRNA between LV groups displaying low, medium

or high inflammation.

Low versus High			Low versus [Medium or High]		
miRNA	Fold regulation	t-test (p)	miRNA	Fold regulation	t-test (p)
let-7a-5p	**-188.20**	**0.0389**	**let-7g-5p**	**-2.39**	**0.0308**
let-7b-5p*	**-11.42**	**0.0006**	**let-7i-5p**	**-2.02**	**0.0130**
let-7c-5p*	**-6.43**	**0.0003**	miR-545-3p	-2.77	0.0739
let-7d-5p*	**-5.06**	**0.0009**	miR-93-5p	1.77	0.0856
let-7e-5p	-2.67	0.0802			
let-7f-5p	**-3.20**	**0.0378**			
let-7g-5p	-2.99	0.0969			
let-7i-5p	**-2.23**	**0.0328**			
miR-15b-5p	-5.24	0.0696			
miR-23a-3p	**-3.44**	**0.0401**			
miR-23b-3p	**-15.17**	**0.0152**			
miR-98-5p	-4.63	0.0823			

Listed are miRNA that showed a fold-change ±1.8, with subsequent t-test with p<0.1, with

miRNA displaying a significant difference of p<0.05 highlighted in bold.

* = Bonferroni correction p<0.0017

Similar trends were observed in the differential miRNA expression identified in the inflamed

LVs and the differential miRNA expression in LVs collected from patients who underwent

relapse during the study. Four significantly differentially expressed miRNA (p<0.05) were

identified in relapse LVs (let-7c-5p, let-7b-5p, miR-23b-3p and miR-186-5p), with three of

these (let-7c-5p, let-7b-5p, miR-23b-3p), also significantly differentially expressed in highly

inflamed LVs (Table 5). Differential expression of miRNA with correction criteria of

p<0.002 was not identified. Of the 17 miRNA that were down-regulated >1.8 fold in relapse

vessels, 8 were also down-regulated in inflamed LVs, and 11 in the highly-inflamed LVs,

including 5 and 6 members of the let-7 family respectively and miR-23b-3p (S2, S3 and S5

Table).

Table 5: Differential expression of miRNA between cancer-infiltrated versus non-cancer

infiltrated LVs and LVs from relapse versus non-relapse patients within 13.2±6 months.

Cancer-infiltrated LVs			LVs from patients that relapsed		
miRNA	Fold regulation	t-test (p)	miRNA	Fold regulation	t-test (p)
miR-144-3p*	**14.02**	**0.0010**	miR-144-3p	5.07	0.0906
miR-181c-5p	**10.63**	**0.0170**	**miR-186-5p**	**4.93**	**0.0294**
miR-101-3p	5.08	0.0740	miR-497-5p	2.29	0.0672
miR-381-3p	**4.07**	**0.0140**	miR-34a-5p	2.09	0.0828
miR-497-5p	**2.9**	**0.0260**	miR-98-5p	-3.43	0.0673
miR-93-5p*	**2.25**	**0.0010**	let-7d-5p	-3.68	0.0645
miR-16-5p	**2.08**	**0.0310**	**let-7c-5p**	**-4.33**	**0.0385**
let-7i-5p	**-2.08**	**0.0210**	**let-7b-5p**	**-6.94**	**0.0383**
let-7b-5p	-2.27	0.0700	**miR-23b-3p**	**-8.53**	**0.0278**
let-7c-5p	-2.56	0.0780			

Listed are miRNA that showed a fold-change ±1.8, with subsequent t-test with p<0.1, with

miRNA displaying a significant difference of p<0.05 highlighted in bold.

* = Bonferroni correction p<0.002

In the cancer-infiltrated LVs, most miRNA expression changes involved miRNA up-

regulation such as miR-16-5p, miR-93-5p and miR-497-5p, miR-381-3p, miR-144-3p and

miR-181a-5p (p<0.05) (Table 5). Of these, up-regulation of miR-93-5p and miR-144-3p

remained significant with application of a Bonferroni correction (p<0.002). In contrast,

inflamed LVs presented with primarily down-regulated miRNA, in particular, significant

down-regulation of members of the let-7 family along with miR-23a-3p, miR-23b-3p and

tumour suppressor miR-15-5p [28]. In the cancer-containing LVs, only 7 miRNA showed a

>1.8 fold down-regulation, 6 of which were members of the let-7 family (S4 Table). Of these

six, five were also down-regulated >1.8-fold in the LVs of patients who relapsed during the

study period.

Similar trends in the differential miRNA expression in cancer-infiltrated LVs and highly

inflamed LVs were also identified, as 15 miRNA were found to undergo an expression

change of >1.8 fold (up- or down-regulated) in both of the comparisons (S2 and S4 Table).

Six of the 7 miRNA that were down-regulated in both groups belonged to the let-7 family.

This common trend in the fold-regulation difference in the highly inflamed and cancer-

infiltrated LVs may be reflective of the fact that 50% of the cancer-infiltrated LVs also

presented with high inflammation. We also noted that the number of differentially expressed

miRNA was higher when comparing LVs presenting with high or low inflammation than

when comparing vessels with or without cancer-infiltration. Expression of several miRNA

were consistently below qPCR detection levels and thus omitted from further analysis.

Pathway analysis of miRNA found to be dysregulated in inflamed

or cancer-infiltrated LVs

Pathway analysis indicated that significantly dysregulated miRNAs are involved in

maintaining LV integrity. The 7 miRNAs that were significantly upregulated in cancer-

infiltrated LVs, and the 7 miRNAs that were significantly down-regulated in inflamed LVs

have known involvement in various pathways related to the TGFβ-pathway, fatty acid

biosynthesis, proteoglycans in cancer, glycosaminoglycan synthesis and glycan biosynthesis

as well as the extracellular matrix receptor interactions pathway and adherens junction

pathway (Table 6). No significantly up-regulated miRNAs were identified in highly inflamed

LVs and only one significantly down-regulated miRNA was identified in cancer-infiltrated

LVs. Pathway analysis carried out with the 4 significantly dysregulated miRNAs in the

relapse LVs showed a range of potential target pathways including adherens junction, ECM-

receptor interactions and mucin type O-Glycan biosynthesis (S6 Table).

Table 6: Pathway analysis using MiRNA that were significantly up-regulated in cancer-

infiltrated LVs (left-hand side) and miRNA significantly down-regulated >1.8 fold

(right-hand side) in highly inflamed LVs.

Upregulated miRNA in cancer-infiltrated LVs				Down-regulated miRNA in highly inflamed LVs			
KEGG pathway	**p-value**	**Genes**	**miRNA**	**KEGG pathway**	**p-value**	**Genes**	**miRNA**
TARBASE				TARBASE			
Proteoglycans in cancer	3.02E-15	96	5	Adherens junction	5.86E-13	52	8
Adherens junction	3.09E-09	44	5	Proteoglycans in cancer	5.86E-13	107	8
Prion diseases	3.09E-09	14	5	TGF-β signaling pathway	3.29E-09	49	8
Viral carcinogenesis	1.64E-08	91	5	Viral carcinogenesis	5.77E-09	99	8
Hippo-signaling pathway	2.37E-08	68	5	Cell cycle	2.79E-08	74	8
TGF-β signaling pathway	9.56E-08	41	5	Hippo signaling pathway	1.24E-07	74	8
TARGETSCAN				TARGETSCAN			

Fatty acid biosynthesis	7.90E-38	2	2	Fatty acid biosynthesis	2.92E-53	3	1
Fatty acid metabolism	8.15E-15	3	2	Fatty acid metabolism	1.03E-21	4	1
Metabolism of xenobiotics by cytochrome P450	7.48E-06	2	2	Signaling path-ways regulating pluri-potency of stem cells	2.58E-09	17	7
Signaling pathways regulating pluri-potency of stem cells	6.26E-05	11	3	Metabolism of xenobiotics by cytochrome P450	8.99E-07	3	3
TGF-β signaling pathway	0.0387	5	3	N-Glycan biosynthesis	0.00466	5	3
Micro-CT-DS				Micro-CT-DS			
Fatty acid biosynthesis	7.90E-38	5	2	Fatty acid biosynthesis	7.39E-26	6	9
Fatty acid metabolism	8.15E-15	11	2	ECM-receptor interaction	3.21E-07	17	9
Signaling pathways regulating pluri-potency of stem cells	3.74E-12	47	3	Signaling pathways regulating pluri-potency of stem cells	1.01E-06	43	9
Hippo signaling pathway	2.06E-06	29	2	Mucin type O-Glycan biosynthesis	5.38E-06	9	9
Proteoglycans in cancer	2.70E-06	59	4	Proteoglycans in cancer	2.34E-05	53	9
				p53 signaling pathway	3.13E-05	27	9

Principle component analysis of miRNA expression in LV samples.

PCA showed that differential miRNA expression identified in inflamed and cancer-infiltrated LVs could also be identified without grouping the samples. This can be seen when assessing the top 16 miRNA whose expression contributed the most to the first and second Principle Components (PC1 and PC2). These components capture factors responsible for the most and second most variation in the data set, respectively. Samples with the most similar miRNA expression profile cluster together and clusters may vary over PC1/PC2 or both (S5 Fig). Twelve miRNA identified in the fold-change analysis of highly inflamed or cancer-infiltrated LVs, were among the top 20 miRNA that contributed to PC1, with 5 of those showing significant differential expression. Furthermore, 8 miRNAs of the top 20 miRNA that contributed to PC2 were all identified as significantly downregulated >1.8 fold in highly

426 inflamed LVs. The remaining miRNA varied greatly between the patient samples, e.g., miR-

427 20, but were not identified by analysis of fold change between groups. The expression levels

428 of these miRNA were therefore spread equally between the compared groups. Overall,

429 variation over PC2 differentiates between highly-inflamed and non-inflamed LVs, supporting

430 the suggestion that miRNA expression could distinguish LV states (S5 Fig).

Application of machine-learning algorithms to predict LV state or clinical outcome based on miRNA expression

Patterns of miRNA expression showed some potential in predicting clinical outcomes by

applying classification algorithms. When predicting patient relapse (Table 7), all of the

classification methods presented with quite high accuracy values and McNemar's

probabilities of $p < 0.05$, meaning that no significant difference was found between the

predicted grouping, based on miRNA expression, and the grouping based on the known

clinical data. Of the methods applied, use of k-Nearest Neighbours and logistic regression

produced the highest classification accuracy with similar prediction accuracy achieved

between methods. Classification of cancer-cell infiltrated LVs based on miRNA expression

(Table 8) showed a similar prediction accuracy compared to classification of relapse LVs, but

the Random Forest classifier achieved the best accuracy. Classification of inflamed and

<=Stage IV samples did not show a high accuracy (S7 Table). As a further evaluation, we

examined the effect of keeping as predictors only those miRNAs that have significant

associations with the outcome variables. Analysis was performed on classifiers build using

only significantly differentially expressed miRNAs ($p < 0.05$). In the cancer-infiltrated LVs,

the differentially expressed miRNAs enhanced the accuracy of the predictive algorithm to 80-

100% (S8 Table). Prediction of relapse based on miR-23b-3p and miR-186-5p expression,

improved the accuracy of classification with all methods by 10%, except when using the

Support Vector Machine, where no improvement was observed (see S8 Table & S1 File). The

addition of miR-let-7b-5p and let-7c-5p improved the accuracy of the Support Vector

Machine from 80% to 90% but decreased the accuracy of Logistic Regression from 80% to

70%. However, the use of miR-23b and miR-186-5p only showed a bias towards classifying

the samples as non-relapsed when using Logistic Regression, which decreased when using all

4 miRNA. These results suggest that the miRNAs with significant associations to the

outcome variables have a capacity to be used as predictors for predicting the outcomes of

interest in new patients.

Table 7: Classification analysis performance for various supervised classification

algorithms for the prediction of patient relapse based on LV miRNA expression. This

illustrates the accuracy of the method (percentage of correct predictions), the per-group

accuracy (No Relapse, Yes Relapse), Cohen's Kappa of inner agreement between predicted

and actual outcome, and McNemar's test of significance of equality of predicted probability

(inner accuracy) between groups for each outcome.

Method	Accuracy	McNemar's Test-pval	Cohen's Kappa	No Relapse	Yes Relapse
Logistic Regression	70%	0.999	0.375	67 %	75 %
K-Nearest Neighbours	80%	0.289	0.583	100 %	50 %
Support Vector Machines	80%	0.289	0.583	100 %	50 %
Random Forests Classifier	50%	0.375	-0.042	67 %	25 %
Gaussian Naive Bayes	60%	0.031	0.167	100 %	0 %

471 **Table 8: Classification performance of various supervised classification algorithms for**

472 **the prediction of cancer infiltration based on LV miRNA expression.** Details are similar

473 to Table 7.

Method	Accuracy	McNemar's Test-pval	Cohen's Kappa	No Cancer	Yes Cancer
Logistic Regression	70%	0.44	0.38	83 %	50 %
K-Nearest Neighbours	50%	0.37	-0.04	67 %	25 %
Support Vector Machines	70%	0.13	0.37	100 %	25 %
Random Forests Classifier	100%	1.00	1.00	100 %	100 %
Gaussian Naive Bayes	50%	0.37	-0.04	67 %	25 %

Discussion

To our knowledge, this is the first analysis of inflammatory and miRNA alterations in tumour

draining LVs in ovarian cancer. In this feasibility study, we aimed to provide insights into LV

features that might be associated with patient relapse. Our results may contribute to a better

prediction of patient relapse, and could guide future experimental work seeking to improve

patient outcome via therapeutic targeting of miRNA.

Macroscopically normal LVs, remote from the tumour site that exhibited high or medium

inflammation correlated with cancer relapse. Despite cancer-cell infiltration in 66% of these

vessels, a moderate but non-significant association was identified between inflamed and

cancer-infiltrated LVs (p=0.07). The staging of LNs is routinely used to quantify cancer

progression but the status of the connecting LVs has not previously been investigated. While

case numbers are small, the presence of high inflammation in the LVs showed the strongest

significant correlation with cancer relapse (p=0.03 **κ**=0.783). The Kaplan Meier estimates of

non-relapse also showed that relapse became less likely without the presence of LV

inflammation or cancer-cell infiltration, and age did not increase likelihood of relapse (S Fig

2 and 3). Whether inflammation is a consequence of factors that contribute to relapse, or the

inflammation-induced changes directly affect relapse, is an area for future work.

Analysis of the effect of cancer cell infiltration of the LVs, without considering LV

inflammatory state, identified a trend towards a non-significant association with relapse

($p=0.07$, $\kappa=0.783$). This suggests that cancer-infiltration of LVs alone may not necessarily

predict relapse, but rather accompanying changes in vessel inflammation may increase the

likelihood of relapse. Furthermore, cancer-cell infiltration did not show a significant

association with high inflammation. The lack of a significant correlation in both cases may

also be simply due to the possible scenario that cancer cells are just transiting through these

vessels, so detecting their presence becomes a matter of timing. Overall, this study suggests

that nearby LVs appear to play an important role in the immune response to cancer and the

association between inflammatory lymphatic changes and relapse deserves further

exploration while potentially representing a therapeutic target.

The use of miRNA expression classification algorithms also showed potential to predict

relapse, with relatively high accuracy for these 10 patients when applying a k-Nearest

Neighbours algorithm or a Support Vector Machine model (80%), rising to 90% when using

significantly differentially expressed miRNA only (Table 7, S8). Further studies spanning a

larger number of patients should therefore be performed. Clinically, the prediction of a

patient's outcome based on miRNA expression of tissue collected during surgery could be of

considerable value. Despite the presence of inflammation showing a moderate agreement

with relapse and high inflammation displaying a significant correlation with relapse, a

classifier based on miRNA expression in high or medium inflamed LVs did not show a high

accuracy (Tables 2 and 3, S7 and S8 Table). Furthermore, due to the limited number of

highly inflamed LVs, it was not possible to build a reliable classifier based on miRNA

expression in LVs with high inflammation versus miRNA expression in LVs with low or

medium inflammation. Indeed, all the classifiers are limited by the number of samples

required to train the predictive algorithms. Therefore, whilst providing insight, these

observations require increased sample input to maintain confidence in miRNA classification.

Analysis of miRNA expression of highly inflamed LVs compared to LVs with no or low

inflammation, showed down-regulation of 6 members of the let-7 family, 3 of which

remained significant with a Bonferroni correction (p<0.0017). A similar trend was observed

when comparing miRNA expression of LVs presenting with high or medium inflammation to

those with low inflammation. In this case 6 members of the let-7 family showed down-

regulation >1.8 fold but only two members showed significant down-regulation (p<0.05),

which may be reflective of less difference in mean LV inflammatory state between the two

groups compared. The change in let-7 miRNA expression was large enough that without

designating groups, the overall variability in sample miRNA expression due to let-7

expression could be detected using PCA (S5 Fig). In ovarian tumour tissue, let-7 expression

is down-regulated and associated with poor prognosis and chemo-resistance [17,29,30]. Let-7

family members target oncogenes HMAG2 and RAS, and loss of let-7 is also associated with

the development of aggressive cancers [31]. Chemo-sensitivity of tumours may also be

affected by let-7 expression. Manipulation of let-7e in EOC cells *in-vitro* and *in-vivo*

confirmed that expression is reduced in cisplatin-resistant cell lines and application of let-7e

mimics alongside cisplatin, reduced tumour growth in mice more than with cisplatin alone

[32]. Furthermore, chimeric-mediated delivery of let-7i miRNA to EOC cells *in-vitro*

reversed paclitaxel-induced chemo-resistance [33]. Expression of let-7 is also associated with

control of the Cdk-dependent cell cycle network, that drives the cell transformation network,

allowing cells to undergo an epigenetic switch between non-transformed and a malignant

transformed state [34].

The implications of let-7 down-regulation in LVs include changes in immune response and

LV permeability. Members of the let-7 family can target inflammatory cytokines, such as IL-

6 by let-7a and IL-13 by let-7f, shown to contribute to cancer-cell apoptosis and anti-

inflammatory action respectively [35,36]. In addition to affecting immune cells in transit,

drainage of cytokines to LNs could impact immune response at the LN, potentially

contributing to the loss of immuno-suppression observed in draining LNs [37]. Cytokines

may also affect LV transport, as IL-6 can increase LV permeability while intradermal

administration of cytokines IL-1β, TNF-α, and IL-6 have been shown to decrease lymphatic

contraction frequency [38,39]. Down-regulation of let-7 in vascular endothelial cells is

associated with increased TGF-β signalling leading to promotion of epithelial-to-

mesenchymal transition and thus increased vessel permeability, but the effects in LVs are

currently unknown [40]. The observed up-regulation of several miRNA (miR-381-3p, miR-

93-5p, miR-16-5p and miR-497-5p) in cancer-infiltrated LVs may inhibit Vascular

Endothelial Growth Factor (VEGF) expression or VEGF receptors, both directly and

indirectly via Bcl-2 inhibition [41-44]. The inhibition of VEGF secretion in tumours has been

a target of several drug trials but manipulation of VEGF secretion in connecting LVs may

present as a future therapeutic area [45]. Expression of miR-144 has been shown to target

RUNX-1 in EOC lines and SRF in HUVECs, both contributing to anti-proliferative action. A

more extensive list of potential targets is provided in Table 9. A logical progression of this

study would be to confirm targets in LECs.

Table 9: Confirmed targets of miRNA in primarily endothelial and cancer cell lines

identified as significantly differentially regulated between groups.

miRNA	Target	Cell type	Pathway/Function	Ref.
let7 family	RAS	EOC lines	tumour/Ras/MAPK-suppressor	[46]
	HMAG2	EOC lines	anti-apoptotic, tumour-suppressor	[47]
	indirect	HUAECs	inhibits TGFβ pathway, pro-EMT	[40]
let7a-5p	IL6	Epithelial cells	pro-cancer-cell survival	[35]
	TGFBR3	HUVECs	anti-angiogenenic, anti-tube formation	[48]
	LOX-1	HUVECs	anti-apoptotic, pro NO synthesis	[49]
let7b-5p	LOX-1	HUVECs	anti-apoptotic, pro NO synthesis	[49]
let7c-5p	Bcl-xl	HUVECs	pro-apoptotic, via ox-LDL induced apoptosis	[50]
let7d-5p	IFI44L	HUVECs	anti-proliferation/migration	[51]
let7e-5p	Ilββ	HUVECs	pro-Nfκ βpathway	[52]
let7f-5p	IL10	CD4+ T cells	pro-inflammatory	[53]
	IL3	Lymphocytes	anti-inflammatory	[36]
let7g-5p	LOX-1	VSCMs	anti-apoptotic	[54]
	TGFBR3/ SMAD2 /THBS	HUVECs	pro-angiogenic, targets TFβ pathway	[55]
	IL10	TH1/TH17 T cells	pro-inflammatory	[56,57]
miR-93-5p	EPLIN	HUVECS	pro-proliferative/migration/angiogenesis	[58]
	PTEN	OC lines	anti-apoptotic miR93-5p/PTEN/pAkt	[59]
	RhoC	EOC lines	pro-apoptotic, anti-migration	[60]
	IL8	Neuroblastoma	anti-angiogenic	[42]
	VEGF	Neuroblastoma	anti-angiogenic	[42]
miR-144-3p	Meis1	Embryotic (Zebrafish)	decreases runx-1,c-myc, haematoepeoeisis	[61]
	SRF	HUVECs	anti-proliferative, pro-apoptotic	[62]
	RUNX-1	EOC lines	anti-proliferation, anti-migration	[63]
miR-23a-3p	ZO-2 & JAM-C	HUVECs	inhibit permeability	[64]
	RUNX2	HUVECs	anti-angiogenic, suppresses VEGF-A	[65]
miR-23b-3p	JAM-C & ZO-2	HUVECs	pro-angiogenic, increase permeability	[64]
	TAB2,TAB3, IKKa	Lymphocytes	suppresses IL-17-associated inflammation	[66]

EOC=Epithelial ovarian cancer

EMT=Epithelial-to-Mesenchymal Transition

HUVEC=Human umbilical vascular endothelial cells

HUAEC=Human umbilical arterial endothelial cells

Further studies of the relationship between inflammatory markers in LVs and differential

miRNA expression would increase the potential for diagnostic capabilities. Jones et al. and

Yee et al, analyzed LEC responses to inflammation and identified a total of 16 differentially

expressed miRNA, one of which (miR-17-5p) overlapped with the panel of miRNAs

screened in our study [18,19]. Chakraborty et al, identified differentially expressed miRNA

associated with autoimmunity and inflammation in rat LECs, with different sets of miRNAs

identified at 2 hours, 48 hours and 96 hours after induction of inflammation [13]. Our results

showed no overlap with the miRNAs regulated at the 2-hour time-point but did show similar

upregulation of miR-19a detected at 48 hours and induction of miR-497, miR-19b and miR-

19a observed at 96 hours [13]. Additionally, there was no evidence of dysregulation of the

let-7 family in the rat LECs. Taken together, chronic inflammation of LVs in EOC patients

may impact miRNA expression differently, aligning more closely with the later time-points

post-inflammatory induction in the controlled rat experiments.

The pathway analysis of the dysregulated miRNA identified involvement in transforming-

growth factor TGF-β2 signalling, as well as glycoproteins in cancer and molecules that

contribute to the glycocalyx of the LV lumen (Table 6). LECs possess a glycocalyx

composed of glycoprotein side chains, a backbone that includes adhesion molecules, and a

proteoglycan core. Therefore, changes to glycoproteins or adherens junctions may affect

integrity of the LV wall [67]. Furthermore, a study by Zolla *et al* showed aged collecting LVs

in rats presented with a thinner endothelial cell glycocalyx and a loss of gap junction proteins

[68]. Functionally this resulted in hyper-permeability of the LVs that allowed pathogens to

escape into surrounding tissue. This may suggest that the miRNA changes that accompany

cancer cell infiltration and inflammation may also alter LV permeability and barrier function.

This in turn could aid cancer cell infiltration but may also impact similar movements of

immune cells such as macrophages and dendritic cells [69]. Further studies to assess the

impact of inflammation-induced miRNA expression changes on LV integrity may identify

new avenues to prevent transmigration of cancer cells across LV walls.

The response to cancer may also be mediated by inflammatory cells normally present in the

LV. Analysis of rat mesenteric, pre-nodal and post-nodal LVs has shown the presence of

innate inflammatory immune cells accumulating on the walls of the LVs including

neutrophils, monocytes and macrophages [69]. Microarray analysis of mesenteric collecting

vessels has also shown significant accumulation of MHCII+ antigen-presenting cells [70].

Furthermore, both antigen-presenting cells and T cells are transported in LVs, with

proportions depending on the status of the upstream tissues [71]. We have not assessed the

status of all of the tissues drained by these LVs, but in any case we would expect some

immune cells to be captured in our images. However, the overall proportion of T cells to LV

wall structural cells should be low enough not to impact overall miRNA expression.

The expression of the enzyme myeloperoxidase (MPO) was selected as an inflammatory

marker due to evidence supporting MPO as a reliable inflammatory marker in a range of

pathogenesis including cancer, rheumatoid arthritis and cardiovascular disease in addition to

tissue injury [72]. The expression of MPO has also been associated with ovarian epithelial

carcinoma cells in early stage carcinomas [73]. Presence of MPO is also specifically

associated with vascular vessel inflammation [74,75]. Furthermore, MPO has been used as a

reliable indicator of lymphatic vessel inflammatory status to quantify LPS-induced lymphatic

vessel inflammation in rats in conjunction with Masson Trichome staining to indicate fibrotic

vessel areas [13]. Other indicators of inflammatory state could be used, perhaps benefitting

the prognostic value for relapse.

A significant limitation of our feasibility study is the small sample size. Although we have

successfully identified associations with high potential for outcomes prediction, further

samples could both strengthen the trends observed and uncover more obscure alterations

associated with inflammation and/or relapse. Additionally, it is possible that increasing

sample size could result in the identification of inflamed but non-cancer infiltrated LVs.

However, there is no guarantee that increasing sample size would lead to the emergence of

this group. The sample size also contributed to the decision to perform real time PCR with a

relatively small number of pre-selected targets. A broader microarray study would require

many more samples to avoid false positives. We screened for miRNA with known

involvement in inflammatory and autoimmunity due to the previous characterization of

inflammatory vessel state and pathophysiological changes associated with inflammation and

lymphatic function in rat lymphatic vessels [13].

A further limitation to this study is the lack of samples from healthy controls. Harvesting

equivalent lymphatic tissues from healthy patients would not be ethical. However, this

means that the changes in LV status noted here are more directly relevant to identification of

factors that aid prognosis than identifying systemic EOC changes compared to healthy

subjects. Epithelial ovarian cancer surgical techniques and types of cytotoxic treatments do

not differ, as of yet, between the various histological subtypes. We therefore included all

available epithelial subtypes, even the clear cell case, to have a broader representation of the

clinical reality. Although our results support a link between systemic inflammation and

cancer cell dissemination, a larger study is warranted to validate our findings and to allow

further analysis to better identify miRNA patterns using predictive algorithms with greater

confidence.

Conclusions

In conclusion, we have demonstrated that LVs appear to show quantifiable inflammatory and infiltrative alterations in patients with advanced EOC and that a high inflammatory LV state significantly correlates with higher risk of post-surgical relapse. Macroscopically normal lymphatic LVs show accompanying alterations in miRNA expression that may impact LV integrity and secretion of VEGF and inflammatory cytokines. Whether LV inflammation is a cause or consequence of cancer-cell infiltration is yet to be determined, but our results suggest systemic inflammation as a new area to target.

Supporting information

S1 File. Supplementary information regarding building supervised classification algorithms. The techniques and assumptions when building classification algorithms and assessing the impact of individual miRNA.

S1 Table. Additional clinical data (patient age and surgery type).

S1 Fig. Experimental design for LV miRNA analysis from sample collection to analysis. Each vessel dissected from one sample is divided into 2 and processed for IHC and miRNA expression analysis.

S2 Fig. Kaplan Meier estimates of non-relapse (survival). A. According to inflammation (MPO) (high vs. medium/low) B. According to inflammation (MPO) (high/medium vs. low). C. According to cancer cell infiltration (WT1 or Pax.8)

S3 Fig. Kaplan Meier estimates of non-relapse (survival) according to multiple patient

age thresholds. A. Under 55 years B. Under 60 years C. Under 65 years . D Under

70years. E Under 75years.

S2 Table. Differential expression of miRNA between LVs displaying high or low

inflammation. We compared expression in LVs with high versus low inflammation (n=7)

Listed are miRNA that showed a fold-regulation change ±1.8 with those showing a

significant difference between groups highlighted (t-test p>0.05). * = miRNA that remained

below a Bonferroni correction of p<0.00208.

S3 Table. Differential expression of miRNA between LVs displaying high or low

inflammation. We compared expression in LVs with high or medium inflammation versus

low inflammation (n=10) Listed are miRNA that showed a fold-regulation change ±1.8 with

those showing a significant difference between groups highlighted (t-test p>0.05). * =

miRNA that remained below a Bonferroni correction of p<0.00161.

S4 Table. Differential expression of miRNA between cancer-infiltrated LVs and non-

cancer-infiltrated LVs. Listed are miRNA that showed a fold-regulation change ±1.8 with

those showing a significant difference between groups highlighted (t-test p>0.05). * =

miRNA that remained below a Bonferroni correction of p<0.002.

S5 Table. Differential expression of miRNA in LVs from patients that relapsed versus

LVs from patients that didn't relapse within 13.2±6 months. Listed are miRNA that

showed a fold-regulation change ±1.8 with those showing a significant difference between

groups highlighted (t-test p>0.05). Bonferroni correction of p<0.002.

S4 Fig. The KDE estimations of the distributions of inner groups for the individual miRNA expressed at statistically significant levels (p<0.05) between LVs with high inflammation and LVs with low inflammation.

S5 Fig. A. Principle component analysis showed that inflamed and non-inflamed samples show variance across Principle Component (PC) 2. Samples with most similar miRNA expression profile cluster together. Unit variance scaling was applied to rows; SVD with imputation was used to calculate principal components. Prediction ellipses are such that with probability 0.95, a new observation from the same group will fall inside the ellipse. N = 10 data points. Figures produced with ClustVis [26] B. The miRNA expression driving PC1 & 2 contributed to the greatest variation between sample groups with several miRNA identified as differentially regulated when comparing inflamed or cancer-infiltrated LVs (bold =>±1.8, *=p<0.05). C. A Scree plot showing the amount of variation described by each PC confirmed that the two primary PCs accounted for much of the variation between all sample.

S6 Table. Table of classification analysis for various supervised classification algorithms to predict LV inflammation (low or medium/high) or patient stage from LV miRNA expression. Depicted is the accuracy of the algorithm to predict LV inflammation or patient relapse (percentage of correct predictions), the per-group accuracy (No Inflam., Yes Inflam. / Stage < IV, Stage >= IV), Cohen's Kappa and thus the agreement between predicted and actual states, and McNemar's test significance of equality of predicted probability (inner accuracy) between groups for each outcome.

S7 Table. Classifier predictions of the examined states (LV inflammation, LV cancer-
cell infiltration, patient relapse or patient stage) based on the expression of significantly
dysregulated miRNA. Accuracy, Cohen's Kappa, Mc Nemar p-value of equality for inner
grouped probabilities and classification are reported for the predicted classes. The expression
of the most significantly differentially expressed miRNA were added one by one, or in pairs
if significance was equal, into the parameter set used to build the classifiers. When only
significantly differentially expressed miRNA identified in inflamed LVs (high and medium
inflammation versus low) were used to build the classifier, the accuracy of subsequent
predictions increased by 20-40% compared to classifiers based on all available miRNA
expression (S6 Table). Similar improvements were found in the accuracy of classifiers
predicting relapse and LV cancer-infiltration (Tables 7 and 8).

S8 Table: Pathway-analysis performed with the 3 significantly up-regulated and single
significantly down-regulated miRNA identified in the relapse versus non-relapse
groupings.

Acknowledgements

The authors gratefully acknowledge the provision of tissue samples by the Imperial College
Healthcare NHS Trust Tissue Bank. Other investigators may have received samples from
these same tissues. The research was supported by the Experimental Cancer Medicine Centre
(ECMC, Hammersmith Campus) and National Institute for Health Research (NIHR)
Biomedical Research Centre based at Imperial College Healthcare NHS Trust and Imperial
College London. The views expressed are those of the author(s) and not necessarily those of
the NHS, the NIHR or the Department of Health.

References

1. UK CR. Cancer Research UK. 2018. Available from:
    https://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics- by-
    cancer-type/ovarian-cancer.

2. Bachmann C, Bachmann R, Kraemer B, Brucker SY, Staebler A, Fend F, et al.
    Prevalence and distribution pattern of nodal metastases in advanced ovarian cancer.
    Molecular and Clinical Oncology. 2016;5(4):483–487.

3. Bachmann C, Bachmann R, Fend F, Wallwiener D. Incidence and Impact of Lymph
    Node Metastases in Advanced Ovarian Cancer: Implications for Surgical Treatment.
    Journal of Cancer. 2016;7(15):2241–2246.

4. Wimberger P, Lehmann N, Kimmig R, Burges A, Meier W, Bois AD. Prognostic
    factors for complete debulking in advanced ovarian cancer and its impact on survival.
    An exploratory analysis of a prospectively randomized phase III study of the
    Arbeitsgemeinschaft Gynäekologische Onkologie Ovarian Cancer Study Group
    (AGO-OVAR). Gynecologic Oncology. 2007;106(1):69 - 74.

5. Harter P, Sehouli J, Lorusso D, Reuss A, Vergote I, Marth C, et al. A Randomized
    Trial of Lymphadenectomy in Patients with Advanced Ovarian Neoplasms. N Engl
    J Med. 2019;380(9):822-832

6. Padera TP, Kadambi A, Tomaso E, Carreira CM, Brown EB, Boucher Y, et al.
    Lymphatic Metastasis in the Absence of Functional Intratumor Lymphatics. Science.
    2002;296(5574):1883–1886.

7. Leu AJ, Berk DA, Lymboussaki A, Alitalo K, Jain RK. Absence of Functional
    Lymphatics within a Murine Sarcoma: A Molecular and Functional Evaluation.
    Cancer Research. 2000;60(16):4324–4327.

8.   Jones D, Pereira ER, Padera TP. Growth and Immune Evasion of Lymph Node

Metastasis. Frontiers in oncology. 2018;8:36;36–36.

9.   Stachura J, Wachowska M, Kilarski WW, Güc E, Golab J, Muchowicz A. The dual

role of tumor lymphatic vessels in dissemination of metastases and immune response

development. Oncoimmunology. 2016;5(7): e1182278.

10. Dieterich LC, Ikenberg K, Cetintas T, Kapaklikaya K, Hutmacher C, Detmar M.

Tumor- Associated Lymphatic Vessels Upregulate PDL1 to Inhibit T-Cell Activation.

Frontiers in immunology. 2017;8:66.

11. Lund A, Duraes F, Hirosue S, Raghavan V, Nembrini C, Thomas S, et al. VEGF-C

Promotes Immune Tolerance in B16 Melanomas and Cross-Presentation of Tumor

Antigen by Lymph Node Lymphatics. Cell Reports. 2012;1(3):191-199.

12. Dubrot J, Duraes FV, Potin L, Capotosti F, Brighouse D, Suter T, et al. Lymph node

stromal cells acquire peptide–MHCII complexes from dendritic cells and induce

antigen-specific CD4+ T cell tolerance. Journal of Experimental Medicine.

2014;211(6):1153–1166.

13. Chakraborty S, Zawieja DC, Davis MJ, Muthuchamy M. MicroRNA signature of

inflamed lymphatic endothelium and role of miR-9 in lymphangiogenesis and

inflammation. American Journal of Physiology - Cell Physiology.

2015;309(10):C680–C692.

14. Liao S, Cheng G, Conner DA, Huang Y, Kucherlapati, RS, Munn, LL, et al. Impaired

lymphatic contraction associated with immunosuppression. Proceedings of the

National Academy of Sciences of the United States of America. 2011;108(46):18784–

18789.

15. Calin GA, Croce CM. MicroRNA signatures in human cancers. Nature Reviews

Cancer. 2006;6: 857-866.

16. Deb B, Uddin A, Chakraborty S. miRNAs and ovarian cancer: An overview. Journal

of Cellular Physiology. 2018;233(5):3846-3854.

17. Dahiya N, Sherman-Baust CA, Wang TL, Davidson B, Shih I, Zhang Y, et al.

MicroRNA Expression and Identification of Putative miRNA Targets in Ovarian

Cancer. PLOS ONE. 2008;3(6):1-11.

18. Jones D, Li Y, He Y, Xu Z, Chen H, Min W. Mirtron MicroRNA-1236 Inhibits

VEGFR- 3 Signaling during Inflammatory Lymphangiogenesis. Arteriosclerosis,

Thrombosis, and Vascular Biology. 2012;32(3):633–642.

19. Yee D, Shah KM, Coles MC, Sharp TV, Lagos D. MicroRNA-155 induction via

TNF- and IFN-suppresses expression of programmed death ligand-1 (PD-L1) in

human primary cells. The Journal of Biological Chemistry. 2017;292(50):20683–

20693.

20. Sasahira T, Kurihara M, Bhawal UK, Ueda N, Shimomoto T, Yamamoto K, et al.

Downregulation of miR-126 induces angiogenesis and lymphangiogenesis by

activation of VEGF-A in oral cancer. British Journal of Cancer. 2012;107:700-706.

21. Hu J, Cheng Y, Li Y, Jin Z, Pan Y, Liu G, et al. microRNA-128 plays a critical role in

human non-small cell lung cancer tumourigenesis, angiogenesis and

lymphangiogenesis by directly targeting vascular endothelial growth factor-C.

European Journal of Cancer. 2014;50(13):2336-2350.

22. Yang B, Jing C, Wang J, Guo X, Chen Y, Xu R, et al. Identification of microRNAs

associated with lymphangiogenesis in human gastric cancer. 2013;16.

23. Xu J, Zhang C, He Y, Wu H, Wang Z, Song W, et al. Lymphatic endothelial cell-

secreted CXCL1 stimulates lymphangiogenesis and metastasis of gastric cancer.

International Journal of Cancer. 2012;130(4):787-797.

24. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al.

Fiji: an open-source platform for biological-image analysis. Nature Methods.

2012;9:676-682.

25. Vierra A, Garret J. Understanding Interobserver Agreement: The Kappa Statistic.

Research Series. 2005;37(5).

26. Metsalu T, Vilo J. ClustVis: a web tool for visualizing clustering of multivariate data

using Principal Component Analysis and heatmap. Nucleic Acids Research.

2015;43(W1):W566-W570.

27. Vlachos I.S, Hatzigeorgiou AG. DIANA-miRPath v3. 0: deciphering microRNA

function with experimental support. Nucleic acids research. 2015;43:W460–W466.

28. Bhattacharya R, Nicoloso M, Arvizo R, Wang E, Cortez A, Rossi S, et al. MiR-15a

and MiR-16 Control Bmi-1 Expression in Ovarian Cancer. Cancer Research.

2009;69(23):9090–9095.

29. Petrillo M, Zannoni GF, Beltrame L, Martinelli E, DiFeo A, Paracchini L, et al.

Identification of high-grade serous ovarian cancer miRNA species associated with

survival and drug response in patients receiving neoadjuvant chemotherapy: a

retrospective longitudinal analysis using matched tumor biopsies. Annals of

Oncology. 2016;27(4):625-634.

30. Sorrentino A, Liu CG, Addario A, Peschle C, Scambia G, Ferlini C. Role of

microRNAs in drug-resistant ovarian cancer cells. Gynecologic Oncology.

2008;111(3):478 - 486.

31. Boyerinas B, Park SM, Hau A, Murmann AE, Peter ME. The role of let-7 in cell

differentiation and cancer. MicroRNA. 2012;1(1):34-39.

32. Cai J, Yang C, Yang Q, Ding H, Jia J, Guo J, et al. Deregulation of let-7e in epithelial

ovarian cancer promotes the development of resistance to cisplatin. Oncogenesis.

2013;2:e75

33. Liu N, Zhou C, Zhao J, Chen Y. Reversal of Paclitaxel Resistance in Epithelial

Ovarian Carcinoma Cells by a MUC1 Aptamer-let-7i Chimera. Cancer Investigation.

2012;30(8):577-582.

34. Gérard C, Lemaigre F, Gonze D. Modeling the Dynamics of Let-7-Coupled Gene

Regulatory Networks Linking Cell Proliferation to Malignant Transformation.

Frontiers in Physiology. 2019;10:848.

35. Meng F, Henson R, Wehbe-Janek H, Smith H, Ueno Y, Patel T. The MicroRNA let-

7a Modulates Interleukin-6-dependent STAT-3 Survival Signaling in Malignant

Human Cholangiocytes. Journal of Biological Chemistry. 2007;282(11):8256-8264.

36. Kumar M, Ahmad T, Sharma A, Mabalirajan U, Kulshreshtha A, Agrawal A, Ghosh

B. Let-7 microRNA-mediated regulation of IL-13 and allergic airway inflammation.

Journal of Allergy and Clinical Immunology. 2011;128(5):1077–1085.e10.

37. Choudhury S, Li Y. miR-21 and let-7 in the Ras and NF-IκβPathways. MicroRNA.

2012;1(1):65-69.

38. Cromer WE, Zawieja SD, Tharakan B, Childs EW, Newell MK, Zawieja DC. The

effects of inflammatory cytokines on lymphatic endothelial barrier function.

Angiogenesis. 2014;17(2):395–406.

39. Aldrich MB, Sevick-Muraca EM. Cytokines are systemic effectors of lymphatic

function in acute inflammation. Cytokine. 2013;64(1):362–369.

40. Chen PY, Qin L, Barnes C, Charisse K, Yi T, Zhang X, et al. FGF Regulates TGF-ˆI Signaling and Endothelial-to-Mesenchymal Transition via Control of let-7 miRNA Expression.Cell Reports. 2012;2(6):1684-1696.

41. Sun CY, She XM, Qin Y, Chu ZB, Chen L, Ai LS, et al. miR-15a and miR-16 affect the angiogenesis of multiple myeloma by targeting VEGF. Carcinogenesis. 2012;34.

42. Metsalu T, Vilo J. MicroRNA miR-93-5p regulates expression of IL-8 and VEGF in neuroblastoma SK-N-AS cells. Oncol Rep. 2016;35(5):2866-72.

43. Tzeng HE, Chang AC, Tsai CH, Wang SW, Tang CH. Basic fibroblast growth factor promotes VEGF-C-dependent lymphangiogenesis via inhibition of miR-381 in human chondrosarcoma cells. Oncotarget. 2016;7(25):38566–38578.

44. Yang G, Xiong G, Cao Z, Zheng S, You L, Zhang T, et al. miR-497 expression, function and clinical application in cancer. Oncotarget. 2016;7(34):55900–55911.

45. Giacca M, Zacchigna S. VEGF gene therapy: therapeutic angiogenesis in the clinic and beyond. Gene Therapy. 2012;19:622-629.

46. Park SM, Shell S, Radjabi AR, Schickel R, Feig C, Boyerinas B, et al. Let-7 Prevents Early Cancer Progression by Suppressing Expression of the Embryonic Gene HMGA2. Cell Cycle. 2007;6(21):2585–2590.

47. Johnson SM, Grosshans H, Shingara J, Byrom M, Jarvis R, Cheng A, et al. RAS Is Regulated by the let-7 MicroRNA Family. Cell. 2005;120(5):635 – 647[48] Wang S, Zhou H, Wu D, Ni H, Chen Z, Chen C, et al. MicroRNA let-7a regulates angiogenesis by targeting TGFBR3 mRNA. Journal of Cellular and Molecular Medicine. 2019;23(1):556–567.

48. Wang S, Zhou H, Wu D, Ni H, Chen Z, Chen C, et al. MicroRNA let-7a regulates angiogenesis by targeting TGFBR3 mRNA. Journal of Cellular and Molecular Medicine. 2019;23(1):556–567

49. Bao Mh, Zhang Yw, Lou Xy, Cheng Y, Zhou Hh. Protective Effects of Let-7a and

Let-7b on Oxidized Low-Density Lipoprotein Induced Endothelial Cell Injuries.

PLOS ONE. 2014 09;9(9):1–10.

50. Bing Q, Xiao B, Liang D, Li Y, Jiang T, Yang H. MicroRNA Let-7c Inhibits Bcl- Xl

Expression and Regulates Ox-LDL-Induced Endothelial Apoptosis. BMB Reports.

2012;45(8):464–69.

51. Ji X, Hua H, Shen Y, Bu S, Yi S. Let-7d modulates the proliferation, migration,

tubulogenesis of endothelial cells. Molecular and Cellular Biochemistry.

2019;462(1):75– 83.

52. Lin Z, Ge J, Wang Z, Ren J, Wang X, Xiong H, et al. Let-7e modulates the

inflammatory response in vascular endothelial cells through ceRNA crosstalk.

Scientific Reports. 2017 02;7:42498

53. Swaminathan S, Suzuki K, Seddiki N, Kaplan W, Cowley MJ, Hood CL, et al.

Differential Regulation of the Let-7 Family of MicroRNAs in CD4+ T Cells Alters

IL- 10 Expression. The Journal of Immunology. 2012;188(12):6238–6246.

54. Ding Z, Wang X, Schnackenberg L, Khaidakov M, Liu S, Singla S, et al. Regulation

of autophagy and apoptosis in response to ox-LDL in vascular smooth muscle cells,

and the modulatory effects of the microRNA hsa-let-7g. International Journal of

Cardiology. 2013;168(2):1378 – 1385.

55. Liao YC, Wang YS, Guo YC, Lin WL, Chang MH, Juo SHH. Let-7g Improves

Multiple Endothelial Functions Through Targeting Transforming Growth Factor-Beta

and SIRT- 1 Signaling. Journal of the American College of Cardiology.

2014;63(16):1685 – 1694.

56. Kagawa T, Watanabe M, Inoue N, Otsu H, Saeki M, Katsumata Y, et al. Increases of

911   microRNA let-7e in peripheral blood mononuclear cells in Hashimotorsquo;s disease.

912   Endocrine Journal. 2016;63(4):375–380.

913   57. Guan H, Fan D, Mrelashvili D, Hao H, Singh N, Singh U, et al. MicroRNA let-7e is

914   associated with the pathogenesis of experimental autoimmune encephalomyelitis.

915   European journal of immunology. 2013 01;43.

916   58. Liang L, Zhao L, Zan Y, Zhu Q, Ren J, Zhao X. MiR-93-5p enhances growth and

917   angiogenesis capacity of HUVECs by down-regulating EPLIN. Oncotarget. 2017

918   11;8(63):107033–107043.

919   59. Fu X, Tian J, Zhang L, Chen Y, Hao Q. Involvement of microRNA-93, a new regula-

920   tor of PTEN/Akt signaling pathway, in regulation of chemotherapeutic drug cisplatin

921   chemosensitivity in ovarian cancer cells. FEBS Letters;586(9):1279–1286.

922   60. Chen X, Chen S, Xiu YL, Sun KX, Zong ZH, Zhao Y. RhoC is a major target of

923   microRNA-93-5P in epithelial ovarian carcinoma tumorigenesis and progression.

924   Molecular cancer. 2015 02;14(1):31; 31–31.

925   61. Su Z, Si W, Li L, Zhou B, Li X, Xu Y, et al. MiR-144 regulates hematopoiesis and

926   vascular development by targeting meis1 during zebrafish development. The

927   International Journal of Biochemistry Cell Biology. 2014;49:53 – 63.

928   62. Honglei Y, Lihua Z, Junbiao Z, Yan Z. miR-144 regulates the proliferation of human

929   vascular endothelial cells through targeting SRF. Int J Clin Exp Pathol;9(3):3357–

930   3364.

931   63. Han S, Zhu J, Zhang Y. miR-144 Potentially Suppresses Proliferation and Migration

932   of Ovarian Cancer Cells by Targeting RUNX1. Medical science monitor basic

933   research. 2018 02;24:40–46.

64. Li J, Zhao Y, Lu Y, Ritchie W, Grau G, Vadas MA, et al. The Poly-cistronic miR-23-

27- 24 Complexes Target Endothelial Cell Junctions: Differential Functional and

Molecular Effects of miR-23a and miR-23b. Molecular Therapy - Nucleic Acids.

2016;5:e354.

65. Wu XD, Guo T, Liu L, Wang C, Zhang K, Liu HQ, et al. MiR-23a targets RUNX2

and suppresses ginsenoside Rg1-induced angiogenesis in endothelial cells.

Oncotarget. 2017;8(35):58072–58085.

66. Zhu S, Pan W, Song X, Liu Y, Shao X, Tang Y, et al. The microRNA miR-23b

suppresses IL-17-associated autoimmune inflammation by targeting TAB2, TAB3

and IKK-. Nature Medicine. 2012;18(7):1077–1086.

67. Reitsma S, Slaaf DW, Vink H, Zandvoort MAMJ, Egbrink MGA. The endothelial

glycocalyx: composition, functions, and visualization. European journal of

physiology. 2007;454(3):345–359.

68. Zolla V, Nizamutdinova IT, Scharf B, Clement CC, Maejima D, Akl T, et al. Aging-

related anatomical and biochemical changes in lymphatic collectors impair lymph

transport, fluid homeostasis, and pathogen clearance. Aging cell. 2015;14(4):582–

594.

69. Chakraborty S, Zawieja SD, Wang W, Lee Y, Wang YJ, von der Weid P, et al.

Lipopolysaccharide modulates neutrophil recruitment and macrophage polarization on

lymphatic vessels and impairs lymphatic function in rat mesentery. American journal

of physiology. Heart and circulatory physiology. 2015;309(12):H2042–H2057

70. Bridenbaugh EA, Wang W, Srimushnam M, Cromer WE, Zawieja SD, Schmidt SE,

et al. An immunological fingerprint differentiates muscular lymphatics from arteries

and veins. Lymphatic research and biology. 2013 09;11(3):155–171.

71. Hunter MC, Teijeira A, Halin C. T Cell Trafficking through Lymphatic Vessels.

Frontiers in Immunology. 2016;7:613.

72. Khan AA, Alsahli MA, Rahmani AH. Myeloperoxidase as an Active Disease

Biomarker: Recent Biochemical and Pathological Perspectives. Medical sciences

(Basel, Switzer- land). 2018 04;6(2):33.

73. Castillo-Tong DC, Pils D, Heinze G, Braicu I, Sehouli J, Reinthaller A, et al.

Association of myeloperoxidase with ovarian cancer. Tumour biology: the journal of

the International Society for Oncodevelopmental Biology and Medicine. 2014

01;35(1):141– 148.

74. Klinke A, Nussbaum C, Kubala L, Friedrichs K, Rudolph TK, Rudolph V, et al.

Myeloperoxidase attracts neutrophils by physical forces. Blood. 2011;117:1350–135.

75. Tiruppathi C, Naqvi T, Wu Y, Vogel SM, Minshall RD, Malik AB. Albumin mediates

the transcytosis of myeloperoxidase by means of caveolae in endothelial cells.

Proceedings of the National Academy of Sciences of the United States of America.

2004 05;101(20):7699–7704.

A Segment to be isolated for RNA extraction

Segment for IHC, maintaining surrounding tissue

25 mm

B

C

D MPO 1.0%

E DAPI 6.3%