Coevolution of retroviruses with *SERINC*s following whole-genome duplication divergence

Pavitra Ramdas[1], Vipin Bhardwaj[1], Aman Singh[1], Nagarjun Vijay[2], Ajit Chande[1*]

[1]*Molecular Virology Laboratory &* [2]*Computational Evolutionary Genomics Lab from the Department of Biological Sciences, Indian Institute of Science Education and Research (IISER) Bhopal, India.*

Correspondence: ajitg@iiserb.ac.in

Abstract

The *SERINC* gene family comprises of five paralogs in humans of which *SERINC3* and *SERINC5* inhibit HIV-1 infectivity and are counteracted by Nef. The origin of this anti-retroviral activity, its prevalence among the remaining paralogs, and its ability to target retroviruses remain largely unknown. Here we show that despite their early divergence, the anti-retroviral activity is functionally conserved among four human *SERINC* paralogs with *SERINC*2 being an exception. The lack of activity in human *SERINC*2 is associated with its post-whole genome duplication (WGD) divergence, as evidenced by the ability of pre-WGD orthologs from yeast, fly, and a post-WGD-proximate *SERINC*2 from coelacanth to inhibit nef-defective HIV-1. Intriguingly, potent retroviral factors from HIV-1 and MLV are not able to relieve the *SERINC*2-mediated particle infectivity inhibition, indicating that such activity was directed towards other retroviruses that are found in coelacanth (like foamy viruses). However, foamy-derived vectors are intrinsically resistant to the action of *SERINC*2, and we show that a foamy virus envelope confers this resistance. Despite the presence of weak arms-race signatures, the functional reciprocal adaptation among *SERINC*2 and *SERINC*5 and, in response, the emergence of antagonizing ability in foamy virus appears to have resulted from a long-term conflict with the host.

Keywords: arms-race, *SERINC*, restriction factors, anti-retroviral, whole-genome duplication

Introduction

Viruses have been exploiting the host machinery for their persistence, and, in response, the host has continued to evolve increasingly intricate antiviral defense strategies. As part of this ongoing arms-race, while viruses have relied on acquisition and fusion of diverse genes (1), the host-defense mechanism has been made possible by the functional divergence of gene copies following duplication of genes as well as whole-genomes (2–7). Restriction factors being at the forefront of this long-term conflict (8–10), show clear molecular signatures of arms-race (11,12). In fact, the presence of these signatures has been proposed to be a hallmark of restriction factors (8,13), and has been used as a screening strategy to identify potential candidates(2,14). In contrast, the recently identified anti-retroviral host inhibitors *SERINC*5 and

*SERINC*3 display an uneventful evolutionary history (15). This is counterintuitive, because

distant retroviruses, with wide host-range, encode anti-*SERINC*5 virulent factors (16–18). We,

therefore, sought to trace the origins of this antiretroviral activity and its relevance for retroviral

inhibition. Our analysis to comprehend the evolutionary origins of the antiretroviral activity in

*SERINC*s, identifies an active *SERINC*2 with a hitherto unknown interaction with a foamy virus.

Results

Antiretroviral activity among human *SERINC* paralogs

Analysis of sequence similarity and gene structure conservation reveals that *SERINC*5 and

*SERINC*4 share a recent ancestry **(Fig-S1)**. Similarly, *SERINC*3 and *SERINC*1 paralogs are

most similar to each other (~60% identity). Despite having the lowest sequence similarity to

either of the established anti-viral *SERINC* paralogs, *SERINC*2 is relatively similar to *SERINC*3

(50%) and *SERINC*5 (37%). Given such levels of sequence similarity and conserved membrane

topology **(Fig-S2)**, we checked the activity of other paralogs to inhibit HIV-1 infectivity in addition

to *SERINC*5 and *SERINC*3. From the native versions of *SERINC*1 and *SERINC*4 we failed to

express the proteins, so we decided to generate codon-optimized versions of these genes.

Having found the proteins being expressed upon codon optimization (**Fig1A**), we used these

conditions to check the inhibitory activity of *SERINC*1 and *SERINC*4. Consistent with previous

reports (19,20), we find that *SERINC*2 is the only *SERINC* to have no activity against HIV-1

while all the other *SERINC*s inhibit the virus infectivity to a varying degree (**Fig-1A and 1B**).

Despite less-prominent membrane localization, *SERINC*4 has a potent activity and is second-

most powerful in inhibiting the virus (**Fig-1C**).

In contrast to other restriction factors, *SERINC*5 and *SERINC*3 genes have a rather uneventful

history that is distinct from the traditional signatures of recurrent selection seen in genes that

are part of an arms-race (21). We investigated the generality of whether the absence of

signatures of positive selection was prevalent across all the *SERINC* paralogs. Towards this,

we compared the evolutionary signature of *SERINC* genes with previously identified restriction

factors, genes showing recurrent positive selection and a few functionally characterized genes

that can act as controls **(Fig-1D)**. Within *SERINC* paralogs there is heterogeneity in the patterns

of positive selection, with *SERINC*1 and *SERINC*5 showing the least number of positively

selected residues while, *SERINC*3 and *SERINC*4 have relatively a greater number of residues

under episodic diversifying selection **(Fig-1D; FigS3, S4;** arms-race signature inference is

found to be sensitive to the multiple sequence alignment tool used, see **Fig-S5)**. Functionally

though, *SERINC*4 and *SERINC*5 are the most potent antiretroviral inhibitors compared to the

other three *SERINC*s, suggesting that merely the fraction of sites that are positively selected is

not an indicator of its ability to inhibit the retrovirus. Although the arms race signatures of few

restriction factors such as BST-2 and EIAF2AK2 were well correlated, the other newly identified

restriction factors, including *SERINC*s did not show any consistent pattern of clustering **(Fig-**

1D). One of the prime features of restriction factors has been their ability to get augmented

upon interferon stimulation. While the genes which formed a cluster are interferon responsive

(IFN responsive genes were obtained from ref (22)), this property did not explain the overall
pattern of clustering **(Fig-1D)**. Arms race signatures are not prominent in several interferon-
inducible genes and innate immune genes, including TLRs **(Fig-1D)**. Hence the lack of arms
race signatures in *SERINC*s is probably not very surprising. The ability to restrict HIV-1 among
human *SERINC*1 and *SERINC*4 suggests that this is an evolutionarily conserved feature
despite the early divergence of these paralogous copies. Hence, we further decided to
investigate the evolutionary origins of this anti-retroviral activity.
88
Anti-retroviral activity in an ancient feature among *SERINC*5 orthologs
*SERINC*s plausibly shaped retroviral evolution, as indicated by the parallel emergence of anti-
*SERINC*5 activity among diverse retroviral genomes. *SERINC* gene family of restriction factors
is conserved across eukaryotic species with evident topological similarity **(Fig-S6)**, so we asked
to what extent the ability of *SERINC* orthologs to restrict HIV-1 is conserved. Unicellular
eukaryotes and invertebrates have a single copy of the *SERINC* gene (*TMS*1); all these
orthologs could restrict *nef*-defective HIV-1 (**Fig-1E and 1F**), albeit the yeast ortholog having
modest activity (2-fold) in comparison to all other orthologs (fold inhibition from 4-36). This lower
potency compared to all other orthologs could be attributed to inadequate localization on the
membrane or heterologous host expression for yeast and fly *TMS*1 in conditions where the
remaining orthologs were adequately expressed on the membrane (**Fig-1G**). Western blot
analysis confirmed the expression of C-terminal HA-tagged TMS-1 and other *SERINC*5
orthologs (**Fig-1E,** lower panel).
102
Coelacanth *SERINC*2 restricts HIV-1
Two rounds of WGD in the ancestor of chordates led to a large increase in the number of genes
leading to the acquisition of new functions. This repertoire of genes also provides greater
flexibility due to their mutually compensatory functions. Hence post-WGD paralogous copies
tend to diversify. A single copy of *SERINC* ortholog (*TMS*1) is present in pre-WGD species and,
post- WGD, the number of copies has increased to five in tetrapods and six in bony fishes
(FigS1). Our investigation of the pre-WGD ortholog of *SERINC* genes from the yeast (*S.*
cerevisiae) and the fly (*D. melanogaster*) *TMS1,* found evidence of anti-retroviral activity (**Fig-**
1E). Functional data on the human paralogs also suggests an ancestrally conserved activity of
*SERINC*s in restricting HIV-1, *SERINC*2, however, being an exception (**Fig1A**). Hence, the
human *SERINC*2 paralog may have lost the ability to restrict HIV sometime after the WGD
(~700 to 400 MYA)(23). It is plausible that the species proximal to the WGD-event might retain
a version of *SERINC*2 with antiviral activity **(Fig-2A)**. We, therefore, decided to systematically
screen *SERINC*2 orthologs from post-WGD species at varying levels of sequence divergence
from human *SERINC*2. This exercise revealed that Coelacanth *SERINC*2 restricts HIV-1 in
conditions where *SERINC*2 isoforms from human did not show any activity (**Fig-2B**). A shorter
isoform of the Human *SERINC*2 (Human-201) was found to be topologically similar (**Fig-S7**) to

that of Coelacanth *SERINC*2 but lacked the activity to restrict HIV-1 concurrent with the longer

isoform.

The gradual loss of antiretroviral activity in *SERINC2*

Upon further assessment of anti-HIV-1 activity of post-WGD *SERINC*2 orthologs, we find that

while Coelacanth *SERINC*2 reduced the infectivity by ~7-fold, Xenopus *SERINC*2 exhibited a

modest inhibition (~2-fold). The activity was completely lost in chicken *SERINC*2 ortholog

onwards, this is despite the topology being conserved and the levels of virus-incorporation for

chicken ortholog were also appreciable **(Fig-2C, S8 and S9)**. This lack of activity, however, is

persistent in mouse, horse and human *SERINC*2 **(Fig-2C and 2D)**. Coelacanth *SERINC*2

restricted HIV-1 to a lesser extent compared to human *SERINC*5; however, it was more potent

than human *SERINC*3 and *SERINC*1. Further, reduced restriction potency can be attributed to

its inadequate membrane presence **(Fig-2E)**, reminiscent of localization of the pre-WGD

orthologs from yeast and fly TMS-1 **(Fig-1G)**. We see that there is a dose-dependent inhibition

when coelacanth *SERINC*2 was expressed from the plasmids carrying the promoters of

different strengths (**Fig-S10**). Western blot of C-terminal HA-tagged *SERINC* expressors

verified that each of these *SERINC*2 orthologs is expressed from the synthetic genes **(Fig-2C,**

lower panel**)**. Loss of human *SERINC*2 antiviral activity could have been associated with

changes in pathogen repertoire or simply gain of a new function. To experimentally test if this

was in response to change in the pathogen repertoire, we asked if the counteraction of Human

*SERINC*5 by known retroviral factors, HIV-1 Nef[1] and MLV GlycoGag[1], was analogous to that

of Coelacanth *SERINC*2. In conditions where Nef and GlycoMA efficiently counteracted the

potent restriction imposed by Human *SERINC*5 and the partial restriction of Xenopus *SERINC*2,

we find that counteraction of Coelacanth *SERINC*2 restriction by these virulent factors was not

appreciable **(Fig-3A and 3B).** We also inspected if the nef alleles from human and non-human

primate lentiviruses showed a similar phenotype and found that nef alleles did not rescue the

infectivity comparable to that of *SERINC*5 **(Fig-3C)**. The ability of *SERINC*5 to restrict HIV-1

inhibition varies with the envelope glycoproteins used for pseudotyping. We checked if this is

the case with coelacanth SERINC2 as well. Coelacanth *SERINC*2 action indeed phenocopied

to that of human *SERINC*5 in terms of the envelope sensitivity **(Fig-3D)**. However, the retroviral

factors still failed to rescue this analogous activity of coelacanth *SERINC*2.

Functional evidence for arms-race dynamics

Three distinct genera of retroviruses were reported to have independently come up with

antagonizing factors to elude the inhibition by *SERINC*5. Having no activity in Nef and Glycogag

against coelacanth *SERINC*2 **(Fig-3A and 3B)** whether represents a new case of functional

arms-race dynamics is what we wanted to check next (16–18,24) **(Fig-3E)**. We learnt that

coelacanth fish has an endogenous foamy virus (25) the genome organization of which

resembled the prototype foamy virus (**Fig-4A**). The activity of coelacanth *SERINC*2 may have

been linked to a foamy virus infection of the host. To experimentally test the idea of whether

*SERINC*2 has in-fact evolved to restrict foamy viruses, we checked the presence of anti-foamy

virus activity in human *SERINC* paralogs as well as the *SERINC*2 orthologs. Unexpectedly, we

find that foamy is insensitive to any of the human *SERINC* paralogs tested as well as the

coelacanth *SERINC*2, in conditions where HIV-1 was consistently inhibited **(Fig-4B)**. We

argued that this might have been associated with an intrinsic ability of a Foamy virus-encoded

factor to counter the inhibition. To dissect this, we co-expressed foamy genes individually to

check its ability to rescue, now, *nef*-defective HIV-1 such that the insensitivity of foamy and the

presence of antagonizing factor can be revealed. Surprisingly, with none of the foamy

components expressed in-*trans* we find an ability to rescue the *nef*-defective HIV-1 in the

presence of Human *SERINC*5 **(Fig-4C)**. The inhibition exerted by coelacanth *SERINC*2,

however, was antagonized by the Envelope glycoprotein of Foamy virus using an equivalent

amount of nef expressing plasmid **(Fig-4D)**, as well as in a dose-dependent manner **(Fig-S11)**.

We interrogated if this rescue was an effect of cross-packaging of the HIV-1 core by the foamy-

virus envelope glycoprotein. By using a pseudoparticle comprising of an HIV core

complemented with the foamy-virus envelope, we show that this was not an effect of cross

packaging **(Fig-S12)**, and that this rescue by the envelope glycoprotein was specific to its ability

to antagonize coelacanth *SERINC*2 **(Fig-4D)**. To then understand the specificity of this

counteraction, we checked the expression of Coelacanth *SERINC*2 at protein level when co-

expressed with Foamy envelope. We checked the effect with both low expression (PBJ6) and

high expression(pcDNA) of Coelacanth SERINC2 and observed that the steady state levels of

Coelacanth SERINC2 is affected in the producer cells when Foamy envelope was co-

expressed **(Fig-4E)**. Further, we questioned as to whether the foamy virus was at all sensitive

to the effects of coelacanth *SERINC*2. To answer this, we pseudotyped a foamy-virus core,

now, with an envelope glycoprotein of HIV-1 lacking a c-terminal domain, for packaging of a

heterologous foamy core since the native, full-length, envelope glycoprotein of HIV-1 was not

capable of the packaging it. After successful packaging using c-terminally deleted HIV

envelope, we show that intrinsically the foamy core is sensitive coelacanth SERINC2 restriction

(Fig 4F). This advocates that foamy-virus is indeed sensitive to coelacanth *SERINC*2 and has

developed a mechanism to antagonize the inhibitory effect through the envelope glycoprotein

likely as part of an ongoing arms-race.

Discussion

The use of comparative evolutionary genetic approaches has continued to enrich our

understanding of restriction factor biology for more than a decade(26–30). Reciprocal loss of

duplicated genes in different species has been shown to contribute to species-specific

differences in susceptibility to pathogens(31–34). Similar to reciprocal gene loss, we find that

*SERINC*2 and *SERINC*5 show reciprocal functional adaptation against divergent retroviruses.

The ability of these orthologous factors to restrict HIV-1 is remarkable, even in the heterologous

hosts **(Fig1E, 2C and Fig-S13)** suggestive of a topology associated feature that remained

conserved and contribute to virus restriction. Further, the *SERINC*s from a human can restrict

divergent viruses, like MLV(16,17), EIAV(15,18), this only indicates a long-term interaction of

these host factors which exerted a strong selective pressure in order to shape retrovirus
evolution. While exogenous retroviruses have not yet been reported from pre-WGD species
that we considered, the range of targets that a pre-WGD *SERINC* (TMS-1) will have remains
to be identified. The restriction on mobility of genetic elements in yeast has earlier been shown
conserved in APOBEC(35). While genes such as CCR4 and DHH1 that physically interact with
TMS-1 have been implicated in yeast retrotransposon activity (36,37) the mechanism that TMS-
1 would manifest on retroelements remains to be investigated. TMS-1 in fly may still be required
to regulate the mobility of gypsy retroelements as the gypsy envelope has been shown to
pseudotype MoMLV-based vector to efficiently infect fly cells (38,39). Moreover, the analogous
mechanism that Rous sarcoma virus uses to produce Pr180gag-pol has been found in yeast
Ty-1 transposons (40), makes such existence of retroelement interactions for other host-factors
like TMS-1 more probable. Post-WGD, *SERINC* has five copies and this may have been
associated with tackling increasing diversity of pathogens during speciation and while doing so
retaining its core function. As demonstrated here, with cross-packaging studies we show that
*SERINC*2 ortholog, which was thought to be inactive, may potentially have constituted a critical
barrier to foamy viruses early on, leading to their divergence over more than 450 MY (25,41).
Loss of activity in other *SERINC*2 orthologs may have been associated with
neofunctionalization: as suggested by localized sequence divergence **(Fig-S14)**, presence of
HNF4alpha binding enhancer **(Fig-S15)** and change in tissue-specific expression patterns **(Fig-**
S16) exemplified by expression of *SERINC*2 in the liver of primates. Interestingly, Coelacanth
*SERINC*2 inhibited HIV-1 and remained invisible to the most powerful retroviral virulent factors
nef (MAC239) and to the glycoGag. Further studies can provide more insights into the lack of
activity in these retroviral factors.
The ability of foamy virus envelope to counteract the restriction is first such evidence where a
canonical gene product is employed to evade *SERINC* restriction by these atypical retroviruses.
Foamy viruses co-diverged with their hosts, and this consistently is also visible in their ability to
introduce variations in envelope glycoproteins, among other documented genomic alterations.
Such changes perhaps are required for its efficient association with cognate entry receptors or
prevent host mechanisms from inhibiting the virus propagation. The nature of many such
interactions remains to be clarified due to the unavailability of reagents to study the phenotype
in its native context. The counteraction phenotype associated with foamy envelope
demonstrated here is not unexpected due to following the reason. It has been earlier suggested
that *SERINC*5-mediated particle inhibition varies with the HIV-1 envelope from various isolates
(16,42,43); indicating envelope as a determinant for *SERINC*5-sensitivity to particle restriction
as also recently demonstrated in (ref 38). Moreover, in the case of EIAV and MLV envelope the
ability to resist *SERINC*5 action has been observed (16,18). Constant engagement of *SERINC*5
by the host seems to have therefore acted as a driving force during the evolution of these
retroviruses. Having found foamy envelope-dependent activity among coelacanth *SERINC*2 is
therefore not surprising and is in agreement with previous reports (44,45) suggesting an ability
of envelope to interact with SERINC. As a result, an early challenge by this host factor leading

to subsequent divergence of foamy envelope towards insensitivity to *SERINC* restriction. The
change in foamy envelope glycoproteins among coelacanth endogenous foamy virus and a
prototype foamy attest these facts that there may have been a selective pressure imposed by
*SERINC*2. Different SERINC paralogs might have specialized to restrain specific viruses
leading to coevolution of viruses in response to specific paralog.

We foresee that weaker signatures, exemplified by *SERINC*s **(Fig-1D, S3)** despite the constant
challenge from viruses, could be due to the native functions endowed to such transmembrane
proteins where the adaptation through diversification in-response to the pathogen would result
in a loss of a core function (46). The poor signatures could also be because the antiretroviral
activity is spread out over multiple host genes e.g. as a recent report shows that a Nef-sensitive
TIM1 activity is potentiated by *SERINC*5 (47). The host, therefore, can afford such redundancy
without having to diversify much. Another example is TLRs where the extracellular domain
shows signatures of recurrent positive selection in contrast to the conserved membrane-
spanning region (9). Intriguingly, virus specific TLRs are under stronger purifying selection than
non-viral TLRs (48) potentially due to the larger number of PAMPs associated with non-viral
pathogens. This constraint however may be more pronounced in SERINCs as they are multi-
pass transmembrane proteins known to inhibit only retroviruses. This nevertheless is indicative
of more *SERINC*-like restriction factors that display poor signatures of arms-race but are
functionally active. The core function of *SERINC*s in eukaryotes awaits independent
observations nevertheless, the most recent effort in providing structural insights into the fly and
human ortholog (44) opens up avenues for structure-inspired biological perturbations.

Despite being conserved, SERINCs have managed to diversify their function against distinct
retroviruses. In order to explain these dynamics of *SERINC*-retrovirus evolution, we first
collated all the known interactions of SERINCs with retroviral factors (**Fig-5A**). Since SERINC1
and SERINC4 are not translated from the native transcripts in the conditions that have been
probed, we reasoned that their presence might have been associated with extinct retroviral
factors. Investigation of SERINC1 and SERINC4 orthologs in diverse species, might be able to
decipher the proposed interactions (**Fig-5A**). In contrast to SERINC1 and SERINC4, we find
evidence for SERINC2 being counteracted by retroviral factor from foamy virus. Having
investigated all the available data, we decided to propose a model of co-evolution between
SERINCs and retroviruses that could explain the diversification of function despite weak
signatures of arms-race. Host-driven changes to the viral genome sequence is similar to niche-
filling models used to explain evolution of phenotypic traits (49). Hence, we reasoned that
divergence of SERINC paralogs following WGD would result in divergence of the niches a virus
can occupy. Not surprisingly, we find that viruses have evolved two types of retroviral
antagonists to resist SERINC restriction. These types reflect the distinct niches that viruses can
occupy. In contrast to previously proposed host-driven models of virus evolution characterized
by host adaptation and host-switches, our model suggests the creation of host-niches within
the same organism due to divergence of paralogous gene copies. We illustrate our model using

the example of post-WGD diversification of interaction between SERINCs and distinct types of
retroviral factors (**Fig-5B**).

In conclusion, evolution-guided analysis for tracking the origin of anti-viral activity in the
SERINC genes and the dynamics following whole-genome duplication have identified the
presence of antiretroviral activity in the only *SERINC* thought to be deficient. The ancient
antiretroviral activity among *SERINC*s seems to have challenged retroviruses for more than
450 MY, as suggested by the presence of an evasion strategy in a spumavirus representative
to target coelacanth *SERINC*2. The parallel emergence of anti-*SERINC* strategies among
divergent retroviruses thus indicates a fundamental role of these host factors in shaping
retrovirus evolution.

Materials and Methods

Plasmids and reagents

All the plasmids used were isolated using MN NucleoBond® Xtra Midi EF according to the
manufacturer's instructions to minimize endotoxin content. The list of plasmids and reagents
used in the experiments are tabulated in ***Supplementary Table I and II***.

Cell lines

All cell lines used were assessed for mycoplasma contamination and were found to be negative.
HEK293T and TZM-GFP indicator cell lines (previously described in Ref (16) were grown in
Dulbecco's Modified Eagle's Medium (DMEM) with 10% heat-inactivated Fetal Bovine Serum),
2mM L-Glutamine and 1% Penicillin-Streptomycin. Jurkat-TAg$^{SERINC3/5KO}$were reported
previously (18), maintained in RPMI 1640 with 10% FBS and 2mM L-Glutamine. All the cultures
were maintained at 37^{0}C and 5% CO2 in a humidified incubator.

Viruses and infectivity assay

HIV-1 was produced from HEK293T by calcium phosphate transfection and were limited to
single-cycle replication. For a 10 cm plate, 7ug NL4-3 env defective and nef defective, 1ug env
expressing plasmid and 100ng of pcDNA vectors expressing *SERINC* paralogs and orthologs
or equivalent empty vector and 1ug of PBJ6 *SERINC*5 HA was used for virus production. For
production of foamy viruses, transient transfection of 0.736 ug pCIES, 1.5ug pCIPS, 11.84ug
pCIGS, 11.84 ug pΔΦ and 100ng of pcDNA vectors expressing *SERINC* paralogs and
orthologs or equivalent control vector in HEK293T(10 cm plate) was done (50). The viral
particles were collected 48 hours post transfection and were centrifuged at 300g for 5 min and
quantified by SG-PERT reverse transcription assay (51). Following this, viruses were serially
diluted 3, 9 and 27-fold for infection in TZM-GFP reporter cells that were seeded in a 96-well
plate, 24 hours prior to infection. HIV-1 Infectivity was quantified by scoring the GFP positive
cells using Spectra max MiniMax^{TM} 300 imaging Cytometer (Molecular devices, USA). Foamy
Virus infectivity was examined by the number of GFP positive cells indicating the population
transduced by Foamy virus. The total cell population was analyzed by nuclei staining with

Hoechst 33342 and visualized on CellInsight CX7 High Content Screening platform (Thermo
Scientific). The acquired values were normalized with the virus titre obtained from SG-PERT
assay as previously described (16) Results are expressed as a percentage of an internal control
sample.

Immunofluorescence

For electroporation in JTag$^{SERINC3/5KO}$, cells were grown, and fresh medium was added a day
before such that the culture concentration did not exceed 10^6 cells/ml. Cells in their exponential
growth phase were pellet down (10^7 cells/sample) at 300xg for 10 mins. Prior to adding Opti-
MEM, the cells were washed with 1X Phoshphate Buffered Saline (PBS (1X) pH 7.0) to remove
residual serum and other cell debris. Each sample was resuspended in 200ul warm
OptiMEM. 5ug of constructs expressing HA-tagged *SERINC*2 orthologs or equivalent control
vector were then added to the cells. The cells and DNA mixture are added to a 0.2cm gap
electroporation cuvette. The cells were pulsed at 140V and 1000uF with exponential decay on
BioRAD GenePulser Xcell. 600ul warm RPMI with 20% FBS was immediately added to the
electroporated cells, following which they were transferred to a plate containing RPMI with 10%
FBS. 48 hours post transfection cells were pellet down at 300g for 5 mins and resuspended
in100ul RPMI and laid on poly-L-lysine coated coverslips and fixed with 4% paraformaldehyde.
After fixation, the cells were washed twice with 1X PBS. The cells were permeabilized using
BD™Perm/wash followed by detection of HA tag with Purified anti-HA.11 Epitope Tag Antibody
and Alexa 488 fluorescent tagged secondary antibody. The coverslips were transferred to a
glass slide and mounted using ProLong™ Glass Antifade reagent. Images were acquired after
12 hours on Zeiss confocal microscope (LSM740).

Virus incorporation

Viruses were produced by calcium phosphate transfection in a 10cm plate of HEK293T as
mentioned above. The media was replaced with DMEM containing 2% FBS after 12-15 hours
of transfection. After 48 hours, the virus containing supernatant was collected and centrifuged
at 500xg for 10 mins to exclude any cell fragments. Following this, the viruses were filtered
using a 0.22µm syringe filter. This was overlaid on 25% sucrose cushion and concentrated at
50000xg for 2 hours at 4 degrees using a Beckmann Coulter ultracentrifuge. After the spin, the
supernatant was aspirated off and the pellet was lysed in Laemmli buffer with TCEP.

Western blotting

After collection of viruses from producer cells (HEK293T), the cells were collected in 1X ice-
cold PBS. They were washed twice at 500g for 5 minutes. The PBS was aspirated until the
pellet was completely dry. The pellets were either processed for lysis or stored in -80 until
further application. The cell pellets were lysed in DDM Lysis buffer (100mM NaCl, 10mM
HEPES pH 7.5, 50mM Tris(2-carboxyethyl) phosphine hydrochloride (TCEP), 1% n-Dodecyl-

β-D-maltoside (DDM), 2xcOmplete™, EDTA-free Protease inhibitor cocktail and rocked on ice
for 30 minutes. Following this, the lysates were centrifuged at 10000g for 15 mins and the
supernatant was collected and mixed with 4x Laemmli buffer with 10mM TCEP.
SDS-PAGE was used for resolving cell pellets and virions for analysis by western blot. The
proteins from the gel were transferred onto a PVDF membrane on a semi-dry transfer unit
(Hoefer TE77) for 75 minutes with 125 Amp constant current. The membrane containing the
proteins was then blocked with 2x Odyssey Blocking Buffer that was diluted with 1x TBS to
make 1x Blocking buffer. After 20 minutes of blocking, Anti-HA.11 Epitope Tag Antibody
(mouse) and anti-actin(rabbit) (diluted 1: 1000 in 1x blocking buffer) was used to probe the
membrane to detect the *SERINC* orthologs and paralogs that were tagged with C-terminal HA
tag. After incubation with the primary antibody for 1 hour, the membrane was washed thrice
with 1X TBST for 5 mins. Goat anti-mouse 680 and goat anti-rabbit 800 Li-Cor antibodies were
used to detect infra-red dyes signal from the membrane. The western blot was analysed on
Odyssey imager system.
373

Reconstruction of coelacanth *SERINC*5 sequence
The *SERINC*5 from coelacanth was found to be shorter than its annotation from its orthologs.
Upon closer inspection it was found that the exons orthologous to human exons1-3 were
missing in the coelacanth *SERINC*5 genomic sequence. We found that the upstream region of
coelacanth *SERINC*5 had a gap in the genomic sequence. It was also found that the transcribed
regions upstream of the gap did not correspond to *SERINC*5 exons. Hence, we decided to use
a chromosome walking strategy to reconstruct the missing exons, such that at least at
sequence level the topology of the protein could be studied (suppl material for reconstruction
of *SERINC*5).
383

384   **Data and code availability**
385   The multiple sequence alignments are available for download under GNU license from the
386   github repository https://github.com/ceglab/RestrictionFactorsArmsRace.
387

388   **Genome correction and quality check**
389
In order to rule out the possibility of artefacts in evolutionary analysis that arise from errors in
the sequence of genome assemblies (true even for high quality reference genomes found on
ensemble(52)) we systematically assessed the quality of each of the primate genomes used in
this study. Genome assemblies of 15 primate species were downloaded from ensembl release
98 through the ftp site. Whole genome sequencing datasets corresponding to each of these
species were obtained from the Short Read Archive (SRA) with the criteria that they should
have >~30x coverage. Details of the genome assemblies used and the corresponding raw read
data from each species are provided in GitHub repository. The raw read data were mapped to
the corresponding genomes using the bwa mem read mapper(53) with default settings. The

alignment files obtained from the mapping step were used to generate the genotype likelihood
estimates using the program angsd (54). The genotype likelihood estimates were provided to
the program referee(55) to assign quality scores and perform genome correction. Overall, we
found that in all the primate genomes considered, less than 1% of the bases were corrected by
the program referee. The sequencing data used for performing genome correction are not from
the same individual that was used for genome assembly. Hence, it is possible that many of the
corrected positions are merely nucleotide polymorphisms.
406

Manual curation and multiple sequence alignment
The manually curated open reading frame multi-fasta files consisting of 70 genes from ~15
primate species were collected from Ensembl. We extend our previous multiple sequence
alignment strategy(56) by including additional multiple sequence alignment programs to
generate 8 independent alignments for each gene. Several multiple sequence alignment tools
were used to ensure that the inferred patterns of sequence evolution are not restricted to the
alignment strategy used. The choice of the actual multiple sequence alignment tools used was
based on the performance-based classification of algorithms (57).
415

Use of FUBAR to find evolutionary fingerprints
Traditional approaches that endeavour to find arms race signatures in genes look for recurrent
occurrence of positive selection in the same gene in different evolutionary lineages. A more
recent approach has been to use the full joint distribution of synonymous (α) and
nonsynonymous (β) rates as an evolutionary fingerprint of a gene(21). The program
FUBAR(58) is available as part of the HyPhy package(59). Pairs of conditionally dependent
sites were identified using the program BGM(60). Estimates of selection coefficients are
inferred from the multiple sequence alignments for pairs of α- β combinations that form a
discretized grid. We calculated the distance measure defined by Murrel(21) to perform
hierarchical clustering of the selection signatures obtained from FUBAR.
426
427

Acknowledgements
This work is supported by IYBA fellowship (BT/010/IYBA/2017/01), a grant
(BT/PR26013/GET/119/191/2017) from the Department of Biotechnology, Government of
India, and the Wellcome Trust/DBT India Alliance Fellowship [grant number IA/I/18/2/504006
awarded to AC]. PR, VB and AS are supported by a fellowship from the MHRD, CSIR and DBT
respectively. The computational analyses were performed on the Har Gobind Khorana
Computational Biology cluster. Authors thank Jeremy Luban, and the NIH AIDS Reagent
Program for the reagents and cell lines. Authors are grateful to Massimo Pizzato for his critical
inputs and reagent support.

References

1.   Koonin E V., Dolja V V., Krupovic M. Origins and evolution of viruses of eukaryotes:
The ultimate modularity. Virology. 2015.

2.   Malfavon-Borja R, Sawyer SL, Wu LI, Emerman M, Malik HS. An evolutionary screen
highlights canonical and noncanonical candidate antiviral genes within the primate
TRIM gene family. Genome Biol Evol. 2013;

3.   Paparisto E, Woods MW, Coleman MD, Moghadasi SA, Kochar DS, Tom SK, et al.
Evolution-Guided Structural and Functional Analyses of the HERC Family Reveal an
Ancient Marine Origin and Determinants of Antiviral Activity. J Virol. 2018;

4.   Daugherty MD, Schaller AM, Geballe AP, Malik HS. Evolution-guided functional
analyses reveal diverse antiviral specificities encoded by ifit1 genes in mammals. Elife.
2016;

5.   Sackton TB, Lazzaro BP, Clark AG, Wittkopp P. Rapid expansion of immune-related
gene families in the house fly, musca domestica. Mol Biol Evol. 2017;

6.   Kasahara M. Genome Duplication and T Cell Immunity. In: Progress in molecular
biology and translational science. 2010. p. 7–36.

7.   Dehal P, Boore JL. Two Rounds of Whole Genome Duplication in the Ancestral
Vertebrate. Holland P, editor. PLoS Biol. 2005 Sep;3(10):e314.

8.   Duggal NK, Emerman M. Evolutionary conflicts between viruses and restriction factors
shape immunity. Nat Rev Immunol. 2012;12(10):687–95.

9.   Wlasiuk G, Khan S, Switzer WM, Nachman MW. A history of recurrent positive
selection at the toll-like receptor 5 in primates. Mol Biol Evol. 2009;

10.   Daugherty MD, Young JM, Kerns JA, Malik HS. Rapid Evolution of PARP Genes
Suggests a Broad Role for ADP-Ribosylation in Host-Virus Conflicts. PLoS Genet.
2014;

11.   Boso G, Buckler-White A, Kozak CA. Ancient Evolutionary Origin and Positive
Selection of the Retroviral Restriction Factor Fv1 in Muroid Rodents . J Virol. 2018;

12.   Compton AA, Malik HS, Emerman M. Host gene evolution traces the evolutionary
history of ancient primate lentiviruses. Philosophical Transactions of the Royal Society
B: Biological Sciences. 2013.

13.   Harris RS, Hultquist JF, Evans DT. The restriction factors of human immunodeficiency
virus. Journal of Biological Chemistry. 2012.

14.   McLaren PJ, Gawanbacht A, Pyndiah N, Krapp C, Hotter D, Kluge SF, et al.
Identification of potential HIV restriction factors by combining evolutionary genomic
signatures with functional analyses. Retrovirology. 2015;

15.   Firrito C, Bertelli C, Vanzo T, Chande A, Pizzato M. SERINC5 as a New Restriction
Factor for Human Immunodeficiency Virus and Murine Leukemia Virus. Annu Rev
Virol. 2018;

16.   Rosa A, Chande A, Ziglio S, De Sanctis V, Bertorelli R, Goh SL, et al. HIV-1 Nef
promotes infection by excluding SERINC5 from virion incorporation. Nature.
2015;526(7572):212–7.

17.   Usami Y, Wu Y, Göttlinger HG. SERINC3 and SERINC5 restrict HIV-1 infectivity and
are counteracted by Nef. Nature. 2015;526(7572):218–23.

18.   Chande A, Cuccurullo EC, Rosa A, Ziglio S, Carpenter S, Pizzato M. S2 from equine
infectious anemia virus is an infectivity factor which counteracts the retroviral inhibitors
SERINC5 and SERINC3. Proc Natl Acad Sci U S A. 2016;113(46).

19.   Schulte B, Selyutina A, Opp S, Herschhorn A, Sodroski JG, Pizzato M, et al.
Localization to detergent-resistant membranes and HIV-1 core entry inhibition
correlate with HIV-1 restriction by SERINC5. Virology. 2018;

20.   Sood C, Marin M, Chande A, Pizzato M, Melikyan GB. SERINC5 protein inhibits HIV-1
fusion pore formation by promoting functional inactivation of envelope glycoproteins. J
Biol Chem. 2017;292(14).

21.   Murrell B, Vollbrecht T, Guatelli J, Wertheim JO. The Evolutionary Histories of
Antiretroviral Proteins SERINC3 and SERINC5 Do Not Support an Evolutionary Arms
Race in Primates. J Virol. 2016 Jun 29;

22.   Shaw AE, Hughes J, Gu Q, Behdenna A, Singer JB, Dennis T, et al. Fundamental
properties of the mammalian innate immune system revealed by multispecies
comparison of type I interferon responses. PLoS Biol. 2017;

23.   Lynch M. The Age and Relationships of the Major Animal Phyla. Evolution (N Y).
1999;

24.   Ahmad I, Li S, Li R, Chai Q, Zhang L, Wang B, et al. The retroviral accessory proteins
S2, Nef, and glycoMA use similar mechanisms for antagonizing the host restriction
factor SERINC5. J Biol Chem. 2019;

25.   Han GZ, Worobey M. An endogenous foamy-like viral element in the coelacanth
genome. PLoS Pathog. 2012;

26.   Sawyer SL, Wu LI, Emerman M, Malik HS. Positive selection of primate
TRIM5 identifies a critical species-specific retroviral restriction domain. Proc Natl
Acad Sci. 2005;

27.   Newman RM, Hall L, Connole M, Chen G-L, Sato S, Yuste E, et al. Balancing
selection and the evolution of functional polymorphism in Old World monkey TRIM5 .
Proc Natl Acad Sci. 2006;

28.   McLaughlin RN, Gable JT, Wittkopp CJ, Emerman M, Malik HS. Conservation and
Innovation of APOBEC3A Restriction Functions during Primate Evolution. Mol Biol
Evol. 2016;

29.   Johnson WE. Rapid adversarial co-evolution of viruses and cellular restriction factors.
Curr Top Microbiol Immunol. 2013;

30.   Blanco-Melo D, Venkatesh S, Bieniasz PD. Origins and Evolution of tetherin, an
Orphan Antiviral Gene. Cell Host Microbe [Internet]. 2016 Aug 10 [cited 2017 Jul
5];20(2):189–201. Available from:
http://linkinghub.elsevier.com/retrieve/pii/S193131281630258X

31.   Meng X, Zhang F, Yan B, Si C, Honda H, Nagamachi A, et al. A paralogous pair of
mammalian host restriction factors form a critical host barrier against poxvirus
infection. PLoS Pathog. 2018;

32.   Browne EP, Littman DR. Species-Specific Restriction of Apobec3-Mediated
Hypermutation. J Virol. 2008;

33.   Mitchell PS, Patzina C, Emerman M, Haller O, Malik HS, Kochs G. Evolution-guided
identification of antiviral specificity determinants in the broadly acting interferon-
induced innate immunity factor MxA. Cell Host Microbe. 2012;

34.   Mitchell PS, Young JM, Emerman M, Malik HS. Evolutionary Analyses Suggest a
Function of MxB Immunity Proteins Beyond Lentivirus Restriction. PLoS Pathog. 2015;

35.   Dutko JA, Schäfer A, Kenny AE, Cullen BR, Curcio MJ. Inhibition of a yeast LTR
retrotransposon by human APOBEC3 cytidine deaminases. Curr Biol. 2005;

36. Irwin B, Aye M, Baldi P, Beliakova-Bethell N, Cheng H, Dou Y, et al. Retroviruses and
yeast retrotransposons use overlapping sets of host genes. Genome Res. 2005;

37. Miller JE, Zhang L, Jiang H, Li Y, Pugh BF, Reese JC. Genome-wide mapping of
decay factor-mRNA interactions in yeast identifies nutrient-responsive transcripts as
targets of the deadenylase Ccr4. G3 Genes, Genomes, Genet. 2018;

38. Llorens J V., Clark JB, Martínez-Garay I, Soriano S, De Frutos R, Martínez-Sebastián
MJ. Gypsy endogenous retrovirus maintains potential infectivity in several species of
Drosophilids. BMC Evol Biol. 2008;

39. Teysset L, Burns JC, Shike H, Sullivan BL, Bucheton A, Terzian C. A Moloney Murine
Leukemia Virus-Based Retroviral Vector Pseudotyped by the Insect Retroviral gypsy
Envelope Can Infect Drosophila Cells. J Virol. 1998;

40. Mellor J, Fulton SM, Dobson MJ, Wilson W, Kingsman SM, Kingsman AJ. A retrovirus-
like strategy for expression of a fusion protein encoded by yeast transposon Ty1.
Nature. 1985;

41. Aiewsakun P, Katzourakis A. Marine origin of retroviruses in the early Palaeozoic Era.
Nat Commun. 2017;

42. Beitari S, Ding S, Pan Q, Finzi A, Liang C. The effect of HIV-1 Env on SERINC5
antagonism. J Virol [Internet]. 2016;(December):JVI.02214-16. Available from:
http://jvi.asm.org/lookup/doi/10.1128/JVI.02214-16

43. Sood C, Marin M, Chande A, Pizzato M, Melikyan GB. SERINC5 protein inhibits HIV-1
fusion pore formation by promoting functional inactivation of envelope glycoproteins. J
Biol Chem [Internet]. 2017 Apr 7 [cited 2017 May 26];292(14):6014–26. Available
from: http://www.ncbi.nlm.nih.gov/pubmed/28179429

44. Pye, V.E., Rosa, A., Bertelli C et al. A bipartite structural organization defines the
SERINC family of HIV-1 restriction factors. Nat Struct Mol Biol. 2020;27:78–83.

45. Beitari S, Ding S, Pan Q, Finzi A, Liang C. Effect of HIV-1 Env on SERINC5
Antagonism. J Virol. 2017;91(4).

46. Sojo V, Dessimoz C, Pomiankowski A, Lane N. Membrane Proteins Are Dramatically
Less Conserved than Water-Soluble Proteins across the Tree of Life. Mol Biol Evol.
2016;

47. Li M, Waheed AA, Yu J, Zeng C, Chen HY, Zheng YM, et al. TIM-mediated inhibition
of HIV-1 release is antagonized by Nef but potentiated by SERINC proteins. Proc Natl
Acad Sci U S A. 2019;

48. Liu G, Zhang H, Zhao C, Zhang H. Evolutionary History of the Toll-Like Receptor
Gene Family across Vertebrates. Enard D, editor. Genome Biol Evol [Internet]. 2020
Jan 1 [cited 2020 Feb 7];12(1):3615–34. Available from:
https://academic.oup.com/gbe/article/12/1/3615/5652095

49. Simmonds P, Aiewsakun P, Katzourakis A. Prisoners of war — host adaptation and its
constraints on virus evolution. Nat Rev Microbiol. 2019;

50. Trobridge G, Josephson N, Vassilopoulos G, Mac J, Russell DW. Improved foamy
virus vectors with minimal viral sequences. Mol Ther. 2002;6(3):321–8.

51. Pizzato M, Erlwein O, Bonsall D, Kaye S, Muir D, McClure MO. A one-step SYBR
Green I-based product-enhanced reverse transcriptase assay for the quantitation of
retroviruses in cell culture supernatants. J Virol Methods [Internet]. 2009 Mar [cited
2017 Jun 19];156(1–2):1–7. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/19022294

52. Mittal P, Jaiswal SK, Vijay N, Saxena R, Sharma VK. Comparative analysis of
corrected tiger genome provides clues to its neuronal evolution. Sci Rep [Internet].
2019 Dec 5 [cited 2020 Jan 13];9(1):18459. Available from:
http://www.nature.com/articles/s41598-019-54838-z

53. Li H. [Heng Li - Compares BWA to other long read aligners like CUSHAW2] Aligning
sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv Prepr
arXiv. 2013;

54. Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of Next Generation
Sequencing Data. BMC Bioinformatics. 2014;

55. Thomas GWC, Hahn MW. Referee: Reference assembly quality scores. Genome Biol
Evol. 2019;

56. Shinde SS, Teekas L, Sharma S, Vijay N. Signatures of Relaxed Selection in the
CYP8B1 Gene of Birds and Mammals. J Mol Evol. 2019;

57. Blackburne BP, Whelan S. Class of multiple sequence alignment algorithm affects
genomic analysis. Mol Biol Evol. 2013;

58. Murrell B, Moola S, Mabona A, Weighill T, Sheward D, Kosakovsky Pond SL, et al.
FUBAR: A fast, unconstrained bayesian AppRoximation for inferring selection. Mol
Biol Evol. 2013;

59. Kosakovsky Pond SL, Frost SDW, Muse S V. HyPhy: Hypothesis testing using
phylogenies. Bioinformatics. 2005;

60. Avino M, Poon AFY. Detecting Amino Acid Coevolution with Bayesian Graphical
Models. In: Methods in Molecular Biology. 2019.

Figure Legends

Figure 1. The activity of *SERINC* paralogs and orthologs on HIV-1 infectivity

A) The activity of Human *SERINC* paralogs (*SERINC*1-*SERINC*5) on nef-defective HIV-
1 Infectivity. (n=4, mean ± s.d., unpaired t-test, (*)p<0.05 (**)p<0.01 (***)p<0.001, ns-
not significant). Lower panel, Western blot showing expression of c-terminally HA-
tagged *SERINC* and the corresponding B-actin from cell lysate (HEK293T). Values
obtained from the empty vector control was normalized to 100% for comparison with
SERINC expressors
B) Representative images of TZM-GFP cells infected with viruses produced for Fig-1A.
Cells are stained with Hoechst and captured at 10x magnification on CellInsight CX7
High Content Screening platform. Scale-100μm
C) Immunofluorescence assay for the indicated HA-tagged human *SERINC*1-5
transfected in JTAg *SERINC*5/3[-/-] cells and visualized using Alexa 488 secondary
antibody. Scale- 10μm
D) Hierarchical clustering of the arms-race signatures in primate genes aligned using
ClustalW aligner and used as input for FUBAR. Interferon-induced genes identified by
(Shaw et al 2017) are color-coded as red (upregulated), blue(downregulated), green
(*SERINC* paralogs), grey (other select genes)
E) The activity of pre-and-post-WGD *SERINC5* orthologs on nef-defective HIV-1
infectivity. (n=4, mean ± s.d., unpaired t-test, (*)p<0.05 (**)p<0.01 (***)p<0.001, ns-
not significant). Lower panel, Western blot showing expression of indicated orthologs
tagged with HA and the corresponding B-actin from cell lysate (HEK293T)
F) Representative images of TZM-GFP cells infected with viruses produced for Fig-1E.
Cells are stained with Hoechst and captured at 10x magnification on Cell Insight CX7
High Content Screening platform. Scale- 100μm

G) Immunofluorescence assay for indicated HA-tagged *SERINC* orthologs transfected in
626   JTAg *SERINC*5/3$^{-/-}$ cells and visualized using Alexa 488 secondary antibody. Scale-
627   10μm

Figure 2- Coelacanth *SERINC*2 inhibits HIV-1 infectivity

A. A Schematic timeline depicting the sequence of events during the course of evolution
with species depicted from left to right being *S. cerevisiae, D. melanogaster, L.*
chalumnae, X. tropicalis, G.gallus, E. caballus, M. musculus and *H. sapiens*
B. The activity of *SERINC*2 from coelacanth and human *SERINC*2 isoforms on nef-
defective HIV-1 infectivity. Values obtained from the empty vector control was
normalized to 100% for comparison with *SERINC* expressors. (n=4, mean ± s.d.,
unpaired t-test, (*) $p<0.05$ (**)$p<0.01$ (***)$p<0.001$, ns-not significant). Lower panel,
Representative images of TZM-GFP cells infected with viruses produced for top
panel. Cells are stained with Hoechst and captured at 10x magnification on
CellInsight CX7 High Content Screening platform.Scale-100μm
C. The activity of *SERINC*2 orthologs from coelacanth, xenopus, gallus, mus, equus and
human on nef-defective HIV-1 infectivity. Values obtained from the empty vector
control was normalized to 100% for comparison with *SERINC* expressors. (n=4,
mean ± s.d., unpaired t-test, (*)$p<0.05$ (**)$p<0.01$ (***)$p<0.001$, ns-not significant).
Lower panel, western blot showing expression of *SERINC*2 orthologs tagged with HA
and the corresponding B-actin from cell lysate (HEK293T).
D. Representative images of TZM-GFP cells infected with viruses produced for Fig-2C.
Cells are stained with Hoechst and captured at 10x magnification on Cell Insight CX7
High Content Screening platform. Scale-100μm
E. Immunofluorescence from for indicated HA-tagged *SERINC* orthologs transfected in
JTAg *SERINC*5/3$^{-/-}$ cells and visualized using Alexa 488 secondary antibody. Scale-
10μm

Figure 3- The ability of retroviral factors to antagonize *SERINC*s

A. The ability of Nef and glycoGag to counteract indicated *SERINC*2 orthologs. Human
*SERINC*5 was served as control for Nef and glycoGag counteraction of the
restriction. Values obtained from the empty vector control without Nef or glycoGag
was normalized to 100% for comparison with *SERINC* expressors
B. Representative images of TZM-GFP cells infected for viruses produced for Fig-3A.
Cells are stained with Hoechst and captured at 10x magnification on Cell Insight CX7
High Content Screening platform. Scale-100μm
C. The ability of nef alleles to antagonize indicated *SERINC*2 orthologs. Human
*SERINC*5 is used as a control
D. Susceptibility of HIV-1 clade-B (HXB2 and JR-FL), clade-C (ZM109F, CAP210) and
VSV envelope glycoprotein (VSV-G) to inhibition of infectivity by coelacanth
*SERINC*2 with human *SERINC*2 and *SERINC*5 as controls.
E. Phylogenetic tree depicting the relationship among various retroviruses. Unknown
retroviral interactions with *SERINC*s are shown in the black ellipse while the known
ones are in blue. Branches colored red are complex lentiviruses, blue are gamma
retroviruses and green are foamy viruses.

Figure 4- The effect of Coelacanth *SERINC*2 on Foamy virus infectivity and antagonism by
the envelop glycoprotein

A. Genome organization of an endogenous foamy virus (CoeFV) from coelacanth and
the prototype foamy virus (FV)

B. Infectivity of foamy virus-derived vector and a nef-defective HIV-1 in the presence of
indicated human *SERINC* paralogs and *SERINC*2 orthologs. Foamy virus and HIV-1
infectivity obtained from the empty vector control was normalized to 100% for
comparison with *SERINC* expressors

C. Nef-defective HIV-1 infectivity obtained from the empty vector control, normalized to
100% for comparison, and with human *SERINC*5. Each foamy virus plasmids

expressing different components (gag, pol and env) along with the transfer vector
(pδφ) were used for its ability to counteract human *SERINC*5. SIV MAC239 Nef was
used as a control. Lower panel, representative images of TZM-GFP cells infected
with viruses produced for Fig-4C. Cells are stained with Hoechst and captured at 10x
magnification on CellInsight CX7 High Content Screening platform. Scale-100μm
683

D. Nef-defective HIV-1 infectivity obtained from the empty vector control, normalized to
100% for comparison, and with Coelacanth *SERINC*2. Each foamy virus plasmids
expressing different components (gag, pol and env) along with the transfer vector
(pδφ) were used for its ability to counteract Coelacanth *SERINC*2. Lower panel,
representative images of TZM-GFP cells infected with viruses produced for Fig-4D.
Cells are stained with Hoechst and captured at 10x magnification on CellInsight CX7
High Content Screening platform. Scale-100μm

E. Western blot showing varying expression of Coelacanth *SERINC*2-HA (from PBJ6
and pcDNA vectors) and the corresponding B-actin from cell lysate (HEK293T) in the
presence and absence of Foamy virus envelope expressor.

F. The sensitivity of the foamy core to the action of coelacanth *SERINC*2. The foamy
core was trans-complemented with either Foamy envelope, HIV-1 envelope or HIV-1
envelope lacking a c-terminal domain and indicated SERINC was co-expressed to
check the ability of individual SERINC to restrict foamy virus. (n = 4, mean ± s.d.,
unpaired t-test).

Figure-5. Schematic representation of *SERINC*-retrovirus interactions

A. *SERINC* antagonists (Glycogag, Nef and S2) from distant retroviruses. An
interaction between known and hypothesized virulent factors with *SERINC* paralogs.
Solid lines show experimentally demonstrated interactions while the dotted lines
represent hypothesized interactions. Color-coding for extinct and extant virulent
factors (VF). Extinct VFs are shown in light blue, the extant VFs (Type-1) are in
purple and Type-2 are in black.

B. A proposed model for adaptive co-evolution among retroviral factors with SERINCs.
The evolution of patterns of interactions between viral factors and SERINCs have
diversified following post-whole-genome duplication.

Supplementary Figure Legends

Figure S1- Phylogenetic analysis of SERINC paralogs**.** *S. cerevisiae, C. elegans, and D.*
melanogaster has a single copy of this gene. Whole-genome duplication that occurred in an
early ancestor of mammals gave rise to a cluster of five genes divided into clusters of
SERINC1, SERINC2, SERINC3, and SERINC4 and SERINC5. Colors denote different
branches of each SERINC in chordates. The lengths of the triangles are proportional to the
number of nucleotide substitutions that have taken place in a particular branch. The tree was
generated using Ensembl.

Figure S2- Topology of human SERINC Paralogs predicted using TOPCONS

Figure S3- Arms-race signatures of primate SERINC genes inferred using FUBAR

Figure S4- Distribution of sites under different selection regimes across the SERINC genes.
The colors are inner (Blue), outer (Pink), helix (Red). Black dots are estimates of dS and red

dots are dN. Pairs of conditionally dependent sites identified by the program BGM (Bayesian
Graphical Model) are connected by a dotted line (see SERINC1 and SERINC4). Vertical
green lines correspond to the sites under purifying selection.

Figure S5- Effect of different aligners on hierarchical clustering of arms-race signatures of
primate genes. Interferon induced genes identified by (Shaw et al 2017) are color coded as
red (upregulated), blue(downregulated), green (SERINC paralogs), grey (other select genes)

Figure S6- Topology of SERINC5 orthologs predicted using Topcons. Coelacanth SERINC5
was reconstructed from the RNAseq data (*) see supplemental material for details

Figure S7- Topological features of Human SERINC2 splice isoforms and coelacanth
SERINC2 predicted using TOPCONS

Figure S8- Topology of SERINC2 orthologs predicted using TOPCONS

Figure S9- Incorporation of SERINC2 orthologs in the virus particles and detection by
indicated antibodies against target proteins

Figure S10- Effect of a dose-dependent expression of coelacanth SERINC2 on HIV-1
infectivity

Figure S11- Counteraction of coelacanth SERINC2 restriction by a Foamy virus envelope
glycoprotein

Figure S12- Reciprocal-packaging of a foamy envelope with an HIV core and sensitivity to
SERINCs

Figure S13- Infectivity of HIV-1 produced from JTAgSERINC5/3KO having ectopic
expression of the indicated SERINC

Figure S14- Multiple sequence alignment of SERINC2 orthologs; highlighted regions showing
the sites of sequence divergence

Figure S15- Evidence for the presence of HNF4a binding site in the intron of human
SERINC2 gene

Figure S16- Gene expression evolution of SERINC genes in vertebrates

Figure 1

Figure 2

A)

Whole Genome Duplication

~MY1000   800   500   450   400   350   300   96 90   40   0.08

TMS-1   SERINC1-5

B)

Infectivity (%)

*** ns ns ***

Vector   Coelacanth   Human   Human-201   Human SERINC5

SERINC2

Vector   Coelacanth   Human   Human-201   Human SERINC5

SERINC2

C)

Infectivity (%)

*** ** ns ns ns ns ***

Vector   Coelacanth   Xenopus   Gallus   Mus   Equus   Human-201   Human SERINC5

SERINC2

55kDa

15kDa

HA

Actin

D)

Vector   Coelacanth   Xenopus   Gallus   Mus   Equus   Human   Human SERINC5

SERINC2

E)

Coelacanth   Xenopus   Gallus   Mus   Equus   Human

SERINC2

Figure 3

Figure 4

Figure 5