IFN-λ4 is associated with increased risk and earlier occurrence of gastrointestinal,

respiratory and malarial infections in Malian children

[1]Ludmila Prokunina-Olsson, [2]Robert D. Morrison, [1]Adeola Obajemu, [3]Almahamoudou

Mahamar, [4]Sungduk Kim, [3]Oumar Attaher, [1]Oscar Florez-Vargas, [3]Youssoufa Sidibe,

[1]Olusegun O. Onabajo, [5]Amy A. Hutchinson, [5]Michelle Manning, [6]Jennifer Kwan, [7]Nathan

Brand, [3]Alassane Dicko, [2]Michal Fried, [4]Paul S. Albert, [8]Sam M. Mbulaiteye, [2]Patrick E.

Duffy

[1]Laboratory of Translational Genomics, Division of Cancer Epidemiology and Genetics,
National Cancer Institute, National Institutes of Health, Bethesda, 20892-4605, MD, USA
[2]Laboratory of Malaria Immunology and Vaccinology, National Institute of Allergy and
Infectious Diseases, National Institutes of Health, Rockville, MD 20852, USA
[3]Malaria Research & Training Center, Faculty of Medicine, Pharmacy and Dentistry,
University of Sciences Techniques and Technologies of Bamako, Bamako, Mali
[4]Biostatistics Branch, Division of Cancer Epidemiology and Genetics, National Cancer
Institute, National Institutes of Health, Bethesda, 20892-9776, MD, USA
[5]Cancer Genomics Research Laboratory, Frederick National Laboratory for Cancer
Research, Frederick, 21702, MD, USA
[6]Laboratory of Clinical Immunology and Microbiology, National Institute of Allergy and
Infectious Diseases, NIH, Bethesda, 20892, MD, USA.
[7]Department of Surgery, University of California San Francisco, San Francisco, 94143, CA,
USA
[8]Infections and Immunoepidemiology Branch, Division of Cancer Epidemiology and
Genetics, National Cancer Institute, National Institutes of Health, Bethesda, 20892-9776,
MD, USA
25

* Correspondence to:
Ludmila Prokunina-Olsson, PhD
Laboratory of Translational Genomics,
Division of Cancer Epidemiology and Genetics,
National Cancer Institute,
9615 Medical Center Dr,
Bethesda, MD 20892-9776
Tel:+1 240-760-6531; Fax: +1 240-541-4442
E-mail: prokuninal@mail.nih.gov
ORCID: 0000-0002-9622-2091

ABSTRACT

Genetic polymorphisms within the *IFNL3/IFNL4* genomic region, which encodes type III interferons, have been strongly associated with impaired clearance of hepatitis C virus (HCV) infection. We hypothesized that type III interferons might be important for the immune response to other pathogens as well. In a cohort of 914 Malian children, we analyzed episodes of malaria, gastrointestinal and respiratory infections using information for 30,626 clinic visits from birth through up to 5 years of follow-up. Genetic polymorphisms *IFNL4*-rs368234815 and *IFNL3*-rs4803217 that functionally affect type III interferons were genotyped with TaqMan assays. For both genetic variants and each infection, we evaluated time-to-first episode and calculated odds ratios (ORs) for the risk of an infection episode during follow-up, controlling for relevant covariates. Compared to children with the rs368234815-TT/TT genotype (IFN-λ4-Null), each copy of the rs368234815-dG allele was associated with an earlier first episode of a gastrointestinal infection (p=0.003) and respiratory infection (p=0.045). The risk of experiencing an infection episode during the follow-up was also significantly increased with each copy of the rs368234815-dG allele – for gastrointestinal infections (OR=1.53, 95%CI (1.13-2.07), p=0.005) and malaria (OR=1.30, 95%CI (1.02-1.65), p=0.033). *IFNL4*-rs368234815 and *IFNL3*-rs4803217 were in moderate linkage disequilibrium in this population (r^2=0.78), and all the associations for rs4803217 were weaker and lost significance after adjusting for rs368234815, implicating IFN-λ4 and not IFN-λ3 as the primary cause of these associations. We conclude that the ability to produce IFN-λ4 may have broad health-related implications by negatively affecting the immune response and clinical outcomes of several common infections.

Key words: IFNL4, IFN-λ4, interferon, immune response, infections

INTRODUCTION

The outcome of any infection is determined by a complex interplay between host and pathogen factors. One of the main host factors is the immune system, which can be modified by genetic polymorphisms in immune response genes. For example, variable ability to clear hepatitis C virus (HCV) infection spontaneously or after treatment has been significantly associated with polymorphisms in the genetic locus that encodes the family of type III interferons (IFNs) [1-5]. Type III IFNs provide a localized immune response at the sites of pathogen entry, such as at the epithelial surfaces of the respiratory and gastrointestinal tracts, as well as in the liver [6,7]. If this first line of defense provided by type III IFNs is successful, it decreases the necessity of a much stronger and systemic immune response provided by type I IFNs (IFNα and IFNβ) and type II IFN (IFN$\square$).

Humans have four type III IFNs, all encoded within the 70 kb region on chromosome 19. Three of these IFNs (IFN-λ1-3) can be inducibly generated in all the individuals, while IFN-λ4 is produced only in about 50% of the world population, including 90% of individuals of African ancestry, versus only 50% of Europeans and 10% of Asians [5,8]. The ability to produce IFN-λ4 is an ancestral trait, which appears to be under negative selection in humans [9]. The reasons for this selection are unclear but could be related to poor clearance of some deadly infections in the past. Thus, historic infections might have shaped the genetic landscape of this locus. On the other hand, the variable production of IFN-λ4 might have influenced and still be influencing susceptibility to infections and clinical outcomes.

The production of IFN-λ4 is controlled by a frameshift dinucleotide polymorphism, rs368234815-dG, within the first exon of the *IFNL4* gene. The open reading frame for IFN-λ4 protein is created by the dG allele, whereas the TT allele results in the truncated, non-functional protein (IFN-λ4-Null) [5]. Another polymorphism, rs12979860, also known as the *IL28B* (*IFNL3*) marker, is located within the first intron of *IFNL4* and often used as a proxy for the functional rs368234815 due to population-dependent moderate to high linkage disequilibrium between these variants. Although rs368234815-dG is generally considered the causal variant due to its direct functional effect on the production of IFN-λ4 protein [5], additional polymorphisms in the *IFNL3/IFNL4* region may also demonstrate significant associations because of considerable linkage disequilibrium. One of these linked variants is

a single nucleotide polymorphism (SNP) rs4803217, which is located within the 3'UTR of

IFNL3; this variant was proposed to affect IFN-λ3 protein by modulating the stability of

IFNL3 mRNA [10].

The strong associations between the *IFNL3*/*IFNL4* genetic variants and HCV clearance have

been well established and replicated by multiple studies. Additionally, a recent study

reported an association between the *IFNL4* genetic polymorphisms (rs12979860 and

rs368234815) and the clearance of respiratory RNA viruses in children from Rwanda [11]. To

further explore the hypothesis that type III IFNs might impact the immune response to

diverse pathogens, we analyzed detailed health records of 914 Malian children and

genotyped their DNA for the functional polymorphisms *IFNL4*-rs368234815 and *IFNL3*-

rs4803217. For these children, information for 30,626 scheduled and walk-in clinic visits

from birth through 1-5 years of follow-up was available for analysis.

We observed that compared to children genetically unable to produce IFN-λ4 (carriers of the

rs368234815-TT/TT genotype, IFN-λ4-Null), those with rs368234815-dG allele were

significantly more likely to have experienced at least one episode of gastrointestinal

infection (p=0.005) and malaria (p=0.033) during up to 5 years of follow-up, with an earlier

first episode of gastrointestinal infections (p=0.003) and respiratory infections (p=0.045).

The associations for *IFNL3*-rs4803217 were weaker and not significant after adjustment for

IFNL4-rs368234815. These results suggest the role of type III IFNs, and IFN-λ4,

specifically, in genetic control of the immune response to diverse pathogens of clinical

significance.

METHODS

The Mali longitudinal birth cohort

We used cord blood samples and data collected by the longitudinal birth cohort study

conducted in Ouelessebougou, Mali, between 2010 and 2016. The goal of the cohort was to

explore host and parasite factors that influence susceptibility to malaria infection and

disease during pregnancy and early childhood. The study enrolled pregnant women aged 15-

45 years presenting for antenatal consultations at the Ouelessebougou Community Health Center (CESCOM). The eligibility criteria included a residency in the district of Ouelessebougou for at least one year before enrollment and absence of a chronic, debilitating illness. Malaria history or diagnosis was not an exclusion criterion. Newborn children of the participating women were enrolled at delivery and were followed-up for 1-5 years. Before enrollment, written informed consent was obtained from the parents/guardians on behalf of their children after receiving written and oral study explanations from a study clinician in their native language.

The protocol and study procedures were approved by the institutional review board of the National Institute of Allergy and Infectious Diseases (NIAID) at the US National Institutes of Health (ClinicalTrials.gov ID NCT01168271), and the Ethics Committee of the Faculty of Medicine, Pharmacy and Dentistry at the University of Bamako, Mali.

The study site was located 80 km south of Bamako, an area of intense but highly seasonal malaria transmission. The follow-up protocol included visits to the clinic scheduled monthly during the high malaria transmission season (June-December) and every two months during the dry season (January-May), as well as *ad hoc* walk-in clinic visits for any acute symptoms. Clinical information was collected during all visits by project clinicians and recorded using standardized forms.

Per study protocol, malaria infection was tested at each visit and was defined as positivity for *P. falciparum* malaria parasite on thick blood smear microscopy, with or without clinical symptoms. Clinical malaria was defined as fever $>37.5^0$C, presence of malaria-related symptoms, and either a positive Rapid Diagnostic Test (RDT) for malaria antigens or visualized malaria parasites on thick blood smear microscopy. Severe malaria (SM) was defined as *P. falciparum* parasitemia and at least one of the following World Health Organization (WHO) criteria for SM - 2 or more convulsions in the past 24 hours, prostration (inability to sit unaided or in younger infants inability to move/feed), hemoglobin <5 g/dl, respiratory distress (hyperventilation with deep breathing, intercostal recessions and/or irregular breathing), or coma (Blantyre score <3). The presence of SM symptoms but with negative tests for malaria antigens or parasites was attributed to other diseases. All other parasitemia episodes, including those that were

asymptomatic, were classified as non-severe malaria (NSM). For ever/never analyses, the

NSM group included children with malaria history who never had an SM episode, while the

SM group included children with at least one SM episode, regardless of the number of NSM

episodes.

Gastrointestinal infections (GII) were defined by diarrhea; respiratory infections (RI) were

defined by cough and chest-related signs and symptoms (sputum production, dyspnea,

tachypnea, abnormal auscultatory findings). The etiology of these infections (bacterial, viral,

parasitic, fungal) was not determined. All children received treatment as clinically indicated

for their symptoms. Episodes of the same infection were considered independent if they

occurred more than 28 days apart.

DNA extraction and genotyping

Genomic DNA was extracted from cord blood clots using the QIAsymphony automation

system with protocol Blood_200_V7_DSP and the DSP DNA Mini Kit (Qiagen) by the

Cancer Genome Research Laboratory of the DCEG/NCI. The DNA samples were tested by

microsatellite fingerprinting (AmpFLSTR Indentifiler, ThermoFisher) to confirm gender

concordance with study records and exclude any mixed up or contaminated samples.

DNA samples were genotyped for *IFNL4*-rs368234815 and *IFNL3*-rs4803217 by custom

TaqMan genotyping assays as previously described [5,12], (**Figure S1**). *HBB* polymorphisms

rs334-T/A, *HBB*-Glu7Val (HbS, encoding hemoglobin variants A/A, A/S, and S/S) and

rs33930165, *HBB*-Glu7Lys (HbC, encoding hemoglobin variants A/A, A/C, and C/C) were

genotyped by Sanger sequencing as previously described [13], (**Figure S2**). Approximately

10% of all samples were genotyped in technical duplicates, which were 100% concordant

for all markers. The distributions of rs368234815 and rs4803217 did not deviate from

Hardy-Weinberg equilibrium (HWE). All genotyping and sequencing was done in the

Laboratory of Translational Genomics (LTG, DCEG, NCI) blinded to demographic and

clinical data. Of the 993 initial samples, 21 (2.1%) were excluded due to poor

quantity/quality of the extracted DNA. Of the remaining 972 samples, 38 samples (3.9%)

were excluded due to discordant *HBB* status between the study records (determined based

on gel electrophoresis) and DNA testing, 18 samples (1.9%) were excluded due to sex

discordance between study records and Identifiler profiles and 2 samples (0.2%) – due to

failed rs368234815 genotyping. The final set included 914 children with available genotype,

demographic and clinical data for 30,626 clinic visits; 447 (48.9%) and 467 (51.1%) of

children were females and males, respectively. The children were followed-up from birth for

an average of 147 weeks (range 52-264 weeks), with an average of 33.5 visits per child

(range 5-81 visits).

Statistical analysis

Haplotypes and linkage disequilibrium metrics (D' and r^2) between the markers were

analyzed and plotted using Haploview 4.2 [14]. The associations for *IFNL4*-rs368234815 and

IFNL3-rs4803217 were tested based on additive and genotypic genetic models, using the

TT/TT genotype or TT allele (IFN-λ4-Null) for rs368234815 and G/G genotype or G allele

for rs4803217 as reference groups.

The occurrence of infection episodes (never/ever) was analyzed for both polymorphisms

separately and in the same model using binary logistic regression models and adjusting for

the follow-up time (weeks). *HBB* status coded as 0, 1 and 2 for AA (wild-type) vs. sickle-

cell trait (A/S or A/C) vs. sickle-cell disease (S/S, S/C or C/C) was used as a covariate for

all malaria-related analyses; sex and birth month (as a proxy for seasonal effects) were

tested but not significantly associated with any outcomes (data not shown). Logistic

regression models were used to evaluate the associations of *IFNL4*-rs368234815 with

infections of interest adjusting for the occurrence of other infections (ever/never) to examine

whether these associations were specific to particular infection types.

Kaplan-Meyer plots and Cox proportional hazards regression models were used to analyze

time-to-first episode of each infection, with P-values calculated per rs368234815-dG and

rs4803217-T alleles. The number of infection episodes during follow-up was analyzed using

Poisson models with overdispersion (Quasi-likelihood), adjusting for the length of follow-

up as an off-set in the model (offset = log(LastVisitAge)).

Since the study did not have permission to perform genome-wide analysis, we used self-

reported paternal tribe as a proxy to control for possible population stratification. Of the 21

tribes, four tribes accounted for 92.5% of all the children: Bamana - 72.3%, Peulh/Fulani -

9.2%, Soninke - 6.0%, and Malinke - 5.0%. The reported paternal tribe (for all 21 tribes)

was used as a random effect variable for logistic regression analyses (glmer package), Cox

regression analyses with gamma frailty, and the analyses of infection episode counts per

year (glmmPQL package). All the analyses were performed using specified packages in R

(https://cran.r-project.org/). The results were not adjusted for multiple testing.

RESULTS

Analysis of infection episodes in the Malian children

We analyzed information for 914 children based on health records for each of the 30,626

routine and walk-in clinic visits from birth through up to 5 years of follow-up. During

follow-up, 76.1% of children had a malaria episode, including 11.1% children that had at

least one SM episode (regardless of NSM episodes) and 65.0% of children that had only an

NSM episode (and never an SM episode); 88.1% and 97.7% children experienced a GII and

RI episode, respectively.

Episodes of any infection (malaria, GII or RI) were reported at 12.7% of the routine and

76.3% of the walk-in visits (**Table 1**). Thus, most infection episodes were acute and

sufficiently severe to require an emergency visit to the clinic without waiting for the next

regular scheduled check-up visit. The overall frequency of malaria infection at any visit was

10.6% (5.4% at routine and 19.9% at walk-in visits); GII was reported at 8.4% of all visits,

(1.8% at routine and 20.2% at walk-in visits), and RI was reported at 21.0% of all visits

(6.5% at routine and 47.4% at walk-in visits). No infection was reported at most visits

(n=19,808, 64.7%), while one and two infection types were reported at 9,407 (30.7%) and

1,380 (4.5%) visits, respectively. Only 31 visits (0.1%) reported all three infection types at

the same visit.

The most likely co-infection combination was malaria and RI, with 5.3% of routine and

8.2% of walk-in visits, and the least likely co-infection combination was malaria and GII,

with 0.8% of routine and 1.6% of walk-in visits (**Table S1**). Logistic regression analyses

showed that infections tended to be non-redundant, and any infection significantly decreased

the risk of any other infection reported at the same visit (**Table S2**).

 Distribution of the HBB, IFNL4 and IFNL3 polymorphisms in the Malian children and the

populations of the 1000 Genomes Project

The *HbB*-A/A genotype, representing wild-type hemoglobin, was present in 80.5% of

children; sickle cell trait (SCT, HbS-A/S or HbC-A/C) – in 18.6%, and sickle cell disease –

(SCD, S/S, S/C or C/C) - in 0.9% of the children. The *IFNL4*-rs368234815 alleles (dG-

72.2% and TT-28.8%) and *IFNL3*-rs4803217 alleles (G-66.7% and T-33.3%) were in

moderate linkage disequilibrium (LD) in the total set of 914 Malian children (D'=0.98 and

r^2=0.78), which was also reflected in the haplotype frequencies of these variants (**Figure**

S3). The allele and genotype frequencies of *HBB, IFNL3* and *IFNL4* polymorphisms were

comparable to those in the populations of African ancestry in the 1000 Genomes Project,

especially to the populations from West-Africa (**Table S3**). The distribution of

alleles/genotypes of these variants in the four main tribes in Mali is also presented (**Table**

S3).

Associations between IFNL4 and IFNL3 polymorphisms and infection episodes in the

Malian children

We analyzed the risk of an infection episode reported at any clinic visit from birth through

up to 5 years of follow-up (ever/never analysis), and present the results adjusted for the

tribes; unadjusted results are also presented in corresponding tables/figures. The

rs368234815-dG allele was associated with an increased risk of GII (with a per-allele

OR=1.53, 95%CI (1.13-2.07), p=0.005), malaria overall (OR=1.30, 95%CI (1.02-1.65),

p=0.033), NSM (OR=1.28, 95%CI (1.00-1.63), p=0.047), but not of RI (OR=1.20, 95%CI

(0.63-2.29, p=0.58) (**Table 2**).

Gastrointestinal and respiratory symptoms might overlap with clinical manifestations of

malaria. However, the associations of *IFNL4*-rs368234815 with these infections appeared

independent, as mutual conditioning on these infections did not significantly alter the results

(**Table S4**). We also analyzed the association between *IFNL4*-rs368234815 and time-to-

first episode of each infection, which represents the earliest occurrence of symptoms of

sufficient severity to require a visit to the clinic. The most significant association was

observed for the *IFNL4*-rs368234815 and the time to the first episode of GII (per allele

p=0.003) (**Figure 1**). The association was marginally significant for RI (p=0.045), but not

for malaria overall and SM (**Figure 1**). The duration of follow-up did not differ significantly

for children with different *IFNL4* genotypes (**Table S5**).

The number of episodes reported for each infection during follow-up was the lowest for

children with the IFN-λ4-Null genotype, but this was significant only in the dominant

genetic model for GII (p=0.03, **Table S6**), suggesting that the increased susceptibility is

related to infection overall and not its recurrence. All the associations for *IFNL3*-rs4803217

were weaker than for rs368234815 (**Table S7**, **Figure S4**) and lost significance in the

models that included both variants, while most of the associations for rs368234815

remained significant (**Table 3**).

DISCUSSION

In 2015, 5.9 million children died globally before the age of five, with 30% of these deaths

being caused by pneumonia, malaria and diarrhea [15]. Thus, it is critical to identify and

understand the factors affecting susceptibility and the host response to these infections. We

identified the genetic ability to produce IFN-λ4, a type III IFN, in carriers of the *IFNL4*-

rs368234815-dG allele, as a factor most significantly increasing the risk of gastrointestinal

infections and earlier occurrence of these infections in Malian children under five. We also

observed evidence for higher risk of malaria infection and earlier episodes of respiratory

infections in carriers of the same allele.

Episodes of acute gastroenteritis (diarrhea and vomiting) are commonly caused by

infections with rotavirus (RoV) or norovirus (NoV). Rotavirus is considered the most

important global cause of severe diarrhea in children under five. In 2016 alone, despite more

than 28,000 deaths considered to be averted by vaccination for rotavirus, still, an estimated

128,500 children under five died of this infection [16]. In Rwanda, rotavirus was detected in

rectal swabs of 36.9% of children under five seeking clinical help for acute gastroenteritis [17].

Norovirus is the second leading cause of acute gastroenteritis, accounting for up to 20% of

all acute infections in all age groups worldwide [18]. Although these infections are ubiquitous

and in healthy individuals spontaneously clear within days, viral clearance may take longer

in younger children, especially in those affected by malnutrition and other infections.

Extended severe dehydration due to diarrhea can be fatal, especially in infants. Human

enteric viruses are difficult to study *in vitro*, but recent advances in human intestinal

organoid models [19,20] should facilitate these studies. In contrast to humans that can produce

either 3 or 4 type III IFNs (IFN-λ1, 2, 3 in all and IFN-λ4 - in a subset of individuals), the

mouse genome invariably encodes only IFN-λ2 and IFN-λ3, which are collectively referred

to as IFN-λ. Although most conclusions on RoV and NoV infections have been drawn using

murine viruses and based on mouse models that differ from human cells by the IFN-λ

repertoire, these studies proposed that intact signaling through type III IFNs is important for

the control of GII. These conclusions were supported by the functional impact caused by the

elimination of IFNLR1, which is a receptor expressed by epithelial cells and specific to the

type III IFN family [21-26]. However, the IFN-λ-related immune responses to enteric viruses

appeared to be viral-strain and mouse-model specific [22], and could also be affected by the

intestinal microbiota [26,27]. Overall, clearance or persistence of enteric viruses likely results

from the combination of factors contributed both by the host and the pathogen.

Although data on GII etiology was not available in our study, a large subset of these acute

gastroenteritis episodes are likely to be caused by RoV and NoV - RNA viruses already

linked with type III IFNs in animal models or *in vitro* studies. A higher risk and earlier

episodes of clinical GII in children with *IFNL4*-rs368234815-dG allele might indicate an

impaired immune response in these children due to the production of IFN-λ4. However, if

diarrhea can be considered as a mechanism to purge those pathogens from the GI tract, these

results might also indicate an enhanced innate immune response, as has been suggested

based on mouse studies [16] and Kotenko, personal communication. The immune response

altered in children with *IFNL4*-rs368234815-dG allele might manifest in more severe

episodes od diarrhea but our study did not have information on the duration and the intensity

of each infection episode. Either impaired or enhanced, the innate immune response

associated with the genetic ability to produce IFN-λ4 appears to have negative health

consequences. With the associations for GII being in the same direction as was previously

reported for HCV and respiratory viruses, GII can be added to the list of outcomes

negatively affected by the ability to produce IFN-λ4.

Acute RI cause 1.9 million deaths annually in children under five, with 70% of these deaths

occurring in Africa and Southeast Asia [28]. Several studies have explored the role of type III

IFNs in the immune response to the respiratory syncytial virus (RSV) and influenza in the

human and murine nasal epithelium. These studies suggested that type III IFNs represent the

first line of defense from respiratory RNA viruses, and treatment with recombinant IFN-λs

can protect from RI and stop the viral spread from upper to lower respiratory tract [29-32].

Recently, a genetic association was reported for *IFNL4*-rs12979860-T and rs368234815-dG

alleles with impaired clearance of respiratory RNA viruses in nasal swabs of children from

Rwanda [11]. The observed effect was in the same direction as for clearance of HCV reported

in many studies, suggesting that the negative impact of IFN-λ4 is relevant for the clearance

of respiratory RNA viruses as well. In our study, we observed a borderline significant

association (p=0.045) with an earlier occurrence of RI in children with rs368234815-dG

allele but the association with increased susceptibility to RI in these children did not reach

statistical significance. This could be because respiratory symptoms recorded during clinic

visits in our study were very common (were experienced by 97.7% of children during

follow-up), and occurred very early in life, with the median timing of the first RI episode

being at 16 weeks of age. Furthermore, RI episodes were defined very broadly, without

stratifying by specific viral etiology, as was done in the study in Rwanda [11]. In addition to

bacterial and viral etiology, respiratory symptoms might also be caused by helminths

(pinworms, roundworms, etc) transiting through the lungs during their life cycle. A more

specific definition of RI etiology and surveillance of clearance is needed to clarify the

association with genetic variants in the *IFNL3*/*IFNL4* region.

Our results showing the association between type III IFNs and malaria infection are also

novel and, if validated, might present new mechanisms. Although III IFNs are mostly

recognized for their antiviral role, the involvement of IFN-λs in bacterial and fungal

infections have also been reported [33]. Malaria is caused by plasmodium, an intracellular

parasite, which first invades hepatocytes and then propagates in the liver where type III

IFNs can be expressed after induction by different stimuli. However, these results were

attenuated by adjusting for the tribe, suggesting possible differences between these tribes,

such as malaria exposure and protection, that should be carefully considered as confounding

factors in genetic association studies. Most of our results remained significant after

adjustment for the tribe, warranting further replication studies.

While rs4803217 was previously suggested as a functional variant affecting the stability of

IFNL3 mRNA and possibly accounting for the association in the *IFNL3/IFNL4* region [10],

this SNP appeared to be associated with all the outcomes in our study weaker than

rs368234815, supporting *IFNL4*-rs368234815 as a primary associated and functional

genetic variant in this region.

The strength of our study is the longitudinal analysis of a cohort of children who were

regularly examined from birth to up to 5 years of age, a large sample size with 30,626 clinic

visits that captured most, if not all, episodes of common infections. Longitudinal studies of

transient but recurrent infectious diseases with high spontaneous clearance are more

informative than cross-sectional studies, which may miss a significant proportion of events

not detectable at the time of the study. The limitations of our study include a broad

definition of gastrointestinal and respiratory symptoms without specifying their infectious

etiology. Due to the regular medical assistance provided to the children at the time of the

routine and walk-in visits, the negative health impacts of these infections were likely to be

diminished, which makes our findings conservative. Even though the rs368234815-dG allele

was associated with a higher risk of these infections in early childhood, our study did not

explore the long-term health consequences of these associations. For example, we were not

able to test whether earlier exposure to infections would affect immune fitness later in life.

Additionally, our results do not explain why the ability to produce IFN-λ4 could be

beneficial in Africa, accounting for its very high frequency, while this allele being selected

against outside of Africa. Our exploratory study tested associations between multiple

infections and models, which might result in some of these results being false-positive due

to multiple testing and unadjusted confounding factors. Further studies in comparable

cohorts are needed to confirm our findings and refine the etiology of these and other

infections in different environmental and socio-economic settings.

In conclusion, our study suggests that type III IFNs, and IFN-λ4, specifically, negatively

affect the immune response to infections in young children in Africa. Clearance of several

RNA viruses likely impaired by IFN-λ4 (HCV, respiratory viruses, RoV and NoV) suggests

that response to other clinically relevant RNA viruses, such a coronavirus, including SARS-

CoV-2 that causes COVID-19, might also be impaired in carriers of the *IFNL4-*

rs368234815-dG allele. These functional associations might contribute to the variable

severity of clinical manifestations and outcomes in the infected patients, warranting further

investigations.

ACKNOWLEDGMENTS

We thank the study participants, the populations and health staff in the study area for their

cooperation. We also thank the Mali Service Center for providing administrative support to

the project, Drs. Richard Sakai and Souleymane Karambe for logistical support. We thank

Dr. Sergei Kotenko, Rutgers University and Thomas O'Brien, Infections and

Immunoepidemiology Branch, Division of Cancer Epidemiology and Genetics, National

Cancer Institute for critical comments and discussions.

Declaration of interests

L.P.-O. is a co-inventor on IFN-λ4-related patents issued to NCI/NIH, and receives

royalties for monoclonal antibodies for IFN-λ4 detection. Other authors have no conflict of

interest to declare.

Funding sources

The study was supported by the Intramural Research Program of the Division of Cancer

Epidemiology and Genetics, National Cancer Institute (NCI), the Intramural Research

Program, National Institute of Allergy and Infectious Diseases (NIAID), National Institutes

of Health, US Department of Health and Human Services. The content of this publication

does not necessarily reflect the views or policies of the Department of Health and Human

Services, nor does mention of trade names, commercial products, or organizations imply

endorsement by the US Government. The content of this manuscript is the sole

responsibility of the authors. The sponsors had no role in the study design, data collection,

analysis, interpretation, writing of the manuscript, or the decision to submit the paper for

publication. The corresponding author had full access to all the data in the study and had

final responsibility for the decision to submit for publication.

Author contributions

LPO, SMM and PED conceived the idea and supervised the study; AM, OA, YS, JK, AD,

RDM, MF and PED provided samples and data; AO, AAH and MM performed DNA

extraction and genotyping of the samples; SK, PSA, RDM, OFV and LPO analyzed the data

and generated figures and tables; LPO drafted the manuscript; RDM, OOO, NB, AD, SMM

and PED helped with critical editing. All authors contributed to the interpretation of the

results and read and approved the final manuscript.

 TABLES

Table 1. Distribution of clinic visits for 914 children from the Mali birth cohort study

	Clinic visits, n			
	All visits#	Routine	Walk-in	Other
	30,626	18,654	10,884	1,088
Visits reporting any infection, n,	10,818	2,377	8,303	138
% of all visits	35.3	12.7	76.3	12.7
Visits with malaria, n,	3,260	1,015	2,167	78
% of malaria visits	100.0	31.1	66.5	2.4
% of all infections	10.6	5.4	19.9	7.2
Visits with GII, n	2,563	342	2204	17
% of GII visits	100.0	13.3	86.0	0.7
% of all infections	8.4	1.8	20.2	1.6
Visits with RI, n	6,437	1,221	5,161	55
% of RI visits	100.0	19.0	80.2	0.9
% of all infections	21.0	6.5	47.4	5.1

GII – gastrointestinal infections; RI – respiratory infections; Malaria – positivity by blood smear test with/without additional clinical symptoms, includes both severe (SM) and non-severe malaria (NSM).

Table 2. Associations between *IFNL4*-rs368234815 and infections in 914 children from the Mali birth cohort study

Infection	Total % of total	rs368234815 genotypes				rs368234815 alleles		Ever/never*		Ever/never adj$	
		Total n, %	TT/TT n , %	TT/dG n , %	dG/dG n , %	TT n, %	dG n, %	OR	P-val	OR	P-val
		914	77	373	464	527	1301				
			8.4	40.8	50.8	28.8	71.2				
Malaria#	Never 23.9	218	24 11.0	99 45.4	95 43.6	147 33.7	289 66.3	Ref		Ref	
	Any 76.1	696	53 7.6	274 39.4	369 53.0	380 27.3	1012 72.7	**1.32 (1.04-1.68)**	**0.022**	**1.30 (1.02-1.65)**	**0.033**
	SM 11.2	102	4 3.9	44 43.1	54 52.9	52 25.5	152 74.5	**1.53 (1.01-2.30)**	**0.044**	1.48 (0.98-2.24)	0.061
	NSM 65.0	594	49 8.3	230 38.7	315 53.0	328 27.6	860 72.4	**1.30 (1.02-1.65)**	**0.035**	**1.28 (1.00-1.63)**	**0.047**
GII	Never 11.9	109	16 14.7	49 45.0	44 40.4	81 37.2	137 62.8	Ref		Ref	
	Ever 88.1	805	61 7.6	324 40.3	420 52.2	446 27.7	1164 72.3	**1.50 (1.11-2.03)**	**0.0079**	**1.53 (1.13-2.07)**	**0.0054**
RI	Never 2.3	21	4 19.1	7 33.3	10 47.6	15 35.7	27 64.3	Ref		Ref	
	Ever 97.7	893	73 8.2	366 41.0	454 50.8	512 28.7	1274 71.3	1.20 (0.63-2.29)	0.58	1.20 (0.63-2.28)	0.58

*Binary logistic regression analysis based on additive genetic models and adjusting for the duration of follow-up (weeks); #results for malaria are also adjusted for *HBB* status; $ - additionally adjusting for tribes; GII – gastrointestinal infections; RI – respiratory infections; any malaria – positivity by blood smear test with/without additional clinical symptoms; SM – severe malaria; NSM – non-severe malaria; in bold – significant associations.

Table 3. Associations between *IFNL4*-rs368234815 and *IFNL3*-rs4803217 with infections in 914 children from the Mali birth cohort study

Infection		*IFNL4*-rs368234815		*IFNL3*-rs4803217		Models with both variants$			
						rs368234815		rs4803217	
		OR, 95%CI	P-val	OR, 95%CI	P-val	OR, 95%CI	P-val	OR, 95%CI	P-val
Malaria #	Never	Ref		Ref		Ref		Ref	
	Any	**1.30 (1.02-1.65)**	**0.033**	1.19 (0.94-1.50)	0.15	**1.73 (0.96-3.10)**	**0.068**	0.74 (0.42-1.31)	0.30
	SM	**1.48 (0.98-2.24)**	**0.061**	**1.52 (1.01-2.27)**	**0.044**	1.04 (0.37-2.88)	0.94	1.46 (0.54-3.98)	0.45
	NSM	**1.28 (1.00-1.63)**	**0.047**	1.15 (0.91-1.46)	0.25	**1.87 (1.03-3.38)**	**0.039**	0.67 (0.37-1.19)	0.17
GII	Never	Ref							
	Ever	**1.53 (1.13-2.07)**	**0.0054**	**1.35 (1.00-1.82)**	**0.047**	**2.34 (1.06-5.19)**	**0.036**	0.63 (0.29-1.39)	0.26
RI	Never	Ref							
	Ever	1.2 (0.63-2.28)	0.58	1.24 (0.66-2.33)	0.51	0.94 (0.23-3.90)	0.93	1.31 (0.32-5.31)	0.71

*Binary logistic regression analysis based on additive genetic models and adjusting for the duration of follow-up (weeks); in bold – significant associations; #results for malaria are adjusted for *HBB* status, $ - results for models including both variants; any malaria – positivity by blood smear test with/without additional clinical symptoms; SM – severe malaria; NSM – non-severe malaria. $-additionally adjusting for tribes.

Figure 1. Time-to-first episode of infections in relation to *IFNL4*- rs368234815 polymorphism in 914 children from the Mali birth cohort study. Malaria – positivity by blood smear test with/without additional clinical symptoms, includes both severe (SM) and non-severe malaria; GII – gastrointestinal infections; RI – respiratory infections. The plots are for Kaplan-Meyer analysis, P-values are for Cox proportional hazards regression models with per-allele linear trends, for malaria – adjusting for *HBB* status. Padj-values are additionally adjusting for tribes.

REFERENCES

1. Ge, D. *et al.* Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature* **461**, 399-401 (2009).
2. Thomas, D.L. *et al.* Genetic variation in IL28B and spontaneous clearance of hepatitis C virus. *Nature* **461**, 798-801 (2009).
3. Suppiah, V. *et al.* IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet* **41**, 1100-4 (2009).
4. Tanaka, Y. *et al.* Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat Genet* **41**, 1105-9 (2009).
5. Prokunina-Olsson, L. *et al.* A variant upstream of IFNL3 (IL28B) creating a new interferon gene IFNL4 is associated with impaired clearance of hepatitis C virus. *Nat Genet* **45**, 164-71 (2013).
6. Wack, A., Terczynska-Dyla, E. & Hartmann, R. Guarding the frontiers: the biology of type III interferons. *Nat Immunol* **16**, 802-9 (2015).
7. Lazear, H.M., Nice, T.J. & Diamond, M.S. Interferon-lambda: Immune Functions at Barrier Surfaces and Beyond. *Immunity* **43**, 15-28 (2015).
8. Prokunina-Olsson, L. Genetics of the Human Interferon Lambda Region. *J Interferon Cytokine Res* (2019).
9. Key, F.M. *et al.* Selection on a variant associated with improved viral clearance drives local, adaptive pseudogenization of interferon lambda 4 (IFNL4). *PLoS Genet* **10**, e1004681 (2014).
10. McFarland, A.P. *et al.* The favorable IFNL3 genotype escapes mRNA decay mediated by AU-rich elements and hepatitis C virus-induced microRNAs. *Nat Immunol* **15**, 72-9 (2014).
11. Rugwizangoga, B. *et al.* IFNL4 Genotypes Predict Clearance of RNA Viruses in Rwandan Children With Upper Respiratory Tract Infections. *Front Cell Infect Microbiol* **9**, 340 (2019).
12. O'Brien, T.R. *et al.* Comparison of functional variants in IFNL4 and IFNL3 for association with HCV clearance. *J Hepatol* **63**, 1103-10 (2015).
13. Legason, I.D. *et al.* Evaluating the Causal Link Between Malaria Infection and Endemic Burkitt Lymphoma in Northern Uganda: A Mendelian Randomization Study. *EBioMedicine* **25**, 58-65 (2017).
14. Barrett, J.C., Fry, B., Maller, J. & Daly, M.J. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* **21**, 263-5 (2005).
15. Liu, L. *et al.* Global, regional, and national causes of under-5 mortality in 2000-15: an updated systematic analysis with implications for the Sustainable Development Goals. *Lancet* **388**, 3027-3035 (2016).
16. Troeger, C. *et al.* Rotavirus Vaccination and the Global Burden of Rotavirus Diarrhea Among Children Younger Than 5 Years. *JAMA Pediatr* **172**, 958-965 (2018).
17. Kabayiza, J.C. *et al.* Diarrhoeagenic microbes by real-time PCR in Rwandan children under 5 years of age with acute gastroenteritis. *Clin Microbiol Infect* **20**, O1128-35 (2014).
18. Ahmed, S.M. *et al.* Global prevalence of norovirus in cases of gastroenteritis: a systematic review and meta-analysis. *Lancet Infect Dis* **14**, 725-730 (2014).
19. Saxena, K. *et al.* A paradox of transcriptional and functional innate interferon responses of human intestinal enteroids to enteric virus infection. *Proc Natl Acad Sci U S A* **114**, E570-E579 (2017).
20. Ettayebi, K. *et al.* Replication of human noroviruses in stem cell-derived human enteroids. *Science* **353**, 1387-1393 (2016).

21. Nice, T.J. *et al.* Interferon-lambda cures persistent murine norovirus infection in the absence of adaptive immunity. *Science* **347**, 269-73 (2015).

22. Lin, J.D. *et al.* Distinct Roles of Type I and Type III Interferons in Intestinal Immunity to Homologous and Heterologous Rotavirus Infections. *PLoS Pathog* **12**, e1005600 (2016).

23. Hernandez, P.P. *et al.* Interferon-lambda and interleukin 22 act synergistically for the induction of interferon-stimulated genes and control of rotavirus infection. *Nat Immunol* **16**, 698-707 (2015).

24. Pott, J. *et al.* IFN-lambda determines the intestinal epithelial antiviral host defense. *Proc Natl Acad Sci U S A* **108**, 7944-9 (2011).

25. Baldridge, M.T. *et al.* Expression of Ifnlr1 on Intestinal Epithelial Cells Is Critical to the Antiviral Effects of Interferon Lambda against Norovirus and Reovirus. *J Virol* **91**(2017).

26. Baldridge, M.T. *et al.* Commensal microbes and interferon-lambda determine persistence of enteric murine norovirus infection. *Science* **347**, 266-9 (2015).

27. Walker, F.C. & Baldridge, M.T. Interactions between noroviruses, the host, and the microbiota. *Curr Opin Virol* **37**, 1-9 (2019).

28. Williams, B.G., Gouws, E., Boschi-Pinto, C., Bryce, J. & Dye, C. Estimates of world-wide distribution of child deaths from acute respiratory infections. *Lancet Infect Dis* **2**, 25-32 (2002).

29. Okabayashi, T. *et al.* Type-III interferon, not type-I, is the predominant interferon induced by respiratory viruses in nasal epithelial cells. *Virus Res* **160**, 360-6 (2011).

30. Klinkhammer, J. *et al.* IFN-lambda prevents influenza virus spread from the upper airways to the lungs and limits virus transmission. *Elife* **7**(2018).

31. Davidson, S. *et al.* IFNlambda is a potent anti-influenza therapeutic without the inflammatory side effects of IFNalpha treatment. *EMBO Mol Med* **8**, 1099-112 (2016).

32. Galani, I.E. *et al.* Interferon-lambda Mediates Non-redundant Front-Line Antiviral Protection against Influenza Virus Infection without Compromising Host Fitness. *Immunity* **46**, 875-890 e6 (2017).

33. Kotenko, S.V., Rivera, A., Parker, D. & Durbin, J.E. Type III IFNs: Beyond antiviral protection. *Semin Immunol* **43**, 101303 (2019).