TITLE: Predicting *Vibrio cholerae* infection and disease severity using metagenomics in a

prospective cohort study

AUTHORS: Inès Levade[1], Morteza M. Saber[1], Firas Midani[2,3,4], Fahima Chowdhury[5], Ashraful

I. Khan[5], Yasmin A. Begum[5], Edward T. Ryan[6,7,9], Lawrence A. David[2,3,4,8], Stephen B.

Calderwood[6,7,10], Jason B. Harris[6,11], Regina C. LaRocque[6], Firdausi Qadri[5], B. Jesse

Shapiro[1,12,13*], Ana A. Weil[14]

[1]Department of Biological Sciences, University of Montreal, Montreal, QC, Canada.

[2]Program in Computational Biology and Bioinformatics, Duke University, Durham, NC, USA

[3]Center for Genomic and Computational Biology, Duke University, Durham, NC, USA

[4]Department of Molecular Genetics and Microbiology, Duke University, Durham, NC, USA

[5]Center for Vaccine Sciences, International Centre for Diarrhoeal Disease Research, Dhaka,

Bangladesh

[6]Division of Infectious Diseases, Massachusetts General Hospital, Boston, MA, USA

[7]Department of Medicine, Harvard Medical School, Boston, MA USA

[8]Department of Biomedical Engineering, Duke University, Durham, NC, USA

[9]Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public

Health, Boston, MA, USA

[10]Department of Microbiology, Harvard Medical School, Boston, MA USA

[11]Department of Pediatrics, Harvard Medical School, Boston, MA, USA

[12]Department of Microbiology and Immunology, McGill University, Montreal, QC, Canada

[13]McGill Genome Centre, Montreal, QC, Canada

[14]Division of Allergy and Infectious Diseases, University of Washington, Seattle, WA, USA

CORRESPONDING AUTHOR : jesse.shapiro@mcgill.ca

RUNNING TITLE: Stool metagenomics predicts cholera

SUMMARY: Cholera infection and disease severity can be predicted using metagenomic

sequencing of the gut microbiome pre-infection in a prospective cohort, and suggests potentially

protective bacterial species and genes.

ABSTRACT

Background: Susceptibility to *Vibrio cholerae* infection is impacted by blood group, age, and pre-existing immunity, but these factors only partially explain who becomes infected. A recent study used 16S rRNA amplicon sequencing to quantify the composition of the gut microbiome and identify predictive biomarkers of infection with limited taxonomic resolution.

Methods: To achieve increased resolution of gut microbial factors associated with *V. cholerae* susceptibility and identify predictors of symptomatic disease, we applied deep shotgun metagenomic sequencing to a cohort of household contacts of patients with cholera.

Results: Using machine learning, we resolved species, strains, gene families, and cellular pathways in the microbiome at the time of exposure to *V. cholerae* to identify markers that predict infection and symptoms. Use of metagenomic features improved the precision and accuracy of prediction relative to 16S sequencing. We also predicted disease severity, although with greater uncertainty than our infection prediction. Species within the genera *Prevotella* and *Bifidobacterium* predicted protection from infection, and genes involved in iron metabolism also correlated with protection.

Conclusion: Our results highlight the power of metagenomics to predict disease outcomes and suggest specific species and genes for experimental testing to investigate mechanisms of microbiome-related protection from cholera.

KEYWORDS: *Vibrio cholerae,* cholera, microbiome, machine learning, metagenomics

INTRODUCTION

Cholera is an acute diarrheal disease caused by *Vibrio cholerae*. It is a major public health threat worldwide that continues to cause major outbreaks, such as in Yemen, where over 1.7 million cases have been reported since 2016 (1,2). Transmission of *V. cholerae* between household members commonly occurs through shared sources of contaminated food or water or through fecal-oral spread (3,4). The clinical spectrum of disease ranges from asymptomatic infection to severe watery diarrhea that can lead to fatal dehydration (5). Host factors such as age, innate immune factors, blood group, or prior acquired immunity partially explain why some people are more susceptible to *V. cholerae* infection than others, but a substantial amount of the variation remains unexplained (6).

The gut bacterial community can protect against enteropathogenic infections (7), and may explain some of the variation in *V. cholerae* susceptibility. Several studies have identified commensal bacteria and mechanisms that could be protective against *V. cholerae*. For instance, a species enriched in the gut microbiota of patients recovering from cholera, *Blautia obeum,* was found to interfere with *V. cholerae* pathogenicity through quorum-sensing inhibition in a mouse model (8). Other experiments have demonstrated that alteration of commensal-derived metabolite levels influenced host susceptibility by affecting *V. cholerae* growth or colonization (9-13).

Studies of *V. cholerae* and the gut microbiota often focus on a few bacterial species, or involve patients who already have symptomatic cholera (8,14). One study recently characterized the gut microbiome of healthy individuals exposed to *V. cholerae* (15). In this study, Midani *et al*. developed a machine learning model to predict susceptibility based on 16S rRNA gene amplicon sequencing of the gut microbiota in a group known to have high risk of infection: household contacts of confirmed cholera patients (4). Midani *et al* showed that microbiome composition at the time *V. cholerae* exposure to can predict infection with similar or better

accuracy as commonly measured host factors known to impact susceptibility. However, 16S

rRNA sequencing has limited taxonomic resolution and does not identify the genetic mechanisms

of protection.

Here we used shotgun metagenomics to analyze an expanded prospective cohort of

persons exposed to *V. cholerae* in Bangladesh. Our metagenomic analysis yielded improved

outcome predictions compared to 16S rRNA sequencing, and identified bacterial genes

associated with remaining uninfected after exposure to *V. cholerae*. We are also able to predict

disease severity among infected contacts, albeit with lower power and precision than

susceptibility. Finally, we highlight several microbiome-encoded metabolic functions associated

with protection against cholera.

METHODS

Sample collection, clinical outcomes and metagenomic sequencing

As described in (15), household contacts were enrolled within 6 hours of the presentation of an index cholera case at the icddr,b (International Center for Diarrheal Disease Research, Bangladesh) Dhaka Hospital. Index patients with severe acute diarrhea, a stool culture positive for *V. cholerae*, age between 2 and 60 years old, and no major comorbid conditions were recruited (4,6). A clinical assessment of symptoms in household contacts was conducted daily for the 10-day period after presentation of the index case, and repeated on day 30. We collected demographic information, rectal swabs, and blood samples for ABO typing and vibriocidal antibody titers as described in the Supplementary Methods. During the observation period, contacts were determined to be infected if any rectal swab culture was positive for *V. cholerae* and/or if the contact developed diarrhea and a 4-fold increase in vibriocidal titer during the follow-up period (4,6). Contacts with positive rectal swabs developing watery diarrhea were categorized as symptomatic and those without diarrhea were considered asymptomatic (**Figure 1**). *V. cholerae* positive contacts (by culture or deep 16S amplicon sequencing (15)) at the time of enrollment were excluded, in addition to contacts who reported antibiotic use or diarrhea during the week prior to enrollment. DNA extraction was performed for the selected samples and used for shotgun metagenomics sequencing. Details on cohorts, sequencing methods and sample processing are described in Supplementary Methods.

Figure 1. Study cohort in Dhaka, Bangladesh. After presentation of a *V. cholerae* culture-positive index case to the hospital on day 1, household contacts were enrolled on day 2. The expanded cohort includes the Midani 2018 cohort (15), with an addition of 33 samples from infected individuals (13 asymptomatic and 2 symptomatic).

Taxonomic/functional profiling and predictive model construction

We used MetaPhlAn2 (version 2.9) (16) for taxonomic profiling and HUMAnN2 (17) to

profile cellular pathways (from MetaCyc) and gene families (identified using the PFAM

database). For identification of biomarkers of susceptibility and disease severity, we used

MetAML (18) to apply a random forests (RF) classifier on species, pathways and gene-family

relative abundances, as well as strain-specific markers presence/absence. Models constructed

using each of these features types were compared to a random dataset with shuffled labels, and to

a model constructed with clinical/demographic data, using two-sample, two-sided *t*-tests over 20

replicate cross-validation (18). We used a stratified 3-fold cross validation approach, splitting our

dataset into validation and training sets (1/3 and 2/3 of samples, respectively) with the same

infected:uninfected ratio. We used an embedded feature selection strategy to identify the most

useful features and improve model accuracy. Feature relative importance was computed using the

mean decrease in impurity strategy, which calculates importance of each feature as the sum of the

number of nodes (across all trees) that use the feature, proportional to the number of samples

each of these nodes splits (18). Further details are described in the Supplementary Methods.

Data availability. After removal of human reads (Supplementary Methods), the sequence data

has been deposited in NCBI under BioProject PRJNA608678.

RESULTS

Metagenomic sequencing of the gut microbiome in household contacts exposed to *V.*

cholerae

We performed metagenomic sequencing of the gut microbiome in 65 contacts of cholera cases

from a cohort described by Midani *et al*. (15), from which sufficient DNA remained. Of these 65

contacts, referred to as the Midani 2018 cohort, 20 developed infection during the follow-up

period, and 45 remained uninfected (**Figure 1**). Among the 20 contacts who became infected, 10

had no symptoms during the 30-day follow-up period, and were classified as asymptomatic, and

10 developed symptoms (Supplementary Methods). To increase our sample size, we surveyed an

expanded cohort (**Table S1a**) by adding 33 samples, including 10 additional pre-infection

samples from timepoints of contacts in the Midani 2018 cohort, and 23 samples from 16 newly

enrolled contacts from the same place and time (2012-2014, Dhaka, Bangladesh). We used pre-

infection samples in order to identify predictive features of disease outcomes in the Midani 2018

cohort, upon which we base the majority of our analyses. We also performed exploratory

analyses on the expanded cohort to determine the potential for predictive models to be

generalized to larger samples. We used the shotgun metagenomic DNA sequence reads from

these samples to characterize four features of the microbiome: 1) relative abundances of

microbial species, 2) the presence/absence of sub-species-level strains, 3) metabolic pathway

relative abundances, and 4) gene family relative abundances (**Table 1**).

		RF – Cohort from Midani et al				RF – Expanded			
		Species abundance	Strain markers	Gene families	Pathways	Species abundance	Strain markers	Gene families	Pathways
	#features	705	54953	6810	443	807	62965	7514	461
Infected Vs Uninfected	Accuracy	0.73 (±0.02)	0.71 (±0.02)	0.76 (±0.02)	0.72 (±0.02)	0.76 (±0.03)	0.69 (±0.03)	0.80 (±0.02)	0.80 (±0.03)
	Precision	0.71 (±0.06)	0.68 (±0.06)	0.77 (±0.04)	0.70 (±0.05)	0.76 (±0.03)	0.70 (±0.03)	0.81 (±0.02)	0.81 (±0.03)
	F1	0.66 (±0.02)	0.64 (±0.03)	0.71 (±0.03)	0.66 (±0.03)	0. 75 (±0.03)	0.68 (±0.03)	0.80 (±0.02)	0.80 (±0.03)
	AUC	0.61 (±0.05)	0.62 (±0.04)	0.74 (±0.04)	0.71 (±0.04)	0.83 (±0.02)	0.76 (±0.03)	0.87 (±0.02)	0.88 (±0.02)
Shuffled	F1	0.55 (±0.04)	0.56 (±0.04)	0.56 (±0.04)	0.56 (±0.05)	0.40 (±0.03)	0.45 (±0.03)	0.48 (±0.03)	0.44 (±0.03)
	AUC	0.40 (±0.04)	0.57 (±0.04)	0.50 (±0.05)	0.50 (±0.04)	0.39 (±0.03)	0.52 (±0.03)	0.51 (±0.03)	0.46 (±0.03)
Asymptomatic vs Symptomatic vs Uninfected	Accuracy	0.70 (±0.02)	0.70 (±0.02)	0.69 (±0.01)	0.69 (±0.01)	0.68 (±0.01)	0.60 (±0.03)	0.69 (±0.02)	0.67 (±0.03)
	Precision	0.53 (±0.03)	0.53 (±0.03)	0.60 (±0.02)	0.59 (±0.02)	0.60 (±0.02)	0.53 (±0.03)	0.61 (±0.02)	0.59 (±0.02)
	F1	0.60 (±0.02)	0.59 (±0.02)	0.57 (±0.02)	0.57 (±0.02)	0.62 (±0.02)	0.55 (±0.03)	0.64 (±0.02)	0.62 (±0.02)
	AUC	NA	NA	NA	NA	NA	NA	NA	NA
Shuffled	F1	0.48 (±0.04)	0.49 (±0.04)	0.46 (±0.03)	0.55 (±0.03)	0.41 (±0.03)	0.35 (±0.03)	0.44 (±0.04)	0.37 (±0.03)
	AUC	NA	NA	NA	NA	NA	NA	NA	NA

Table 1. Assessment of prediction performance for a random forest (RF) model applied to the Midani 2018 and expanded cohorts. Species abundances, strain-specific markers presence/absence, relative abundance of Pfam-grouped gene families, and MetaCyc pathways were used as features. For each dataset, we applied a binary (uninfected vs. infected contacts) and a multi-class (asymptomatic vs. symptomatic vs. uninfected contacts) classifier and reported performance metrics for each dataset. Metrics obtained by the same classifier applied to the same datasets with shuffled class labels (random assignment of labels to samples) are also reported (shuffled). The margins of errors (95% confidence intervals) are reported in parenthesis.

Predicting susceptibility to *V. cholerae* infection with Random Forests

We first used an RF model to predict *V. cholerae* susceptibility (developing infection or

remaining uninfected) from baseline microbiome features (**Figure 1**). In the Midani 2018 cohort,

functional pathways and gene families predicted infection significantly better than random (Two-

sample *t*-tests comparing area under the curve [AUC] across 20 replicate 3 fold cross-validations;

$p < 0.05$) compared to data with shuffled (randomized) labels, and also predicted infection better

than species or strain features (**Tables 1 and S2**). Pathways and gene families had significantly

higher mean AUCs (0.71 and 0.74, respectively) compared to species or strains (0.61 and 0.62) (*p*

< 0.05; **Table 1**; **Figure S1, Table S3**).

To determine the minimum number of metagenomic features required for prediction, we

repeated the analysis using smaller subsets of features. Using only 30 species, 60 gene families or

pathways, or 200 strains achieved similar cross-validation AUC values (**Figure S2**). We then

trained an RF model on this reduced number of selected features, yielding improved predictions

for all feature types (**Figure S1; Table S4**). This suggests that only a limited number of strains,

species, genes and pathways in the gut microbiome at the time of exposure are sufficient to

predict *V. cholerae* susceptibility. For example, prediction using strain-level markers after feature

selection yielded an AUC of 0.95 (**Table S4**). However, such high AUC values should be treated

with caution because the models can be overfit when a supervised feature selection step is applied

on the same data used to train the model (18). Because we did not have a fully independent

validation cohort (*e.g.* from another continent) to test our model, we decided to use the features

selected from the Midani cohort to make predictions on the Expanded dataset. Using the same

features selected from the Midani 2018 training dataset, we made predictions on the Expanded

cohort and achieved AUCs between 0.89 and 0.93 for prediction of infection using the four types

of features (**Table S4**). Again, because the expanded cohort partly overlaps with the Midani

cohort, and includes some repeated samples from the same individuals over time, these results

could also be prone to overfitting, but they demonstrate the potential for generalized predictions.

Finally, we repeated the RF analysis using all features in the expanded dataset, whichh

increased predictive performance relative to the original Midani cohort (**Figure S1**). Once again,

genes and pathways outperformed species and strains according to all metrics, with AUC

reaching ~0.88 using cellular pathways (**Table 1**). This improvement in the expanded cohort also

highlights the importance of using larger, more balanced datasets as input to predictive models.

Improved prediction compared to known factors impacting susceptibility

To put the metagenomic predictions in context, we compared their predictive power and

accuracy to clinical and demographic factors (**Table S1a**). Three of these factors (age, baseline

vibriocidal antibodies and blood group) are known to impact susceptibility to *V. cholerae*

infection (6,15) and we used them to train RF models (**Table S5**). As expected, contacts who

became infected tended to be younger and have lower baseline antibody titers than those who

remained uninfected (**Table S1b**), but these small differences were not sufficient to train a

significantly predictive model. An RF model trained on the seven clinical and demographic

factors did not perform better than a random model with shuffled labels (AUC=0.60, p=0.66;

Figure 2). Predictions were not improved using all species-level metagenomic features present at

the time of exposure to *V. cholerae* (AUC=0.61), but significantly improved using a selected

number of species (AUC=0.80, $p < 1 \times 10^{-7}$). The use of all gene families or a selected number of

genes showed an increased predictive performance (AUC=0.74 and AUC=0.89 respectively;

Figure 2) compared to species-level or clinical and demographic contact data ($p < 1 \times 10^{-7}$ for all

comparisons). We again note the caveat that models with selected features may be overfit and

represent an upper bound for predictive power. Even without feature selection, we found that

gene families clearly provide superior predictions, and adding clinical data did not improve the

predictions based on microbiome features alone (**Figure 2**). Together, these results demonstrate

that gene families present in the gut microbiome at the time of exposure contain more

information about *V. cholerae* susceptibility compared to species-level or clinical and

demographic contact data.

Figure 2. Metagenomic features predict *V. cholerae* infection better than clinical and
demographic features. Random forest prediction of infection status was applied to 7 clinical and
demographic features, and compared with all species and all gene families (top row), as well as
30 selected species features from metagenomes and 60 selected gene family features (bottom
row), or a combination of clinical, demographic and metagenomic features. Plots show receiver
operating characteristic (ROC) curves (average across cross-validations) for the Midani 2018
dataset. Shuffled labels represent the prediction run on a dataset with a random assignment of
infection outcomes. AUC = area under the curve.

Disease severity is more difficult to predict than likelihood of infection

To predict symptomatic disease among infected individuals (**Figure 1**), we divided
samples into uninfected, symptomatic and asymptomatic groups and again applied the RF
approach. We used the F1 score as a performance metric since it is well suited for uneven class
distributions in our uninfected/symptomatic/asymptomatic comparison. Applied to the Midani
2018 cohort, this model predicted outcomes significantly better than random (shuffled labels)
using species, strains or pathway data, but not gene families (**Table 1;** see **Table S3** for *p*-
values). However, the F1 scores for the symptomatic/asymptomatic predictions were
systematically lower (mean scores ranging from of 0.57 to 0.60) than for the infected/uninfected
prediction (means ranging from 0.64 to 0.71). Using the expanded cohort, the scores were
improved only slightly (**Table 1**). These results suggest that disease severity is predictable in
principle, but with greater uncertainty than the infection outcome.

Taxonomic biomarkers of disease susceptibility and severity

Predictive features in the gut microbiome identified to a species/strain or gene level allow
the possibility of experimental follow-up to investigate mechanisms of the associations we
observed. We characterized the most predictive species, pathways, and gene families (**Tables S6-
S9**). The most common discriminating species in individuals that remained uninfected during the
follow-up period were *Eubacterium rectale, Campylobacter hominis, Ruminococcus gnavus,
Bacteroides vulgatus, Veillonella parvula* and members of the *Prevotella* and *Eubacterium*
genera (**Figures 3A**, **S3A and S4A**). These species are ranked by their importance score, which
is effectively their relative weighting in the RF model. Several species associated with contacts
that developed *V. cholerae* infection belonged to the genera *Bifidobacterium, Actinomyces* or
Collinsella, and many of the species were also associated with asymptomatic infection (**Figures**

3B, **S3B and S4B**), including three species of *Bifidobacterium.* The top predictive species in

contacts who developed symptomatic infection were *Clostridium ventriculi* (formerly *Sarcina*

ventriculi), *Streptococcus parasanguinis* and members of *Veillonella*. *Shigella* species were also

associated with the gut microbiome of persons who developed symptomatic *V. cholerae*

infection, although persons enrolled in this study were *Shigella* stool-culture negative. *Shigella*

identified by DNA presence in stool may be the result of recent or resolving infection, or may be

present at subclinical levels due to ingestion of contaminated water. The features identified by the

multivariate RF model were confirmed using univariate statistics for the uninfected/infected

prediction (**Figure S5)**, but the overlap was poorer for the uninfected/symptomatic/asymptomatic

prediction (**Figure S6)**. This is consistent with the difficulty of predicting disease severity.

In general, the most important species were selected by the model because of differences

in relative abundance at baseline among uninfected/symptomatic/asymptomatic outcomes

(**Figure S7, S8**). In rare cases, species presence/absence information was predictive. For

example, *Ruminococcus gnavus,* is absent (near or below limit of detection) in most of the

individuals who become infected, but present in many (but not all) of those who remain

uninfected (**Figure S7**). Thus, there is no single, strong predictor of infection outcomes, but

rather a probabilistic combination of many species, each of relatively modest predictive value.

A

Top discriminating species associated with contacts
who remained uninfected or became infected

Mitsuokella multacida
Catenibacterium sp CAG 290
Burkholderia pyrrocinia
Eubacterium rectale
Prevotella sp 885
Bifidobacterium longum
Prevotella sp TF12 30
Roseburia sp CAG 471
Bifidobacterium adolescentis
Faecalibacterium prausnitzii
Prevotella sp CAG 5226
Campylobacter hominis
Slackia isoflavoniconvertens
Bifidobacterium bifidum
Firmicutes bacterium CAG 83
Dialister sp CAG 486
Eubacterium sp CAG 202
Actinomyces odontolyticus
Clostridiales bacterium KLE1615
Ruminococcus gnavus
Prevotella copri
Shigella flexneri
Veillonella parvula
Burkholderia stabilis
Bacteroides vulgatus

Feature importance ranking

B

Top discriminating species associated with contacts
who remained uninfected or became infected
symptomatic or infected asymptomatic

Collinsella massiliensis
Prevotella sp 885
Burkholderia pyrrocinia
Enorma massiliensis
Catenibacterium sp CAG 290
Veillonella parvula
Eubacterium rectale
Collinsella aerofaciens
Clostridium ventriculi
Escherichia coli
Gemmiger formicilis
Bifidobacterium bifidum *
Roseburia faecis
Bifidobacterium adolescentis *
Shigella sonnei
Faecalibacterium prausnitzii
Shigella boydii
Streptococcus parasanguinis
Bifidobacterium longum *
Eubacterium sp CAG 146
Prevotella sp AM42 24
Roseburia sp CAG 471
Veillonella atypica
Veillonella infantium
Prevotella sp TF12 30

Figure 3. Most important discriminating species of the gut microbiome at the time of exposure to *V. cholerae* identified in the Midani 2018 dataset, classified by clinical outcome. (A) Species associated with contacts that became infected (red) or remained uninfected (black) during follow-up. (B) Species associated with contacts who remained uninfected (black), or became infected asymptomatic (green), or symptomatic (red) during follow-up. The top 25 most important features for discriminating between classes in the RF model are shown here; see Table S6 for the full list. Yellow lines connect species associated with uninfected individuals in both (A) and (B); red lines connect species associated with infection in (A) and symptomatic disease in (B); grey lines connect species associated with infection in (A) but asymptomatic infection in (B). Three species of *Bifidobacterium* are marked with an asterisk.

Identification of functional biomarkers of disease susceptibility and severity

We also identified gene families in the gut microbiome of persons who remained

uninfected during follow-up (**Figures S9 and S10**), with some of the top gene families involved

in DNA repair, transmembrane transporter activity, iron metabolism (indicated with asterisks in

Figure 4), and genes of unknown function (**Table S8**). Long-chain fatty acid biosynthesis

pathways (*e.g.* cis-vaccenate, gondoate and stearate) were associated with individuals who

remained uninfected, while amino acid biosynthesis and catabolic pathways were associated with

individuals who developed infection (**Figures S11 and S12, Table S9**). We identified three iron-

related genes associated with remaining uninfected: (1) the ferric uptake regulator Fur, a major

regulator of iron homeostasis, (2) thioredoxin, a redox protein involved in adaptation to oxidative

and iron-deficiency stress, and (3) the TonB/ExbD/TolQR system, a ferric chelate transporter

(19-21). In individuals who became infected but asymptomatic, two genes involved in the

conversion of riboflavin into catalytically active cofactors, the riboflavin kinase and the FAD

synthetase, were found as the first and the third most discriminant features (**Figure 4, Table S8**).

We next asked which taxa in the microbiome likely encoded these genes. In some cases,

specific taxonomic groups corresponded to discrete gene functions. For example, several iron

metabolism-related gene families tend to be encoded by *Prevotella* genomes (**Figure S14**). In

other cases, the major contributors to protective gene families were unclassified (**Figures 5 and**

S13). These results partly explain why gene families or pathway features tend to outperform

species-level features in predicting infection status – because predictive gene families are

distributed across many species, including several with poor taxonomic annotation or families

lacking representation in taxonomic databases.

302
303
Figure 4. Most important discriminating gene families of the gut microbiome at the time of
exposure to *V. cholerae* identified in the Midani 2018 dataset, classified by clinical outcome.
(A) Genes families associated with contacts that became infected (red) or remained uninfected
(black) during follow-up. (B) Genes families associated with contacts who remained uninfected
(black), or became infected asymptomatic (green), or symptomatic (red) during follow-up. The
top 25 most important features for discriminating between classes in the RF model are shown
here; see Table S8 for the full list. Yellow lines connect species associated with uninfected
individuals in both (A) and (B). Asterisks indicate genes involved in redox or iron metabolism.
312

Top discriminating pathways in contacts who became infected (symptomatic)

Top discriminating pathways in contacts who became infected (asymptomatic)

Top discriminating pathways in contacts who remained uninfected

Figure 5. Top predictive cellular pathways of the gut microbiome at the time of exposure to *V. cholerae* in the Midani 2018 cohort, annotated by their taxonomic contributors. The four top-ranked pathways associated with uninfected contacts (left column), contacts who developed asymptomatic infection (middle), and contacts who developed symptomatic infection (right column) are shown. Total bar height reflects log10-scaled community relative abundance of each pathways. The contributions of each genus to encoding these pathways are shown as stacked colors within each bar, linearly scaled within the total. See Table S9 for the complete list of pathways

DISCUSSION

The gut microbiome is a potentially modifiable host risk factor for cholera, and identification of specific genes and strains correlated with susceptibility is needed for experimental testing to understand the mechanisms of observed correlations. Compared to a previous study using a single marker gene, shotgun metagenomics provides this degree of resolution, potentially to the species and strain level, and to the level of individual genes and cellular functions. We found that gene families in the gut microbiome at the time of exposure to *V. cholerae* were more predictive of susceptibility compared to taxonomic or clinical and demographic information. Selecting a subset of the most informative features further improved predictions, but using these selected features may lead to overfitting. This suggests an upper limit to predictive power that requires validation in larger, independent cohorts.

Most of the top predictive biomarkers were associated with remaining uninfected after exposure to *V. cholerae*. An example is the genus *Prevotella,* including several strains within *Prevotella* sp. 885, identified only at the genus level in a previous study (15). *Prevotella* species are hypothesized to be beneficial members of the microbiota in healthy individuals in non-Westernized countries, and this species is a potential candidate for follow-up experimental studies in *V. cholerae* susceptibility (14,22,23).

Several species known to ferment mucin glycans into short chain fatty acids (SCFAs) correlated with remaining uninfected, including *Eubacterium rectale*, *Ruminococcus gnavus* and *Bacteroides vulgatu*s (24,25). This finding is consistent with experiments of SCFAs applied to animal models. *B. vulgatus* has been shown to inhibit *V. cholerae* colonization in mice, an effect that was dependent upon SCFA production (13). SCFAs are known to impact immune cell

development and attenuate inflammation by inhibiting histone deacetylases and other

mechanisms of altering gene expression (26-29).

All three *Bifidobacterium* species associated with contacts that developed infection were

also associated with asymptomatic rather than symptomatic disease, and prior work on this genus

supports several hypotheses for this relationship. First, *Bifidobacteria* are known to produce the

SCFA acetate that can protect against enteric infection in mice (30,33,34). SCFAs are also known

to inhibit cholera toxin-related chloride secretion in the mouse gut, reducing water and sodium

loss, and have been observed to increase cholera toxin-specific antibody responses (31-33).

Bifidobacteria are also major producers of lactate, a metabolite that has been shown to impair *V.*

cholerae biofilm formation, a function that can impact virulence (12). Lastly, *B. bifidum* and *B.*

adolescentis are known to reduce the activity of *V. cholerae* type VI secretion systems through

modification of bile acids (9).

Metagenomics also allowed us to identify bacterial functions that could impact the ability

of *V. cholerae* to compete and colonize the gut. For example, several gene families involved in

iron transport, iron regulation, and riboflavin conversion appeared among the top twenty features

associated with uninfected and asymptomatic individuals, suggesting that competition for iron

might be a protective mechanism of the gut microbiota against *V. cholerae*, as in other pathogens

(7). Iron is often a limiting redox cofactor in the gut, and bacteria have evolved strategies to

solubilize and internalize iron (34,35). Riboflavin (another major redox cofactor in bacteria) and

iron levels are reciprocally regulated in *V. cholerae*, and riboflavin may allow *V. cholerae* to

overcome iron limitation in the gut (34,36). A gut microbiota more competitive for iron could

therefore help resist *V. cholerae* colonization or reduce its virulence. Further work is thus needed

to understand mechanisms of how the enrichment of these genes may protect people after

exposure to *V. cholerae*.

Our results are currently not generalizable beyond the study cohort in Dhaka, Bangladesh,

since a similar cohort in another geographic location is not available. As with any association-

based study (37), it is unknown if any of the metagenomic features that correlate with protection

from *V. cholerae* infection are causal, and many may be markers of clinical or environmental

factors that themselves impact susceptibility. Despite our deep sequencing and collection of

standard cholera risk factors, our study was unable to measure all potentially relevant

environmental or clinical risk factors. In line with recent studies in Dhaka, we assume that *V.*

cholerae transmission occurs mainly within households (3) and did not consider how the mode of

transmission (*e.g.* waterborne or not) might affect outcomes. It has also been noted that

microbiome-disease associations may be poorly portable across human populations (37). For

instance, we identified species of *Prevotella* as protective features in Bangladesh, but *Prevotella*

is much less abundant and less diverse in Western countries (22). It thus remains to be seen if

protective gene features (*e.g.* iron metabolism) are encoded in other species of the microbiome

outside endemic areas like Bangladesh, or if people outside these areas are simply at greater risk

for cholera. Further experimental characterization of metagenomic features correlated with

protection from infection or symptoms are needed to understand if factors we identified impact *V.*

cholerae pathogenesis or host responses to infection. Ultimately, the strains and functionalities

identified have the potential to inform microbiota-based therapeutics to ameliorate or prevent

disease. Our results show the power of metagenomic data from the gut microbiome to predict

health outcomes such as susceptibility to infection and disease severity.

FUNDING INFORMATION

This study was supported by CIHR (Canadian Institutes of Health Research) and the Canada Research Chairs program (BJS), The icddr,b: Centre for Health and Population Research, the Alfred P. Sloan Fellowship (LAD), grants AI099243 (J.B.H and L.C.I), AI103055 (J.B.H and F.Q), AI106878 (E.T.R and F.Q.), AI058935 (E.T.R, S.B.C and F.Q.), T32AI070611976 and K08AI123494 (A.A.W.) from the National Institutes of Health, and the Robert Wood Johnson Foundation Harold Amos Medical Faculty Development Program (R.C.C.).

ACKNOWLEDGEMENTS

We thank Meti Debela for technical assistance. Finally, we are grateful to the people of Dhaka where our study was undertaken; to the field, laboratory and data management staff who provided tremendous effort to make the study successful; and to the people who provided valuable support in our study. The icddr,b gratefully acknowledges the Government of the People's Republic of Bangladesh; Global Affairs Canada (GAC); Swedish International Development Cooperation Agency (Sida) and the Department for International Development, (UKAid). We declare that we have no competing financial interest.

ETHICAL STATEMENT

The Ethical and Research Review Committees of the icddr,b and the Institutional Review Board of MGH reviewed the study. All adult subjects provided informed consent and parents/guardians of children provided informed consent. Informed consent was written.

CONFLICTS OF INTEREST

The authors declare that there are no conflicts of interest.

Supplementary tables S1-S9 are available at:
https://figshare.com/articles/Supplementary_Tables_-_Levade_et_al_2020/12440417

REFERENCES

1.   Ali M, Nelson AR, Lopez AL, Sack DA. Updated Global Burden of Cholera in Endemic Countries. PLoS Negl Trop Dis. **2015**; 9(6):e0003832.

2.   Camacho A, Bouhenia M, Alyusfi R, et al. Cholera epidemic in Yemen, 2016-18: an analysis of surveillance data. Lancet Glob Health. **2018**; 6(6):e680–e690.

3.   Domman D, Chowdhury F, Khan AI, et al. Defining endemic cholera at three levels of spatiotemporal resolution within Bangladesh. Nat Genet. **2018**; :1–10.

4.   Weil AA, Khan AI, Chowdhury F, et al. Clinical Outcomes in Household Contacts of Patients with Cholera in Bangladesh. Clin Infect Dis. **2009**; 49(10):1473–1479.

5.   Nelson EJ, Harris JB, Morris JG, Calderwood SB, Camilli A. Cholera transmission: the host, pathogen and bacteriophage dynamic. Nature. **2009**; :1–10.

6.   Harris JB, LaRocque RC, Chowdhury F, et al. Susceptibility to *Vibrio cholerae* Infection in a Cohort of Household Contacts of Patients with Cholera in Bangladesh. PLoS Negl Trop Dis. **2008**; 2(4):e221–8.

7.   Ubeda C, Djukovic A, Isaac S. Roles of the intestinal microbiota in pathogen protection. Clinical & Translational Immunology. **2017**; 6(2).

8.   Hsiao A, Ahmed AMS, Subramanian S, et al. Members of the human gut microbiota involved in recovery from *Vibrio cholerae* infection. Nature. **2014**; 515(7527):423–426.

9.   Bachmann V, Kostiuk B, Unterweger D, Diaz-Satizabal L, Ogg S, Pukatzki S. Bile Salts Modulate the Mucin-Activated Type VI Secretion System of Pandemic *Vibrio cholerae*. PLoS Negl Trop Dis. **2015**; 9(8):e0004031.

10.   Yoon MY, Min KB, Lee K-M, et al. A single gene of a commensal microbe affects host susceptibility to enteric infection. Nature Communications. **2016**; 7:1–11.

11.   Mao N, Cubillos-Ruiz A, Cameron DE, Collins JJ. Probiotic strains detect and suppress cholera in mice. Science Translational Medicine. **2018**; 10(445).

12.   Kaur S, Sharma P, Kalia N, Singh J, Kaur S. Anti-biofilm Properties of the Fecal Probiotic Lactobacilli Against Vibrio spp. Front Cell Infect Microbiol. **2018**; 8.

13.   You JS, Yong JH, Kim GH, et al. Commensal-derived metabolites govern *Vibrio cholerae* pathogenesis in host intestine. Microbiome.. **2019**; 7(1):1–18.

14.   David LA, Weil A, Ryan ET, et al. Gut Microbial Succession Follows Acute Secretory Diarrhea in Humans. mBio **2015**; 6(3):e00381–15.

450   15.   Midani FS, Weil AA, Chowdhury F, et al. Human Gut Microbiota Predicts Susceptibility
451   to *Vibrio cholerae* Infection. J Infect Dis. **2018**; 218(4):645–653.

452   16.   Truong DT, Franzosa EA, Tickle TL, et al. MetaPhlAn2 for enhanced metagenomic
453   taxonomic profiling. Nature Methods. **2015**; 12(10):902–903.

454   17.   Franzosa EA, McIver LJ, Rahnavard G, et al. Species-level functional profiling of
455   metagenomes and metatranscriptomes. Nature Methods. **2018**; 15(11):962–968.

456   18.   Pasolli E, Truong DT, Malik F, Waldron L, Segata N. Machine Learning Meta-analysis of
457   Large Metagenomic Datasets: Tools and Biological Insights. PLoS Comput Biol. **2016**;
458   12(7):e1004977.

459   19.   Fillat MF. The FUR (ferric uptake regulator) superfamily: diversity and versatility of key
460   transcriptional regulators. Arch Biochem Biophys. **2014**; 546:41–52.

461   20.   Wang B-Y, Huang H-Q, Li S, et al. Thioredoxin H (TrxH) contributes to adversity
462   adaptation and pathogenicity of *Edwardsiella piscicida*. Vet Res. **2019**; 50(1):1–13.

463   21.   Noinaj N, Guillier M, Barnard TJ, Buchanan SK. TonB-dependent transporters: regulation,
464   structure, and function. Annu Rev Microbiol. **2010**; 64:43–60.

465   22.   Tett A, Huang KD, Asnicar F, et al. The Prevotella copri Complex Comprises Four
466   Distinct Clades Underrepresented in Westernized Populations. Cell Host and Microbe.
467   **2019**; 26(5):666–679.e7.

468   23.   Kovatcheva-Datchary P, Nilsson A, Akrami R, et al. Dietary Fiber-Induced Improvement
469   in Glucose Metabolism Is Associated with Increased Abundance of Prevotella. Cell Metab.
470   **2015**; 22(6):971–982.

471   24.   Crost EH, Tailford LE, Le Gall G, Fons M, Henrissat B, Juge N. Utilisation of mucin
472   glycans by the human gut symbiont *Ruminococcus gnavus* is strain-dependent. PLoS ONE.
473   **2013**; 8(10):e76341.

474   25.   Tailford LE, Crost EH, Kavanaugh D, Juge N. Mucin glycan foraging in the human gut
475   microbiome. Front Genet. **2015**; 6.

476   26.   Sun Y, O'Riordan MXD. Regulation of bacterial pathogenesis by intestinal short-chain
477   Fatty acids. Adv Appl Microbiol. **2013**; 85:93–118.

478   27.   Koh A, De Vadder F, Kovatcheva-Datchary P, Bäckhed F. From Dietary Fiber to Host
479   Physiology: Short-Chain Fatty Acids as Key Bacterial Metabolites. Cell. **2016**;
480   165(6):1332–1345.

481   28.   Louis P, Flint HJ. Formation of propionate and butyrate by the human colonic microbiota.
482   Environmental Microbiology. **2017**; 19(1):29–41.

29. Fachi JL, Souza Felipe J de, Pral LP, et al. Butyrate Protects Mice from *Clostridium*
difficile-Induced Colitis through an HIF-1-Dependent Mechanism. Cell Reports. **2019**. p.
750–761.e7.

30. Fukuda S, Toh H, Hase K, et al. Bifidobacteria can protect from enteropathogenic
infection through production of acetate. Nature. **2011**; 469(7331):543–547.

31. Canani RB, Costanzo MD, Leone L, Pedata M, Meli R, Calignano A. Potential beneficial
effects of butyrate in intestinal and extraintestinal diseases. World J Gastroenterol. **2011**;
17(12):1519–1528.

32. Yang W, Xiao Y, Huang X, et al. Microbiota Metabolite Short-Chain Fatty Acids
Facilitate Mucosal Adjuvant Activity of Cholera Toxin through GPR43. The Journal of
Immunology. **2019**; 203(1):282–292.

33. Rabbani GH, Albert MJ, Rahman H, Chowdhury AK. Short-Chain Fatty Acids Inhibit
Fluid and Electrolyte Loss Induced by Cholera Toxin in Proximal Colon of Rabbit In
Vivo. Dig Dis Sci. **1999** 44(8):1547–1553.

34. Sepúlveda Cisternas I, Salazar JC, García-Angulo VA. Overview on the Bacterial Iron-
Riboflavin Metabolic Axis. Front Microbiol. **2018**; 9:1478.

35. Rivera-Chávez F, Mekalanos JJ. Cholera toxin promotes pathogen acquisition of host-
derived nutrients. Nature. **2019**; 572(7768):244–248.

36. Sepúlveda Cisternas I, Aguirre LL, Flores AF, de Ovando IVS, García-Angulo VA.
Transcriptomics reveals a cross-modulatory effect between riboflavin and iron and outlines
responses to riboflavin biosynthesis and uptake in *Vibrio cholerae*. Scientific Reports.
2018; 8(1):1–14.

37. Schmidt TSB, Raes J, Bork P. The Human Gut Microbiome: From Association to
Modulation. Cell. **2018**; 172(6):1198–1215.