Mice lacking paternal expression of imprinted *Grb10* are risk-takers

Claire L. Dent[a], Kira D. A. Rienecker[a], Andrew Ward[b], Jon F Wilkins[c] Trevor Humby[d], &

Anthony R. Isles[a]*

[a]Behavioural Genetics Group, MRC Centre for Neuropsychiatric Genetics and Genomics,

Neuroscience and Mental Health Research Institute, Cardiff University, Hadyn Ellis Building,

Maindy Road, Cardiff, United Kingdom

[b]Department of Biology and Biochemistry, University of Bath, Bath, United Kingdom

[c]Ronin Institute, Montclair, NJ 07043, USA

[d]Behavioural Genetics Group, School of Psychology, Cardiff University, Tower Building,

Cardiff, United Kingdom

*Author for correspondence: Anthony R. Isles, E-mail IslesAR1@cardiff.ac.uk

ABSTRACT

The imprinted genes *Grb10* and *Nesp* influence impulsive behavior on a delay discounting

task in an opposite manner. A recently developed theory suggests that this pattern of

behavior may be representative of predicted effects of imprinted genes on tolerance to risk.

Here we examine whether mice lacking paternal expression of *Grb10* show abnormal

behavior across a number of measures indicative of risk-taking. Although $Grb10^{+/p}$ mice

show no difference from wild type littermates in their willingness to explore a novel

environment, their behavior on an explicit test of risk-taking, namely the Predator Odour

Risk-Taking task, is indicative of an increased willingness to take risks. Follow-up tests

suggest that this risk-taking is not simply due to a general decrease in fear, or a general

increase in motivation for a food reward, but reflects a change in the trade-off between cost

and reward. These data, coupled with previous work on the impulsive behaviour of $Grb10^{+/p}$

mice in the delayed reinforcement task, and taken together with our work on mice lacking

maternal *Nesp*, suggest that maternally and paternally expressed imprinted genes oppositely

influence risk-taking behaviour as predicted.

Introduction

The imprinted genes *Grb10* and *Nesp* affect impulsive choice behavior in opposing
directions (Dent et al., 2016, 2018). Mice lacking paternal *Grb10* (*Grb10*$^{+/p}$) prefer a larger,
but delayed reward to a smaller, but more immediate reward in the delayed reinforcement
task (DRT) (Dent et al., 2018). In contrast, mice lacking maternal *Nesp* (*Nesp*$^{m/+}$) choose a
more immediate, smaller reward over a larger, but delayed reward in the DRT (Dent *et al.*,
2016). These behavioral findings, coupled with co-localisation of expression of *Nesp* and
Grb10 in a number of brain regions (Garfield *et al.*, 2011, Plagge *et al.*, 2005) and cell types
(Dent et al., 2018), has led to the suggestion that they may have an antagonistic effect on
the control of behavior (Dent *et al.*, 2018, Dent & Isles, 2014). This fits with the general idea
that genomic imprinting evolved as a consequence intragenomic conflict between maternal-
and paternally-derived alleles arising as a consequence of kin-selection (Moore & Haig,
1991, Wilkins *et al.*, 2016).

Recently a theoretical basis for how imprinted genes may influence risk-taking behavior has
been proposed (Wilkins & Bhattacharya, 2019). Here, intragenomic conflict between
maternal and paternal genomes arises not because of kinship differences, but as a
consequence of differences in reproductive variance between males and females. The
prediction is that when reproductive variance is higher in males (as it is for most mammals)
then paternally expressed imprinted genes will promote risk-averse, variance-reducing
behaviors, and maternally expressed imprinted genes will promote risk-tolerant, variance-
increasing behaviors. The authors argue that the opposing pattern of behavior shown by
Grb10$^{+/p}$ and *Nesp*$^{m/+}$ mice in the DRT supports this idea. The DRT is generally considered a
measure of impulsive choice; *Grb10*$^{+/p}$ are more likely to choose the larger reward and are
interpreted as less impulsive. However, in a naturalistic environment, delay introduces the
possibility a reward will be lost to a competitor (loss of opportunity), or that the individual will
be exposed to risk of predation before receipt of the reward (cost in death or injury) (Benson
& Stephens, 2015, Green & Myerson, 1996, Kacelnik & Bateson, 1997, Schonberg *et al.*,
2011). Choosing a delayed, larger reward in the DRT may therefore indicate not only less
impulsive, but also more risky behavior (Holt *et al.*, 2003).

The idea that the choice of the more immediate, but smaller reward, displayed by *Nesp*$^{m/+}$
mice in the DRT may reflect less risk-taking is supported by their behavior in other tasks.
Nesp$^{m/+}$ mice show altered reactivity to, and are less willing to explore, novel environments
(Plagge *et al.*, 2005). The propensity to explore a novel environment is regarded as a good
index of risk-taking behavior and has been used as such in a number of studies (Blanchard
et al., 2009). However, the data for the behavior of *Grb10*$^{+/p}$ mice being indicative of an
increased willingness to take risk are more equivocal. Here we address this, by examining
the behavior of *Grb10*$^{+/p}$ mice in a novel environment. We also directly tested risk-taking

behavior using the Predator Odor Risk-Taking (PORT) task (Dent, Isles, & Humby, 2014). The PORT task was developed by us (Dent et al., 2014), and has been used by others in both mice (Blomeley *et al.*, 2018, Viola *et al.*, 2019) and rats (Schneider *et al.*, 2015), to specifically assess "real-life" risky situations where there is a trade-off between a cost (risk of predation) and a food reward. Although $Grb10^{+/p}$ mice show no difference from wild type animals in their willingness to explore a novel environment, their behavior on the PORT task is indicative of increased risk-taking. Follow-up tests suggest that this risk-taking is not simply due to a general decrease in fear, or a general increase in motivation for a food reward, but reflects a change in the trade-off between cost and reward.

MATERIALS AND METHODS

Animals

All procedures were conducted in accordance with the UK Animals (Scientific Procedures) Act 1986. The $Grb10$ null line was maintained on an F1-hybrid (C57Bl/6J X CBA/Ca) background. Due to potentially confounding metabolic phenotypes associated with $Grb10^{m/+}$ mice (Smith *et al.*, 2007), comparisons of $Grb10^{+/p}$ were made with wild-type (WT) littermate controls. Subjects were male mice aged between 3-6 months during testing. Standard laboratory chow was available *ad libitum*, but during the progressive ratio experiment water was restricted to 2h access per day. This regime maintained the subjects at ≈90% of free-feeding body weight and motivated the animals to work for the food reward used in the task. Two separate cohorts were used. Cohort 1 (WT, n=11; $Grb10^{+/p}$, n=13) undertook locomotor activity, novelty place preference, milk consumption test, PORT task, acoustic startle. Cohort 2 (WT, n=14; $Grb10^{+/p}$, n=11) undertook milk consumption testing and the progressive ratio. All animals within a cohort were used throughout testing but a number of animals were removed or lost (due to an inability to perform training stages or death) as testing progressed.

Locomotor Activity

Locomotor activity (LMA) was measured using an apparatus consisting of a battery of 12 cages, each measuring 21 x 21 x 36cm. The activity cages were clear Perspex boxes containing two transverse infrared beams 10mm from the floor, spaced equally along the length of the box, linked to an Acorn computer using ARACHNID software (Cambridge Cognition Ltd., Cambridge, UK). Activity was measured for two-hour sessions in the dark, over 3 consecutive days; testing took place at the same time every day. The cages were thoroughly cleaned after each animal, using 1% acetic acid solution. The main measure was 'runs', recorded when the animal broke the two infrared beams consecutively.

Novelty Place Preference

Novelty place preference (NPP) was assessed using an apparatus consisting of two adjacent boxes with an opening in the middle, which could be occluded by a door. These two arenas were made distinct by the colour (black or white) and the texture of the floor (smooth plastic or sandpaper). In the first stage of the test the door was closed and the animal was placed in one side of the box for one hour and allowed to habituate to that side. Then the mouse was taken out, the door removed and the mouse put back in the same side and allowed to explore both the habituated side and the novel side for a total of 30 minutes. The side to which the animal was habituated, and in which the sand-paper was placed, was pseudo-randomly allocated to avoid confounding the results. The arena was thoroughly cleaned after each animal, using 1% acetic acid solution.

The movement of each subject was tracked using a camera mounted approximately two metres above the test arena, connected to EthoVision Observer software (Noldus Information Technology, Netherlands). Behavioral measures, obtained automatically, included the duration of time spent in each zone, frequency of entries, and latency of first entrance into the novel zone. The time spent in and the number of entries into the novel compartment was measured automatically by a video tracking system, using Noldus software.

Predator Odor Risk-Taking (PORT) task

The PORT task was conducted using the same methods and apparatus as previously described (Dent et al., 2014) and full details can be found in the Supplementary Information. Briefly, following habituation to the apparatus, animals were trained to leave the start chamber, traverse the middle chamber, and to collect a food reward in the third chamber of the apparatus. For training trials, clean standard mouse bedding (wood shavings) was distributed evenly over the floor of the middle chamber. Trial length was set to 10 minutes but was terminated when the subject had traversed the apparatus and was observed to have collected the reward. The mouse was then removed and placed in a holding box until the start of the next trial. In the test trials the middle chamber bedding was mixed with either 'self-odor bedding' (bedding taken from the mouse's home cage), or 'predator-odor bedding' (wood shavings mixed with a synthetic predator cue, 2,4,5-trimethylthiazoline, TMT; Contech Inc., Canada). The main measurements taken in each trial was the latency to leave the start chamber.

Predator odor enhanced acoustic startle response (POE-ASR)

The POE-ASR was assessed in two separate test sessions, a week apart,

immediately following a 10-min exposure to either untainted wood-shaving bedding (control

condition) or fox odor-tainted bedding, mixed at the same concentration of TMT as used in

the PORT task (see above). The order of odor presentation was counter-balanced between

mice. ASR was monitored using SR-Lab apparatus (San Diego Instruments, USA),

according to the previous method used (Dent et al., 2014) (see Supplementary Information

for full details).

Condensed Milk Test

In order to increase motivation and performance of the mice in the progressive ratio task

mice were placed on a schedule of water restriction. Water was maintained at a 2hr regime

for the duration of the experiment, and food was available *ad libitum* at all times (apart from

when in chambers)*.* Body weight was monitored prior to water restriction, and throughout the

first 10 days of restriction. Once body weight had stabilised (>90% free-feeding weight) the

mice were habituated to the condensed milk reward used in the operant tasks. This was

carried out using the Condensed Milk Test (CMT) to check for preference of condensed milk

over water as described previously (Dent *et al.*, 2014) (see Supplementary Information for

full details). The amount drunk and preference for condensed milk over water was measured

on five successive days. This was to prevent mice having a neophobic reaction to the reward

during the experiment and also to test for any differences in consumption and/or acquisition

of a preference.

Progressive ratio

All the sessions of the Progressive Ratio (PR) task were performed in 9-hole operant

chambers (Cambridge Cognition Ltd, U.K) modified for use in mice, as described previously

(Humby *et al.*, 1999). For the PR task, only the central nose-poke hole was used. The mice

were presented with a visual stimulus (light) recessed into the holes and were trained to

respond to this stimulus with a nose-poke as recorded by infra-red beams spanning the hole.

Reward was presented in a recessed compartment on the wall opposite to the nose-

poke/stimulus array.   The control of the stimuli and recording of the responses were

managed by an Acorn Archimedes computer with additional interfacing by ARACHNID

(Cambridge Cognition Ltd). For all operant testing, animals were maintained on a restricted

water access schedule, water provided for two hours immediately after testing.

During training and testing, a nose-poke in the illuminated central hole resulted in the

presentation of 20 µl of an 10% condensed milk (Nestle) reward. Collection of this reward

initiated a subsequent trial. Conditioned reinforcement (CRf - one nose poke required for

reward delivery) was carried out for five days. Following this, a progressive ratio (PR)

schedule was carried out. Here, the number of nose pokes required to receive a reward

ascended linearly every four trials for three (FR4) sessions. FR4 sessions were followed by

three FR2 sessions, where the number of nose pokes required to receive a reward

ascended linearly every two trials. These PR sessions were followed by three CRf sessions.

A number of measures were taken, including number of rewards received, latency to first

nose-poke and latency to collect the reward. During the PR sessions, an additional measure,

the maximum number of nose pokes an animal was willing to make to receive a reward, was

deemed the "breakpoint" and was the main indication of the animal's motivation to work for a

reward.

Data analysis and statistics

All behavioral data were analysed using SPSS 20 (SPSS, USA). Data were assessed for

normality and then analysed by Student's t-test or mixed ANOVA, with between-subjects

factors of GENOTYPE ($Grb10^{+/p}$ vs. WT), and within-subject factors BIN; DAY; CHAMBER

(start, middle or reward chamber of PORT task); ODOR (control or fox odors in PORT and

POE-ASR); DAY (day of testing on CMT); SESSION (CRf, FR4 or FR2 session in PR task).

For repeated-measures analyses, Mauchly's test of sphericity of the covariance matrix was

applied; significant violations from the assumption of sphericity were subject to the Huynh–

Feldt correction to allow more conservative comparisons by adjusting the degrees of

freedom. Preference for exploring the novel environment in the NPP test was tested using

Kolmogolov Smirnov test. All significance tests were performed at an alpha level of 0.05.

RESULTS

$Grb10^{+/p}$ mice show normal reactivity to novel environments

We used two measures to assess the reactivity of $Grb10^{+/p}$ and WT mice to novel

environments. Firstly, locomotor activity was measured in activity chambers over three

consecutive days. On day 1 of testing, $Grb10^{+/p}$ and WT mice showed robust levels of

activity that reduced over the course of the two hour session (Figure 1A; main effect of BIN,

$F_{23,506}=11.55$, $P<0.001$, partial $\eta^2=0.344$). This habituation to the environment over time was

Figure 1 Locomotor activity and novelty place preference behavior in in *Grb10*^{+/p} and WT littermates. As the LMA session progress, activity reduces (**A**), a pattern also seen across consecutive days (**B**). However, there were no activity differences detected between *Grb10*^{+/p} and WT mice. Similarly, in the NPP test (**C**) all animals showed a preference in the proportion of time spent in the novel environment, but there were no differences detected between *Grb10*^{+/p} and WT mice. This was supported by other measures in the NPP test, including absolute time (**D**), number of entries into (**E**) and latency to first enter (**F**) the novel environment. Data are mean values ±SEM. # (P<0.05) and ## (P<0.01) indicate within subject (factors BIN or DAY) differences .

also seen over consecutive daily sessions, with the total activity levels being highest on Day

1 and reducing with consecutive daily sessions (Figure 1B; main effect of DAY,

$F_{1.30,25.92}=7.70$, P=0.006, partial $\eta^2=0.278$). However, there were no significant differences

between *Grb10*^{+/p} and WT mice either in total levels of activity across the first two hour

session (Figure 1A; main effect of GENOTYPE, $F_{1,22}=0.48$, P=0.496, partial $\eta^2=0.021$), or the

three days (Figure 1B; main effect of GENOTYPE, $F_{1,20}=0.004$, P=0.952, partial $\eta^2=0.0002$),

or in the degree of habituation, as inferred from the rate of change in activity and indicated

by the lack of a significant interaction between GENOTYPE and BIN on Day 1 (Figure 1A; $F_{23,506}$=1.08, P=0.363, partial η^2=0.047), and GENOTYPE and DAY across all locomotor activity sessions (Figure 1B; $F_{1.30,25.92}$=0.42, P=0.572, partial η^2=0.021).

We then explicitly measured the willingness of $Grb10^{+/p}$ and WT mice to explore a novel environment using a novelty place preference test (NPP). During the test phase, both WT and $Grb10^{+/p}$ mice spent significantly more time than by chance in the novel chamber (approximately 60%; Figure 1C; Kolmongolov Smirnov test, WT P=0.003, $Grb10^{m/-}$ P=0.048). Analysis of absolute measures suggested there was no distinction between $Grb10^{+/p}$ and WT mice in the total exploration of the novel chamber, with no significant differences in total time (Figure 1D; t=-0.14, P=0.89), number of entries (Figure 1E; t=0.77, P=0.45) or latency to first enter (Figure 1F; t=-0.16, P=0.87) the novel chamber.

$Grb10^{+/p}$ mice show increased risk-taking in the PORT task

There were no differences between $Grb10^{+/p}$ and WT mice during the 20-min session of habituation to the PORT apparatus (Supplementary Information, Table S1). $Grb10^{+/p}$ and WT mice spent equivalent amounts of time (main effect of GENOTYPE, $F_{1,22}$ = 0.85, P=0.37 partial η^2=0.037) and made the same number of entries to each chamber of the PORT apparatus (main effect of GENOTPYE, $F_{1,25}$ = 1.13, P=0.30 partial η^2=0.049). More entries were made in the middle chamber (main effect of CHAMBER, $F_{1.52,\ 33.47}$=120.2, P = 0.0001, partial η^2=0.85), as might be expected as the mice traversed the apparatus, and this was also reflected in an increase in the amount of time spent in this chamber (main effect of CHAMBER, $F_{1.24,\ 33.47}$=856.0, P = 0.0001, partial η^2=0.98). During task acquisition, the mice were trained to cross the apparatus to collect a reward, passing through the middle chamber which had plain wood shavings on the floor (Supplementary Results Figure S1). Both $Grb10^{+/p}$ and WT mice spontaneously demonstrated this behavior, and there was no difference between $Grb10^{+/p}$ and WT mice (main effect of GENOTYPE, $F_{1,22}$ = 3.78, P=0.072 partial η^2=0.20). These data suggest that habituation and task acquisition were equivalent between $Grb10^{+/p}$ and WT mice.

As expected (Dent *et al.*, 2014, Schneider *et al.*, 2015, Viola *et al.*, 2019), the introduction of a predator odor into the middle chamber of the test arena significantly increased the overall latency to leave the start chamber, relative to the presence of a control odor (Figure

Figure 2. Wild type and *Grb10*$^{+/p}$ behavior in PORT task. **A** All but one wild type animal showed an increase in latency to leave the Start chamber when a predator odor (fox) was introduced into the middle chamber of the apparatus, relative to a control odor (wood shavings from the mouse's home cage). **B** In contrast, only 5/10 *Grb10*$^{+/p}$ mice showed an increase upon introduction of the predator odor, and the overall magnitude of change in latency to leave the Start chamber was reduced. Representative traces from single trials of a wild type mouse with control bedding (**C**) and fox odor (**D**); and a *Grb10*$^{+/p}$ mice with control bedding (**E**) and fox odor (**F**). Videos of these trials can be found in the Supplementary material.

2a and b, main effect of ODOR, $F_{1,22}$=14.73, P=0.001, partial η^2=0.40). However, this effect was more pronounced in WT mice (significant interaction between GENOTYPE and ODOR (ANOVA, $F_{1,22}$=6.75, P=0.016, partial η^2=0.24). *Post-hoc* pairwise comparisons indicated that whilst latency to leave the start chamber was equivalent in control conditions (*P*=0.751), in the presence of the fox odor *Grb10*[+/p] mice had significantly reduced latencies compared to WT mice (*P*=0.016), being on average 85s (60%) quicker (Figure 2b). Furthermore, whilst all but one of the 14 WT animals showed an increase in the time in the Start chamber following the introduction of fox odor, only half (5/10) of *Grb10*[+/p] mice showed a similar slowing of the latency to leave the Start chamber.

As previously (Dent et al., 2014), we confirmed that the presence of the predator odor induced an equivalent fear response in the *Grb10*[+/p] and WT mice. Exposure to fox odor caused a significantly enhanced (24% increase) acoustic startle response (ASR) relative to prior exposure to control bedding (Figure 3a and b, main effect of ODOR, $F_{1,22}$=11.34, P=0.003, partial η^2=0.34), an increase that was equivalent in *Grb10*[+/p] and WT mice (main effect of GENOTYPE, $F_{1,22}$=2.07, P=0.164, partial η^2=0.09).

Figure 3. Wild type and *Grb10*[+/p] behavior in PORT task. Acoustic startle response in both wild types (**A**) and *Grb10*[+/p] mice (**B**) showed an equivalent increase following pre-exposure predator odor in comparison to control odor.

Motivation for the food reward is not altered in Grb10[+/p]* mice*

We performed the CMT to examine consumption and preference for a novel palatable foodstuff, namely 10% condensed milk. The total volume of milk consumed increased across successive days (Figure 4A; main effect of DAY, $F_{3.10,68.2}$=33.28, P<0.001, partial η^2=0.602), but did not differ between *Grb10*[+/p] and WT mice (main effect of GENOTYPE, $F_{1,22}$=0.34, P=0.565, partial η^2=0.015). All animals showed an initial aversion to the novel foodstuff

(preference of less than 50%), but with subsequent exposures acquired a preference for the

milk reward over water (preference of approximately 75%) (Figure 4B; main effect of DAY,

$F_{2.66,58.6}$=25.63, P<0.001, partial η^2=0.54). However, there was no difference between

$Grb10^{+/p}$ and WT mice in either their overall preference (main effect of GENOTYPE,

$F_{1,22}$=1.79, P=0.194, partial η^2=0.075) nor in the rate at which their preference was acquired

(interaction between GENOTYPE and DAY, $F_{2.66,58.6}$=1.23, P=0.305, partial η^2=0.53). These

data are from the animals that went on to the progressive ratio test (Cohort 2), but a similar

pattern of results was seen in animals who went on to be tested in the PORT task (Cohort 1,

see Supplementary Information, Figure S2).

We then examined the motivation to work for a palatable solution using a progressive ratio

task. Mice were initially trained to respond on a conditioned reinforcement (CRf) schedule for

5 sessions. During the CRf sessions, subjects were able to carry out a maximum of 100

trials, which is equal to 100x 22µl rewards within each 30min test session. Analysis of the

initial CRf stage revealed that both genotypes achieved the required level of performance

per session, showing no effect of genotype ($t_{1,19}$=0.80, p=0.44; data not shown). Following

CRf training, subjects were switched to the two progressive ratio (PR) schedules: first FR4,

followed by FR2. The breakpoint (BP) was defined as the maximum number of nose pokes

an animal was willing to make to receive a reward and is an indication of the animal's

motivation to work for a reward. To demonstrate the effects of the imposition of the PR

schedule, performance during the PR sessions were compared to the average of the three

CRf sessions following PR testing (Figure 3C). Imposition of the PR schedule led to a

significant reduction in the number of rewards earned within a session (main effect of

SESSION, $F_{1.24,23.6}$=160.36, P=8.80E-13, partial η^2=0.89). There were no differences

between $Grb10^{+/p}$ and WT mice (main effect of GENOTYPE, $F_{1,19}$=0.45, P=0.510, partial

η^2=0.023). This decrease in rewards earned was not due to mice running out of time to

collect all the available rewards, as the average PR session did not run for the full 30

minutes and there were no significant differences in session duration between the PR (FR4

18.2mins ±1.8; FR2 19.1mins ±1.7) and CRf (21.0mins ±1.4) sessions (main effect of

SESSION $F_{1.44,27.28}$=1.36, P=0.267, partial η^2=0.067). Although $Grb10^{+/p}$ appeared to have a

higher breakpoint, the main PR measure, in both FR4 and FR2 (Figure 4D), this did not

reach significance (main effect of GENOTYPE, $F_{1,19}$=1.15, P=0.296, partial η^2=0.057). This

suggests an equivalent level of motivation to work for the food reward between the $Grb10^{+/p}$

and WT mice. This finding was underlined by no differences in latency measures, such as

latency to first nose-poke (WT 7.97s ±1.05, $Grb10^{+/p}$ 9.97s ±1.21; main effect of

GENOTYPE, $F_{1,19}$=1.56, P=0.227, partial η^2=0.076) and latency to collect reward (WT 1.48s

±0.28, *Grb10*^{+/p} 1.84s ±0.32; main effect of GENOTYPE, $F_{1,19}=0.73$, $P=0.403$, partial $\eta^2=0.037$).

Figure 4 Palatable food consumption and progressive ratio behavior in *Grb10*^{+/p} and WT littermates. Consumption (**A**) and preference (**B**) for 10% condensed milk increased with successive sessions but was not different between *Grb10*^{+/p} and WT littermates. In the PR task imposition of the FR4 and FR2 reduced the total number of trials relative to CRf, but there was no difference between *Grb10*^{+/p} and WT littermates (C). Similarly, the BPs at FR4 and FR2 were also equivalent between *Grb10*^{+/p} and WT mice. ## indicates within subject (factor DAY) differences P<0.01.

DISCUSSION

Grb10 is currently a unique example of an imprinted gene in which the different parental alleles show distinct patterns of expression and have distinct physiological functions (Garfield *et al.*, 2011). We have previously demonstrated that mice lacking a paternal copy of *Grb10* have altered behavior (Rienecker *et al.*, 2020), including a higher tolerance of delayed rewards in a delay reinforcement task (DRT) (Dent *et al.*, 2018). One suggestion is that these changes reflect a role for *Grb10* in regulating risk-taking behavior broadly (Wilkins & Bhattacharya, 2019). We examined this in a number of tests showing that, although *Grb10*^{+/p} mice explore a novel environment to the same extent as their wild-type littermates, they are

more risk-taking on the Predator Odor Risk-taking (PORT) task. This is the direction of effects predicted from our previous analysis on the DRT (Dent *et al.*, 2018) and by evolutionary theory (Wilkins & Bhattacharya, 2019), and taken together these data suggest that *Grb10* normally influences the cost vs. risk assessment and acts to reduce risk-taking.

To test whether loss of paternal *Grb10* expression would influence risk-taking we took a broad approach. Firstly, we examined the propensity of *Grb10*[+/p] mice to explore novelty, both in terms of basic locomotor activity and habituation to a new environment, and also an explicit test of investigation of a novel environment. In both tests, the behaviour of *Grb10*[+/p] mice was equivalent to wild-type littermates. These data were supported by the condensed milk test, which was used to assess consumption and preference for a palatable substance but can also be regarded as a measure of food neophobia. The prediction from the behavior of *Grb10*[+/p] mice on the DRT (Dent *et al.*, 2018) would be that they will be more willing to take risks and therefore explore a novel environment more (or more quickly). It is possible that these tests are not sensitive enough, or that there is a ceiling effect, to detect such a 'positive' change. Nevertheless, it seems that across novelty domains, *Grb10*[+/p] mice appear to behave normally.

We also used an explicit test of risk-taking to examine the behavior of *Grb10*[+/p] mice, namely the PORT task. This task was developed by us (Dent *et al.*, 2014), and has been used by others to assess risk-taking in both mice (Blomeley *et al.*, 2018, Viola *et al.*, 2019) and rats (Schneider *et al.*, 2015). The PORT task examines a more ecologically valid aspect of risk-taking, namely the trade-off between a food reward and the risk of predation in obtaining that reward. Behavior in the task is sensitive to changes that affect this balance, such as the presence of a predator odor, or a reduction in the value of the reward (Dent *et al.*, 2014). In the PORT task, *Grb10*[+/p] mice are quicker than wild-type littermates to leave the start chamber in the presence of a predator odor (fox) in order to obtain the food reward. There were no differences in habituation or acquisition of the task, and no difference in latencies in control trials, suggesting that the *Grb10*[+/p] mice are more willing to take risks. Importantly, the difference in behavior in the PORT task is not due to changes in either fear, or motivation for palatable food alone, as predator odor enhanced acoustic startle and behavior in a progressive ratio task were equivalent between *Grb10*[+/p] and wild-type mice. This suggests that loss of paternal *Grb10* alters the point of trade-off between a obtaining a food reward and the risk of predation in obtaining that reward.

The direction of effects in the PORT task, where *Grb10*[+/p] mice show increased risk-taking, is consistent with findings from the DRT. Here, *Grb10*[+/p] mice were more willing to wait for a large, but delayed, food reward. Although not developed as a direct test, behavior in discounting tasks such as the DRT have been correlated with (Hayden & Platt, 2007), and have been used as a proxy measure for, risk-taking (Bateson & Kacelnik, 1997, Holt *et al.*,

2003, Kacelnik & Bateson, 1996). Taken together, these data suggest that in the brain paternal *Grb10* normally influences the cost vs. risk decision-making, and acts to make mice more risk-averse. Moreover, this is also in line with the predicted direction of effects of how the paternal genome will influence sensitivity to risk (Wilkins & Bhattacharya, 2019). Maternal and paternal genomes are expected to have conflicting influences on risk-tolerance as a consequence of differences in reproductive variance between males and females. When reproductive variance is higher in males (as it is for most mammals) then paternally expressed imprinted genes will promote risk-averse, variance-reducing behaviors, and maternally expressed imprinted genes will promote risk-tolerant, variance-increasing behaviors. This pattern is not only seen in $Grb10^{+/p}$ mice, but also in mice lacking maternal *Nesp* that show reduced exploration of a novel environment (Plagge *et al.*, 2005) and decreased tolerance of delay in the DRT (Dent *et al.*, 2016). This suggests opposite effects of *Grb10* and *Nesp* on risk-taking behaviors. Interestingly, these genes show a strong degree of co-localisation (Dent *et al.*, 2018), including in neurons of the Dorsal Raphé Nucleus (DRN) and Locus Coeruleus (LC), two brain areas known to modulate risk-taking behaviours (Sara *et al.*, 1995, Xu *et al.*, 2017).

Here we test the idea that the imprinted gene *Grb10* is involved in modulating risk-taking behavior. Although not true across all domains, mice lacking paternal *Grb10* do show increased risk-taking on the PORT task. This, coupled with previous work on a delayed reinforcement task, suggest that this idea is broadly correct. Taken together with our previous work on another imprinted gene, *Nesp*, these data suggest that maternally and paternally expressed imprinted genes oppositely influence risk-taking behaviour. This "parliament of the mind" (Haig, 2000), caused by opposing parental genomes pulling impulsive choice and risk-taking in different directions, is an additional factor that should be taken into consideration when considering apparently irrational or sub-optimal choice behaviors (Fawcett *et al.*, 2014, Hayden, 2019).

REFERENCES

Bateson, M. & Kacelnik, A. (1997) Starlings' preferences for predictable and unpredictable delays to food. *Anim Behav,* **53,** 1129-1142.

Benson, K.E. & Stephens, D.W. (2015) Interruptions, Tradeoffs, and Temporal Discounting1. *American Zoologist,* **36,** 506-517.

Blanchard, M.M., Mendelsohn, D. & Stamp, J.A. (2009) The HR/LR model: Further evidence as an animal model of sensation seeking. *Neuroscience and biobehavioral reviews,* **33,** 1145-1154.

Blomeley, C., Garau, C. & Burdakov, D. (2018) Accumbal D2 cells orchestrate innate risk-avoidance according to orexin signals. *Nature neuroscience,* **21,** 29-32.

Dent, C.L., Humby, T., Lewis, K., Plagge, A., Fischer-Colbrie, R., Wilkins, J.F., Wilkinson, L.S. & Isles, A.R. (2016) Impulsive choices in mice lacking imprinted Nesp55. *Genes, brain, and behavior,* **15,** 693-701.

Dent, C.L., Humby, T., Lewis, K., Ward, A., Fischer-Colbrie, R., Wilkinson, L.S., Wilkins, J.F. & Isles, A.R. (2018) Impulsive Choice in Mice Lacking Paternal Expression of Grb10 Suggests Intragenomic Conflict in Behavior. *Genetics,* **209,** 233-239.

Dent, C.L. & Isles, A.R. (2014) Brain-expressed imprinted genes and adult behaviour: the example of Nesp and Grb10. *Mammalian genome : official journal of the International Mammalian Genome Society,* **25,** 87-93.

Dent, C.L., Isles, A.R. & Humby, T. (2014) Measuring risk-taking in mice: balancing the risk between seeking reward and danger. *The European journal of neuroscience,* **39,** 520-530.

Fawcett, T.W., Fallenstein, B., Higginson, A.D., Houston, A.I., Mallpress, D.E., Trimmer, P.C., McNamara, J.M. & Modelling Animal Decisions, G. (2014) The evolution of decision rules in complex environments. *Trends in cognitive sciences,* **18,** 153-161.

Garfield, A.S., Cowley, M., Smith, F.M., Moorwood, K., Stewart-Cox, J.E., Gilroy, K., Baker, S., Xia, J., Dalley, J.W., Hurst, L.D., Wilkinson, L.S., Isles, A.R. & Ward, A. (2011) Distinct physiological and behavioural functions for parental alleles of imprinted Grb10. *Nature,* **469,** 534-538.

Green, L. & Myerson, J. (1996) Exponential versus hyperbolic discounting of delayed outcomes: Risk and waiting time. *American Zoologist,* **36,** 496-505.

Haig, D. (2000) Genomic imprinting, sex-biased dispersal, and social behavior. *Ann N Y Acad Sci,* **907,** 149-163.

Hayden, B.Y. (2019) Why has evolution not selected for perfect self-control? *Philosophical transactions of the Royal Society of London. Series B, Biological sciences,* **374,** 20180139.

Hayden, B.Y. & Platt, M.L. (2007) Temporal discounting predicts risk sensitivity in rhesus macaques. *Current biology : CB,* **17,** 49-53.

Holt, D.D., Green, L. & Myerson, J. (2003) Is discounting impulsive?. Evidence from temporal and probability discounting in gambling and non-gambling college students. *Behav Process,* **64,** 355-367.

Humby, T., Laird, F.M., Davies, W. & Wilkinson, L.S. (1999) Visuospatial attentional functioning in mice: interactions between cholinergic manipulations and genotype. *European Journal of Neuroscience,* **11,** 2813-2823.

Kacelnik, A. & Bateson, M. (1996) Risky theories - The effects of variance on foraging decisions. *American Zoologist,* **36,** 402-434.

Kacelnik, A. & Bateson, M. (1997) Risk-sensitivity: crossroads for theories of decision-making. *Trends in cognitive sciences,* **1,** 304-309.

Moore, T. & Haig, D. (1991) Genomic imprinting in mammalian development - a parental tug-of-war. *Trends in Genetics,* **7,** 45-49.

Plagge, A., Isles, A.R., Gordon, E., Humby, T., Dean, W., Gritsch, S., Fischer-Colbrie, R., Wilkinson, L.S. & Kelsey, G. (2005) Imprinted nesp55 influences behavioral reactivity to novel environments. *Mol Cell Biol,* **25,** 3019-3026.

Rienecker, K.D.A., Chavasse, A.T., Moorwood, K., Ward, A. & Isles, A.R. (2020) Detailed analysis of paternal knockout Grb10 mice suggests effects on stability of social behavior, rather than social dominance. *Genes, brain, and behavior,* **19,** e12571.

Sara, S.J., Dyon-Laurent, C. & Herve, A. (1995) Novelty seeking behavior in the rat is dependent upon the integrity of the noradrenergic system. *Brain Res Cogn Brain Res,* **2,** 181-187.

Schneider, M., Kasanetz, F., Lynch, D.L., Friemel, C.M., Lassalle, O., Hurst, D.P., Steindel, F., Monory, K., Schafer, C., Miederer, I., Leweke, F.M., Schreckenberger, M., Lutz, B., Reggio, P.H., Manzoni, O.J. & Spanagel, R. (2015) Enhanced Functional Activity of the Cannabinoid Type-1 Receptor Mediates Adolescent Behavior. *The Journal of neuroscience : the official journal of the Society for Neuroscience,* **35,** 13975-13988.

Schonberg, T., Fox, C.R. & Poldrack, R.A. (2011) Mind the gap: bridging economic and naturalistic risk-taking with cognitive neuroscience. *Trends in cognitive sciences,* **15,** 11-19.

Smith, F.M., Holt, L.J., Garfield, A.S., Charalambous, M., Koumanov, F., Perry, M., Bazzani, R., Sheardown, S.A., Hegarty, B.D., Lyons, R.J., Cooney, G.J., Daly, R.J. & Ward, A. (2007) Mice with a disruption of the imprinted Grb10 gene exhibit altered body composition, glucose homeostasis, and insulin signaling during postnatal life. *Molecular and cellular biology,* **27,** 5871-5886.

Viola, T.W., Wearick-Silva, L.E., Creutzberg, K.C., Kestering-Ferreira, E., Orso, R., Centeno-Silva, A., Albrechet-Souza, L., Marshall, P.R., Li, X., Bredy, T.W., Riva, M.A. & Grassi-Oliveira, R. (2019) Postnatal impoverished housing impairs adolescent risk-assessment and increases risk-taking: A sex-specific effect associated with histone epigenetic regulation of Crfr1 in the medial prefrontal cortex. *Psychoneuroendocrinology,* **99,** 8-19.

Wilkins, J.F. & Bhattacharya, T. (2019) Intragenomic conflict over bet-hedging. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences,* **374,** 20180142.

Wilkins, J.F., Ubeda, F. & Van Cleve, J. (2016) The evolving landscape of imprinted genes in humans and mice: Conflict among alleles, genes, tissues, and kin. *Bioessays,* **38,** 482-489.

Xu, S., Das, G., Hueske, E. & Tonegawa, S. (2017) Dorsal Raphe Serotonergic Neurons Control Intertemporal Choice under Trade-off. *Current biology : CB,* **27,** 3111-3119 e3113.