An Effective CTL Peptide Vaccine for Ebola Zaire Based on Survivors' CD8+ Targeting of a Particular Nucleocapsid Protein Epitope with Potential Implications for COVID-19 Vaccine Design

Herst CV[a], Burkholz S[a], Sidney J[i], Sette A[i], Harris PE[g], Massey S[b], Brasel T[b], Cunha-Neto E[c,d,e], Rosa DS[d,f], Chao WCH[j], Carback R[a], Hodge T[a], Wang L[a], Ciotlos S[a], Lloyd P[a], Rubsamen R[a,h,*]

[a]*Flow Pharma, Inc., 3451 Vincent Road, Pleasant Hill, CA 94523*
[b]*University of Texas, Medical Branch,301 University Blvd, Galveston, TX 77555*
[c]*Laboratory of Clinical Immunology and Allergy-LIM60, University of São Paulo School of Medicine, São Paulo, Brazil*
[d]*Institute for Investigation in Immunology (iii) INCT, São Paulo, Brazil*
[e]*Heart Institute (Incor), School of Medicine, University of São Paulo, São Paulo, Brazil*
[f]*Department of Microbiology, Immunology and Parasitology, Federal University of São Paulo (UNIFESP/EPM), São Paulo, Brazil*
[g]*Endocrinology Division, Department of Medicine, School of Medicine, Columbia University, New York, New York, USA*
[h]*Massachusetts General Hospital, Department of Anesthesia, Critical Care and Pain Medicine, 55 Fruit St, Boston, MA 02114*
[i]*La Jolla Institute for Allergy and Immunology, 9420 Athena Circle La Jolla, CA 92037*
[j]*University of Macau, E12 Avenida da Universidade, Taipa, Macau, China*

Abstract

The 2013-2016 West Africa EBOV epidemic was the biggest EBOV outbreak to date. An analysis of virus-specific CD8+ T-cell immunity in 30 survivors showed that 26 of those individuals had a CD8+ response to at least one EBOV protein. The dominant response (25/26 subjects) was specific to the EBOV nucleocapsid protein (NP). It has been suggested that epitopes on the EBOV NP could form an important part of an effective T-cell vaccine for Ebola Zaire. We show that a 9-amino-acid peptide NP44-52 (YQVNNLEEI) located in a conserved region of EBOV NP provides protection against morbidity and mortality after mouse adapted EBOV challenge. A single vaccination in a C57BL/6

[*]Corresponding author
    Email address: `reidrubsamen@alum.mit.edu` (
Rubsamen R)

mouse using an adjuvanted microsphere peptide vaccine formulation containing NP44-52 is enough to confer immunity in mice. Our work suggests that a peptide vaccine based on CD8+ T-cell immunity in EBOV survivors is conceptually sound and feasible. Nucleocapsid proteins within COVID-19 contain multiple class I epitopes with predicted HLA restrictions consistent with broad population coverage. A similar approach to a CTL vaccine design may be possible for that virus.

Keywords: Ebola Zaire vaccine, CTL Vaccine, controller, YQVNNLEEI, COVID-19, Flow Focusing

1. Introduction

Development of safe and effective vaccines for some viruses such as HIV and EBOV has been challenging [15]. Although vaccine development has been almost exclusively focused on eliciting a humoral immune response in the host
5 through inoculation with whole protein antigen [40][56][46][24], CTL peptide vaccines producing a T-cell response may offer an important alternative approach [19]. For HIV and EBOV and influenza in particular, the potential of CTL vaccines has been discussed [17][6][44]. Although computational prediction alone has been used for T-cell vaccine design [2][11], we saw a unique
10 opportunity to see if a preventative EBOV T-cell vaccine could be successfully designed based on the specific epitopes targeted by survivors of documented EBOV infection.

The notion of HLA restricted HIV control has been described [45]. Pereyra-Heckerman conducted an analysis of virus-specific CD8+ T-cell immunity in
15 individuals living with HIV [33]. They reported that HIV controllers, individuals living with HIV not undergoing treatment who do not progress to AIDS, have CD8+ cells targeting different HLA restricted class I epitopes on HIV compared with progressors, individuals with HIV who progress to AIDS in the absence of therapy. Pereyra-Heckerman suggested that this observation could guide the
20 *in-silico* development of a CTL vaccine for HIV and other diseases.

Acquired immunity has been documented after EBOV infection [3]. Antibody as well as T-cell responses have been described [34]. Sakebe et al have shown that of 30 subjects surviving the 2013-2016 EBOV outbreak in West Africa, CD8+ T-cells from 26 of those survivors responded to at least one
[25] EBOV antigen, with 25 of the 26 responders targeting epitopes on EBOV NP [39]. One of the most commonly targeted EBOV eptitopes on EBOV NP in the survivor group (targeted by CD8+ cells from four survivors) was NP41-60 (IPVYQVNNLEEICQLIIQAF). They also suggested that a CTL vaccine for could be designed using epitopes targeted by CD8+ T-cells identified in these
[30] EBOV controllers.

Human pathogen-derived peptide antigens that are also recognized by C57BL/6 T-cells have been previously described. These include peptides from vesicular stomatitis virus (VSV) RGYVYQGL [55], and human immunodeficiency virus (HIV) RGPGRAFVTI [4]. The existence of such epitopes makes possible a
[35] range of pre-clinical vaccine experiments without having to rely on non-human primates and expensive and complex-to-manage humanized mouse models. Wilson et al showed that the EBOV nucleoprotein (NP) is an immunogen that provides protective, CTL-mediated immunity against EBOV in a C57BL/6 mouse model and that this protection was conferred by a peptide sequence within Ebola
[40] Zaire: NP43-53 (VYQVNNLEEIC) [59]. Wilson et al came to this conclusion based on studying splenocytes harvested from mice vaccinated with Ebola Zaire NP using a Venezuelan equine encephalitis (VEE) vector. Their experiments showed that splenocytes from the vaccinated mice re-stimulated with NP43-53 had high levels of cytotoxic activity against target cells loaded with the EBOV
[45] NP peptide. Remarkably, NP43-53 also happens to be an 11 amino acid subsequence of the epitope identified by Sakebe et al, as most commonly favored for T-cell attack by survivors of the 2013-2016 EBOV outbreak in West Africa.

We set out to see if we could drive CTL expansion directed against NP43-53 to occur after vaccinating C57BL/6 mice with Ebola Zaire NP43-53 (VYQVNNLEEIC),
[50] and to subsequently conduct an *in-vivo* EBOV challenge study to see if this peptide was protective.

We fabricated adjuvanted microspheres for this study as a room temperature stable dry powder using the Flow Focusing process to be $11\mu M$ in diameter so as to prevent more than one microsphere from being phagocytosed by any given antigen presenting cell (APC) at the same time [29]. By loading only one peptide sequence per microsphere, we maximized the peptide payload and mitigated the possibility of multiple, different peptide sequences being delivered to the APC simultaneously, which could possibly result in competitive inhibition at the motif which could interfere with antigen presentation and subsequent T-cell expansion (Supplementary Material).

2. Results

We used a previously described biodegradable dry powder, PLGA microsphere, synthetic vaccine platform adjuvanted with TLR-4 and TLR-9 ligands [37] to immunize C57BL/6 mice with NP43-53 11-mer, the CTL+ class I peptide antigen from the Ebola Ziare NP protein identified by Wilson et al [59]. Microspheres containing NP43-53 and CpG were prepared as a dry powder formulation and suspended before use in a PBS injectate solution containing MPLA, and administered intradermally via injection at the base of the tail into mice as described in a previous publication [37]. Splenocytes from eight mice receiving the active vaccine, and eight mice receiving microspheres and adjuvants only, were harvested on day 14 and subjected to ELISPOT analysis evaluating $IFN - \gamma$ release in response to stimulation with the peptide used in the vaccination. As illustrated in Supplementary Material Figure 1, there was no statistically significant difference between the ELISPOT data for the vaccinated mice *versus* the response seen in the negative control group.

Wilson reported that protection seen in her experiment was due to a peptide sequence within NP-43-53. We hypothesized that the NP43-53 epitope was inefficiently processed into MHC binding sub-sequences during antigen presentation. In order to explore possible H2-D^b matches for peptide sequences contained within Ebola Zaire NP43-53 (VYQVNNLEEIC), we prepared three

9mer Subsequences of VYQVNNLEEIC
Evaluated for Immune Response

Peptide Label	Peptide Sequence	Description
NP43-53	VYQVNNLEEIC	Ebola Zaire NP 11mer peptide not H2-D^b matched
NP43-51	VYQVNNLEE	Sub-sequence 9mer VYQVNNLEEIC
NP44-52	YQVNNLEEI	Sub-sequence 9mer VYQVNNLEEIC
NP45-53	QVNNLEEIC	Sub-sequence 9mer VYQVNNLEEIC

Table 1: Class I peptides used in the study. NP43-53 is the class I 11mer described by Wilson et al. which we found not to produce an immune response in a C57BL/6 mouse model. NP43-51, NP 44-52 and NP 45-53 are the three possible 9mer sub-sequences of NP43-53.

peptide vaccine formulations, each containing one of the three possible 9mer sub-sequences within NP43-53. These sequences are shown in Table 1. We then vaccinated, via intradermal (tail) injection, three groups of mice with microspheres containing one of the three 9mer sub-sequences of NP43-53 (6 per group). ELISPOT analysis was performed, stimulating harvested splenocytes with the three possible 9mer sub-sequences. Splenocytes from mice receiving the NP44-52 sub-sequence had a statistically higher ELISPOT response than mice vaccinated with the other two possible sub-sequence 9mers (P < 0.0001) as shown in Figure 1. This is consistent with the predicted H2-D^b binding affinity of YQVNNLEEI as shown in Supplementary Material Table 3.

We then loaded one population of adjuvanted microspheres with NP44-52 and a second population of adjuvanted microspheres loaded with VG19 from EBOV Zaire NP 273-291 (VKNEVNSFKAALSSLAKHG), a Class II epitope predicted to be relevant to NP43-53 based on the TEPITOPE algorithm using a technique described by Cunha-Neto at al [11]. This peptide has a predicted favorable H2-I^b binding affinity as shown in Supplementary Material Table 5.

We vaccinated 16 mice with those two populations of microspheres via intradermal (tail) administration and an 12 additional mice with two populations of

microspheres, one population containing NP43-53 and the other VG19. $IFN-\gamma$ release as quantified by ELISPOT after the spleens were harvested on day 14 after immunization showed that the immune response to the 9mer NP44-52 was higher than the immune response after vaccination with NP43-53 and that this difference was statistically significant (P < 0.0001) Figure 2.

We conducted a pilot study demonstrating that intraperitoneal injection produced a statistically superior immune response by ELISPOT compared with intradermal tail injection in C57BL/6 mice (Supplementary Material). Based on the data from that study, we chose to proceed with intraperitoneal administration for the challenge portion of this study.

We dosed three groups of mice, ten mice per group, with the adjuvanted microsphere vaccine formulation containing NP44-52 and VG-19, with each peptide in a distinct microsphere population, and challenged these mice 14 days after vaccine administration with escalating IP administered doses of mouse adapted EBOV (maEBOV) (Group 3 - 100 PFU, Group 5 - 1000 PFU and Group 7 - 10,000 PFU). The composition of the vaccine used for the exposure study is described in Supplementary Material. A second set of three control groups of mice (groups 2, 4 and 6), ten mice per group (mock groups), received PBS buffer solution alone and served as control animals for the study and were similarly challenged with maEBOV. Group 1 animals served as study controls and received no PBS buffer, vaccine or maEBOV injections. All mice were sourced from Jackson Labs and were 6-8 weeks of age and 15-25 grams at the time of vaccination. The dosing regimen is outlined in Table 2.

Peak mortality across all groups tested was seen in mice challenged with 1,000 PFU maEBOV *versus* PBS buffer control as shown in the survival curve in Figure 3a. Clinical observation data shown in Figure 3b and Figure 3c and daily weight data shown in Figure 3d and Figure 3e show protection from morbidity in all active vaccinated mice exposed to 1,000 PFU maEBOV.

PBS buffer mock-vaccinated mice showed mortality increasing from the 100 PFU to 1,000 PFU as shown in Figure 4a and Figure 3a. We saw a paradoxical effect in control animals with survival increasing between 1,000PFU (Figure 3a)

130 and 10,000 PFU (Figure 5a). We believe this was caused by innate immunity
triggered by the very large maEBOV challenge. All mice in all vaccinated
groups across both experiments survived and showed no morbidity by clinical
observation scores and weight data.

For each of the three challenge levels, the difference between the number of
135 survivors in the vaccinated group *versus* the PBS control group was statistically
significant by chi square (100 PFU P = 0.001; 1000 PFU P = 0.0003; 10,000
PFU P = 0.003)

Serum samples from sacrificed animals exposed to EBOV who did not receive
vaccine were quantitatively assayed for various cytokines using BioPlex plates.
140 Animals having unwitnessed demise did not have serum samples collected. A
Pearson Correlation Analysis was performed to assess relationships between
specific cytokine levels and survival. The results are shown in Table 3.

We observed low levels of IL-6 in surviving mice. NHPs infected with EBOV
have been determined by other researchers to have elevated levels of IL-6 in
145 plasma and serum [23][14]. EBOV infected humans have also shown elevated
IL-6 levels and these elevated levels have been associated with increased morality
[58].

Similarly, we observed low levels of MCP-1, IL-9 and GM-CSF in survivors.
Increased serum and plasma levels of MCP-1 have been observed in EBOV
150 infected NHPs [18][23][14] and elevated levels of MCP-1 were associated with
fatalities in EBOV infected human subjects [58]. Human survivors of EBOV
have been found to have very low levels of circulating cytokines IL-9 and elevated
levels of GM-CSF have been associated with fatality in humans exposed to
EBOV [58].

155 We saw increased levels of $IFN - \gamma$ in survivors. Other vaccine studies have
associated $IFN - \gamma$ with protection [57][30].

Dosing Table

Vaccinated Animals *versus* PBS Controls

100, 1000, and 10,000 PFU maEBOV Challenge

Group	N	Active / Control	Formulation	Route	Challenge
1	4	Control	N/A	N/A	N/A
2	10	Control	PBS	400μl IP	100PFU maEBOV
3	10	Active	10mg Adjuvanted Microspheres with NP44-52	400μl IP	100 PFU maEBOV
			10mg Adjuvanted Microspheres with VG-19		
4	10	Control	PBS	400μl IP	1,000PFU maEBOV
5	10	Active	10mg Adjuvanted Microspheres with NP44-52	400μl IP	1,000 PFU maEBOV
			10mg Adjuvanted Microspheres with VG-19		
6	10	Control	PBS	400μl IP	10,000PFU maEBOV
7	10	Active	10mg Adjuvanted Microspheres with NP44-52	400μl IP	10,000 PFU maEBOV
			10mg Adjuvanted Microspheres with VG-19		

Table 2: C7BL/6 maEBOV challenge study dosing regimen with PBS (buffer) controls. All challenges were done with Ebola virus M. musculus/COD/1976/Mayinga-CDC-808012 (maE-BOV) delivered IP. Mice in Group 1 received no injections.

3. Summary and Discussion

Most preventative vaccines are designed to elicit a humoral immune response, typically via the administration of whole protein from a pathogen. In contrast, a T-cell vaccine is meant to elicit a cellular immune response directing CD8+ cells to expand and attack cells presenting the HLA Class I restricted pathogen-derived peptide antigen. Difficulty in obtaining a reliable immune response from peptide antigens and the HLA restricted nature of CTL vaccines have limited their utility to protect individuals from infectious disease [61]. However, observations derived from individuals able to control HIV infection [33] and EBOV infection [39] demonstrating that control may be associated with specific CTL targeting behavior, suggest that there may be an important role for HLA restricted peptide vaccines for protection against infectious disease for which development of an effective traditional whole protein vaccine has proved to be difficult. The adjuvanted microsphere peptide vaccine platform described here incorporates unmodified peptides making possible rapid manufacture and deployment to respond to a new viral threat.

NP44-52 is located within one of the EBOV nucleocapsid proteins considered

essential for virus replication. This epitope resides in a sequence conserved across multiple EBOV strains as shown in Supplementary Material Figure 7. A 7.3Å structure for NP and VP24 is shown for context in Figure 6a [54]. A 1.8Å resolution structure rendering for EBOV NP shown in Figure 6b illustrates that NP44-52 is a buried structural loop, which is likely to be important to the structural integrity of the EBOV NP protein [13]. This structural role of NP44-52 likely explains its conservation across EBOV strains.

CTL targeting of the EBOV NP protein has been described [38][20]. Nucleocapisid proteins are essential for EBOV replication [48]. Recent advances in T-cell based vaccines have focused on avoiding all variable viral epitopes and incorporating only conserved regions [6][21]. EBOV NP may more conserved than nucleocapsid proteins VP35 and VP24 making it more suitable as a CTL vaccine target [7][59]. The nucleocapsid proteins in SARS-CoV are also essential for that virus to function normally [8]. This suggests that a CTL vaccine targeting coronavirus nucleocapsid could be effective against SARS-CoV or COVID-19.

We have shown that an H2-D^{b} restricted Class I peptide exists within the NP41-60 epitope identified by Sakebe et al as the most commonly favored NP epitope for CD8+ attack by survivors of the 2013-1016 EBOV outbreak in West Africa. We have demonstrated, when delivered in conjunction with a predicted-matched Class II epitope using an adjuvanted microsphere peptide vaccine platform, NP44-52 protection against mortality and morbidity for the maEBOV challenge doses tested in a C57BL/6 mouse model. We accomplished this with an adjuvanted, microsphere-based, synthetic CTL peptide vaccine platform producing a protective immune response 14 days after a single administration.

We saw what appears to be an innate immune response at the 10,000 PFU EBOV exposure level. It has been suggested that EBOV can mediate an innate immunity response through stimulation of TLR-4 [28]. Because the adjuvanted microsphere vaccine used in this experiment incorporates a TLR-4 agonist, we dosed 10 mice with adjuvanted microspheres without peptides and found the level of protection after exposure to 100 PFU EBOV to be statistically no different from that seen in PBS buffer controls. We conclude that level of protection

conferred by the adjuvanted vaccine described in this study is dependant on delivering peptides with the microspheres.

EBOV can cause severe pulmonary problems in exposed subjects [31]. These problems can be especially severe when the virus is delivered by aerosol [12][26]. Interaction of EBOV specific antibody, NHP lung tissue and EBOV delivered to NHPs via aerosol can produce a more lethal effect than in NHPs without circulating anti-EBOV antibody exposed to aerosolized EBOV (unpublished conference presentation). This suggests that a CTL vaccine may be more effective for prophylaxis against filovirus protection than an antibody vaccine if the anticipated route of EBOV exposure is via aerosol.

Sakebe et al identified A*30:01:01 as the only HLA type common to all four survivors in their study with CD8+ targeting of NP41-60. The A*30 supertype is relatively common in West Africa: 13.3% for Mali, 15.4% for Kenya, 16.3% for Uganda, and 23.9% for Mozambique [27]. Although peptide vaccines are by their nature HLA restricted, it may be possible to create a CTL vaccine directed against EBOV for use alone or in conjunction with a whole protein vaccine to produce an antibody response in tandem, by incorporating additional Class I peptides from epitopes targeted by controllers to broaden the HLA coverage of the vaccine. MHC binding algorithms hosted by the IEDB predict that YQVNNLEEI will bind strongly to the MHC of HLA-A*02:06, HLA-A*02:03 and HLA-A*02:01 individuals (Supplementary Material Table 2) [52]. HLA-DR binding database analysis also suggests that VKNEVNSFKAALSSLAKHG demonstrates sufficiently promiscuous binding characteristics cover that same population (Supplementary Material Table 4) [52]. Taken together, a peptide vaccine based on YQVNNLEEI and VKNEVNSFKAALSSLAKHG could produce a cellular immune response in about 50% of the population of the Sudan and about 30% of the population of North America.

The internal proteins located within influenza virus, in contrast to the glycoproteins present on the surface, show a high degree of conservation. Epitopes within these internal proteins often stimulate T-cell-mediated immune responses. As a result, vaccines stimulating influenza specific T-cell immunity

have been considered as candidates for a universal influenza vaccine [53].

This same approach could be applied to COVID-19 which has conserved regions in nucleocapsid which is located within the virus (see multiple sequence alignment in Supplementary Material Figure 8 and Supplementary Material Figure 9). Antigenic escape allows a virus to retain fitness despite an immune response to vaccination [16]. Picking conserved regions for vaccine targeting is an important part of mitigating this problem. Coronavirus spike protein, for example, may be particularly susceptible to mutation meaning that antigenic escape would be likely if the spike protein was targeted by a coronavirus vaccine, making it difficult to achieve durable protection [60]. A recent paper conducted a population genetic analysis of 103 COVID-19 genomes showing that the virus has evolved into two major types: L and S, with changes in their relative frequency after the outbreak possibly due to human intervention resulting in selection pressure [49].

We took all possible 424 9mer peptide sequences from the COVID-19 nucleocapsid protein sequences available and evaluated each peptide for HLA restriction using NetMHC 4.0 and NetMHCpan 4.0 [52][25][1]. We analyzed 9mer peptide sequences because these are often associated with superior MHC binding properties than class I peptides of other lengths [50]. We found 53 unique peptides with predicted binding below 500nM from NetMHC 4.0 and/or NetMHCpan 4.0. These results are shown in Supplementary Material Table 6, Supplementary Material Table 7, Supplementary Material Table 8 and Supplementary Material Table 9.

We proceeded to determine the predicted HLA population coverage of a vaccine incorporating all 53 peptides using median values of the ANN, SMM, NetMHC 4.0 and NetMHCpan 4.0 algorithms hosted by IEDB [52]. These 53 peptides, taken together, had predicted HLA coverage of greater than 97% of the world's population as shown in Supplementary Material Table 10. We also calculated HLA coverage based on alleles specific to populations in China and found that coverage across those individuals could be expected to be within 3% percent of the world wide coverage estimate as shown in Supplementary

Material Table 11. This same population coverage could be achieved with 16 of the 53 unique peptides as shown in Table 4.

Seven of the 53 peptides with a predicted HLA match have been tested *in-*

270 *vitro* for HLA binding affinity by various researchers [52]. These binding affinity assays were originally performed with the SARS virus during a previous outbreak. Specific literature references for these *in-vitro* assays for each peptide sequence are as follows: ASAFFGMSR, LSPRWYFYY, QQQGQTVTK: [42], FPRGQGVPI: [42][22][36][47], GMSRIGMEV: [22][51][10][32][9], KTFPPTEPK:

275 [42][22][35][47][5] and LLLDRLNQL: [32][10][9][51][62]. These seven peptides are shown in red in Supplementary Material Table 6, Supplementary Material Table 7, Supplementary Material Table 8 and Supplementary Material Table 9.

The remaining 46 COVID-19 peptides listed in could also be further qualified as potential vaccine candidates by confirming MHC binding predictions by

280 *in-vitro* binding affinity and/or binding stability studies [43][41][22]. Another approach to evaluating the 53 COVID-19 candidate vaccine peptides though *in-vitro* testing is also possible.

As we have shown in this paper, a peptide targeted by EBOV controllers could form the basis of a preventative vaccine for EBOV. ELISPOT analysis of

285 PBMCs taken from the peripheral blood of COVID-19 controllers and progressors to assess the presence of a differential response to the 53 peptides could lead to a broadly applicable protective CTL vaccine against COVID-19 by incorporating peptides into the vaccine that are more commonly targeted for CD8+ attack by the controllers *versus* the progressors. The extent of the COVID-19

290 outbreak should allow many more controllers to be identified then the 30 individuals studied by Sakabe. Furthermore, Sakebe did not report progressor data perhaps because of the difficulty in obtaining blood samples from those patients. If researchers act now during the COVID-19 outbreak, perhaps controller and progressor blood samples could be collected and prospectively analyzed, quickly

295 creating a database of optimal candidate class I peptides for inclusion into a CTL vaccine with potentially broad HLA coverage for subsequent rapid manufacture and deployment. It would be interesting to see the extent to which the peptides

favored by controllers appear on COVID-19 nucleocapsid, making COVID-19 a second example, across two different viruses, of controllers exhibiting CTL attack preferentially on the nucleocapsid protein.

4. Acknowlegements

All animal handling was done in accordance with NIH and institutional animal care and use guidelines by Aragen Bio-sciences in Morgan Hill California and the University of Texas, Medical Branch, Galveston Texas working in conjunction with the Galveston National Laboratory. The research was funded by Flow Pharma, Inc.

5. Declaration of Interest Statement

CV Herst, Scott Burkholz, Lu Wang, Peter Lloyd and Reid Rubsamen are employees of Flow Pharma, Inc. all receiving cash and stock compensation. Alessandro Sette, Paul Harris, William Chao and Tom Hodge are members of Flow Pharmas Scientific Advisory Board. Alessandro Sette and Paul Harris have received cash and stock compensation as SAB members. Richard Carback and Serban Ciotlos are consultants to Flow Pharma, both receiving cash and stock compensation. John Sidney works with Alessandro Sette at the La Jolla Institute of Allergy and Immunology. Flow Pharma, Inc. has previously contracted with the La Jolla Institute of Allergy and Immunology to support other research not related to this study funded under STTR contract CBD18-A-002-0016. Reid Rubsamen, CV Herst, Scott Burkholz, Lu Wang, Peter Lloyd, Richard Carback, Serban Ciotlos and Tom Hodge are named inventors on various issued and pending patents relating to Flow Pharma's technology. All of the rights to all of these patents have been assigned by each of the inventors to Flow Pharma. Shane Massey, Trevor Brasel, Edecio Cunha-Neto and Daniela Rosa have nothing to declare.

References

₃₂₅ [1] Massimo Andreatta and Morten Nielsen. Gapped sequence alignment using artificial neural networks: application to the mhc class i system. *Bioinformatics*, 32(4):511–517, 2016. URL https://doi.org/10.1093/bioinformatics/btv639.

[2] Mekibib B and Arilen K. Aerosol transmission of filoviruses. *Viruses*, 9(5):
₃₃₀ 148, May 2016. doi: 10.3390/v8050148. URL https://doi.org/10.3390/v8050148.

[3] Steve E Bellan, Juliet R C Pulliam, Jonathan Dushoff, and Lauren Ancel Meyers. Ebola control: effect of asymptomatic infection and acquired immunity. *The Lancet*, 384(9953):1499–1500, October 2014.
₃₃₅ doi: 10.1016/S0140-6736(14)61839-0. URL https://doi.org/10.1016/S0140-6736(14)61839-0.

[4] R Billeskov, Y Wang, S Solaymani-Mohammadi, B Frey, S Kulkarni, P Andersen, E Marie Agger, Y Sui, and J Berzofsky. Low antigen dose in adjuvant-based vaccination selectively induces cd4 t cells with enhanced
₃₄₀ functional avidity and protective efficacy. *J Immunol*, 198(9):3494–3506, 2017. doi: 10.4049/jimmunol.1600965. URL https://doi.org/10.4049/jimmunol.1600965.

[5] Thomas Blicher, Jette Sandholm Kastrup, Soren Buus, and Michael Gajhede. High-resolution structure of hla-a* 1101 in complex with sars
₃₄₅ nucleocapsid peptide. *Acta Crystallographica Section D: Biological Crystallography*, 61(8):1031–1040, 2005. doi: 10.1107/S0907444905013090.

[6] N. Borthwick, T. Ahmed, B. Ondondo, P. Hayes, A. Rose, U. Ebrahimsa, E. J. Hayton, A. Black, A. Bridgeman, M. Rosario, A.V. Hill, E. Berrie, E. Moyle, N. Frahm, J. Cox, S. Colloca, A. Nicosia, J. Gilmour, A.J.
₃₅₀ McMichael, L. Dorrell, and L. Hanke. Vaccine-elicited human t cells recognizing conserved protein regions inhibit hiv-1. *Mol Ther.*, 22(2):464–475,

2014. doi: 10.1038/mt.2013.248. URL https://doi.org/10.1038/mt.2013.248.

[7] BaslerGaya CF and Amarasinghe K. Evasion of interferon responses by ebola and marburg viruses. *Journal of Interferon & Cytokine Research*, 29 (9), September 2009. doi: 10.1089/jir.2009.0076. URL https://doi.org/10.1089/jir.2009.0076.

[8] Chung-ke Chang, Ming-Hon Hou, Chi-Fon Chang, Chwan-Deng Hsiao, and Tai-huang Huang. The sars coronavirus nucleocapsid protein–forms and functions. *Antiviral research*, 103:39–50, 2014. URL https://doi.org/10.1016/j.antiviral.2013.12.009.

[9] Ying-Kit Cheung, Samuel Chak-Sum Cheng, Fion Wan-Yee Sin, Kin-Tak Chan, and Yong Xie. Induction of t-cell response by a dna vaccine encoding a novel hla-a* 0201 severe acute respiratory syndrome coronavirus epitope. *Vaccine*, 25(32):6070–6077, 2007. URL https://doi.org/10.1016/j.vaccine.2007.05.025.

[10] YK Cheung, SC Cheng, FW Sin, KT Chan, and Y Xie. Investigation of immunogenic t-cell epitopes in sars virus nucleocapsid protein and their role in the prevention and treatment of sars infection. *Hong Kong medical journal= Xianggang yi xue za zhi*, 14:27–30, 2008.

[11] E Cunha-Neto, DS Rosa, PE Harris, T Olson, A Morrow, A Ciotlos, CV Herst, and RM Rubsamen. An approach for a synthetic ctl vaccine design against zika flavivirus using class i and class ii epitopes identified by computer modeling. *Frontiers in Immunology*, 2017. doi: doi.org/10.3389/fimmu.2017.00640. URL https://doi.org/10.3389/fimmu.2017.00640.

[12] Alves DA, Glynn AR, Steele KE, Lackemeyer MG, Garza NL, Buck JG, Mech C, and Reed DS. Aerosol exposure to the angola strain of marburg virus causes lethal viral hemorrhagic fever in cynomolgus macaques. *Veterinary Pathology*, 2010. doi: 10.1177/0300985810378597.

[13] Shishang Dong, Peng Yang, Guobang Li, Baocheng Liu, Wenming Wang, Xiang Liu, Boran Xia, Cheng Yang, Zhiyong Lou, Yu Guo, and Zihe Rao. Insight into the ebola virus nucleocapsid assembly mechanism: crystal structure of ebola virus nucleoprotein core domain at 1.8 å resolution. *Protein & Cell*, 6(5):351–362, May 2015. ISSN 1674-8018. doi: 10.1007/s13238-015-0163-3. URL https://doi.org/10.1007/s13238-015-0163-3.

[14] H. Ebihara, B. Rockx, A. Marzi, F. Feldmann, E. Haddock, D. Brining, R.A. LaCasse, D. Gardner, and H. Feldmann. Host response dynamics following lethal infection of rhesus macaques with zaire ebolavirus. *J. Infect. Dis.*, 204:S991–S999, 2011. doi: DOI: 10.1093/infdis/jir336.

[15] Heinz Feldmann, Steven Jones, Hans-Dieter Klenk, and Hans-Joachim Schnittler. Ebola virus: from discovery to vaccine. *Nature Reviews Immunology*, 3:677–685, August 2003. doi: 10.1038/nri1154. URL https://doi.org/10.1038/nri1154.

[16] Steven A Frank and Robin M Bush. Barriers to antigenic escape by pathogens: trade-off between reproductive rate and antigenic mutability. *BMC evolutionary biology*, 7(1):229, 2007. URL https://doi.org/10.1186/1471-2148-7-229.

[17] Rezza G. A vaccine against ebola: Problems and opportunities. *Human Vaccines and Immunotherapeutics*, 11(5):1258–1260, 2014. doi: 10.1080/21645515.2015.1021528. URL https://doi.org/10.1080/21645515.2015.1021528.

[18] T.W. Geisbert, L.E. Hensley, T. Larsen, H.A. Young, D.S. Reed, J.B. Geisbert, D.P. Scott, E. Kagan, and K.J. Jahrling, P.B.; Davis. Pathogenesis of ebola hemorrhagic fever in cynomolgus macaques: Evidence that dendritic cells are early and sustained targets of infection. *Am. J. Pathol.*, 163: 2347–2370, 2003. doi: DOI: 10.1016/S0002-9440(10)63591-2.

[19] S.C. Gilbert. T-cell-inducing vaccines - what's the future? *Immunology*,

135(1):19–26, 2012. doi: 10.1111/j.1365-2567.2011.03517.x. URL https://doi.org/10.1111/j.1365-2567.2011.03517.x.

[20] Manisha Gupta, Patricia Greer, Siddhartha Mahanty, Wun-Ju Shieh, Sherif R. Zaki, Rafi Ahmed, and Pierre E. Rollin. Cd8-mediated protection against ebola virus infection is perforin dependent. *Journal of Immunology*, 174(7):4198–4202, April 2005. doi: 10.4049/jimmunol.174.7.4198. URL https://doi.org/10.4049/jimmunol.174.7.4198.

[21] Scott G. Hansen, Julia C. Ford, Matthew S. Lewis, Abigail B. Ventura, Colette M. Hughes, Lia Coyne-Johnson, Nathan Whizin, Kelli Oswald, Rebecca Shoemaker, Tonya Swanson, Alfred W. Legasse, Maria J. Chiuchiolo, Christopher L. Parks, Michael K. Axthelm, Jay A. Nelson, Michael A. Jarvis, Michael Piatak, Jeffrey D. Lifson, and Louis J. Picker. Profound early control of highly pathogenic siv by an effector memory t-cell vaccine. *Nature*, 473:523–527, May 2011. doi: 10.1038/nature10003. URL https://doi.org/10.1038/nature10003.

[22] Mikkel Harndahl, Kasper Lamberth, Sune Justesen, Gustav Rder, Michael Madsen, Christina Sylvester-Hvid, Morten Nielsen, Claus Lundegaard, Mette Voldby Larsen, Sheila Tang, Sren Brunak, Ole Lund, and Sren Buus. Large scale analysis of peptide-hla class i interactions. *IEDB*, 2006. URL https://www.iedb.org/reference/1000945.

[23] L.E.. Hensley, H.A. Young, P.B. Jahrling, and T.W. Geisbert. Proinflammatory response during ebola virus infection of primate models: Possible involvement of the tumor necrosis factor receptor superfamily. *Immunol. Lett.*, 80:169–179, 2002. URL https://www.ncbi.nlm.nih.gov/pubmed/11803049.

[24] Steven M Jones, Heinz Feldmann, Ute Ströher, Joan B Geisbert, Lisa Fernando, Allen Grolla, Hans-Dieter Klenk, Nancy J Sullivan, Viktor E Volchkov, Elizabeth A Fritz, Kathleen M Daddario, Lisa E Hensley, Peter B Jahrling, and Thomas W Geisbert. Live attenuated recombinant

vaccine protects nonhuman primates against ebola and marburg viruses. *Nature Medicine*, 11:786–790, June 2005. doi: 10.1038/nm1258. URL https://doi.org/10.1038/nm1258.

[25] Vanessa Jurtz, Sinu Paul, Massimo Andreatta, Paolo Marcatili, Bjoern Peters, and Morten Nielsen. Netmhcpan-4.0: improved peptide–mhc class i interaction predictions integrating eluted ligand and peptide binding affinity data. *The Journal of Immunology*, 199(9):3360–3368, 2017. URL https://doi.org/10.4049/jimmunol.1700893.

[26] Leffel K and Reed D. Marburg and ebola viruses as aerosol threats. *Biosecurity and Bioterrorism: Biodefense Strategy, Practice, and Science*, 2(3), September 2004. doi: 10.1089/bsp.2004.2.186.

[27] I Kalidi, Y Fofana, A A Rahly, V Bochu, C Dehay, J Gony, and J Hors. Study of hla antigens in a population of mali (west africa). *Tissue Antigens*, 31:98–102, 03 1988. URL https://www.ncbi.nlm.nih.gov/pubmed/3163860.

[28] Chih-Yun Lai, Daniel P Strange, Teri Ann S Wong, Axel T Lehrer, and Saguna Verma. Ebola virus glycoprotein induces an innate immune response in vivo via tlr4. *Frontiers in microbiology*, 8:1571, 2017. URL https://doi.org/10.3389/fmicb.2017.01571.

[29] LF Martin-Banderas, M Flores-Mosquera, P Riesco-Chueca, A Rodriguez-Gil, A Cebolla, and S Chávez. Flow focusing: a versatile technology to produce size-controlled and specific-morphology microparticles. *Small*, 4, 2005. doi: 10.1002/smll.200500087.

[30] A.K. McElroy, R.S. Akondy, C.W. Davis, A.H. Ellebedy, A.K. Mehta, C.S. Kraft, G.M. Lyon, B.S. Ribner, J. Varkey, and J. Sidney. Human ebola virus infection results in substantial immune activation. *Proc. Natl. Acad. Sci.*, 1122:4719–4724, 2015. doi: doi: 10.1073/pnas.1502619112.

⁴⁶⁵ [31] Charles K. Nfon, Anders Leung, Greg Smith, Carissa Embury-Hyatt, Gary Kobinger, and Hana M. Weingartl. Immunopathogenesis of severe acute respiratory disease in zaire ebolavirus-infected pigs. *PLoS ONE*, 2013. doi: 10.1371/journal.pone.0061904.

[32] Satoshi Ohno, Shunsuke Kohyama, Maiko Taneichi, Osamu Moriya, Hide-⁴⁷⁰ nori Hayashi, Hiroshi Oda, Masahito Mori, Akiharu Kobayashi, Toshitaka Akatsuka, Tetsuya Uchida, et al. Synthetic peptides coupled to the surface of liposomes effectively induce sars coronavirus-specific cytotoxic t lymphocytes and viral clearance in hla-a* 0201 transgenic mice. *Vaccine*, 27(29): 3912–3920, 2009. URL https://doi.org/10.1016/j.vaccine.2009.04.⁴⁷⁵ 001.

[33] F Pereyra, D Heckerman, J M Carlson, C. Kadie, D Z Soghoian, D Karel, A Goldenthal, O B. Davis, C E DeZiel, and T Lin. Hiv control is mediated in part by cd8+ t-cell targeting of specific epitopes. *J Virol*, 88:12937–12948, 2014. doi: 10.1128/JVI.01004-14. URL https://doi.org/10.1128/JVI.⁴⁸⁰ 01004-14.

[34] Julia Ponomarenko, Kerrie Vaughan, Sinu Paul, Bjoern Peters, Alessandro Sette, Maximilian Haeussler, and Sebastian Maurer-Stroh. Ebola: an analysis of immunity at the molecular level. *IEEE*, July 2015. doi: https://doi:10.1109/aisw.2015.7469230. URL https://doi:10.1109/⁴⁸⁵ aisw.2015.7469230.

[35] Michael Rasmussen, Mikkel Harndahl, Anne Bregnballe Kristensen, Ida Kallehauge Nielsen, Kasper W Jorgensen, Anette Stryhn, Morten Nielsen, and Sren Buus. Large scale analysis of peptide - hla-i stability. *IEDB*, 2014. URL https://www.iedb.org/reference/1028287.

⁴⁹⁰ [36] Michael Rasmussen, Mikkel Harndahl, Anne Bregnballe Kristensen, Ida Kallehauge Nielsen, Kasper W Jorgensen, Anette Stryhn, Morten Nielsen, and Sren Buus. Large scale analysis of peptide - hla-i stability. *IEDB*, 2014. URL https://www.iedb.org/reference/1028291.

[37] R. M. Rubsamen, C. V. Herst, P. M. Lloyd, and D. E. Heckerman. Eliciting cytotoxic t-lymphocyte responses from synthetic vectors containing one or two epitopes in a c57bl/6 mouse model using peptide-containing biodegradable microspheres and adjuvants. *Vaccine*, 32:4111–4116, 2014. doi: 10.1016/j.vaccine.2014.05.071. URL https://doi.org/10.1016/j.vaccine.2014.05.071.

[38] Jain S and Baranwal M. Conserved peptide vaccine candidates containing multiple ebola nucleoprotein epitopes display interactions with diverse hla molecules. *Medical Microbiology and Immunology*, 208(2):227–238, April 2019. doi: 10.1007/s00430-019-00584-y. URL https://doi.org/10.1007/s00430-019-00584-y.

[39] Saori Sakabe, Brian M. Sullivan, Jessica N. Hartnett, Refugio Robles-Sikisaka, Karthik Gangavarapu, Beatrice Cubitt, Brian C. Ware, Dylan Kotliar, Luis M. Branco, Augustine Goba, Mambu Momoh, John Demby Sandi, Lansana Kanneh, Donald S. Grant, Robert F. Garry, Kristian G. Andersen, Juan Carlos de la Torre, Pardis C. Sabeti, John S. Schieffelin, and Michael B. A. Oldstone. Analysis of cd8+ t cell response during the 2013–2016 ebola epidemic in west africa. *Proceedings of the National Academy of Sciences*, 2018. ISSN 0027-8424. doi: 10.1073/pnas.1806200115. URL http://www.pnas.org/content/early/2018/07/17/1806200115.

[40] M.D. Selidji T. Agnandji, M.D. Angela Huttner, M.D. Madeleine E. Zinser, M.Med. Patricia Njuguna, Ph.D. Christine Dahlke, M.D. José F. Fernandes, M.Sc. Sabine Yerly, M.D. Julie-Anne Dayer, Ph.D. Verena Kraehling, Ph.D. Rahel Kasonta, Ph.D. Akim A. Adegnika, M.D., and Ph.D. Marcus Altfeld, M.D. Phase 1 trials of rvsv ebola vaccine in africa and europe. *The New England Journal of MedicineNew England*, 374:1647–1660, April 2016. doi: 10.1056/NEJMoa1502924. URL https://doi.org/10.1056/NEJMoa1502924.

[41] Alessandro Sette, John Sidney, Marie-France del Guercio, Scott Southwood, Jörg Ruppert, Carol Dahlberg, Howard M Grey, and Ralph T Kubo. Peptide binding to the most frequent hla-a class i alleles measured by quantitative molecular binding assays. *Molecular immunology*, 31(11):813–822, 1994. doi: https://doi.org/10.1016/0161-5890(94)90019-1.

[42] John Sidney, Jason Botten, Benjamin Neuman, Michael Buchmeier, and Alessandro Sette. Sars iedb entries. *IEDB*, 2006. URL https://www.iedb.org/reference/1000425.

[43] John Sidney, Scott Southwood, Carrie Moore, Carla Oseroff, Clemencia Pinilla, Howard M Grey, and Alessandro Sette. Measurement of mhc/peptide interactions by gel filtration or monoclonal antibody capture. *Current protocols in immunology*, 100(1):18–3, 2013. URL https://doi.org/10.1002/0471142735.im1803s100.

[44] S. Sridhar. Heterosubtypic t-cell immunity to influenza in humans: Challenges for universal t-cell influenza vaccines. *Front Immunol.*, 7(195), 2016. doi: doi: 10.3389/fimmu.2016.00195. URL https://doi.org/10.3389/fimmu.2016.00195.

[45] The International HIV Controllers Study et al. The major genetic determinants of hiv-1 control affect hla class i peptide presentation. *Science (New York, NY)*, 330(6010):1551, 2010. URL https://doi.org/10.1126/science.1195271.

[46] Nancy J. Sullivan, Anthony Sanchez, Pierre E. Rollin, Zhi yong Yang, and Gary J. Nabel. Development of a preventive vaccine for ebola virus infection in primates. *Nature*, 408:605–609, 2000. doi: 10.1038/35046108. URL https://doi.org/10.1038/35046108.

[47] C Sylvester-Hvid, Morten Nielsen, K Lamberth, G Røder, S Justesen, C Lundegaard, P Worning, H Thomadsen, O Lund, S Brunak, et al. Sars

ctl vaccine candidateshla supertype, genome-wide scanning and biochemical validation. *Scandinavian Journal of Immunology*, 59(6):632–632, 2004. URL https://doi.org/10.1111/j.0001-2815.2004.00221.x.

[48] Yuki Takamatsu, Larissa Kolesnikova, and Stephan Becker. Ebola virus proteins np, vp35, and vp24 are essential and sufficient to mediate nucleocapsid transport. *Proceedings of the National Academy of Sciences*, 115(5): 1075–1080, 2018. URL https://doi.org/10.1073/pnas.1712263115.

[49] Xiaolu Tang, Changcheng Wu, Xiang Li, Yuhe Song, Xinmin Yao, Xinkai Wu, Yuange Duan, Hong Zhang, Yirong Wang, Zhaohui Qian, et al. On the origin and continuing evolution of sars-cov-2. *National Science Review*, page nwaa036, 2020. URL https://doi.org/10.1093/nsr/nwaa036.

[50] Thomas Trolle, Curtis P McMurtrey, John Sidney, Wilfried Bardet, Sean C Osborn, Thomas Kaever, Alessandro Sette, William H Hildebrand, Morten Nielsen, and Bjoern Peters. The length distribution of class i–restricted t cell epitopes is determined by both peptide supply and mhc allele–specific binding preference. *The Journal of Immunology*, 196(4):1480–1487, 2016. URL https://doi.org/10.4049/jimmunol.1501721.

[51] Yeou-Ping Tsao, Jian-Yu Lin, Jia-Tsrong Jan, Chih-Hsiang Leng, Chen-Chung Chu, Yuh-Cheng Yang, and Show-Li Chen. Hla-a 0201 t-cell epitopes in severe acute respiratory syndrome (sars) coronavirus nucleocapsid and spike proteins. *Biochemical and biophysical research communications*, 344(1):63–71, 2006. URL https://doi.org/10.1016/j.bbrc.2006.03.152.

[52] R Vita, JA Overton, JA Greenbaum, J Ponomarenko, JD Clark, JR Cantrell, DK Wheeler, JL Gabbard, D Hix, A Sette, and B Peters. The immune epitope database (iedb) 3.0. *Nucleic Acids Res.*, 2014.

[53] Olivia A Vogel and Balaji Manicassamy. Broadly protective strategies against influenza viruses: Universal vaccines and therapeutics. *Frontiers*

in Microbiology, 11:135, 2020. URL https://doi.org/10.3389/fmicb.2020.00135.

[54] William Wan, Larissa Kolesnikova, Mairi Clarke, Alexander Koehler, Takeshi Noda, Stephan Becker, and John AG Briggs. Structure and assembly of the ebola virus nucleocapsid. *Nature*, 551(7680):394–397, 2017. URL https://doi.org/10.1038/nature24490.

[55] F Wang, T Ono, A Kalergis, W Zhang, T DiLorenzo, and K Lim. On the defing rules for interactions between the t cell receptor and its ligand: a critical role for a specific amino acid residue of the t cell receptor b chain. *Proc Natl Acad Sci USA*, 95(6), 1998. doi: 10.1073/pnas.95.9.5217.

[56] Kelly L. Warfield, John M. Dye, Jay B. Wells, Robert C. Unfer, Frederick W. Holtsberg, Sergey Shulenin, Hong Vu, Dana L. Swenso, Sina Bavari, and M. Javad Aman. Homologous and heterologous protection of nonhuman primates by ebola and sudan virus-like particles. *PLoS ONE*, 10(3):e0118881, March 2015. doi: 10.1371/journal.pone.0118881. URL https://doi.org/10.1371/journal.pone.0118881.

[57] K.L. Warfield, B.C. Bosio, C.M. Welcher, and E.M. Deal. Ebola virus-like particles protect from lethal ebola virus infection. *Proc. Natl. Acad. Sci.*, 100(26):15889–15894, 2003. doi: 10.1073/pnas.2237038100.

[58] P. Wauquier, N.and Becquart, C.; Padilla, S. Baize, and E.M. Leroy. Human fatal zaire ebola virus infection is associated with an aberrant innate immunity and with massive lymphocyte apoptosis. *Negl. Trop. Dis.*, 4 (e837), 2010. doi: doi: 10.1371/journal.pntd.0000837.

[59] JA Wilson and MK Hart. Protection from ebola virus mediated by cytotoxic t lymphocytes specific for the viral nucleoprotein. *J Virol*, 75 (6):2660–2664, 2001. doi: 10.1128/JVI.75.6.2660-2664.2001. URL https://doi.org/10.1128/JVI.75.6.2660-2664.2001.

[60] Guang Wu and Shaomin Yan. Reasoning of spike glycoproteins being more vulnerable to mutations among 158 coronavirus proteins from different species. *Journal of molecular modeling*, 11(1):8–16, 2005. URL https://doi.org/10.1007/s00894-004-0210-0.

[61] L Zhao, M Zhang, and H Cong. Advances in the study of hla-restricted epitope vaccines. *Human Vaccines and Immunotherapeutics*, 9:2566–2577, 2018. doi: 10.4161/hv.26088. URL https://doi.org/10.4161/hv.26088.

[62] Minghai Zhou, Dongping Xu, Xiaojuan Li, Hongtao Li, Ming Shan, Jiaren Tang, Min Wang, Fu-Sheng Wang, Xiaodong Zhu, Hua Tao, et al. Screening and identification of severe acute respiratory syndrome-associated coronavirus-specific ctl epitopes. *The Journal of Immunology*, 177(4):2138–2145, 2006. URL https://doi.org/10.4049/jimmunol.177.4.2138.

9mer Epitope ELISPOT Data

Figure 1: ELISPOT data from mice vaccinated with NP44-52 in 2mg of adjuvanted microspheres administered via intradermal (tail) injection was statistically different from the response after vaccination with NP43-51 and NP45-53.

Figure 2: ELISPOT data from mice vaccinated with Class I peptides NP44-52 compared with NP43-53, each delivered adjuvanted microspheres adminstered via intradermal (tail) injection with with Class II help peptide VG19 in separate microspheres.

(a) Post infection survival curves for 1,000 PFU challenged mice comparing mice vaccinated with microspheres containing the class I epitope sequence and different microspheres containing the class II epitope sequence versus PBS buffer control 14 days before maEBOV challenge. The difference between the number of survivors in the vaccinated group *versus* the PBS control group was statistically significant by chi square (P = 0.0003).

(b) Clinical observations (control). (c) Clinical observations (active).

(d) Daily weights (control). (e) Daily weights (active).

Figure 3: 1000 PFU post-challenge data (active *versus* PBS buffer solution) collected beginning 14 days after vaccination.

(a) Survival curve *versus* PBS buffer control. The difference between the number of survivors in the vaccinated group *versus* the PBS control group was statistically significant by chi square (P = 0.001).

(b) Clinical observations (control).   (c) Clinical observations (active).

(d) Daily weights (control).   (e) Daily weights (active).

Figure 4: 100 PFU post-challenge data (active *versus* PBS buffer solution) collected beginning 14 days after vaccination.

(a) Survival curve *versus* PBS buffer control. The difference between the number of survivors in the vaccinated group *versus* the PBS control group was statistically significant by chi square (P = 0.003).

(b) Clinical observations (control).   (c) Clinical observations (active).

(d) Daily weights (control).   (e) Daily weights (active).

Figure 5: 10,000 PFU post-challenge data (active *versus* PBS buffer solution) collected beginning 14 days after vaccination.

(a) EBOV nucleocapsid proteins NP and VP24 shown.

(b) NP44-52 is a conserved structural loop (red) buried inside the NP structure. The conservation and the location of NP44-52 suggest that residues 44-52 are important for the structural integrity of the EBOV NP.

Figure 6: The class I epitope used for this study is located within NP. Nucleocapsid proteins NP and VP24 are shown together in (a). A detailed view of NP with the study epitope position highlighted in shown in (b).

Cytokine / Survival Correlations for Control Groups

Cytokine	p-Value	Correlation
Mo IL-6	0.050	Decreased with survival
Mo MCP-1	0.019	Decreased with Survival
Mo IL-9	0.015	Increased with survival
Mo MIP-1b	0.009	Decreased with survival
Mo IL-12(p40)	0.006	Increased with Survival
Mo G-CSF	0.005	Decreased with Survival
Mo IL-1b	0.005	Increased with Survival
Mo IFN-g	0.003	Increased with Survival
Mo GM-CSF	0.002	Increased with Survival
Mo IL-12(p70)	0.001	Increased with Survival
Mo TNF-a	0.001	Increased with Survival
Mo IL-17	0.000	Increased with Survival
Mo IL-10	0.000	Decreased with Survival

Table 3: Cytokines with statistically significant (positive or negative) correlation with survival in non-vaccinated mice are shown here along with (Pearson Correlation Analysis) p-values.

COVID-19 Nucleocapsid Top Candidate Peptides with Associated
Predicted HLA Restricted Binding Affinities

Peptide	Start Position	Allele	NetMHC 4.0 pIC$_{50}$nM	NetMHCpan 4.0 pIC$_{50}$nM	SARS Same?
LSPRWYFYY	104	HLA-A*01:01	48.64	76.9	YES
LLLDRLNQL	222	HLA-A*02:01	14.81	11.3	YES
KTFPPTEPK	361	HLA-A*03:01	20.8	18.8	YES
KTFPPTEPK	361	HLA-A*11:01	6.8	7.7	YES
KHIDAYKTF	355	HLA-A*23:01	134.12	778.7	YES
YYRRATRRI	86	HLA-A*24:02	74.89	322	NO
NTASWFTAL	48	HLA-A*26:01	1113.04	122.6	YES
IGYYRRATR	84	HLA-A*33:03	N/A	57.8	YES
FPRGQGVPI	66	HLA-B*07:02	3.82	4.7	YES
SPRWYFYYL	105	HLA-B*08:01	13.77	42.1	YES
KAYNVTQAF	266	HLA-B*15:01	40.35	19	NO
RRIRGGDGK	92	HLA-B*27:05	65.94	72.5	NO
NTASWFTAL	48	HLA-B*39:01	47.87	353.3	YES
MEVTPSGTW	322	HLA-B*44:02	11.48	14.2	YES
LPNNTASWF	45	HLA-B*53:01	19.03	25.7	YES
KAYNVTQAF	266	HLA-B*58:01	12.51	17.7	NO
KAYNVTQAF	266	HLA-C*03:04	N/A	12.7	NO
YRRATRRIR	87	HLA-C*07:01	112.27	8786.2	NO
QRNAPRITF	9	HLA-C*07:02	112.27	237.8	NO
FAPSASAFF	307	HLA-C*08:01	N/A	280.1	YES

Table 4: This set of 16 unique peptides represents the minimum number required to achieve >
95% world-wide population coverage. The starting position is within the Nucleocapsid. Top
binding affinity predictions chosen via NetMHC 4.0 or NetMHCpan 4.0. Peptide sequences
colored in red have literature references as known *in-vitro* binders to the predicted allele match
(see text).