The evolution of mechanisms to divide labour in microorganisms

Guy Alexander Cooper[*1,2], Ming Liu[2], Jorge Peña[3], and Stuart Andrew West[2]

[1]St. John's College, Oxford, OX1 3JP, United Kingdom

[2]Department of Zoology, University of Oxford, Oxford, OX1 3SZ, United Kingdom

[3]Institute for Advanced Study in Toulouse, University of Toulouse Capitole, Toulouse,

31080 Cedex 6, France

<u>Abstract</u>

Division of labour, where cooperating individuals specialise to perform complementary

tasks, plays a fundamental role at all levels of biological complexity, from genes to

complex animal societies. Different labour-dividing species employ different mechanisms

to determine how tasks are allocated between individuals, including coordinated, random,

and genetic determination of phenotype (caste). It is not clear if this diversity is adaptive—

arising because different mechanisms are favoured in different environments—or is

merely the result of non-adaptive, historical artifacts of evolution. We use theoretical

models to analyse the relative advantages of the two dominant mechanisms employed

for reproductive division of labour in microorganisms: coordinated and random

specialisation. We show that fully coordinated specialisation is more likely to evolve over

random specialisation in well-mixed groups when: (i) social groups are small; (ii)

cooperation has a greater relative impact on fitness (i.e., is more "essential"); and (iii)

there is a relatively low metabolic cost to coordination. We find analogous results when

we allow for spatial structure in a more mechanistic model of growing cyanobacteria

[*] Corresponding author: guy.cooper@zoo.ox.ac.uk

filaments. More generally, our results show that the diversity in mechanisms to divide

labour are likely to have arisen through the action of adaptation by natural selection, and

not as an historical artifact of non-adaptive evolution.

Key words: random specialisation, phenotypic noise, adaptive coin-flipping, coordination,

signalling, cellular differentiation, adaptation, evolutionary theory, cyanobacteria, social

microbes

<u>Introduction</u>

Different species use different mechanisms to divide labour between reproductives and

their helpers (Figure 1) [1–3]. In some cases, there is coordination across individuals to

determine which individual will perform which role (*coordinated specialisation*). This

coordination could use signals, cues or a developmental programme to provide

information about the phenotypes adopted by other individuals in the group. This

happens, for example, when honey bee workers feed royal jelly to larvae to produce

reproductive queens (Figure 1A), or when the local density of a signalling molecule

determines whether cyanobacteria cells develop into sterile nitrogen-fixing heterocysts

(Figure 1B)[4–6]. In other cases, each individual adopts a helper phenotype with a certain

probability, independently and without knowledge of the phenotypes adopted by other

individuals (*random specialisation*). For example, in *Salmonella enterica* co-infections,

random biochemical fluctuations within each cell's cytoplasm are used to determine

whether the cell sacrifices itself to trigger an inflammatory response that eliminates

competitor species (Figure 1D)[7,8]. In yet further cases, phenotype is influenced by the

individual's genotype (genetic control). For instance, in some ant societies whether

individuals develop into queens, major, or minor workers can be determined, in part, by

their genes (Figure 1C)[3,9–11]. Across the tree of life some species employ one mechanism

to divide labour whereas in other species mixed forms exist with a combination of

coordinated specialisation, random specialisation, or genetic control[3,10,12–17].

Figure 1. Different mechanisms for division of labour in nature. A) In honey bee hives (Apis mellifera), larvae develop as sterile workers unless they are fed large amounts of royal jelly by adult workers (coordinated specialisation) [4]. (Photo by Wausberg via the Wikimedia Commons.) B) In *A. cylindrica* filaments (cyanobacteria), some individuals develop into sterile nitrogen fixers (larger, round cells) if the amount of nitrogen fixed by their neighbours is insufficient (coordinated specialisation). This leads to a precise allocation of labour, with nitrogen-fixing cells distributed at fixed intervals along the filament. [5] (Picture taken by Robert Calentine.) C) In the army ant (*Eciton Burchelli*), whether individual ants become a major or minor worker has a genetic component (genetic control) [11]. (Photo by Alex Wild via the Wikimedia Commons, cropped.) D) In *S. enterica* infections (serovar Typhymurium), each cell amplifies intra-cellular noise to determine whether it will self-sacrifice and trigger an inflammatory response that

eliminates competing strains (random specialisation). [8] (Photo by Rocky Mountain
Laboratories, NIAID, NIH via Wikimedia Commons.)

We lack general evolutionary explanations for why different species use different

mechanisms to divide labour between reproductives and their helpers[2,3,18,19]. Previous

work has focused on non-reproductive division of labour in the social insects, and the

proximate mechanisms that lead to different worker castes[11,20–25]. However, the focus in

that literature is on a different question – how different proximate mechanisms can

produce coordinated specialisation – rather than the broader question of whether

coordinated specialisation should be favoured over random specialisation or genetic

control in the first place. It is with reproductive division of labour that these three very

different mechanisms have been observed in different species and for which there is an

absence of evolutionary explanations[2,3,18,19,26].

Bacteria and other social microbes offer excellent opportunities for studying why different

labour-dividing mechanisms are favoured in different species[1,2]. Across microbes, the two

primary mechanisms used to produce reproductive division of labour are coordinated and

random specialisation (Figure 2). Furthermore, while the form of cooperation and life

histories of social microbes share many similarities, they also vary in factors that could

influence the evolution of division of labour, such as social group size [27,28]. We develop

theoretical models to examine whether the relative advantage of random versus

coordinated specialisation can depend upon social or environmental conditions. Our aim

here is to use microbes as a 'test system' to address the broader question of whether

evolutionary models can explain the diversity in the mechanisms that produce division of

labour.

Figure 2. Mechanisms to divide labour in social microorganisms. We examine the
relative advantages and disadvantages of the two key mechanisms for reproductive
division of labour in social microorganisms. (A) Random specialisation occurs when cells
randomly specialise into helpers or reproductives independently of one another. This can
occur when a genetic feedback circuit is used to amplify small molecular fluctuations in
the cytoplasm of each cell (phenotypic noise) [7,29–33]. (B) Coordinated specialisation occurs
when cells interact with one another, and share (or gain) phenotypic information while
they are differentiating. This could occur through the secretion and detection of
extracellular molecules (signals or cues), or with a shared developmental programme
(epigenetics)[1,2].

Results and Discussion

We compare the relative fitness advantages of dividing labour with either coordinated or

random specialisation. Our first aim is to capture the problem in a deliberately simple

model, which is easy to interpret, and can be applied across diverse microbe species[34,35].

We begin by assuming that coordinated specialisation always produces the optimal

proportion of helpers and reproductives (fully coordinated specialisation) and that there

is no within-group spatial structure (well-mixed groups). We then test the robustness of

our results by examining a number of alternate models for different biological scenarios

and by developing a more mechanistic model of growing cyanobacteria filaments that

includes the effects of within-group spatial structure and partial coordination. Throughout,

we assume a form of cooperation that is common in microbes, where some individuals

produce a 'public good' that benefits all cells.

<u>Random specialisation vs fully coordinated specialisation</u>

We assume that a single cell arrives on an empty patch and, through a fixed series of

replications, produces a clonal group of n individuals that consists of k sterile helpers and

$n - k$ pure reproductives $(k \in \{0,1,2, \ldots, n\})$. We denote group fecundity, $g_{k,n}$, as the

reproductive success of a particular group in the absence of mechanism costs. This is

measured as the per capita number of offspring that disperse at the end of the group life

cycle, given by

$$g_{k,n} = \frac{1}{n}(n - k)f_{k,n}, \tag{1}$$

where $n - k$ is the number of reproductives in the group, and $f_{k,n}$ is the fecundity of each

reproductive in the absence of mechanism costs. We assume that $f_{k,n}$ increases with the

number of helpers in the whole group (k).

Expression (1) highlights the trade-off between the number of reproductives in the group

$(n - k)$, which is higher when there are fewer helpers (lower k), and the amount of help

that those reproductives obtain $(f_{k,n})$, which is higher when there are more helpers (higher

k). The balance of this trade-off often results in an optimal number of helpers, k^*, that is

intermediate (i.e., $0 < k^* < n$).

In species that divide labour by coordination, the outcome of individual specialisation

depends on the phenotypes of social group neighbours. Our first model is deliberately

agnostic to the details of how phenotype information is shared between group members

in order to facilitate predictions across different systems. For instance, individuals may

share phenotype information via signalling between cells or with a common

developmental program (Figure 2B)[1,2,36]. We make the simplifying assumption that

individuals coordinate fully, so that coordinated groups always form with precisely the

optimal number of helpers, k^*. The disadvantage of coordinated specialisation is that the

mechanism could incur metabolic costs, such as the production of extracellular signaling

molecules. The fitness of a group of coordinated specialisers is given by:

$$w_C = (1 - c_C)g_{k^*,n} \, , \tag{2}$$

where $g_{k^*,n}$ is the group fecundity with the optimal number of helpers (i.e., $g_{k^*,n} = \max_k g_{k,n}$)

and $0 \leq c_c \leq 1$ is the metabolic cost of coordination, whose form we leave unspecified

but could in principle depend on further factors such as group size. A number of different

models have examined how different proximate mechanisms can produce coordinated

division of labour in specific systems [20,23–25].

In species that divide labour by random specialisation, each individual in the group

independently becomes a helper with a given probability and a reproductive otherwise

(Figure 2A). Hence, the final number of helpers in the group is a binomial random variable.

We assume that the probability of adopting a helper role is equal to the optimal proportion

of helpers ($p^* = k^*/n$). Thus, the expected fitness of a group of random specialisers is

given by:

$$w_R = (1 - c_R) \sum_{k=0}^{n} \binom{n}{k} p^{*k} (1 - p^*)^{n-k} g_{k,n},\qquad(3)$$

where $0 \leq c_R \leq 1$ is the metabolic cost of random specialisation, which we assume is

independent of the number of helpers in the group, k. The potential advantage of random

specialisation is that there may be fewer upfront metabolic costs from, for example,

between cell signaling (i.e., if $c_R < c_C$ holds). The downside of random specialisation is

that groups sometimes form with fewer or more helpers than is optimal (demographic

stochasticity). In principle, the probability of becoming a helper could be regulated by

environmental cues so as to produce on average more or fewer helpers when this is more

favourable. However, throughout our analysis we assume a stable environment and

ignore such regulation.

We need to specify how reproductive fecundity depends on the number of helpers in the

group. We focus here on one of the most common forms of cooperation in microbes,

where individuals secrete factors that provide a benefit to the local population of cells

("public goods") [37]. We assume that the amount of public good in the social group

depends linearly on the number of helpers in the group and is "consumed" by all group

members equally [38,39]. An example of such a public good is found in *Bacillus subtilis*

populations, where only a subset of cells (helpers) produce and secrete proteases that

degrade proteins into smaller peptides, but where these are then re-absorbed as a

nutrient source by all cells [40].

We allow the relative importance of producing public goods to vary between species.

Each reproductive has a baseline fecundity, $b \geq 0$, that is independent of the amount of

public good in the group. The fecundity benefit of helpers scales according to $h \geq 0$ as

the amount of public good in the group increases. When reproductives have no baseline

fecundity ($b = 0$), we say that cooperation is essential. When baseline fecundity is non-

zero ($b > 0$), cooperation is non-essential and the ratio h/b provides a useful metric for

the relative importance of cooperation.

Our assumptions give the following expression for the fecundity of a reproductive:

$$f_{k,n} = b + h\frac{k}{n}. \qquad (4)$$

184   By substituting Equation 4 into Equations 1-3, we can determine when the fitness of

185   coordinated specialisation is greater than the fitness of random specialisation (i.e., $w_C >$

186   w_R), which gives the simplified condition:

$$\gamma < \frac{h-b}{n(h+b)}, \qquad (5)$$

where $\gamma = (c_C - c_R)/(1 - c_R)$ captures the relative change in metabolic costs paid when

switching to coordinated specialisation from random specialisation. If $h < b$, then helpers

are not required and the group is composed of all reproductives ($k^* = 0$). Thus, division

of labour is favoured to evolve only when $h > b$, which we will assume henceforth

(supplementary section C). Condition (5) specifies that coordinated specialisation is

favoured when the relative change in metabolic costs of switching from random

specialisation to coordination (γ), is less than the fecundity benefits gained from doing so

(right-hand side). The condition can be used to predict how key environmental and

ecological factors will influence which labour-dividing mechanism is more likely to evolve

(Figure 3).

Figure 3. Random versus coordinated specialisation. Small group sizes (lower n),
relatively more important cooperation (higher h/b), and lower relative metabolic costs to
coordination (lower γ) favour division of labour by coordinated specialisation (black) over division
of labour by random specialisation (white). Here we have used $\gamma = 2 \times 10^{-3}$ (solid boundary) and
$\gamma = 1 \times 10^{-3}$ (dashed boundary). We note that the limit as the relative importance of
cooperation goes to infinity (very large h/b) illustrates the outcome when cooperation is
essential ($b = 0$).

Prediction 1: Smaller relative metabolic costs of coordination favour coordinated

specialisation.

When the metabolic cost of coordination is smaller (lower c_C) and the metabolic cost of

random specialisation is larger (higher c_R), then the relative cost of switching from random

specialisation to coordinated specialisation is lessened (smaller γ), which favours the

evolution of coordinated specialisation (smaller left-hand side of Condition 5). If the

metabolic costs of random specialisation are equal to or larger than the metabolic costs

of coordination ($c_R \geq c_C \Rightarrow \gamma \leq 0$), then coordinated specialisation is always the favoured

mechanism (Condition 5 always satisfied). Conversely, random specialisation can only

ever be the favoured strategy ($w_R > w_C$; Condition 5 not satisfied) if the metabolic costs

of random specialisation are less than the metabolic costs of coordination ($c_C > c_R \Rightarrow \gamma >$

0; a necessary but not sufficient condition). This arises directly from our starting

assumption that coordinated specialisation always produces groups with the optimal

proportion of helpers whereas random specialisation only does so sometimes.

Larger metabolic costs of coordinated specialisation $(c_R < c_C \Rightarrow \gamma > 0)$ may be a

reasonable assumption for many biological systems. The metabolic costs of random

specialisation are determined by the production costs of the regulatory proteins employed

in the genetic feedback circuit that amplifies intra-cellular noise[29,41–43]. In contrast,

coordinated specialisation requires both an intracellular genetic feedback circuit and

some mechanism by which phenotype is communicated between cells, such as the costly

production and secretion of extra-cellular signaling molecules[1,2,36,44–46].

As a result, in many scenarios, the optimal mechanism to divide labour depends on how

the potentially higher metabolic costs of coordination ($c_C > c \Rightarrow \gamma > 0$) balance against

the benefit of avoiding the stochastic costs of random specialisation (right-hand side of

condition 5). The stochastic costs of random specialisation are determined by: (i) the

relative likelihood that random groups deviate from the optimal proportion of helpers, and

(ii) the degree to which those deviations from the optimal proportion of helpers leads to a

reduced fecundity for the group (supplementary section C). Equation (5) shows how the

importance of these two factors depends upon the size of the group (n) and on the relative

importance of cooperation (h/b).

Prediction 2. Smaller social groups favour coordinated specialisation.

The number of cells in the group has a large impact on the relative likelihood that random

groups deviate from the optimal proportion of helpers (Figure 3). In smaller groups, there

are fewer possible outcomes for the proportion of helpers ($n + 1$ possible allocations of

labour for groups of size n). Consequently, random specialisation can more easily lead to

the formation of groups with a realized proportion of helpers that deviates significantly

from the optimum ($p \ll p^*$ or $p \gg p^*$). In contrast, in larger groups there are more possible

outcomes and the resulting proportion of helpers will be more closely clustered about the

optimal composition with highest fitness (with $p \approx p^*$ for very large group sizes).

This effect of group size on the stochastic cost of random specialisation is a consequence

of the law of large numbers. For example, outcomes close to 50% heads are much more

likely when tossing 100 coins in a row compared to only tossing 4 coins in a row where

no heads or all heads may frequently occur. Our prediction is related to how, when mating

occurs in small groups, small brood sizes select for more precise and less female biased

sex ratios as there would otherwise be a high probability of producing a group containing

no males at all[47–49]. In another analogue, Wahl showed a mechanistically different effect

when division of labour is determined genetically and the number of group founders is

small: groups may sometimes form that do not contain all of the genotypes required to

produce all of the necessary phenotypes in the division of labour [19].

Prediction 3: The higher the relative importance of cooperation, the more coordinated

specialisation is favoured.

When the relative importance of cooperation is larger (higher h/b), the fitness costs

incurred from producing too few helpers increases. In addition, as the relative importance

of cooperation increases (higher h/b), the optimal proportion of helpers increases to 50%

helpers ($p^* \approx \frac{1}{2}$). This increases the variance in the proportion of helpers produced by

random specialisers, and so sub-optimal groups may arise more frequently

(supplementary section C). Thus, a higher relative importance of cooperation increases

both (i) the likelihood that groups deviate from the optimal proportion of helpers and (ii)

the scale of the fitness cost when they do. Both of these effects increase the stochastic

costs of random specialisation (larger right-hand side of Condition 5), and thus favour the

evolution of coordinated specialisation (Figure 3).

<u>Alternative forms of cooperation</u>

The above analysis employs a deliberately simple public goods model, focusing on

factors that are expected to be relevant across many microbial systems. This facilitates

the interpretation of our results and generates broadly applicable predictions that are less

reliant on the specifics of any particular biological species.

In order to test the robustness of our results (predictions 1-3) we also developed a series

of alternative simplified models corresponding to different biological scenarios

(supplementary sections D and E; Figures S1-S3). We examined the possibility that the

public good provided by helpers is: (i) not consumed by its beneficiaries, as may occur

when self-sacrificing *S. enterica* cells enter the gut to trigger an immune response that

eliminates competitors (non-rivalrous or non-congestible collective good); or (ii) only

consumed by the reproductives in the group, as may occur for the fixed nitrogen secreted

by heterocyst cells in *A. cylindrica* filaments (excludible or club good) [7,46,50,51]. We allowed

for reproductive fecundity to depend non-linearly on the proportion of helpers in the group,

for helpers to have some fecundity (non-sterile helpers), and for division of labour to occur

in each generation of group growth. In all of these alternative biological scenarios, we

found broad qualitative agreement across all three predictions of the linear public goods

model.

We found that less specialised helpers (with some fecundity) favours random

specialisation over coordinated specialisation. In contrast to prediction 3, more fecund

helpers can lead to a scenario where a larger relative importance of cooperation (higher

h/b) disfavours coordinated specialisation. This occurs because a high relative

importance of cooperation (higher h/b) can produce groups composed predominantly of

non-sterile helpers ($p^* \approx 1$), where the likelihood that random groups deviate from the

optimal proportion of helpers is significantly diminished (supplementary section E.4).

In supplementary section F we develop an individual based simulation, which also

supports predictions 1-3. In addition, this simulation shows that costly coordination can

evolve incrementally from random specialisation, and that intermediate levels of

coordination can be favoured (Figure S4).

<u>Division of labour in a cyanobacteria filament</u>

We then developed a more mechanistically detailed model of a growing cyanobacteria

filament to investigate the impact of within-group spatial structure (supplementary section

G). When there is insufficient fixed nitrogen (N_2) in the environment, some cyanobacteria

species will facultatively divide labour between reproductive cells (autotrophs) that

photosynthesise light and sterile helper cells (heterocysts) that fix and secrete

environmental N_2 (Figure 1B)[5,52,53]. The fixed N_2 diffuses along the filament where it is

used by reproductives to grow and produce new cells. Division of labour in cyanobacteria

is a canonical example of coordinated specialisation as helpers produce a variety of

signalling molecules that diffuse along the filament to ensure that a regular pattern of

phenotypes develops (Figure 1B)[5,52,53]. Previous models of cyanobacteria focused on

determining the signalling and regulatory network required to recreate the exact pattern

of heterocysts along the filament[52,54–58].

Cyanobacteria spores (hormogonium) tend to contain multiple cells [5,52]. We assumed that

each filament begins as a clonal sequence of two reproductives (R) and two helpers (H)

in the arrangement (H-R-R-H). Over time, the number of cells in the filament increases

as reproductives grow and divide to produce within-filament offspring cells, which become

either helpers or reproductives (Figure 4A). The group life cycle ends when the filament

has reached a maximum size of L cells. At this time, the reproductives in the filament

produce dispersing hormogonia that found filaments in the next generation of the group

life cycle and all remaining cells die (non-overlapping generations) [5].

Reproductives absorb fixed N_2 in order to replicate. Each reproductive receives fixed N_2

from the abiotic environment at a rate of $\phi \geq 0$ units of fixed N_2 per unit time (uniform

background density of fixed N_2)[58]. In addition, each helper in the filament produces fixed

N_2, at a maximum rate of $\bar{\phi} > 0$ units of fixed N_2 per unit time. We assume that the fixed

N_2 produced by a helper diffuses across the filament, decaying by a factor of $0 < \eta \leq 1$

with each cell position that is further away (cooperation range).

A) Filament growth
B) Signal-response norm
C) Random Specialisation
D) Coordinated Specialisation

Figure 4: Division of labour in a cyanobacteria filament. (A) Black cells represent helpers, and grey cells reproductives. When a reproductive replicates, the parent cell produces an offspring cell (white cell) to one side of itself along the filament. The red shading shows the density of the signal molecule produced by the helpers as it diffuses along the filament. (B) When an offspring cell is sensitive to the signal ($v > 0$), a greater (lesser) signal density will decrease (increase) the probability that it becomes a helper ($q = 0.5, d = 5, v = 0, 0.2, 1$). (C) A simulated example of a filament growing that employs random specialisation ($q = 0.33, s = 0, d = 0$, and $v = 0$). (D) A simulated example of a filament growing that employs coordinated specialisation ($q = 0.33, s = 0.1, d=1$ and $v = 1.5$) (Supplementary information G).

Whether a new cell becomes a helper or a reproductive depends on four evolutionary

traits that jointly determine the extent of division of labour and coordination in the filament

$(q, s, d,$ and v; Figure 4B). The baseline probability $(0 \leq q \leq 1)$ is the underlying probability

that a cell becomes a helper in the absence of coordination. The level of signalling ($0 \leq$

$s \leq 1$) is the fraction of resources that a helper commits to the production and secretion

of signalling molecules. The signalling molecules produced by a helper diffuse along the

filament, decreasing in density by a fixed factor for each cell position that is further away,

and which we assume is distinct from the N_2 diffusivity (Figure 4A). The local density of

signalling molecules allows new cells to estimate how close they are to a helper, or how

many helpers there may be nearby.

Empirically, the reason that new cyanobacteria filaments do not simply use the local

density of fixed N_2 as a cue may be that signalling molecules are cheaper and faster to

produce and thus coordination can occur even before helpers begin to fix N_2 [57].

Furthermore, using a dedicated signal could be more reliable than one based on fixed N_2

density alone, which might be biased by transient fluctuations in the background level of

fixed N_2 (ϕ).

Whether and how the new cell responds to the signal depends on the response sensitivity

($v \geq 0$) and the response threshold ($d \geq 0$; Figure 4B). If $v = 0$, then a new cell is

insensitive to the signal and adopts the helper phenotype with the baseline probability q

(random specialisation). If the daughter cell is sensitive to the signal ($v > 0$) then a local

signal density that is greater than the response threshold, d, will lead to that cell being

less likely to adopt the helper phenotype (Figure 4B). A higher signal density than the

threshold leads to the opposite effect. As sensitivity increases (higher v), the response to

the signal becomes more deterministic (Figure 4B).

Increasing levels of coordination (higher v and s), allows for a more precise patterning of

helpers and reproductives in the filament (compare Figures 4C and 4D). However, we

assume that increased coordination is costly. First, as helpers produce more signalling

molecule (higher s), they can produce proportionally less fixed N_2. Second, when cells

are more sensitive to the signal (higher v), we assume an increasing delay in the time it

takes for a replication event to occur.

<u>Simulations</u>

We simulated an evolving population to estimate the strategy that is favoured by natural

selection in different scenarios (q^*, s^*, d^*, v^*) (supplementary section G). We started with

a uniform population that specialises randomly ($s = d = v = 0$), and allowed the helper

probability (q) to mutate and evolve for 500 generations, until an approximate equilibrium

was reached. We then held the baseline helper probability (q) fixed and allowed the

coordination traits (s, d and v) to mutate and evolve for 3500 generations. Each

generation, the mutant strategy successfully replaces the resident strategy if it has a

higher estimated average fitness. We calculate the fitness of individual filaments as the

summed fecundity of reproductives in the last generation of the group life cycle, divided

by the amount of time that it took the filament to grow to L cells. An equilibrium was

generally reached within a few hundred generations.

We found that the degree to which specialising cells evolve to coordinate can depend on

social and environmental factors. In particular, both a smaller background density of fixed

N_2 (low ϕ) and a smaller cooperation range (low η) lead to the evolution of higher signaling

levels (higher s^*; Figure 5A) and higher response sensitivities (higher v^*; Figure 5B). This

produced filaments with more precise allocation of labour across the filament (Figure 5C),

where we calculate the variance in the number of helpers in a contiguous sub-block of 10

cells relative to the variance that would be expected for a binomial random variable of the

same mean (supplementary section G.4.). These results also illustrate that intermediate

coordination can be favoured in certain scenarios.

Figure 5: The optimal level of coordination. We present simulation results for two key
factors that affect the optimal level of coordination (supplementary section G.4.) A lower
background density of fixed N_2 (lower ϕ) and a smaller cooperation range (lower η)
favours both: (A) the evolution of a higher level of signalling (larger s^*); and (B) a higher
response sensitivity to the signal (larger v^*). (C) The combined effect of higher levels of
signalling (larger s^*) and higher response sensitivity (larger v^*) is that groups form with
an estimated lower variance in the allocation of help across the filament, where the
estimated variance is calculated relative to a binomial distribution with the same mean
(supplementary section G).

The predictions of our cyanobacteria model agree broadly with those of our simpler

analytical model. When the benefits of cooperation are shared more locally (low η) this

leads to a smaller social group (analogous to lower *n*). With random specialisation, a

smaller social group can lead to proportions of helpers that show greater deviations from

the optimum, increasing the benefit that can be obtained by coordination (Figure 3). When

the background density of fixed N_2 is small (low ϕ), this increases the relative benefit of

cooperation (analogous to higher *h/b*). With an increased benefit from cooperation there

is a greater advantage from coordinating to produce the optimum proportion of helpers

(Figure 3).

However, care is required when examining factors in mechanistic models that can have

additional effects unaccounted for by their analogues in simpler models. For instance, an

increase in the background density of fixed N_2 (higher ϕ) means that cooperation is

relatively less important (lower h/b), which we have found favours less coordination

(Figure 5). However, relatively less important cooperation (lower h/b) in the mechanistic

model also means that helpers may be willing to dedicate more effort to signal production

(higher s) as there is then a relatively lower fitness cost to producing less of the public

good. Another example is how helpers that produce more fixed N_2 (larger $\bar{\phi}$) leads to

cooperation that is relatively more important (higher h/b) but can also lead to larger

effective social groups sizes (larger n) as the increased good that helpers produce can

then diffuse further along the filament and benefit reproductives that are farther away.

Our simulations show that coordination ($s^* > 0, v^* > 0$) is often favoured over random

specialisation ($s^* = 0, v^* = 0$). In groups with fixed spatial structure and local cooperation

(lower γ), an effective division of labour requires a regular distribution of helpers across

the group. We hypothesize that random specialisation is particularly disadvantageous

when groups grow over time because it is more likely to lead to contiguous groups of

helpers that expand as the whole group grows (compare Figures 4C and 4D). The helpers

within these clumps can neither reproduce to break up the clump, nor are they close

enough to reproductives to provide fixed N_2. Consequently, even a little bit of coordination

($s^* > 0, v^* > 0$) can lead to a substantial fitness advantage by decreasing the chance that

such "dead zones" can form and grow.

<u>Distribution of labour dividing mechanisms in the natural world</u>

Our analyses provide a theoretical framework to help explain why different species of

microorganisms use different mechanisms to divide labour [2]. While testing our

predictions with a formal comparative analysis would require data from more species,

our predictions can help to understand the mechanisms that have evolved in well

studied examples.

There are many reasons why coordinated specialisation was favoured to evolve in

cyanobacteria filaments. First, cyanobacteria only divide labour when fixed N_2 is growth-

limiting and so the relative importance of cooperation is high (low ϕ and high h/b)[5,53,58].

Second, the fixed nitrogen produced by helpers diffuses across the filament,

preferentially aiding nearby reproductives and so the effective social group size is small

(low η and small n)[45,59,60]. Third, the initial costs of coordination may have been quite

small as new cells could use the local level of fixed N_2 as a cue (low γ) [61]. Finally,

cyanobacteria filaments have a rigid linear structure that can grow to very large sizes

(higher L) and thus random specialisation could have lead to the accumulation of large

sterile clumps.

Colonies of *Volvox carteri* and *Dictyostelium discoideum* use coordination to divide

labour, despite the fact that these groups are composed of large numbers of cells (high

n; on the order of 1000s of cells or more) [15,62–64]. However, the initial specialisation of

helpers in each case appears to make use of a shared developmental programme and

so the relative costs of coordination could be quite small for both species (low γ) [62,63,65].

Further, in both cases an absence of helpers would lead to cell death and thus

cooperation is important (higher h/b) [1,2,63,66]. These conflicting effects highlights that no

single factor can fully explain empirical patterns. Furthermore, some details of the

mechanism of division of labour are still not well understood. For instance, it is possible

that there is also an initially random component to pre-stalk specialisation in *D.*

discoideum [63].

There are multiple reasons why random specialisation would have been favoured to

evolve in other well-studied species. In *Salmonella enterica*, the self-sacrificing helper

cells provide a competitive advantage that eliminates other microbes but is not

"essential" to the replication of *Salmonella* cells (lower h/b)[7,8]. Further, the benefits of

cooperation are provided to all cells in the co-infection ($\eta = 1$) and so the effective

social group size is reasonably large (higher n). Finally, division of labour occurs in one

generation of the group life cycle so there is no scope for the accumulation of growing

helper clumps as for cyanobacteria filaments. In *B. subtilis*, a subset of cells become

helpers that produce and secrete protein degrading proteases[40]. However, these helper

cells are not sterile and so the consequence of deviating from the optimal caste ratios is

much smaller (supplementary section E.4).

Conclusion

Most previous work on reproductive division of labour has tended to be either

mechanistic, focusing on how different phenotypes are produced (caste determination),

or evolutionary, focusing on why division of labour is favoured in the first place [2–4,18–

24,26,45,63,67–72]. We have used evolutionary models to explain the broader question of why

different mechanisms are used in different species [2,3,7,18–20]. We have shown that

coordinated specialisation is more likely to be favoured over random specialisation in

small groups, when relative coordination costs are low, and when there are larger fitness

costs to deviating from optimal caste ratios. We have also shown how these patterns can

hold in groups with spatial structure and where different degrees of coordination can be

favoured. These results identify social and environmental factors that could explain the

distribution of mechanisms for division of labour that have been observed in bacteria and

other microbes. Beyond microorganisms, our results also suggest a hypothesis for why

random caste determination has not been widely observed in animal societies: during the

initial evolution of complex animal societies, group sizes were likely to be small and

relative costs of coordination might have been minor compared to each individual's day-

to-day organismal metabolic expenditure.

Acknowledgements

We thank Anna Dewar, Kevin Foster, Andy Gardner, Ashleigh Griffin, Asher Leeks,
Matishalin Patel, Tom Scott, and Daniel Unterwegger for their helpful comments and
suggestions, and St. John's College, Oxford (GAC), the French National Research
Agency (ANR) under the Investments for the Future (Investissements d'Avenir) program
(grant ANR-17-EURE-0010 to the IAST; JP) and the ERC (Horizon 2020 Advanced
Grant; SAW) for funding. The authors would like to acknowledge the use of the
University of Oxford Advanced Research Computing (ARC;
http://dx.doi.org/10.5281/zenodo.22558) facility in carrying out this work.
520

Data and Code Availability

All simulated data was generated using C and Matlab. The codes and generated data

used for this study are available at:

https://github.com/mingpapilio/Codes_DOL_Mechanisms.

Author Contributions

G.A.C., S.A.W., and J.P. conceived the study. G.A.C. and J.P. and designed and

analysed the analytical models, G.A.C. and M.L. designed and analysed the simulation

models. G.A.C. and S.A.W. wrote the first draft. All authors contributed toward writing the

final manuscript.

Declaration of Interests

The authors declare no competing interests.

Supplementary information:

A. Overview

B. Labour dividers and their fitness

C. Linear public goods

D. Alternative modelling assumptions

E. Alternative forms of cooperation

F. The optimal level of coordination

G. Dividing labour in a cyanobacteria filament

Figure S1. Alternative modelling assumptions

Figure S2. Non-linear public goods

Figure S3. Alternative biological scenarios

Figure S4. The optimal level of coordination

Figure S5. Simulation results for other traits

Table S1: Cyanobacteria model

References:

1. Ackermann, M. A functional perspective on phenotypic heterogeneity in microorganisms. *Nat. Rev. Microbiol.* **13**, 497–508 (2015).
2. West, S. A. & Cooper, G. A. Division of labour in microorganisms: An evolutionary perspective. *Nat. Rev. Microbiol.* **14**, 716–723 (2016).
3. Schwander, T., Lo, N., Beekman, M., Oldroyd, B. P. & Keller, L. Nature versus nurture in social insect caste differentiation. *Trends Ecol. Evol.* **25**, 275–282 (2010).
4. Linksvayer, T. A. *et al.* Larval and nurse worker control of developmental plasticity and the evolution of honey bee queen-worker dimorphism. *J. Evol. Biol.* **24**, 1939–1948 (2011).
5. Flores, E. & Herrero, A. Compartmentalized function through cell differentiation in filamentous cyanobacteria. *Nat. Rev. Microbiol.* **8**, 39–50 (2010).
6. Weaver, N. Rearing of Honeybee Larvae on Royal Jelly in the Laboratory. *Science (80-.).* **121**, 509–510 (1955).
7. Ackermann, M. *et al.* Self-destructive cooperation mediated by phenotypic noise. *Nature* **454**, 987–990 (2008).
8. Diard, M. *et al.* Stabilization of cooperative virulence by the expression of an avirulent phenotype. *Nature* **494**, 353–356 (2013).
9. Hughes, W. O. H., Sumner, S., Van Borm, S. & Boomsma, J. J. Worker caste polymorphism has a genetic basis in Acromyrmex leaf-cutting ants. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 9394–9397 (2003).
10. Anderson, K. E., Linksvayer, T. a & Smith, C. R. The causes and consequences of genetic caste determination in ants (Hymenoptera : Formicidae). *Myrmecol. News* **11**, 119–132 (2008).
11. Jaffe, R., Kronauer, D. J. C., Kraus, F. B., Boomsma, J. J. & Moritz, R. F. A. Worker caste determination in the army ant Eciton burchellii. *Biol. Lett.* **3**, 513–516 (2007).
12. Zhang, Z. *et al.* Antibiotic production in Streptomyces is organized by a division of labor through terminal genomic differentiation. *Sci. Adv.* **6**, eaay5781 (2020).
13. Shelton, D. E., Desnitskiy, A. & Michod, R. E. Distributions of reproductive and somatic cell numbers in diverse Volvox (Chlorophyta) species. *Evol. Ecol. Res.* **14**, 707–727 (2012).
14. Griesemer, J. The Units of Evolutionary Transition. *Selection* **1**, 67–80 (2001).
15. Smith, G. M. *A comparative study of the species of Volvox. Transactions of the American Microscopical Society* vol. 63 (1944).
16. Dragoš, A. *et al.* Division of Labor during Biofilm Matrix Production. *Curr. Biol.* (2018) doi:10.1016/j.cub.2018.04.046.
17. Dragoš, A. *et al.* Collapse of genetic division of labour and evolution of autonomy in pellicle biofilms. *Nat. Microbiol.* (2018) doi:10.1038/s41564-018-0263-y.
18. Wahl, L. M. Evolving the division of labour: Generalists, specialists and task allocation. *J. Theor. Biol.* **219**, 371–388 (2002).
19. Wahl, L. M. The Division of Labor: Genotypic versus Phenotypic Specialization. *Am. Nat.* **160**, 135–145 (2002).
20. Duarte, A., Weissing, F. J., Pen, I. & Keller, L. An Evolutionary Perspective on Self-Organized Division of Labor in Social Insects. *Annu. Rev. Ecol. Evol. Syst.*

42, 91–110 (2011).

21.   Oster, G. F. & Wilson, E. O. *Caste and ecology in the social insects. Caste and*
ecology in the social insects (Princeton University Press, 1978).

22.   Wilson, E. O. The ergonomics of caste in the social insects. *Am. Econ. Rev.* **102**,
41–66 (1978).

23.   Duarte, A., Pen, I., Keller, L. & Weissing, F. J. Evolution of self-organized division
of labor in a response threshold model. *Behav. Ecol. Sociobiol.* **66**, 947–957
(2012).

24.   Duarte, A., Scholtens, E. & Weissing, F. J. Implications of behavioral architecture
for the evolution of self-organized division of labor. *PLoS Comput. Biol.* **8**,
e1002430 (2012).

25.   Waibel, M., Floreano, D., Magnenat, S. & Keller, L. Division of labour and colony
efficiency in social insects: Effects of interactions between genetic architecture,
colony kin structure and rate of perturbations. *Proc. R. Soc. B Biol. Sci.* **273**,
1815–1823 (2006).

26.   Bourke, A. F. G. *Principles of Social Evolution. Principles of Social Evolution*
(Oxford University Press, 2011).
doi:10.1093/acprof:oso/9780199231157.001.0001.

27.   Powers, S. T. & Lehmann, L. When is bigger better? The effects of group size on
the evolution of helping behaviours. *Biol. Rev.* (2017) doi:10.1111/brv.12260.

28.   Peña, J. & Nöldeke, G. Group size effects in social evolution. *J. Theor. Biol.*
(2018) doi:10.1016/j.jtbi.2018.08.004.

29.   Smits, W. K., Kuipers, O. P. & Veening, J. W. Phenotypic variation in bacteria:
The role of feedback regulation. *Nat. Rev. Microbiol.* **4**, 259–271 (2006).

30.   Süel, G. M., Garcia-Ojalvo, J., Liberman, L. M. & Elowitz, M. B. An excitable gene
regulatory circuit induces transient cellular differentiation. *Nature* **440**, 545–550
(2006).

31.   Dubnau, D. & Losick, R. Bistability in bacteria. *Mol. Microbiol.* **61**, 564–572
(2006).

32.   Balázsi, G., Van Oudenaarden, A. & Collins, J. J. Cellular decision making and
biological noise: From microbes to mammals. *Cell* **144**, 910–925 (2011).

33.   Delong, E. F. *et al.* Noise in Gene Expression Determines Cell Fate in Bacillus
subtilis. *Science (80-.).* **27**, 526–530 (2007).

34.   Parker, G. A. & Maynard Smith, J. Optimality theory in evolutionary biology.
Nature **348**, 27–33 (1990).

35.   May, R. M. Uses and Abuses of Mathematics in Biology. *Science (80-.).* **303**,
790–793 (2004).

36.   Gestel, J. V. A. N., Vlamakis, H. & Kolter, R. Division of Labor in Bio films : the
Ecology of Cell Differentiation. *Microb. Biofilms* **3**, 1–24 (2015).

37.   West, S. A., Diggle, S. P., Buckling, A., Gardner, A. & Griffin, A. S. The Social
Lives of Microbes. *Annu. Rev. Ecol. Evol. Syst.* **38**, 53–77 (2007).

38.   Frank, S. A. A general model of the public goods dilemma. *J. Evol. Biol.* **23**,
1245–1250 (2010).

39.   West, S. A. & Buckling, A. Cooperation, virulence and siderophore production in
bacterial parasites. *Proc. R. Soc. B Biol. Sci.* **270**, 37–44 (2003).

40.   Veening, J.-W. *et al.* Transient heterogeneity in extracellular protease production

644 by Bacillus subtilis. *Mol. Syst. Biol.* **4**, 184 (2008).

645 41. Veening, J.-W., Smits, W. K. & Kuipers, O. P. Bistability, Epigenetics, and Bet-
646  Hedging in Bacteria. *Annu. Rev. Microbiol.* **62**, 193–210 (2008).

647 42. Davidson, C. J. & Surette, M. G. Individuality in Bacteria. *Annu. Rev. Genet.* **42**,
648  253–268 (2008).

649 43. McAdams, H. H. & Arkin, A. Stochastic mechanisms in gene expression. *Proc.*
650  *Natl. Acad. Sci.* **94**, 814–819 (1997).

651 44. Keller, L. & Surette, M. G. Communication in bacteria : an ecological and
652  evolutionary perspective. *Nat. Rev. Microbiol.* **4**, 249–258 (2006).

653 45. Rossetti, V., Schirrmeister, B. E., Bernasconi, M. V. & Bagheri, H. C. The
654  evolutionary path to terminal differentiation and division of labor in cyanobacteria.
655  *J. Theor. Biol.* **262**, 23–24 (2010).

656 46. Flores, E. & Herrero, A. Compartmentalized function through cell differentiation in
657  filamentous cyanobacteria. *Nat. Rev. Microbiol.* **8**, 39–50 (2010).

658 47. West, S. A. & Herre, E. A. Stabilizing Selection and Variance in Fig Wasp Sex
659  Ratios. *Evolution (N. Y).* **52**, 475–485 (1998).

660 48. Nagelkerke, C. J. & Sabelis, M. W. Precise control of sex allocation in pseudo-
661  arrhenotokous phytoseiid mites. *J. Evol. Biol.* **11**, 649–684 (1998).

662 49. Green, R. F., Gordh, G. & Hawkins, B. A. Precise Sex Ratios in Highly Inbred
663  Parasitic. *Am. Nat.* **120**, 653–665 (1982).

664 50. Dionisio, F. & Gordo, I. The Tragedy of the Commons, the Public Good Dilemma
665  and the Meaning of Rivalry and Excludability in Evolutionary Biology. *Evol. Ecol.*
666  *Res.* **8**, 321–332 (2006).

667 51. Peña, J., Nöldeke, G. & Lehmann, L. Evolutionary dynamics of collective action in
668  spatially structured populations. *J. Theor. Biol.* (2015)
669  doi:10.1016/j.jtbi.2015.06.039.

670 52. Meeks, J. C. & Elhai, J. Regulation of Cellular Differentiation in Filamentous
671  Cyanobacteria in Free-Living and Plant-Associated Symbiotic Growth States.
672  *Microbiol. Mol. Biol. Rev.* **66**, 94–121 (2002).

673 53. Herrero, A., Stavans, J. & Flores, E. The multicellular nature of filamentous
674  heterocyst-forming cyanobacteria. *FEMS Microbiol. Rev.* **40**, 831–854 (2016).

675 54. Wolk, C. P. Formation of one-dimensional patterns by stochastic processes and
676  by filamentous blue-green algae. *Dev. Biol.* **46**, 370–382 (1975).

677 55. Gerdtzen, Z. P. *et al.* Modeling heterocyst pattern formation in cyanobacteria.
678  *BMC Bioinformatics* **10**, 1–13 (2009).

679 56. Torres-Sánchez, A., Gómez-Gardeñes, J. & Falo, F. An Integrative Approach for
680  Modeling and Simulation of Heterocyst Pattern Formation in Cyanobacteria
681  Filaments. *PLoS Comput. Biol.* **11**, 1–18 (2015).

682 57. Muñoz-García, J. & Ares, S. Formation and maintenance of nitrogen-fixing cell
683  patterns in filamentous cyanobacteria. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 6218–
684  6223 (2016).

685 58. Brown, A. I. & Rutenberg, A. D. Heterocyst placement strategies to maximize the
686  growth of cyanobacterial filaments. *Phys. Biol.* **9**, (2012).

687 59. Flores, E. & Herrero, A. Compartmentalized function through cell differentiation in
688  filamentous cyanobacteria. *Nature Reviews Microbiology* (2010)
689  doi:10.1038/nrmicro2242.

60. Co, A. D., van Vliet, S., Kiviet, D. J., Schlegel, S. & Ackermann, M. Short-range
interactions govern the dynamics and functions of microbial communities. *Nat.*
Ecol. Evol. **4**, 366–375 (2020).

61. Brown, J., Sanderson, J. & Michod, E. Evolution of Social Behavior by
Reciprocation. 319–339 (1982).

62. Kirk, D. L. Germ-soma differentiation in Volvox. *Dev. Biol.* **238**, 213–223 (2001).

63. Strmecki, L., Greene, D. M. & Pears, C. J. Developmental decisions in
Dictyostelium discoideum. *Dev. Biol.* **284**, 25–36 (2005).

64. Maree, A. F. M. & Hogeweg, P. How amoeboids self-organize into a fruiting body:
Multicellular coordination in Dictyostelium discoideum. *Proc. Natl. Acad. Sci.* **98**,
3879–3883 (2002).

65. Kirk, D. L. Asymmetric division, cell size and germ-soma specification in Volvox.
Semin. Dev. Biol. (1995) doi:10.1016/S1044-5781(06)80078-2.

66. Koufopanou, V. The Evolution of Soma in the Volvocales. *Am. Nat.* **143**, 907–931
(1994).

67. Kamakura, M. Royalactin induces queen differentiation in honeybees. *Nature* **473**,
478 (2011).

68. Gavrilets, S. Rapid transition towards the division of labor via evolution of
developmental plasticity. *PLoS Comput. Biol.* **6**, e1000805 (2010).

69. Ispolatov, I., Ackermann, M. & Doebeli, M. Division of labour and the evolution of
multicellularity. *Proc. R. Soc. B Biol. Sci.* **279**, 1768–1776 (2012).

70. Cooper, G. A. & West, S. A. Division of labour and the evolution of extreme
specialization. *Nat. Ecol. Evol.* **2**, 1161–1167 (2018).

71. Linksvayer, T. A. Ant species differences determined by epistasis between brood
and worker genomes. *PLoS One* **2**, e994 (2007).

72. Walsh, J. T., Warner, M. R., Kase, A., Cushing, B. J. & Linksvayer, T. A. Ant
nurse workers exhibit behavioural and transcriptomic signatures of specialization
on larval stage. *Anim. Behav.* **141**, 161–169 (2018).