**Optogenetic control of gut bacterial metabolism to promote longevity**

Lucas A. Hartsough[1], Matthew V. Kotlajich[1], Bing Han[3,7], Chih-Chun J. Lin[3,4], Lauren Gambill[2],

Meng C. Wang*[3,4,5], Jeffrey J. Tabor*[1,2,6]

[1]Department of Bioengineering, [2]Systems, Synthetic, and Physical Biology Program, Rice

University, Houston, TX; [3]Huffington Center on Aging, [4]Department of Molecular & Human

Genetics, Baylor College of Medicine, Houston, TX; [5]Howard Hughes Medical Institute;

[6]Department of Biosciences, Rice University; [7]Current Address: Children's Hospital & Institutes

of Biomedical Sciences, Fudan University, Shanghai, China, 201102

*corresponding authors

**Abstract**

Gut microbial metabolism is associated with host longevity. However, because it requires

direct manipulation of microbial metabolism *in situ*, establishing a causal link between these two

processes remains challenging. We demonstrate an optogenetic method to control gene

expression and metabolite production from bacteria residing in the host gut. We genetically

engineer an *Escherichia coli* strain that synthesizes and secretes colanic acid (CA) under the

quantitative control of light. Using this optogenetically-controlled strain to induce CA production

directly in the *Caenorhabditis elegans* gut, we reveal the local effect of CA in protecting

intestinal mitochondria from stress-induced hyper-fragmentation. We also exploit different

intensities of light to determine that the lifespan-extending effect of CA is positively correlated

with its levels produced from bacteria. Our results show that optogenetic control offers a rapid,

reversible and quantitative way to fine-tune gut bacterial metabolism and uncover its local and

systemic effects on host health and aging.

## Introduction

**Introduction**

Microbiome studies have identified correlations between bacteria and host aging (Kundu

et al., 2017; O'Toole and Jeffery, 2015). For example, 16S rRNA and metagenomic DNA

sequencing are used to associate the presence or abundance of specific bacteria to human

centenarians (Biagi et al., 2016; Claesson et al., 2012, 2010). However, given the complexity and

heterogeneity of the human gut environment, these approaches are unable to elucidate how a

specific microbial species contributes to longevity. The nematode *Caenorhabditis elegans* has a

short and easily-measured lifespan, features that have revolutionized our understanding of the

molecular genetics of aging and longevity (Kenyon, 2010). Studies using *C. elegans* also provide

mechanistic insight into the association between bacterial species and host longevity (Gusarov et

al., 2013; Kim, 2013). Importantly, recent studies have revealed that bacterial metabolism can

produce specific products to directly influence the aging process in the host *C. elegans* or

modulate the effects of environmental cues on *C. elegans* lifespan (Cabreiro et al., 2013; Pryor et

al., 2019; Virk et al., 2016). These findings highlight the significance of bacterial metabolism in

regulating host physiology during the aging process and have inspired interest in directly

manipulating bacterial metabolism *in situ* in the host gastrointestinal (GI) tract.

In several recent studies, researchers have administered antibiotic- or carbohydrate-based

small molecule inducers to modulate gene expression from gut bacteria (Kotula et al., 2014; Lim

et al., 2017; Mimee et al., 2015). While this approach has enabled the *in situ* analysis of a gut

bacterial-host interaction (Lim et al., 2017), chemical effectors may have unwanted side-effects

on host or microbial physiology and subject to slow and poorly-controlled transport and

degradation processes that ultimately limit their precision.

Optogenetics combines light and genetically-engineered photoreceptors to achieve

unrivaled control of biological processes (Olson and Tabor, 2014). Previously, we and others

have engineered bacterial photoreceptors that activate or repress gene expression in response to

specific wavelengths of light (Levskaya et al., 2005; Li et al., 2020; Ohlendorf et al., 2012; Ong

and Tabor, 2018; Ong et al., 2017; Ramakrishnan and Tabor, 2016; Ryu and Gomelsky, 2014;

Schmidl et al., 2014). These photoreceptors have been used to achieve precise quantitative

(Olson et al., 2017, 2014), temporal (Chait et al., 2017; Milias-Argeitis et al., 2016; Olson et al.,

2017, 2014), and spatial (Chait et al., 2017; Levskaya et al., 2005; Ohlendorf et al., 2012; Tabor

et al., 2010) control of bacterial gene expression in culture conditions. They have also been used

to characterize and control transcriptional regulatory circuits (Chait et al., 2017; Olson et al.,

2014; Tabor et al., 2009) and bacterial metabolic pathways (Fernandez-Rodriguez et al., 2017;

Tandar et al., 2019) *in vitro*. Here, we hypothesized that optogenetic control of bacterial gene

expression might provide a new way to manipulate bacterial metabolism *in vivo* in the host GI

tract, with high temporal and spatial precision and no unwanted side-effects.

We address this possibility using the *E. coli-C. elegans* interaction model, a testable

system with known mechanistic links between bacterial metabolism and host longevity and with

complete optical transparency. In particular, CA is an exopolysaccharide synthesized and

secreted from *E. coli*, which can extend the lifespan of the host *C. elegans* through modulating

mitochondrial dynamics (Han et al., 2017). We thus have genetically engineered an *E. coli* strain

to put its biosynthesis of CA under a switchable control between green and red lights, and then

utilized green light to induce CA production from this strain in the gut of the host *C. elegans*. We

discovered that light-induced CA from bacteria residing in the host is sufficient to modulate

mitochondrial dynamics and lifespan, which gives even more potent effects than dietary

supplementation of CA. Furthermore, this optogenetic manipulation allowed us to investigate the

local effect of CA on intestinal cells in a time-controlled manner and its systemic effect on

organisms in a quantitative way, which are not possible with purified CA or CA-overproducing

genetic mutants. This work paves the road for future application of bacterial optogenetics in

understanding bacteria-host interaction with temporal, spatial and quantitative controls and

minimal chemical interference.

**Results**

To demonstrate optogenetic control over gut bacterial gene expression, we first

engineered *E. coli* strain LH01, wherein our previous light-switchable two-component system

CcaSR (Schmidl et al., 2014) controls expression of superfolder green fluorescent protein (*sfgfp*),

and *mcherry* is expressed constitutively to facilitate identification of the bacteria (**Fig. 1a**,

**Supplementary Fig. 1, Supplementary Tables 1-3**). In LH01, green light exposure switches

CcaS to an active state in the presence of chromophore phycocyanobilin (PCB), wherein it

phosphorylates the response regulator CcaR. Phosphorylated CcaR then activates transcription of

*sfgfp* from the P$_{cpcG2\text{-}172}$ output promoter. Red light reverts active CcaS to the inactive form, de-

activating *sfgfp* expression (**Fig. 1a**).

We then reared two groups of *C. elegans* from the larval to the adult stage on plates of

LH01 under red or green light, respectively (**Fig. 1b**). Next, we washed away external bacteria,

applied the paralyzing agent levamisole to prevent expulsion of gut contents, and transferred the

worms to agar pads. Finally, we switched the light color from red to green, or green to red, and

used epi-fluorescence microscopy to image the resulting changes in fluorescence in the gut

lumen over time (**Fig. 1c**). In the red-to-green (step ON) experiment, we observed that sfGFP

fluorescence in the worm gut lumen starts low, begins to increase within 2 hours, and reaches a

saturated high level at 6 hours (**Fig. 1d**). In contrast, in the green-to-red (step OFF) experiment,

sfGFP fluorescence begins high, and decreases exponentially between hours 1-7 (**Fig. 1d**). This

light response is abolished when the PCB biosynthetic operon is removed (ΔPCB) (**Fig. 1d**),

demonstrating that sfGFP levels are specifically controlled by CcaSR.

Next, we used flow cytometry to achieve high-throughput single-cell resolution analysis

of this *in situ* bacterial light response. Specifically, we reared worms in red and green light as

before, but then washed, paralyzed, and placed them into microtubes prior to light switching

(**Fig. 1b**). At several time points over the course of 8 hours, we homogenized the animals,

harvested the gut contents, and sorted bacterial cells and measured fluorescence via cytometry.

This experiment revealed that bacteria in the host gut remain intact (**Supplementary Fig. 2**) and

respond to light in a unimodal fashion (**Fig. 1e-g**). Furthermore, the temporal dynamics of the

gene expression response and dependence on PCB recapitulate our microscopy results

(**Supplementary Fig. 3**). We also confirmed that residual bacteria on the exterior of worms do

not contribute to the flow cytometry measurements (**Supplementary Fig. 4**). Together, these

experiments demonstrate that we can use optogenetics to rapidly, reversibly induce gene

expression of *E. coli* residing in the *C. elegans* gut.

Next, we sought to utilize our optogenetic method to modulate the production of specific

metabolites in bacteria residing in the gut of live hosts. It is well-known that bacterial genes

involved in the same metabolic process are often clustered into operons and co-regulated at the

transcriptional level. We took advantage of this coordinated mode of regulation and chose the

*cps* operon and its transcription activator RcsA for testing optogenetic control of bacterial

metabolism. The *cps* operon in *E. coli* consists of 19 genes that encode enzymes required for the

biosynthesis and secretion of CA (Torres-Cabassa and Gottesman, 1987), and CA-overproducing

bacterial mutants Δ*lon* and Δ*hns* promote longevity in the host *C. elegans* (Han et al., 2017). To

place CA biosynthesis under optogenetic control, we engineered an *E. coli* strain (MVK29)

lacking genomic *rcsA* and expressing a heterologous copy of *rcsA* under the control of CcaSR

(**Fig. 2a, Supplementary Tables 1-3**). We first examined whether MVK29 could respond to

green light and induce CA production and secretion. To this end, we grew the strain in batch

culture under red or green light and quantified supernatant CA levels. In red light, MVK29

secretes CA to concentrations below the limit of detection of the assay, similar to the Δ*rcsA*

mutant (**Fig. 2b**). Green light, on the other hand, induces MVK29 to secrete high levels of CA,

and removal of the PCB biosynthetic operon abolishes this response (**Fig. 2b**). Moreover,

mutation of the CcaS catalytic histidine to a non-functional alanine (H534A), or the CcaR

phosphorylation site from an aspartic acid to a non-functional asparagine (D51N) abolishes

detectable CA production (**Fig. 2b**). Importantly, the level of secreted CA increases sigmoidally

with green light intensity, similar to the response of CcaSR itself (**Fig. 2c**) (Schmidl et al., 2014).

We conclude that we have placed CA production under the control of the CcaSR system, and

that we can use light to tune the production of bacterial metabolites.

We then used this approach to study a gut bacterial metabolite-host interaction pathway

*in vivo*. We first reared worms expressing mitochondrially-localized GFP (mito-GFP)

(**Supplementary Table 3**) on MVK29 with red light. We then continued this red-light exposure

for one group, and switched a second to green, for an additional 6 hours, and immediately

imaged intestinal cell mitochondrial morphology using confocal microscopy (**Fig. 3b**). We found

that mitochondrial fragmentation increases in worms exposed to the bacteria with light-induced

CA secretion (**Fig. 3c**). This result recapitulates the phenotype observed in worms supplemented

with purified CA (Han et al., 2017).

Next, we induced CA production directly from bacteria residing within the host gut. To

this end, we paralyzed the worms carrying mito-GFP, split them into two groups, and treated one

with red and the second with green for 6 hours (**Fig. 3a**). We then analyzed mitochondrial

morphology in intestinal cells using confocal microscopy. We found that the 6-hour levamisole

treatment does not kill worms but leads to mitochondrial hyper-fragmentation (**Fig. 3d**), which

might be due to the inhibitory effect of levamisole on mitochondrial NADH-oxidizing enzymes

(Köhler and Bachmann, 1978). This stress-induced effect resembles mitochondrial decay related

to aging and age-related neurodegenerative diseases (Cho et al., 2009; Exner et al., 2007;

Sebastián et al., 2017). Interestingly, we found that green light exposure counteracts this hyper-

fragmentation in paralyzed worms bearing MVK29 in the gut (**Fig. 3d**). Importantly, we found

no such effects in worms bearing the ΔPCB, CcaS(H534A), CcaR(D51N) or *ΔrcsA* mutant

strains (**Fig. 3d**), suggesting that this protective effective is a result of light-induced CA

overproduction in the gut. These results not only show that optogenetics can be utilized to induce

CA secretion from gut-borne *E. coli in vivo*, but also reveal a local protective effect of CA on

intestinal cells.

Finally, we took the advantage of the quantitative control afforded by optogenetics to

investigate how the lifespan-extending effect of CA relates to CA levels. Beginning at the day-1

adult stage, we exposed worms bearing MVK29 to red, or two green light intensities resulting in

intermediate or high CA secretion, and measured their lifespans. We found that the lifespan

extension increases proportionally with green-light intensity (**Fig. 4a**), revealing the pro-

longevity effect of CA is dose dependent. We also noticed that the extent of lifespan extension

by light is much stronger than that caused by dietary supplementation of purified CA (Han et al.,

2017), suggesting the high efficacy of optogenetic induction to modulate host physiology. As a

control, we repeated the experiment with worms bearing the CA-overproducing Δ*lon* mutant and

showed that the lifespan extension caused by Δ*lon* is independent of light exposure (**Fig. 4b**).

The extent of lifespan extension is similar to the MVK29 intermediate green light condition, but

less than the MVK29 high green light condition (**Fig. 4a, b**), suggesting that MVK29 is capable

of producing higher levels of CA than Δ*lon*. In addition, the lifespan of worms bearing the Δ*rcsA*

mutant is also light-independent and is comparable to that of MVK29 worms under red light

(**Fig. 4a, c**). These results suggest that optogenetic control is sufficient to induce bacterial

production of pro-longevity compounds and improve host health, and can exert stronger

beneficial effects than administration of a bacterial mutant or supplementing purified

compounds. Importantly, unlike the traditional approach of introducing a bacterial mutant,

optogenetic control of bacterial metabolism can modulate a host-level phenotype in a

quantitative manner.

**Discussion**

Our method has broad applications for studying microbe-host interactions *in situ*. For

example, we have identified about two dozen additional *E. coli* genes that are unrelated to CA

biosynthesis and that enhance worm longevity when knocked out (Han et al., 2017), though the

mechanisms by which they act remain largely unclear. By using light to induce their expression

in the gut and measuring acute host responses such as changes in mitochondrial dynamics, the

role of these genes in gut microbe-host interactions could be further explored. In another

example, the quorum-sensing peptide CSF and nitric oxide, both of which are produced by

*Bacillus subtilis* during biofilm formation, have been found to extend worm lifespan through

downregulation of the insulin-like signaling pathway (Donato et al., 2017). We have recently

ported CcaSR into *B. subtilis* and demonstrated that it enables rapid and precise control of gene

expression dynamics (Castillo-Hair et al., 2019). The method we report here should enable *in*

*situ* studies of how gene expression and metabolite production from this important Gram-positive

model bacterium impact longevity as well.

Multiple photoreceptors could also be combined to study more complex microbe-host

interaction pathways. Specifically, we and others have co-expressed CcaSR with independently-

controllable blue/dark and red/far-red reversible light sensors in order to achieve simultaneous

and independent control of the expression of up to three genes in the same bacterial cell

(Fernandez-Rodriguez et al., 2017; Olson et al., 2017; Tabor et al., 2010). Such optogenetic

multiplexing could be performed *in situ* and used to study potential synergistic, antagonistic, or

other higher-order effects of multiple bacterial genes or pathways. A large number of eukaryotic

photoreceptors have also been developed, enabling optical control of many cell- and

neurobiological processes (Deisseroth, 2015; Gautier et al., 2014; Goglia and Toettcher, 2018;

Leopold et al., 2018). Bacterial and eukaryotic photoreceptors could be combined to enable

simultaneous optical manipulation of bacterial and host pathways in order to interrogate whether

or how they interact. Optogenetics could also be used to manipulate bacterial and/or host

pathways at specific locations within the gut to examine location- or tissue-dependent

phenomena.

Finally, our method could be extended to other bacteria or hosts. In particular, it should

be possible to port CcaSR or other bacterial photoreceptors into native *C. elegans* symbionts

(Zhang et al., 2017) or pathogens(Couillault and Ewbank, 2002). Because these strains stably

colonize the host, the use of these bacteria could eliminate the need for paralysis, and facilitate

longer-term experiments. It is likely that light can also be used to control gut bacterial gene

expression in other model hosts such as flies, zebrafish, or mammals. Red-shifted wavelengths

and corresponding optogenetic tools(Ong et al., 2017; Ryu and Gomelsky, 2014) may prove

superior for less optically transparent or larger animals. Overall, by enabling precision control of

bacterial gene expression and metabolism *in situ*, we believe that optogenetics will greatly

improve our understanding of a wide range of microbe-host interactions.

**Acknowledgements**

The LED array used to illuminate *C. elegans* plates was designed by Brian Landry &

Sebastián Castillo-Hair. The mounting hardware for the microscope LED array was designed by

Ravi Sheth. We thank Ravi Sheth for discussions during early stages of the project. We thank Dr.

Joel Moake for the use of his cytometer. This work was supported by the John S. Dunn

Foundation (J.J.T. and M.C.W.) and US National Institutes of Health, 1R21NS099870-01

(J.J.T.), DP1DK113644 (M.C.W.), R01AT009050 (M.C.W.). LAH was supported by a NASA

Office of the Chief Technologist Space Technology Research Fellowship (NSTRF

NNX11AN39H).

**Author Contributions**

JJT and MW conceived of the study. LAH and MVK designed experiments. MVK and LAH

constructed plasmids and strains. LAH, MVK, BH, CJL, LG, and MW performed experiments.

CJL and LG scored single-blinded mitochondrial confocal micrographs. LAH, MVK, and MW

analyzed and interpreted results. LAH, MW, and JJT wrote the manuscript.

## Declaration of Interests

The authors declare no competing interests.

**Figures and legends**

**Figure 1. Optogenetic control of *C. elegans* gut bacterial gene expression.** (a) Strain LH01.

(b) Microscopy and cytometry workflows. (c) Fluorescence microscopy images 0 and 8 h after

green light exposure in the step ON experiment. Scale bar: 10 μm. (d) Response dynamics in the

step ON (green) and step OFF (red) microscopy experiments. Black: ΔPCB strain (step ON

experiment). Individual- (light lines) and multi-worm average (dark lines) data are shown. $n = 7$,

4, 6 worms for green, red, black data sets (measured over 2, 3, 1 days, respectively). Error bars:

SEM. (e-g) Flow cytometry histograms for response dynamics experiments. MEFL: molecules of

equivalent fluorescein.

**Figure 2. Optogenetic control of colanic acid biosynthesis.** (a) Strain MVK29. (b) CA

secretion levels for MVK29 and control strains exposed to red and green light. JW1935-1 is the

*E. coli rcsA* background strain used in this study. N.D.: below assay limit of detection. (c)

Green light intensity versus CA secretion level for MVK29. Data points represent 3 biological

replicates collected on a single day. Dashed line: limit of detection. Error bars indicate standard

deviation of the three biological replicates.

**Figure 3. Light-regulated CA secretion modulates *C. elegans* mitochondrial dynamics.** (a) Schematic of experiment for activating CA biosynthesis *in situ*. (b) Representative images of the mitochondrial network of anterior intestinal cells immediately distal to the pharynx are scored as fragmented, intermediate, or tubular, as previously(Han et al., 2017). Scale bars: 10 μm. (c) Mitochondrial fragmentation profiles of un-paralyzed worms fed MVK29 while exposed to red or green light for 6 h. (d) Fragmentation profiles for worms fed the indicated strain, then paralyzed for 6 h while exposed to red or green light. The number of worms included in each condition is indicated below each bar. The Chi-Squared Test of Homogeneity was used to calculate *p*-values between conditions.

267

268 **Figure 4. Optogenetically-regulated CA biosynthesis extends worm lifespan.** (a) When

269 exposed to green light, worms grown on MVK29 live longer than those exposed to red light, and

270 the magnitude of lifespan extension is proportional to green light intensity (*p<0.0001* green vs.

271 red, log-rank test). (b-c) The lifespans of worms grown on the Δ*lon* (b) or the Δ*rcsA* (c) controls

272 are not affected by light condition (*p>0.1* green vs. red, log-rank test).

## Methods

*E. coli* plasmids, strains, and media

Plasmids used in this study are described in **Supplementary Table 1**. Genbank accession

numbers are given in **Supplementary Table 2**. All plasmids constructed in this study were

assembled via Golden Gate cloning(Engler et al., 2009). Primers were ordered from IDT

(Coralville, IA). Assembled plasmids were transformed into *E. coli* NEB10β (New England

Biolabs) for amplification and screening. All plasmid sequences were confirmed by Sanger

sequencing (Genewiz; S. Plainfield, NJ). To construct pLH401 and pLH405, pSR58.6(Schmidl

et al., 2014) was modified by inserting an *mCherry* expression cassette composed of a

constitutive    promoter    (J23114;    http://parts.igem.org/Promoters/Catalog/Anderson),    RBS

(BBa_B0034; http://parts.igem.org/Part:BBa_B0034), *mCherry*, and a synthetic transcriptional

terminator (L3S1P52 (Chen et al., 2013)). To construct pLH405, pLH401 was further modified

by exchanging the superfolder GFP gene (*sfgfp*) for *gfpmut3\**. pMVK201.2 was built by

modifying pSR58.6 to control expression of *rcsA*.

All *E. coli* strains are described in **Supplementary Table 3.** *ΔrcsA* (JW1935-1) was

obtained from the Coli Genetic Stock Center. *Δlon* (JW0429-1) was obtained from the Keio *E.*

*coli* knockout library(Baba et al., 2006), a gift from the Herman lab. All *E. coli* strains were

maintained in LB media supplemented with appropriate antibiotics (chloramphenicol 34 μg/mL,

spectinomycin 100 μg/mL, kanamycin 100 μg/mL) in a shaking incubator at 37 °C and 250 rpm

unless otherwise noted.

*C. elegans* strains and media

All *C. elegans* strains (**Supplementary Table 3**) were provided by the Caenorhabditis Genetics Center (University of Minnesota), which is funded by the NIH office of Research Infrastructure Programs (P40 OD010440). Worms were grown at 20°C on 1.7% NGM-agar plates in 60 mm Petri dishes inoculated with a lawn of *E. coli* (CGSC str. BW28357), as described in the CGC WormBook (wormbook.org), unless otherwise specified. The common strain *E. coli* OP50 was not used for worm feeding, as it produces CA during normal growth(Han et al., 2017). M9 buffer for *C. elegans* (abbreviated M9Ce to distinguish from *E. coli* M9 media) was composed of 3 g $KH_2PO_4$, 6 g $Na_2HPO_4$, 5 g NaCl, 1 mL 1 M $MgSO_4$, $H_2O$ to 1 L, and sterilized by autoclaving (wormbook.org).

Optogenetic control of CA production

3 mL starter cultures of appropriate *E. coli* strains were inoculated from a -80°C freezer and grown 12 h at 37°C. These starters were diluted to $OD_{600} = 1\times10^{-2}$ in M9 minimal media (1x M9 salts, 0.4% w/v glucose, 0.2% w/v casamino acids, 2 mM $MgSO_4$, 100 µM $CaCl_2$) supplemented with appropriate antibiotics. The M9/cell mixtures were then distributed into 3 mL aliquots in 15 mL clear polystyrene culture tubes and grown at 37°C in a shaking incubator at 250 rpm while illuminated with the appropriate light wavelength and intensity, using the Light Tube Array (LTA) device (Gerhardt et al., 2016). After 22 h, cultures were removed and iced to halt growth and the $OD_{600}$ was measured. Culture samples were collected for CA quantification.

## CA quantification

We adapted a previous CA quantification protocol(DISCHE, 1947; DISCHE and

SHETTLES, 1948) that takes advantage of the fact that it is the only exopolysaccharide

produced in our *E. coli* strains that incorporates fucose. In particular, we quantified the amount

of fucose in cell-derived exopolysaccharides (EPS), and used that value as a proxy CA levels.

First, EPS was liberated from cells by boiling 2 mL of culture for 15 min. in a 15 mL conical

tube. The sample was then centrifuged in 1.5 mL Eppendorf tubes for 15 min. at 21,000 x g.

Then, 0.7 mL of supernatant was dialyzed against water for at least 12 h using Pur-A-Lyzer Midi

3500 dialysis mini-tubes (Sigma-Aldrich, PURD35100-1KT) to remove monomeric fucose from

the sample.

Fucose monomers were then liberated from the EPS polymers by hydrolyzing 0.2 mL of

dialyzed media with 0.9 mL of $H_2SO_4$ solution (6:1 v/v acid:water). This mixture was boiled in a

15 mL conical for 20 min and then cooled to room temperature. The absorbance at 396 nm and

427 nm was measured. Next, 25 µL of 1 M L-cysteine HCl was added and mixed thoroughly by

pipetting. The absorbance at 396 nm and 427 nm was measured again. Simultaneously,

absorbance measurements of L-fucose standards pre- and post-L-cysteine addition were also

recorded. Absorbance change, given by *D* in the formula below, were used to compare the L-

fucose standard samples to the dialyzed culture samples and estimate the L-fucose concentration

in the dialyzed product.

$$D = \left( \left( A_{post}^{396} - A_{pre}^{396} \right) - \left( A_{post}^{427} - A_{pre}^{427} \right) \right)$$

<u>Preparation of NGM-agar plates for worm feeding</u>

3 mL *E. coli* starter cultures were inoculated from -80 °C freezer stocks and grown for 12

h at 37 °C. These starters were then diluted to $OD_{600} = 1$ x $10^{-6}$ in M9 minimal media

supplemented with appropriate antibiotics. The M9/cell mixture was then distributed into 3mL

aliquots in 15 mL clear polystyrene culture tubes and grown at 37 °C in a shaking incubator at

250 rpm while illuminated with the appropriate light in the LTA. Once cultures reached $OD_{600} =$

0.1-0.4, tubes were iced for 10 min and subsequently concentrated to $OD_{600} \sim$ 20 by

centrifugation (4 °C, 4000 rpm, 10 min) and resuspension in fresh M9 media. 400-600 μL of

dense bacterial culture was then applied to sterile NGM-agar plates and allowed to dry in a dark

room, or a room with green overhead safety lights if cultures were preconditioned in green light.

Plates were wrapped in foil and refrigerated at 4 °C for no more than 5 days until needed.

<u>Time-lapse microscopy</u>

To obtain age-synchronized worm cultures, axenized *C. elegans* (strain *glo-1*) eggs were

isolated and allowed to arrest in L1 by starvation in M9 buffer (distinct from M9 media: 3 g

$KH_2PO_4$, 6 g $Na_2HPO_4$, 5 g NaCl, 1 mL 1M $MgSO_4$, and water to 1 L, sterilized by autoclaving

at 121 °C for 20 min) at room temperature for 12-18 h. 10-100 larvae were transferred to a

previously prepared NGM-agar plate containing a lawn of the appropriate bacterial strain. The

plate was then placed in a 20 °C incubator and illuminated with appropriate optogenetic light

provided by a single LED positioned 1cm above the Petri dish. Adult worms were transferred to

a fresh plate as necessary to maintain only a single generation.

Individual worms aged 1-3 days were removed from the dish and prepared for time-lapse

epifluorescence imaging. A 1.5% agar pad was prepared using M9 buffer as previously

described[45], and punched into ½" circles with a hollow punch. A 4 μL droplet of 2 mM

levamisole was deposited on a pad and 5 adult worms were transferred from the NGM plate to

the droplet. An additional 4 μL of levamisole solution was added and the worms were gently

washed to remove external bacteria. Worms were then transferred to a fresh pad with a 4 μL

droplet of levamisole solution, which was allowed to dry, thereby co-localizing and aligning the

worms on the pad. The pad was then inverted and placed into a 13 mm disposable microscopy

petri dish with a #1.5 coverslip on the bottom (Cell E&G; Houston, TX). Another coverslip was

placed on the top of the pad in the dish to curtail evaporation.

The dish was then placed on the stage of a Nikon Eclipse Ti-E inverted epifluorescence

microscope (Nikon Instruments, Inc; Melville, NY). Complete paralysis was induced by

incubating the dish at room temperature (~23 °C) for 30 min. Meanwhile, worms were exposed

to appropriate preconditioning light supplied by a circular array of 8 LEDs (4 x 660 nm, 4 x 525

nm) mounted to the microscope condenser ring, about 2 cm above the Petri dish. Light was then

switched from the preconditioning to the experimental wavelength, and worms were imaged

periodically using 10x, 40x, and 60x objectives. For each time point, the LEDs were turned off

and images acquired in the brightfield (DIC) and fluorescent channels. Afterwards, the LEDs

were turned on again to maintain optogenetic control.

Epifluorescence image analysis

All epifluorescence images were analyzed using the Nikon Elements software package

(Nikon Instruments, Inc; Melville, NY). The mCherry signal was used as a marker for the gut

lumen, and only cells in this region were included in the analysis. Image ROIs were created by

thresholding the sfGFP signal to identify the boundaries of cell populations. Out of focus regions

were eliminated from analysis. The average sfGFP pixel intensity inside the ROIs was calculated

and recorded for each time point.

<u>Flow cytometry</u>

1-3 day old *glo-1* worms were prepared for flow cytometry of the microbiome

constituents by washing, using a protocol adapted from previous work(Portal-Celhay et al.,

2012). Groups of 5 worms were washed 2x in a 5 µL droplet of lytic solution: *C. elegans* M9

buffer containing 2 mM levamisole, 1% Triton X-100, and 100 mg/mL ampicillin. The worms

were then washed 2x in 5 µL droplets of M9 buffer containing 2 mM levamisole only. Finally,

the worms were transferred to clear 0.5 mL Eppendorf tubes containing 50 µL of M9 buffer + 2

mM levamisole, ensuring that 5 worms were deposited in the liquid contained in each tube. Each

tube was then exposed to light by placing it within one well of a 24-well plate (AWLS-303008,

ArcticWhite LLC)  atop a Light Plate Apparatus (LPA) containing green and red LEDs[47] for 8 h

at room temperature. In separate control experiments, we demonstrated that any stray bacteria

that may escape the worms over this period, or which were inadvertently added to the 50 µL of

M9 buffer, are incapable of responding to optogenetic light (**Supplementary Fig. 4**).

At the conclusion of the experiment, tubes were removed from the plate and immediately

chilled in an ice slurry for 10 min in the dark. Worms were homogenized using an anodized steel

probe sterilized between samples via 70% ethanol treatment and flame before being cooled.

Next, we used our previous antibiotic-based fluorescent protein maturation

protocol(Olson et al., 2014) to allow unfolded proteins to mature while preventing the production

of new protein. In particular, 250 µL PBS containing 500 mg/mL Rifampicin was added to the

50 µL homogenized worm samples and transferred to cytometry tubes. These tubes were

incubated in a 37 °C water bath for precisely 1 h, then transferred back to an ice slurry.

These samples were measured on a BD FACScan flow cytometer. For gating, an

FSC/SSC polygon gate was first created using non-fluorescent bacteria grown *in vitro* at 37 °C

(**Supplementary Fig. 2**). Events outside this region were excluded as non-bacterial material. To

isolate the engineered gut bacteria, only events with a high mCherry signal (FL3 > 1200 a.u.,

FL3 gain: 999) were included (**Supplementary Fig. 2**). Samples were measured until 20,000

events were recorded or the sample was exhausted.

<u>Flow cytometry data analysis</u>

All flow cytometry data (FCS format) were analyzed using FlowCal(Castillo-Hair et al.,

2016) and Python 2.7. We wrote a standard cytometry analysis workflow that truncated the

initial and final 10 events to prevent cross-sample contamination, removed events from saturated

detector bins at the ends of the detection range, and added 2D density gate on SSC/FSC retaining

the densest 75% of events (**Supplementary Fig. 2a**). GFPmut3* fluorescence units were

converted into standardized units of molecules of equivalent fluorescein (MEFL) using a

fluorescent bead standard (Rainbow calibration standard; cat. no. RCP-30-20A, Spherotech, Inc.)

as described previously(Castillo-Hair et al., 2016). Finally, to eliminate events associated with *C.*

*elegans* autofluorescence (**Supplementary Fig. 2b**), any events in the region FL1 $\leq$ 1200 MEFL

were discarded.

<u>Mitochondrial Fragmentation Assays</u>

Synchronized L1 worms (strain *ges-1*) were applied to NGM agar plates containing

bacterial strain BW25113 and allowed to develop until adulthood. This parental bacterial strain is

used to allow all worms to develop at the same rate, avoiding any developmental/growth effects

the experimental strains may exert on the worms. All experimental bacterial strains were

preconditioned in red light with optogenetic light provided by a single LED positioned 1 cm

above the Petri dish.

After 3-5 days (between days 1-3 of adulthood), worms allocated for the experiment were

transferred to experimental strains for approximately 60-90 minutes to thoroughly inoculate the

GI tract. In the case of the unparalyzed worms, red or green light was then applied for an

additional 6 hours. For the paralyzed worm experiments, 1.5% low-melt agar pads were prepared

as described above and placed on individual slides. About 15 adult worms were transferred from

the experimental strain Petri dish to an agar pad containing 10 µL of *C. elegans* M9 buffer + 2

mM levamisole (M9Ce+Lev), where worms were gently washed before being transferred to a

fresh pad also containing 10 µL of M9Ce+Lev. The majority of M9Ce+Lev on the pad was

allowed to evaporate, which causes the worms to align longitudinally before a cover slip was

applied. Slides were then exposed to either red or green light by placing them under a single

LED positioned 1 cm above the Petri dish for 6 h. Afterward, the anterior intestinal cells were

imaged using confocal microscopy (Olympus Fluoview 3000) in the brightfield and GFP

channels.

## Confocal Microscopy Image Analysis

All confocal images for an experiment were manually cropped to display only the anterior intestinal cells of a single worm (in the GFP channel). These cropped images were then anonymized, randomized and the mitochondrial networks of each were blindly classified by two researchers independently as either tubular, fragmented, or intermediate. Tubular samples were marked by a high degree of network connectivity throughout. Fragmented samples were composed almost exclusively of isolated clusters of fluorescence with high circularity. Intermediate samples contained regions of both types. Scores were then de-randomized and aggregated. For each experimental strain, the red and green light conditions were compared for statistical significance using the chi-squared test of homogeneity.

## Lifespan experiments

3 mL starter cultures of $\Delta lon$, $\Delta rcsA$ or MVK29 were inoculated from -80ºC freezer stocks into LB supplemented with appropriate antibiotics and grown shaking for 12 h at 37 ºC at 250 rpm. These cultures were diluted to $OD_{600} = 1\times10^{-6}$ in 27 mL M9 media supplemented with appropriate antibiotics. 1.5 mL of each M9/cell mixture was added to each of 18 wells on three 24-well plates and grown in 3 LPA devices under the appropriate light conditions at 37 ºC and 250 rpm. Once cultures reached $OD_{600} = 0.1\text{-}0.4$, all tubes were iced for 10 min and subsequently concentrated 10 times by centrifugation (4ºC, 4000 rpm, 10 min). Approximately 50 μL of this dense bacterial culture was then applied to sterile NGM-agar plates with no antibiotics and allowed to dry in a dark room. The plates were then illuminated with the appropriate light wavelength and intensity for 16 h at room temperature, and immediately used for lifespan assays.

During the longitudinal lifespan assay, exposure to white light is limited to the minimal

level. To reach this goal, the *sqt-3(e2117)* temperature sensitive mutant (**Supplementary Table**

**3**) was used to perform longitudinal analyses at 25 ℃, which avoids time-consuming animal

transfer without interrupting normal reproduction. *sqt-3(e2117)* is a collagen mutant of *C.*

*elegans* that reproduces normally but is embryonically lethal at 26 ℃, and has been used

previously in longitudinal studies(Han et al., 2017; Wang et al., 2014). Worms were age-

synchronized by bleach-based egg isolation followed by starvation in M9 buffer at the L1 stage

for 36 hours. Synchronized L1 worms were grown on BW25113 *E. coli* at 15 ℃ until the L4

stage, when worms were transferred to 24-well plates (~15 worms/well) with $\Delta lon$, $\Delta rcsA$ or

MVK29 (**Supplementary Table 3**). The plates were placed in LPA. The LPA LEDs were

programmed to illuminate wells with constant red (10 μmol/m$^2$/s), low-intensity green (0.25

μmol/m$^2$/s), or high-intensity green light (10 μmol/m$^2$/s). The apparatus was then transferred to a

26 ℃ incubator. The number of living worms remaining in each well was counted every day.

Death was indicated by total cessation of movement in response to gentle mechanical

stimulation. Statistical analyses were performed with SPSS (IBM Software) using Kaplan-Meier

survival analysis and the log-rank test (**Supplementary Table 4**).

**Supplemental Information Titles and Legends**

**Supplementary Figure 1**. *In vitro* **characterization of GFP reporter strains used in this**

**study.** All optogenetic strains carry our previously published CcaSR v2.0 system, which is

encoded on plasmids (a) pSR43.6 and (b) pSR58.6. (c) pLH401 (used for microscopy) and (d)

pLH405 (used for cytometry) genetic device schematics. We replaced *sfgfp* with *gfpmut3\** in

pLH405 as we hypothesized that the latter may be less stable and thus result in faster response

dynamics. However, we observed no difference in dynamics. (e) Batch culture light responses of

CcaSR v2.0 and all GFP reporter strains used in this study. GFP fluorescence was measured by

flow cytometry. The dynamic range (ratio of GFP fluorescence in green versus red light) is

shown above each data set. We note that CcaSR v2.0 exhibits 77-fold dynamic range in the

reference strain BW29655 (Δ*envZ*, Δ*ompR*), which is similar to our previous measurement of

this strain at 120-fold (Schmidl et al., 2014). The calculated fold-change is sensitive to

fluctuations in the measured *E. coli* autofluorescence and red-light expression level, which likely

explains the slight discrepancy. CcaSR v2.0 dynamic range increases slightly to 84.2-fold in

Δ*rcsA* (JW1935-1), which is used throughout this work, due to higher sfGFP expression in green

light. The mCherry cassette in pLH401 decreases dynamic range to 38.3-fold due to higher leaky

sfGFP expression in red light. In worms, the fold-change in response to green light decreases

further to 5.52 ± 2.4-fold (**Fig. 1d**). The use of *gfpmut3\** in pLH405 further decreases dynamic

range to 13.5 ± 0.0-fold for reasons that are not clear, while in worms the fold-change is 8.63 ±

3.6-fold (**Fig. 1e**). Data represent the mean of three independent, autofluorescence-subtracted,

biological replicates acquired on 3 separate days. Error bars: standard deviation.

**Supplementary Figure 2. Flow cytometry gating.** (a) Strain LH05 (**Supplementary Table 3**),

which expresses only mCherry, was fed to worms, isolated, and analyzed by flow cytometry to

quantify *E. coli* autofluorescence through the FL1 (GFP) channel. To eliminate events

corresponding to cytometer noise and autofluorescence of homogenized worm samples, three

gates were then applied: (1) a density gate for the most homogeneous 75% of samples in forward

scatter (FSC) vs side-scatter (SSC) (the area within the bold black line in the plots in Column 1),

(2) a threshold gate on the FL1 (GFP) channel (marked by a red dashed line; **Methods**), and (3) a

threshold gate for events exhibiting high FL3 (mCherry, red dashed line; **Methods**). (b)

Applying these gates to samples from the step ON experiment (**Fig. 1e**) reveals robust isolation

of bacteria with a FSC/SSC profile consistent with our previous *in vitro* experiments[24,47] and an

increase in expression of GFPmut3* in response to green light. Ungated data are shown for

comparison to gated data in all plots.

**Supplementary Figure 3. Gene expression dynamics from flow cytometry experiments.** (a)

LH03 and (b) LH04 (ΔPCB) step ON results. (c) LH03 and (d) LH04 step OFF results. Each

individual trajectory (faint lines) corresponds to a single biological replicate. A biological

replicate comprises the homogenized contents of five worms collected on a single day. Each

biological replicate on a given plot was run on a different day. Population medians taken across

all trajectories are shown in bold (**Methods**). Data are composed of six and eight trajectories in

the step ON and step OFF experiments, respectively. Error bars: SEM.

**Supplementary Figure 4. Bacteria on the exterior of worms do not contribute to the**

**measured light response in flow cytometry experiments.** Worms were suspended in clear

tubes containing M9Ce+Lev media and levamisole in the flow cytometry experiments. To

demonstrate that bacterial cells outside the worm gut (e.g. on the exterior of the worm) do not

respond to light in these conditions, samples of pre-conditioned bacteria from the NGM plates

were suspended in the M9Ce+Lev or *E. coli* M9 media supplemented with casamino acids and

glucose (M9Ec). These samples were then exposed to either green or red light for 8 h, then

measured via flow cytometry after the fluorophore maturation protocol (**Methods**). As expected,

the bacterial populations are only responsive to light when grown in M9Ec and when the PCB

operon is intact (strain JW1935-1/pLH405/pSR43.6, aka LH03). We conclude that the responses

shown in data in **Fig. 1e-f** and **Supplementary Fig. 3** are not due to bacteria outside the worm,

nor to bacteria that escape the worm over the course of the experiment, as such cells do not

respond to light in the experimental buffer (M9Ce). Data represent 7, 6, 3, and 5 replicates (left);

and 8, 3, 8, and 3 replicates (right) over 8 and 10 days, respectively; error bars: SEM. Individual

data points for each condition are overlaid as white markers.

**Supplementary Table 1. Plasmids used in this study**

**Supplementary Table 2. Genbank Accession Numbers**

**Supplementary Table 3. Bacterial and worm strains used in this study**

**Supplementary Table 4. Statistical analysis of worm lifespan experiments**

**References**

Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner

BL, Mori H. 2006. Construction of Escherichia coli K□12 in□frame, single□gene knockout

mutants: the Keio collection. *Mol Syst Biol* **2**:2006.0008. doi:10.1038/msb4100050

Biagi E, Franceschi C, Rampelli S, Severgnini M, Ostan R, Turroni S, Consolandi C, Quercia S,

Scurti M, Monti D, Capri M, Brigidi P, Candela M. 2016. Gut Microbiota and Extreme

Longevity. *Curr Biol* **26**:1480–1485. doi:10.1016/j.cub.2016.04.016

Cabreiro F, Au C, Leung K-Y, Vergara-Irigaray N, Cochemé HM, Noori T, Weinkove D,

Schuster E, Greene NDE, Gems D. 2013. Metformin retards aging in C. elegans by altering

microbial folate and methionine metabolism. *Cell* **153**:228–39. doi:10.1016/j.cell.2013.02.035

Castillo-Hair SM, Baerman EA, Fujita M, Igoshin OA, Tabor JJ. 2019. Optogenetic control of

Bacillus subtilis gene expression. *Nat Commun* **10**:3099. doi:10.1038/s41467-019-10906-6

Castillo-Hair SM, Sexton JT, Landry BP, Olson EJ, Igoshin OA, Tabor JJ. 2016. FlowCal: A

User-Friendly, Open Source Software Tool for Automatically Converting Flow Cytometry

Data from Arbitrary to Calibrated Units. *Acs Synth Biol* **5**:774–80.

doi:10.1021/acssynbio.5b00284

Chait R, Ruess J, Bergmiller T, Tkačik G, Guet CC. 2017. Shaping bacterial population behavior

through computer-interfaced control of individual cells. *Nat Commun* **8**:1535.

doi:10.1038/s41467-017-01683-1

Chen Y-J, Liu P, Nielsen AAK, Brophy JAN, Clancy K, Peterson T, Voigt CA. 2013.

Characterization of 582 natural and synthetic terminators and quantification of their design

constraints. *Nat Methods* **10**:659–664. doi:10.1038/nmeth.2515

Cho D-H, Nakamura T, Fang J, Cieplak P, Godzik A, Gu Z, Lipton SA. 2009. S-nitrosylation of

Drp1 mediates beta-amyloid-related mitochondrial fission and neuronal injury. *Sci New York*

*N Y* **324**:102–5. doi:10.1126/science.1171091

Claesson MJ, Cusack S, O'Sullivan O, Greene-Diniz R, Weerd H de, Flannery E, Marchesi JR,

Falush D, Dinan T, Fitzgerald G, Stanton C, Sinderen D van, O'Connor M, Harnedy N,

O'Connor K, Henry C, O'Mahony D, Fitzgerald AP, Shanahan F, Twomey C, Hill C, Ross

RP, O'Toole PW. 2010. Composition, variability, and temporal stability of the intestinal

microbiota of the elderly. *P Natl Acad Sci Usa* **108 Suppl 1**:4586–91. doi:10.1073/pnas.1000097107

Claesson MJ, Jeffery IB, Conde S, Power SE, O'Connor EM, Cusack S, Harris HMB, Coakley M, Lakshminarayanan B, O'Sullivan O, Fitzgerald GF, Deane J, O'Connor M, Harnedy N, O'Connor K, O'Mahony D, Sinderen D van, Wallace M, Brennan L, Stanton C, Marchesi JR, Fitzgerald AP, Shanahan F, Hill C, Ross RP, O'Toole PW. 2012. Gut microbiota composition correlates with diet and health in the elderly. *Nature* **488**:178–84. doi:10.1038/nature11319

Couillault C, Ewbank JJ. 2002. Diverse Bacteria Are Pathogens of Caenorhabditis elegans. *Infect Immun* **70**:4705–4707. doi:10.1128/iai.70.8.4705-4707.2002

Deisseroth K. 2015. Optogenetics: 10 years of microbial opsins in neuroscience. *Nat Neurosci* **18**:1213–25. doi:10.1038/nn.4091

DISCHE Z. 1947. A new specific color reaction of hexuronic acids. *J Biological Chem* **167**:189–98.

DISCHE Z, SHETTLES LB. 1948. A specific color reaction of methylpentoses and a spectrophotometric micromethod for their determination. *J Biological Chem* **175**:595–603.

Donato V, Ayala FR, Cogliati S, Bauman C, Costa JG, Leñini C, Grau R. 2017. Bacillus subtilis biofilm extends Caenorhabditis elegans longevity through downregulation of the insulin-like signalling pathway. *Nat Commun* **8**:14332. doi:10.1038/ncomms14332

Engler C, Gruetzner R, Kandzia R, Marillonnet S. 2009. Golden gate shuffling: a one-pot DNA shuffling method based on type IIs restriction enzymes. *Plos One* **4**:e5553. doi:10.1371/journal.pone.0005553

Exner N, Treske B, Paquet D, Holmstrom K, Schiesling C, Gispert S, Carballo-Carbajal I, Berg

D, Hoepken H-H, Gasser T, Kruger R, Winklhofer KF, Vogel F, Reichert AS, Auburger G,

Kahle PJ, Schmid B, Haass C. 2007. Loss-of-Function of Human PINK1 Results in

Mitochondrial Pathology and Can Be Rescued by Parkin. *J Neurosci* **27**:12413–12418.

doi:10.1523/jneurosci.0719-07.2007

Fernandez-Rodriguez J, Moser F, Song M, Voigt CA. 2017. Engineering RGB color vision into

Escherichia coli. *Nat Chem Biol* **13**:706–708. doi:10.1038/nchembio.2390

Gautier A, Gauron C, Volovitch M, Bensimon D, Jullien L, Vriz S. 2014. How to control

proteins with light in living systems. *Nat Chem Biol* **10**:533–541. doi:10.1038/nchembio.1534

Gerhardt KP, Olson EJ, Castillo-Hair SM, Hartsough LA, Landry BP, Ekness F, Yokoo R,

Gomez EJ, Ramakrishnan P, Suh J, Savage DF, Tabor JJ. 2016. An open-hardware platform

for optogenetics and photobiology. *Sci Rep-uk* **6**:35363. doi:10.1038/srep35363

Goglia AG, Toettcher JE. 2018. A bright future: optogenetics to dissect the spatiotemporal

control of cell behavior. *Curr Opin Chem Biol* **48**:106–113. doi:10.1016/j.cbpa.2018.11.010

Gusarov I, Gautier L, Smolentseva O, Shamovsky I, Eremina S, Mironov A, Nudler E. 2013.

Bacterial Nitric Oxide Extends the Lifespan of C. elegans. *Cell* **152**:818–830.

doi:10.1016/j.cell.2012.12.043

Han B, Sivaramakrishnan P, Lin C-CJ, Neve IAA, He J, Tay LWR, Sowa JN, Sizovs A, Du G,

Wang J, Herman C, Wang MC. 2017. Microbial Genetic Composition Tunes Host Longevity.

*Cell* **169**:1249-1262.e13. doi:10.1016/j.cell.2017.05.036

Kenyon CJ. 2010. The genetics of ageing. *Nature* **464**:504–512. doi:10.1038/nature08980

Kim DH. 2013. Bacteria and the aging and longevity of Caenorhabditis elegans. *Annu Rev Genet*

**47**:233–46. doi:10.1146/annurev-genet-111212-133352

Köhler P, Bachmann R. 1978. The effects of the antiparasitic drugs levamisole, thiabendazole,

praziquantel, and chloroquine on mitochondrial electron transport in muscle tissue from

Ascaris suum. *Mol Pharmacol* **14**:155–63.

Kotula JW, Kerns SJ, Shaket LA, Siraj L, Collins JJ, Way JC, Silver PA. 2014. Programmable

bacteria detect and record an environmental signal in the mammalian gut. *P Natl Acad Sci*

*Usa* **111**:4838–43. doi:10.1073/pnas.1321321111

Kundu P, Blacher E, Elinav E, Pettersson S. 2017. Our Gut Microbiome: The Evolving Inner

Self. *Cell* **171**:1481–1493. doi:10.1016/j.cell.2017.11.024

Leopold AV, Chernov KG, Verkhusha VV. 2018. Optogenetically controlled protein kinases for

regulation of cellular signaling. *Chem Soc Rev* **47**:2454–2484. doi:10.1039/c7cs00404d

Levskaya A, Chevalier AA, Tabor JJ, Simpson ZB, Lavery LA, Levy M, Davidson EA, Scouras

A, Ellington AD, Marcotte EM, Voigt CA. 2005. Engineering Escherichia coli to see light.

*Nature* **438**:441–442. doi:10.1038/nature04405

Li X, Zhang C, Xu X, Miao J, Yao J, Liu R, Zhao Y, Chen X, Yang Y. 2020. A single-

component light sensor system allows highly tunable and direct activation of gene expression

in bacterial cells. *Nucleic Acids Res*. doi:10.1093/nar/gkaa044

Lim B, Zimmermann M, Barry NA, Goodman AL. 2017. Engineered Regulatory Systems

Modulate Gene Expression of Human Commensals in the Gut. *Cell* **169**:547-558.e15.

doi:10.1016/j.cell.2017.03.045

Milias-Argeitis A, Rullan M, Aoki SK, Buchmann P, Khammash M. 2016. Automated

optogenetic feedback control for precise and robust regulation of gene expression and cell

growth. *Nat Commun* **7**:12546. doi:10.1038/ncomms12546

Mimee M, Tucker AC, Voigt CA, Lu TK. 2015. Programming a Human Commensal Bacterium,

Bacteroides thetaiotaomicron, to Sense and Respond to Stimuli in the Murine Gut Microbiota.

*Cell Syst* **1**:62–71. doi:10.1016/j.cels.2015.06.001

Ohlendorf R, Vidavski RR, Eldar A, Moffat K, Möglich A. 2012. From Dusk till Dawn: One-

Plasmid Systems for Light-Regulated Gene Expression. *J Mol Biol* **416**:534–542.

doi:10.1016/j.jmb.2012.01.001

Olson EJ, Hartsough LA, Landry BP, Shroff R, Tabor JJ. 2014. Characterizing bacterial gene

circuit dynamics with optically programmed gene expression signals. *Nat Methods* **11**:449–

55. doi:10.1038/nmeth.2884

Olson EJ, Tabor JJ. 2014. Optogenetic characterization methods overcome key challenges in

synthetic and systems biology. *Nat Chem Biol* **10**:502–11. doi:10.1038/nchembio.1559

Olson EJ, Tzouanas CN, Tabor JJ. 2017. A photoconversion model for full spectral

programming and multiplexing of optogenetic systems. *Mol Syst Biol* **13**:926.

doi:10.15252/msb.20167456

Ong NT, Tabor JJ. 2018. A Miniaturized Escherichia coli Green Light Sensor with High

Dynamic Range. *Chembiochem* **19**:1255–1258. doi:10.1002/cbic.201800007

Ong NTX, Olson EJ, Tabor JJ. 2017. Engineering an E. coli near-infrared light sensor. *Acs Synth*

*Biol* **7**:240–248. doi:10.1021/acssynbio.7b00289

O'Toole PW, Jeffery IB. 2015. Gut microbiota and aging. *Sci New York N Y* **350**:1214–5.

doi:10.1126/science.aac8469

Portal-Celhay C, Bradley ER, Blaser MJ. 2012. Control of intestinal bacterial proliferation in

regulation of lifespan in Caenorhabditis elegans. *Bmc Microbiol* **12**:49. doi:10.1186/1471-

2180-12-49

Pryor R, Norvaisas P, Marinos G, Best L, Thingholm LB, Quintaneiro LM, Haes WD, Esser D,

Waschina S, Lujan C, Smith RL, Scott TA, Martinez-Martinez D, Woodward O, Bryson K,

Laudes M, Lieb W, Houtkooper RH, Franke A, Temmerman L, Bjedov I, Cochemé HM,

Kaleta C, Cabreiro F. 2019. Host-Microbe-Drug-Nutrient Screen Identifies Bacterial

Effectors of Metformin Therapy. *Cell* **178**:1299-1312.e29. doi:10.1016/j.cell.2019.08.003

Ramakrishnan P, Tabor JJ. 2016. Repurposing Synechocystis PCC6803 UirS-UirR as a UV-

Violet/Green Photoreversible Transcriptional Regulatory Tool in E. coli. *Acs Synth Biol*

**5**:733–40. doi:10.1021/acssynbio.6b00068

Ryu M-H, Gomelsky M. 2014. Near-infrared light responsive synthetic c-di-GMP module for

optogenetic applications. *Acs Synth Biol* **3**:802–10. doi:10.1021/sb400182x

Schmidl SR, Sheth RU, Wu A, Tabor JJ. 2014. Refactoring and Optimization of Light-

SwitchableEscherichia coliTwo-Component Systems. *Acs Synth Biol* **3**:820–831.

doi:10.1021/sb500273n

Sebastián D, Palacín M, Zorzano A. 2017. Mitochondrial Dynamics: Coupling Mitochondrial

Fitness with Healthy Aging. *Trends Mol Med* **23**:201–215.

doi:10.1016/j.molmed.2017.01.003

Tabor JJ, Levskaya A, Voigt CA. 2010. Multichromatic control of gene expression in

Escherichia coli. *J Mol Biol* **405**:315–24. doi:10.1016/j.jmb.2010.10.038

Tabor JJ, Salis HM, Simpson ZB, Chevalier AA, Levskaya A, Marcotte EM, Voigt CA,

Ellington AD. 2009. A synthetic genetic edge detection program. *Cell* **137**:1272–81.

doi:10.1016/j.cell.2009.04.048

Tandar ST, Senoo S, Toya Y, Shimizu H. 2019. Optogenetic switch for controlling the central

metabolic flux of Escherichia coli. *Metab Eng* **55**:68–75. doi:10.1016/j.ymben.2019.06.002

Virk B, Jia J, Maynard CA, Raimundo A, Lefebvre J, Richards SA, Chetina N, Liang Y,

Helliwell N, Cipinska M, Weinkove D. 2016. Folate Acts in E. coli to Accelerate C. elegans

Aging Independently of Bacterial Biosynthesis. *Cell Reports* **14**:1611–20.

doi:10.1016/j.celrep.2016.01.051

Wang MC, Oakley HD, Carr CE, Sowa JN, Ruvkun G. 2014. Gene pathways that delay

Caenorhabditis elegans reproductive senescence. *Plos Genet* **10**:e1004752.

doi:10.1371/journal.pgen.1004752

Zhang F, Berg M, Dierking K, Félix M-A, Shapira M, Samuel BS, Schulenburg H. 2017.

Caenorhabditis elegans as a Model for Microbiome Research. *Front Microbiol* **8**:485.

doi:10.3389/fmicb.2017.00485