Functional remodeling of lysosomes by type I interferon modifies host defense

Hailong Zhang[1,2,3], Abdelrahim Zoued[1,2,3], Xu Liu[1,2,3], Brandon Sit[1,2], Matthew K. Waldor[1,2,3]*

1Division of Infectious Diseases, Brigham and Women's Hospital, Boston, Massachusetts, USA

2Department of Microbiology, Harvard Medical School, Boston, Massachusetts, USA

3Howard Hughes Medical Institute, Boston, Massachusetts, USA

*Correspondence: mwaldor@research.bwh.harvard.edu

SUMMARY

Organelle remodeling is critical for cellular homeostasis, but host factors that control organelle function during microbial infection remain largely uncharacterized. Here, a genome-scale CRISPR/Cas9 screen in intestinal epithelial cells with the prototypical intracellular bacterial pathogen *Salmonella* led us to discover that type I interferon (IFN-I) remodels lysosomes. Even in the absence of infection, IFN-I signaling modified the localization, acidification, protease activity and proteomic profile of lysosomes. Proteomic and genetic analyses revealed that multiple IFN-I-stimulated genes including *Ifitm3*, *Slc15a3*, and *Cnp* contribute to lysosome acidification. IFN-I-dependent lysosome acidification stimulated intracellular *Salmonella* virulence gene expression, leading to rupture of the *Salmonella*-containing vacuole and host cell death. Moreover, IFN-I signaling promoted *in vivo Salmonella* pathogenesis in the intestinal epithelium, where *Salmonella* initiates infection. Our findings explain how an intracellular bacterial pathogen co-opts epithelial IFN-I signaling. We propose that IFN-I control of lysosome function broadly impacts host defense against diverse viral and microbial pathogens.

KEYWORDS

INTRODUCTION

Microbial pathogens have evolved varied virulence strategies to modulate host cell function (Geoghegan and Holmes, 2018; Ribet and Cossart, 2015). A common mechanism, employed by all viral and some bacterial pathogens, is to enter host cells, where they co-opt cellular functions while simultaneously evading extracellular threats such as innate and adaptive immune mechanisms (Hybiske and Stephens, 2008; Lee et al., 2019). Inside cells, intracellular pathogens interact with and exploit host cell organelles to support their proliferation (Omotade and Roy, 2019). As a result of their intimate relationships with and manipulation of varied host cell functions, intracellular pathogens have proven to be outstanding tools to probe basic eukaryotic cell biology (Welch, 2015).

Compared to knowledge of how microbial pathogens interact with phagocytic cells, less is known about the landscape of pathogen-epithelial cell interactions at barrier sites, where most infections originate (Jo, 2019). The human foodborne pathogen *Salmonella enterica* serovar Typhimurium (Stm) is a model intracellular bacterium that initially invades and subsequently kills intestinal epithelial cells (IECs) before spreading systemically via circulating phagocytes (Hurley et al., 2014). Stm's entry into and initial trafficking inside IEC is well-characterized, and a hallmark of Stm infection is the formation of the *Salmonella*-containing vacuole (SCV), a dynamic, lysosome-like compartment that is permissive for Stm replication (Steele-Mortimer, 2008; Tuli and Sharma, 2019). However, the IEC pathways that control Stm-induced cytotoxicity remain incompletely defined.

One striking feature of the host response to Stm is the induction of type I interferons (IFN-Is), which include IFNα and IFNβ (Hess et al., 1989). IFN-Is are cytokines that, once secreted, bind the IFN-I receptor (IFNAR1/2) to activate JAK-STAT signaling, which triggers expression of intracellular anti-microbial transcriptional programs consisting of over 400 IFN-stimulated genes (ISGs) (Schoggins and Rice, 2011). Due to the large size of the "interferome" and the complex interactions of ISGs with thousands of additional cellular proteins (Hubel et al., 2019), knowledge of the full spectrum of IFN-I-mediated changes in cellular function is incomplete. Although IFN-Is are known to have critical roles in antiviral responses, their functions in bacterial infection are less clear, and IFN-I signaling has been reported to be either protective or detrimental to the host depending on the specific bacterial pathogen (Kovarik et al., 2016).

Here, we carried out a genome-scale CRISPR/Cas9 screen to identify the host factors that contribute to Stm's cytotoxicity to IECs. This screen revealed IFN-I signaling as a key susceptibility factor for cytotoxicity in IECs and led to our discovery of a novel role for IFN-I signaling in lysosomal localization and function, including the modification of this organelle's pH, protease activity, and protein content. Organellar proteomics revealed that 11 ISGs were enriched in lysosomes following IFN-I stimulation, several of which were found to directly impact lysosomal pH. IFN-I signaling-dependent lysosomal acidification stimulated Stm virulence gene expression, and *in vivo* studies confirmed a role for epithelial IFN-I signaling in promoting systemic Stm infection. IFN-I signaling mediated control of lysosome function likely contributes to host responses to diverse intracellular pathogens and viruses.

RESULTS

A CRISPR/Cas9 screen identifies IEC factors required for Stm cytotoxicity

As large-scale genetic screens to identify epithelial cell factors that mediate interactions with

intracellular pathogens are lacking, we performed a multi-round, genome-wide CRISPR/Cas9

loss-of-function screen in the human colonic epithelial cell line HT29-Cas9 (Blondel et al.,

2016), to identify IEC genes that confer resistance to Stm cytotoxicity (Figure 1A). HT29 cells

are efficiently invaded and subsequently killed by Stm, providing a strong selective force to

enrich for guide RNAs targeting host factors that modulate cytotoxicity. The screen identified

known pathways that sensitize cells to Stm infection, including those involved in regulation of

actin dynamics (Arp2/3 and Rac), which are important in pathogen invasion (Table S1 and

Figures 1B, C), (Unsworth et al., 2004; Yeung et al., 2019). Genes linked to pathways not

previously directly linked to Stm virulence, including the Fc-gamma receptor-dependent

phagocytic and GPI anchor modification pathways were also enriched (Figures 1B, C).

Strikingly, the top 'hits' of the screen were remarkably coherent - the seven most enriched guide

RNAs from both libraries screened corresponded to genes in the IFN-I signaling pathway,

including the receptor (*Ifnar1*/*Ifnar2*), adaptor (*Jak1*/*Tyk2*), and transcription factor

(*Stat1*/*Stat2*/*Irf9*) components of the system (Figures 1D, E), suggesting that IFN-I signaling is

a major driver of Stm-mediated cytotoxicity in IECs.

IFN-I promotes Stm cytotoxicity in IECs

Stm induces IFN-I production during infection (Hess et al., 1989), but the function of IFN-I

signaling in IECs is unknown. A clonal knockout of *Ifnar2,* the top enriched hit in both libraries,

was constructed in the HT29-Cas9 background (Figure S1A), to validate the screen findings. At

both early and late infection time points, *Ifnar2* KO cells were more resistant to Stm-induced

cell death than the wild-type (WT) parental line (Figure 2A and S1B, C). Priming cells with

IFNβ (a major IFN-I), conditions that mimic the multiple rounds of the original screen, further

sensitized WT but not *Ifnar2* KO cells to death (Figure 2A). To complement these findings, we

treated WT cells with chemical inhibitors of JAK-STAT signaling, the downstream target of

activated IFNAR1/2. Similar to *Ifnar2* KO cells, treatment with the JAK inhibitors ruxolitinib

and pyridone-6 also diminished Stm-induced death in WT cells, indicating active IFN-I

signaling is required for this phenotype (Figure 2B).

Our overall observations that IFN-I promotes Stm-induced IEC death is consistent with

prior data that this cytokine enhances necroptosis in Stm-infected macrophages (Robinson et al.,

2012). However, it is not clear whether macrophage and epithelial cell responses to Stm

infection are similar; furthermore, it is known that Stm-induced cell death in macrophages is

invasion-independent (van der Velden et al., 2000). Thus, we next tested whether

IFN-I-promoted epithelial cell death depended on SPI-1 or SPI-2, critical *Salmonella*

pathogenicity islands that each encode type 3 secretion systems (T3SS) required for cellular

invasion and intracellular survival, respectively (Galan et al., 2014). SPI-1-deficient (*ΔprgH*)

Stm did not induce epithelial cell death in any condition, confirming that in IECs cytotoxicity

requires cell invasion (Figure 2A). In contrast, SPI-2-deficient (*ΔssaV*) Stm led to reduced but

still detectable levels of cytotoxicity that remained sensitive to IFNβ priming, suggestive of

both SPI-2-dependent and independent mechanisms of intracellular Stm-induced cytotoxicity

(Figure 2A).

In support of the population-level LDH assays, flow cytometry of HT29 or HeLa cells

infected with fluorescent Stm and stained with the cell death probe Annexin-V indicated that

IFN-I only influenced cell death in the population of cells that contained intracellular Stm

(Figures 2C and S1D-F). In addition, we found that. IFN-I signaling did not impact Stm

invasion (Figures 2D and S2A), nor did it influence bacterial association with the early

endosomal marker Rab5, late endosomal marker Rab7 or lysosomal marker LAMP1

(Desjardins et al., 1994) (Figures 2E, F and S2B-E). Together, these data suggest that

IFN-I-mediated sensitization of epithelial cells to Stm occurs downstream of cell invasion and

initial SCV formation.

IFN-I regulates lysosome localization and function

During our analyses of SCV formation, we unexpectedly observed that IFN-I signaling alters

the subcellular localization of lysosomes in epithelial cells, even in the absence of infection. In

HeLa cells, lysosomes (identifiable as LAMP1+/Lysotracker+ co-staining organelles) were

scattered throughout the cytoplasm under basal conditions. Following IFNβ stimulation,

lysosomes re-localized to the perinuclear region (Figures 3A, B, Video S1, 2); lysosome

re-localization was not observed in *Ifnar2* KO HeLa cells, confirming that this response was

dependent on IFN-I signaling.

Notably, IFNβ priming led to significantly higher intensities of two fluorescent reporters

(Lysotracker and Lysosensor) of lysosomal pH in WT but not *Ifnar2* KO cells, indicating that

IFN-I signaling lowers lysosome pH (Figures 3C and S3A-C). Moreover, staining with

fluorescent reporters of general lysosomal protease activity (DQ-BSA (Reis et al., 1998)), or

cathepsin D (a major lysosomal protease) activity revealed that their activities were elevated by

IFNβ stimulation in an *Ifnar2*-dependent fashion (Figures 3D, E and S3D). These findings are

consistent with the idea that the activity of most resident lysosomal proteins, including

cathepsins and other degradative enzymes, is positively regulated by acidic pH (Butor et al.,

1995). Staining with fluorescent reporters of endocytic activity (Dextran-568) suggested that

IFN-I signaling does not impact general endocytic trafficking; thus, IFN-I signaling appears to

specifically influence lysosome function and positioning (Figure S3E). IFNβ-induced lysosomal

acidification and protease activation was abolished by the addition of the v-ATPase inhibitor

bafilomycin A1 (Bfa1) (Yoshimori et al., 1991) (Figure 3C, D), demonstrating that IFN-I

signaling primarily relies on the conventional lysosomal acidification machinery.

Together, these observations reveal that IFN-I signaling promotes epithelial cell lysosomal

re-localization, acidification and degradative activity, without broadly affecting intracellular

trafficking. Besides epithelial cells such as HeLa and HT29, IFNβ treatment also reduced

lysosomal pH in monocyte/macrophage-like THP-1 cells (Figure S3F), suggesting that IFN-I

signaling controls lysosomal acidification outside of epithelial cell lineages.

The ISG IFITM3 regulates lysosomal function and Stm cytotoxicity

To begin to dissect the mechanism by which IFN-I signaling regulates lysosome acidification

and function, we first took a candidate-based approach and investigated IFITM3. This

transmembrane ISG has antiviral activity and is thought to reside in the endosomal trafficking

system and to interact with the lysosomal v-ATPase complex (Spence et al., 2019; Wee et al.,

2012), suggesting a potential role for this protein in lysosome function. Immunofluorescence

analysis revealed that IFITM3 co-localized with LAMP1, but not Rab5, confirming that

IFITM3 is a lysosomal protein (Figure 4A). Remarkably, lysosomal pH in *Ifitm3* KO cells was

elevated in both basal and IFNβ-primed conditions relative to WT cells, partially phenocopying

the *Ifnar2* KO, and suggesting that IFITM3 contributes to IFN-I-meditated lysosomal

remodeling (Figures 4B-D). *Ifitm3*'s contribution to basal pH levels are consistent with the tonic

activities of IFNs observed in diverse mammalian cell types (Schoggins et al., 2014). *Ifitm3* KO

cells were more resistant to Stm-induced cell death than the WT parental line both before and

after IFNβ priming (Figure 4E), suggesting that this ISG contributes to IFN-I signaling

augmentation of Stm cytotoxicity.

Discovery of ISGs with novel roles in lysosomal pH regulation

Given that both lysosomal pH and degradative activity in *Ifitm3* KO cells were still somewhat

sensitive to IFNβ priming (Figures 4C, D), we hypothesized that additional ISGs regulate

lysosome function. To identify these factors, we employed organellar proteomics, a powerful

and unbiased affinity-based technique that has not yet been applied to host-pathogen

interactions. We used a recently described lysosomal pulldown system, LysoIP (Abu-Remaileh

et al., 2017), to profile the proteomes of intact lysosomes from WT or *Ifnar2* KO cells in basal

or IFNβ-stimulated states. The purity and integrity of the lysosome samples was confirmed by

verifying the presence of luminal cathepsin D and the absence of cytosolic and Golgi apparatus

markers (Figures 5A and S4). Quantitative profiling revealed that the abundances of ~15

proteins, most of them ISGs, were increased in purified lysosomes upon IFNβ stimulation

(Figure 5B). Spectral counts for IFITM3 were enriched in lysosomes from IFNβ-treated cells,

supporting the imaging above (Figure 4A) and providing validation of the dataset. Immunoblots

of purified lysosomal and cytoplasmic fractions from naïve and IFNβ-treated cells with

antibodies to IFITM3 further corroborated this observation, and immunoblots for the known

cytosolic ISG IFIT3 served as a negative control in this assay (Figure 5A).

We constructed KO cell lines for most of the lysosomally enriched ISGs and assessed their

contributions to the pH of this degradative organelle (Figure 5C). Although most ISGs did not

appear to influence lysosomal pH, we found two additional ISGs (*Slc15a3* and *Cnp*) that like

Ifitm3 contributed to both basal and IFN-I-mediated lysosomal acidification (Figure 5C). In

contrast to *Ifitm3*, *Slc15a3*, a lysosome-resident, proton-coupled histidine and di-tripeptide

transporter (Song et al., 2018), likely does not interact with the v-ATPase, suggesting that both

v-ATPase-dependent and independent mechanisms may underlie IFN-I-induced lysosomal

acidification. Similarly*, Cnp*, the other identified regulator of lysosome pH, is a 2',3'-cyclic

nucleotide 3' phosphodiesterase whose activity has not been linked to the v-ATPase.

IFN-I stimulates intracellular Stm virulence gene expression and facilitates SCV damage

We hypothesized that IFN-I's role in lysosomal acidification might explain why IFN-I signaling

enhances Stm cytotoxicity because acidic pH is known to stimulate expression of

SPI-2-encoded and other virulence genes (Chakraborty et al., 2015; Prost et al., 2007).

Consistent with this idea, IFNβ priming increased intracellular Stm SPI-2 encoded gene

expression (Figures 6A and S5A). These genes were only induced after SCV formation (i.e.

later than one-hour post-infection), and the effect of IFNβ priming was eliminated in *Ifnar2* KO

cells. Furthermore, treatment with Bfa1 abolished IFNβ induction of SPI-2 expression (Figures

6A and S5A), suggesting that diminished SCV acidification is the primary mechanism of

IFN-I-enhanced SPI-2 induction. Analyses of SPI-2 gene expression using flow cytometry and

a fluorescent $P_{sifB}::gfp$ Stm reporter strain (Garmendia et al., 2003), confirmed this phenotype at

single bacterial cell resolution (Figure 6B). Similar expression trends were also observed in

known acid-induced, virulence-associated genes that are not encoded within SPI-2, such as

pagD (Gunn et al., 1995) (Figures 6C and S5B, C). This is consistent with the observation

above (Figure 2A) that SPI-2-deficient Stm retain some cytotoxicity. Importantly, the

expression of SPI-1 genes, which encode invasion-specific functions, was not altered in

infections with IFNβ priming or in *Ifnar2* KO cells (Figures 6D and S5D). Together, these data

indicate that IFN-I-mediated acidification of lysosomes promotes intracellular Stm virulence

gene expression.

The Stm virulence program can lead to the breakage of SCV, exposing the pathogen to the

host cytosol (Roy et al., 2004; Xu et al., 2019). To assess whether the pathogen was

cytosol-exposed, infected cells were treated with high concentrations of gentamicin, an

antibiotic that can penetrate into cells at high concentrations (Myrdal et al., 2005). Stm in

IFNβ-treated WT cells were markedly more sensitive to gentamicin than bacteria in

IFNβ-treated *Ifnar2* KO cells (Figure 6E), suggesting that IFN-I activation of Stm virulence

gene expression promotes SCV rupture and facilitates the pathogen's access to the cytosol.

Consistent with this idea, ~60% of Stm stained positive for galectin-3, a marker of SCV damage

(Thurston et al., 2012), in infected IFNβ-primed WT HeLa cells, whereas <20% of Stm were

galectin-3+ in infected *Ifnar2* KO cells (Figures 6F, G). Together, these data suggest a model

that explains why IFN-I signaling was a hit in the CRISPR/Cas9 screen: IFN-I

signaling-dependent lysosome acidification stimulates intracellular Stm virulence gene

expression, which promotes SCV rupture and subsequent Stm-induced cytotoxicity.

IFN-I promotes epithelial Stm pathogenesis *in vivo*

To understand the function of IFN-I signaling in Stm pathogenesis, we used a more physiologic

culture system - primary human-derived small intestinal organoids. IFNβ priming of organoids

increased cell death associated with Stm infection, whereas treatment of organoids with

pyridone-6 had the opposite effect (Figures 7A, B), supporting the idea that IFN-I signaling

promotes Stm pathogenicity in IECs.

To further dissect the importance of IFN-I signaling in the context of *in vivo* Stm infection,

we used bone marrow transfers to generate chimeric C57BL/6 mice that had *Ifnar1* deleted in

only the hematopoietic compartment or in other bodily tissues, including epithelial surfaces

(Figure S6A-C). Following intraperitoneal delivery of poly (I:C) to induce IFNβ production

(Lauterbach et al., 2010), chimeric mice were oro-gastrically inoculated with Stm to assess the

roles of epithelial and hematopoietic compartments in resistance to infection (Figure 7C).

Strikingly, mice with KO epithelium and WT bone marrow were relatively resistant to oral Stm

infection, with reduced weight loss and distal organ bacterial loads compared to mice that had

WT epithelium and bone marrow (Figures 7D, E), suggesting that IEC IFN-I signaling

enhances Stm pathogenicity during infection. We also observed a similar phenotype in mice

with WT epithelium and KO bone marrow (Figures 7D, E), consistent with previous

observations that immune cell IFN-I signaling also promotes Stm pathogenesis (Robinson et al.,

2012). Mice that had both KO epithelium and bone marrow were more protected than either

chimera (Figures 7D, E), further supporting the idea that Stm takes advantage of IFN-I

signaling in both the gut epithelium as well as in bone marrow-derived cells.

Although histological analyses revealed similar levels of tissue damage in both chimeras

(Figures 7F, G), finer-scale immunofluorescence studies with TUNEL staining to quantify cell

death showed that TUNEL+ (dying) cells tracked with the WT compartment. In chimeric mice

with WT epithelium, TUNEL staining primarily co-localized with E-cadherin-positive IECs

(Figures 7H, I). In contrast, in Stm-infected chimeric mice with WT bone marrow, cell death

was primarily localized to E-cadherin-negative cells in the lamina propria (Figures 7H, I).

Together, these *in vivo* studies suggest that IFN-I signaling in the epithelial compartment

facilitates Stm-induced IEC death and pathogen spread.

DISCUSSION

Our findings underscore the utility of model intracellular pathogens like Stm as probes for

the investigation of fundamental cell processes. The top hits in the genome-scale CRISPR/Cas9

screen that initiated this study were remarkably coherent and revealed that IFN-I signaling

sensitizes epithelial cells to Stm cytotoxicity. IFN-I-dependent lysosome acidification in IECs

stimulated Stm virulence gene expression, heightened SCV damage and exacerbated cell death,

offering a plausible molecular pathway that explains the results of the screen. Importantly, our

work uncovered a fundamental new role for IFN-I signaling. We discovered that this canonical

antiviral signaling pathway, which has been studied for more than 5 decades (Gonzalez-Navajas

et al., 2012), controls the subcellular localization, protein content, pH, and protease activity of

lysosomes. Several ISGs, including *Ifitm3*, *Slc15a3*, and *Cnp,* that were found to localize to

lysosomes, were shown to contribute to lysosomal acidification. Thus, IFN-I signaling controls

the function of an organelle – the lysosome – in addition to directly or indirectly modulating the

expression and activities of hundreds of ISGS and their interaction partners.

The functions of the three ISGs that were identified as participants in IFN-I-mediated

lysosomal acidification suggests that more than one mechanism accounts for this phenotype. It

seems likely that putative v-ATPase-associated proteins such as *Ifitm3* contribute to

acidification processes by directly modulating the proton concentration gradient. However,

ISGs with known non-ATPase-related functions such as *Slc15a3,* which is a proton-coupled

histidine and di-tripeptide transporter, may not play similar roles. While we cannot exclude

v-ATPase-mediated mechanisms for such proteins, we speculate that *Slc15a3* may show

transport preferences for non-neutral dipeptides that could influence lysosome luminal pH.

Previous studies have linked *Cnp* with not only lysosomal, but mitochondrial compartments

(McFerran and Burgoyne, 1997), raising the possibility that ISG function in additional cell

compartments could also indirectly contribute to lysosomal acidification. Interestingly, *Cnp,* a

membrane-bound protein, has additionally been linked to microtubule function, suggesting that

it may also play a role in IFN-I-mediated lysosomal repositioning (Bifulco et al., 2002).

Although a virtually universal antiviral immune signal, the consequences of IFN-I

signaling on bacterial pathogens has remained less clear and is in many cases detrimental to the

host (Kovarik et al., 2016). Several intracellular bacterial pathogens, including *Listeria*

monocytogenes, Mycobacterium tuberculosis and Stm*,* appear to have decreased virulence in

IFNAR1-deficient mice. Studies of the bases for these phenotypes have primarily focused on

immune-mediated explanations (Boxx and Cheng, 2016; Kernbauer et al., 2013; O'Connell et

al., 2004) (bring back previous citations). For Stm*,* we propose that IFN-I signaling contributes

to the outcome of infection at least in part by inducing remodeling of epithelial cell lysosomes

and thereby stimulating the Stm virulence program. Our findings suggest that IFN-I signaling

can modify innate defense in the epithelial as well as the immune compartment and is

complementary and compatible with previously proposed mechanisms for IFN-I-enhanced Stm

infection in bone-marrow derived immune cells. Such mechanisms include elevated

macrophage necroptosis (Hos et al., 2017; Robinson et al., 2012) and transcriptional

reprogramming (Perkins et al., 2015), altered dendritic cell homeostasis (Stefan et al., 2017),

and increased neutrophil-mediated inflammation (Wilson et al., 2019). The role of IFN-I

modulation of lysosome function to Stm infection in non-epithelial cells, such as macrophages,

requires further study. Furthermore, it remains an open question whether Stm purposely

stimulates and exploits IFN-I signaling as part of its pathogenic strategy.

Although IFN-I-induced lysosomal acidification sensitizes cells to an intracellular bacterial

pathogen, our finding that several known ISGs with antiviral properties, such as *Ifitm3*, *Slc15a3*

and *Cnp,* participate in this process leads us to speculate that this mechanism may be protective

against viral threats. IFN-I-mediated lysosomal remodeling may also play a role in

non-infectious pathologies, such as lysosomal cholesterol accumulation (Kuhnl et al., 2018) and

other lysosome-related disorders. It remains unclear whether these effects might be driven by

the tonic IFN-I signaling that occurs in many tissues (Schoggins et al., 2014), or instead require

pathogenic elevations of IFN-I.

Our finding that IFN-I signaling governs the composition and function of the lysosome

provides a new cell biologic perspective for understanding cytokine function. It will be of

interest to determine whether other immune signals (i.e. including other IFNs and cytokines)

can also direct remodeling of lysosomes and other organelles, such as the mitochondria and

endoplasmic reticulum, under homeostasis and in diverse pathogenic contexts. Such activities

may constitute a broadly applicable lens through which to view and enhance our understanding

of the cell biology of innate defense.

AUTHORCONTRIBUTIONS

H.L.Z. and M.K.W. conceived and all authors designed the study. H.L.Z., A.Z., and X.L. performed all experiments and analyzed data. H.L.Z., B.S., and M.K.W. wrote the manuscript and all authors edited the paper.

ACKNOWLEDGEMENTS

We thank members of the Waldor lab for helpful discussions on all aspects of this project, Dr. David Breault at the Harvard Digestive Diseases Center (HDDC) Organoid Core for the primary human small intestine organoids; and Michal Pyzik from Dr. Richard Blumberg's lab for assistance in creation of bone marrow chimeric mice.

Research in the M.K.W. laboratory is supported by HHMI and NIH grant R01 AI-042347. A.Z. was supported by an EMBO long-term fellowship (ALTF 1514-2016) and by a HHMI Fellowship of the Life Sciences Research Foundation.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Figure 1. A CRISPR/Cas9 screen identifies IEC factors required for Stm cytotoxicity

(A) Workflow for CRISPR/Cas9 Stm cytotoxicity screen in HT29-Cas9 cells.

(B) Adjusted p-values for selected enriched Gene Ontology (GO) terms from GO-analyzed hits

in the Stm cytotoxicity screen (upper threshold set at p < 1E-03).

(C) Cytoscape visualization of enriched pathways.

(D) Scatterplots showing normalized read enrichment of specific sgRNAs in two libraries (A

and B) after 4 rounds of Stm infection. Genes involved in IFN-I signaling are delineated by the

dashed red circle.

(E) Overview of IFN-I signaling pathway. Numbers correspond to hit ranks in each library.

Figure 2. IFN-I promotes Stm cytotoxicity in IECs

(A) Survival of IFNβ-primed or unprimed WT or *Ifnar2* KO HT29 cells 4 hours post WT or

mutant Stm infection. Mean ± s.d., n = 3.

(B) Survival of mock or drug-treated WT HT29 cells 20 hours post WT Stm infection. Mean ±

s.d., n = 3.

(C) Flow cytometry of IFNβ-primed or unprimed WT and *Ifnar2* KO HT29 cells 20 hours post

mCherry-Stm infection and stained with Annexin V-FITC. FITC, fluorescein isothiocyanate.

(D) Flow cytometric quantification of invasion of HT29 cells by mCherry-Stm. Mean ± s.d., n

= 3.

(E) Representative images of LAMP1-RFP-expressing HeLa cells 4 hours post GFP-Stm

infection. Boxed insets depict higher magnification showing bacterial colocalization with

LAMP1-RFP. Scale bar, 10μm.

(F) Quantification of LAMP1-RFP-positive Stm from 10 fields. Mean ± s.d., n = 3.

Statistical analysis was performed by two-tailed Student's *t* test (*P < 0.05, **P < 0.01 and

***P < 0.001).

See also Figure S1, 2 and Table S1.

Figure 3. IFN-I signaling regulates lysosomal positioning, acidity, and protease activity

(A) Representative images of lysosome (LAMP1-GFP+/LysoTracker+ compartment)

distribution in WT and *Ifnar2* KO HeLa cells with or without 16 hours of IFNβ stimulation.

Nuclei (blue) were stained with DAPI and actin (purple) was stained with phalloidin. Scale bar,

5μm.

(B) Quantification of perinuclear lysosome indices from 10 cells. Mean ± s.d., n = 3.

(C-D) Flow cytometry of LysoTracker Red (C) and DQ-Green BSA fluorescence (D) in HeLa

cells ±16 hours of treatment with IFNβ or the lysosomal acidification inhibitor Bfa1. Vertical

dashed lines indicate the mean fluorescence value of the mock control in WT (red) or *Ifnar2*

KO (blue) cells.

(E) Relative cathepsin D activity in WT and *Ifnar2* KO HeLa cells ± 16 hours of IFNβ

treatment. Mean ± s.d., n = 5.

Statistical analysis was performed by two-tailed Student's *t* test (***P < 0.001).

See also Figure S3.

Figure 4. The ISG IFITM3 regulates lysosomal function and Stm cytotoxicity

(A) Representative images of LAMP1-RFP or Rab5-RFP–expressing HeLa cells stained with

IFITM3 antibody (GFP). Nuclei (blue) were stained with DAPI. Scale bar, 5μm.

(B) Immunoblotting for IFITM3 in WT and *Ifitm3* KO HeLa cells ± 16 hours of IFNβ

treatment.

(C-D) Flow cytometry of LysoTracker Red (D) and DQ-Green BSA fluorescence (E) in WT,

Ifitm3 KO and *Ifnar2* KO HeLa cells ± 16 hours of IFNβ treatment.

(E) Survival of IFNβ-primed or unprimed WT, *Ifitm3* or *Ifnar2* KO HeLa cells 4 hours post Stm

infection. Mean ± s.d., n = 3.

Statistical analysis was performed by two-tailed Student's *t* test (*P < 0.05, **P < 0.01 and

***P < 0.001).

Figure 5. Discovery of ISGs with novel roles in lysosomal pH regulation

(A) Immunoblotting for known (LAMP1, CTSD) and suspected (IFITM3) lysosomal proteins

in whole-cell lysates (T) and purified lysosomes (IP).

(B) Relative fold change scatterplot of protein abundance in lysosomes purified from WT or

Ifnar2 KO HeLa cells ± 16 hours of IFNβ treatment. Colored dots indicate proteins that are

known ISGs.

(C) Quantification of mean fluorescence intensity from flow cytometry of Lysotracker staining

in WT or ISG KO cells ± 16 hours of IFNβ treatment.

Statistical analysis was performed by two-tailed Student's *t* test (*P < 0.05, **P < 0.01 and

***P < 0.001).

See also Figure S4 and Table S2.

Figure 6. IFN-I signaling promotes Stm virulence gene expression and SCV rupture

Relative induction of SPI-2 (*ssaV*) (A), PhoP-induced virulence gene (*pagD*) (C) and SPI-1

(*prgH*) (D) in intracellular Stm from WT and *Ifnar2* KO HeLa cells ± 16 hours of drug

treatment. Data are normalized to transcript levels in LB-cultured Stm (red). Mean ± s.d., n = 3.

(B) Flow cytometry of intracellular P$_{sifB}$*::gfp* Stm isolated from WT and *Ifnar2* KO HeLa cells

± 16 hours of drug treatment. LB-cultured Stm were used as the mock control.

(E) Intracellular CFU counts from IFNβ-treated WT and *Ifnar2* KO HeLa cells 2 hour post Stm

infection. Infected cells were treated with gentamicin (Gm) at the indicated concentrations

(μg/ml). Data were normalized to the WT+IFNβ Gm 10 group. Mean ± s.d., n = 5.

(F) Representative images of Gal3-GFP-expressing HeLa cells 4 hour post mCherry-Stm

infection. Scale bar, 10μm.

(G) Quantification of the Gal3 positive SCVs from 10 cells. Mean ± s.d., n = 3.

Statistical analysis was performed by two-tailed Student's *t* test (***P < 0.001).

See also Figure S5.

Figure 7. IFN-I signaling in intestinal epithelial cells promotes Stm pathogenesis.

(A) Representative images of IFNβ or pyridone-6-primed or unprimed human small bowel

enteroids 20 hours post WT Stm infection. Propidium iodide (PI) staining was used to detect

cell death. Scale bar, 100 μm.

(B) Enteroid survival 20 hours post WT Stm infection. Mean ± s.d., n = 3.

(C) Timeline of generation (top) and oral Stm infection (bottom) of *Ifnar1* chimeric mice.

(D) Body weights of Stm-infected chimeric mice. Mean ± s.e.m., n = 12 mice.

(E) Liver and spleen Stm CFU burdens from chimeric mice 5 days post-Stm infection. Mean ±

s.d., n = 12 mice.

(F) Representative H&E stained ileal sections from chimeric mice 5 days post-Stm infection.

Scale bars, 100μm.

(G) Average histological scores of chimeric mice 5 days post-Stm infection from 8 fields. Mean

± s.d., n = 4 mice.

(H) Representative images of ileal sections from chimeric mice 5 days post-Stm infection. IECs

were identified with E-cadherin (red), dying cells with TUNEL (green), and nuclei with DAPI

(blue). The white dashed line marks the epithelial surface. Scale bar, 100μm.

(I) Quantification of TUNEL+/E-cadherin+ (red) or TUNEL+/E-cadherin- (blue) cells per field

from 8 fields. Mean ± s.d., n = 4 mice.

Statistical analysis was performed by two-tailed Student's *t* test in (B) and (G). Statistical

analysis was performed by two-tailed Mann-Whitney U-test in (D) and (E). (*$P < 0.05$, **$P <$

0.01 and ***$P < 0.001$).

See also Figure S6.

Figure S1. IFN-I promotes intracellular Stm cytotoxicity, Related to Figure 2

(A) Relative expression of the IFN-I target gene *oas1* in WT and *Ifnar2* KO HT29 cells 16 hours post-IFNβ treatment. Mean ± s.d., n = 3.

(B) Survival of IFNβ-primed or unprimed WT or *Ifnar2* KO HT29 cells 20 hours post WT or mutant Stm infection. Mean ± s.d., n = 3.

(C) Representative bright-field images of WT and *Ifnar2* KO HT29 cells 2 days post Stm infection. Scale bar, 250 μm or 100 μm, respectively.

(D) Quantification of flow cytometry data in Figure 2C. Mean ± s.d., n = 4.

(E) Flow cytometry of IFNβ-primed or unprimed WT and *Ifnar2* KO HeLa cells 20 hours post mCherry-Stm infection and stained with Annexin V-FITC.

(F) Quantification of flow cytometry data from Figure S2E. Mean ± s.d., n = 4.

Statistical analysis was performed by two-tailed Student's *t* test (*P < 0.05, **P < 0.01 and ***P < 0.001).

Figure S2. IFN-I does not affect Stm invasion or SCV formation, Related to Figure 2

(A) Flow cytometry of IFNβ-primed WT and *Ifnar2* KO HT29 cells 4 hours post mCherry-Stm infection. Quantification is shown in Figure 2D.

(B) Representative images of Rab5-RFP-expressing HeLa cells at 4 hours post GFP-Stm infection. Scale bar, 10 μm.

(C) Quantification of Rab5-RFP-positive Stm from 10 fields. Mean ± s.d., n = 3.

(D) Representative images of Rab7-RFP-expressing HeLa cells 4 hours post GFP-Stm infection.

Scale bar, 10 μm.

(E) Quantification of Rab7-RFP-positive Stm from 10 fields. Mean ± s.d., n = 3.

Figure S3. IFN-I signaling regulates lysosomal remodeling in both epithelial cell and

THP1 cells, Related to Figure 3

(A) Quantification of mean fluorescence intensity (MFI) from Figure 3C. Mean ± s.d., n = 3.

(B) Flow cytometry of LysoSensor staining in WT and *Ifnar2* KO HeLa cells ± 16 hours of

IFNβ treatment.

(C) Flow cytometry of LysoTracker staining in HT29 cells ± 16 hours of drug treatment.

(D) Quantification of mean fluorescence intensity from Figure 3D. Mean ± s.d., n = 3.

(E) Flow cytometry of Dextran-568 uptake in WT and *Ifnar2* KO HeLa cells ± 16 hours of

IFNβ treatment.

(F) Flow cytometry of LysoTracker staining in monocytic macrophage-like THP1 cells ± 16

hours of IFNβ treatment.

Statistical analysis was performed by two-tailed Student's *t* test (**P < 0.01 and ***P < 0.001).

Figure S4. Purity of isolated lysosomes and IFITM3 gene KO in HeLa cells, Related to

Figure 5

Immunoblotting for protein markers of indicated subcellular compartments in whole-cell lysates

(T) and purified lysosomes (IP).

Figure S5. Intracellular Stm virulence gene expression, Related to Figure 6

(A-D) Relative induction of individual SPI-2 (A), PhoP-induced (B), SPI-3 (C) or SPI-1 (D)

genes in intracellular Stm from WT and *Ifnar2* KO HeLa cells ± drug treatment. Data are

normalized to transcript levels from LB-cultured Stm (red). Mean ± s.d., n = 3.

Statistical analysis was performed by two-tailed Student's *t* test (***P < 0.001).

Figure S6. Generation of chimeric mice by bone marrow transfer, Related to Figure 7

(A-C) Flow cytometry of peripheral blood CD45.1 and CD45.2+ cells in mock and chimeric

mice 4 weeks after bone marrow transplantation. WT mock mice carry CD45.2 allele but not

CD45.1 (A), which is abolished by irradiation (B, C). 4 weeks later after CD45.1 BM transfer,

the chimeric mice carry CD45.1 allele but not CD45.2 (B, C).

STAR METHODS

 ## KEY RESOURCE TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
LAMP1 (D4O1S) Mouse mAb	Cell Signaling Technology	15665S
Anti-IFIT3/P60 antibody [OTI1G1]	Abcam	ab118045
IFITM3 Antibody	Proteintech	11714-1-AP
Cathepsin D Monoclonal Antibody (CTD-19)	ThermoFisher	MA1-26773
Anti-GM130 antibody [EP892Y] - cis-Golgi Marker	Abcam	ab52649
β-Actin Antibody (AC-15)	Santa Cruz	sc-69879
PE/Cy7 anti-mouse CD45.2	Biolegend	109829
FITC anti-mouse CD45.1	Biolegend	110706
Human/Mouse E-Cadherin Antibody	R&D	AF748
Anti-Rabbit IgG (whole molecule)–Peroxidase antibody	Sigma	A4914
Goat anti-Mouse IgG (H+L) Secondary Antibody, HRP	ThermoFisher	31430
Rabbit anti-Goat IgG (H+L) Cross-Adsorbed Secondary Antibody, HRP	Invitrogen	R-21459
Donkey anti-Goat IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 568	Invitrogen	A-11057
Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 488	Invitrogen	A-11034
Bacterial Strains		
Salmonella Typhimurium SL1344	Dr. Dirk Bumann	
Salmonella Typhimurium SL1344 ΔprgH	This study	
Salmonella Typhimurium SL1344 ΔssaV	This study	
Salmonella Typhimurium SL1344-eGFP	This study	
Salmonella Typhimurium SL1344-mCherry	This study	
Salmonella Typhimurium SL1344-P$_{sifB}$-GFP	Dr. Dirk Bumann	
Salmonella Typhimurium SL1344-mCherry-P$_{sifB}$-GFP	This study	
One Shot Stbl3 Chemically Competent E. coli	Thermo Fisher Scientific	Cat No.C737303
Chemicals and Recombinant Proteins		
IFNβ	Peprotech	Cat No. 300-02BC
pyridine-6	BioVision	Cat No. 2534
Ruxolitinib (NCB018424)	Selleckchem	Cat No.S1378
Polybrene,	Sigma	Cat No. TR-1003-G
Trizol	Invitrogen	Cat No. 15596018

SuperScript III reverse transcriptase	Invitrogen	Cat No. 18080085
Roche 2×SYBR master mix	Roche	Cat No. 04707516001
BfA1	SantaCruz	Cat No. sc-201550
Critical Commercial Assays		
Blood and Cell Culture DNA MaxiKit	QIAGEN	Cat No. 13362
TransIT-LT1	Mirus	Cat No. MIR230
QIAquick Gel Extraction Kit	Qiagen	Cat No. 28704
LDH assay kit	Promega	Cat No. G1780
FITC Annexin V Apoptosis Detection kit	BioLegend	Cat No. 640922
Lipofectamine 3000	ThermoFisher	Cat No. L3000008
Lysotracker	ThermoFisher	Cat No. L7528
Lysosensor	Thermofisher	Cat No. L7545
fluorogenic peptide substrate of cathepsin D	Biovision	Cat No. K143
Dextran 568	ThermoFisher	Cat No. D22912
DQ-Red BSA	ThermoFisher	Cat No. D12050
10% Tris-Glycine gels	ThermoFisher	Cat No. XP00102BOX
nitrocellulose membranes	Invitrogen	Cat No. IB23002
SuperSignal West Pico Enhanced Chemiluminescence kit	ThermoFisher	Cat No. 34577
PureLink Micro-to-Midi total RNA purification system	Invitrogen	Cat No. 12183
Ambion Turbo DNA-free DNase	Invitrogen	Cat No. AM1907
TUNEL kit	ThermoFisher	Cat No. A23210
Experimental Models: Cell Lines		
HeLa	ATCC	CRM-CCL-2, female
HEK293T	ATCC	CRL-3216, female
HT29	ATCC	HTB-38, female
THP-1	ATCC	TIB-202, male
Primary human small intestine organoids	Harvard Digestive Diseases Center Organoid Core	Gift from Dr. David Breault
Experimental Models: Organisms/Strains		
Mouse: C57BL/6J B6(Cg)-Ifnar1tm1.2Ees/J	The Jackson Laboratory	Stock No: 028288
Mouse: B6.SJL-Ptprca Pepcb/BoyJ	The Jackson Laboratory	Stock No: 002014
Oligonucleotides		
qPCR primers, see Table S3		
CRISPR gene KO primers, see Table S4		

Recombinant DNA		
lentiGuide-Puro	Addgene	52963
psPAX2	Addgene	12260
pMD2.G	Addgene	12259
pLJC5-LAMP1-RFP-3xHA	Addgene	102932
pHR-FKBP:mCherry-Rab5a	Addgene	72901
pHR-FKBP:mCherry-Rab7a	Addgene	72903
LAMP1-mGFP	Addgene	34831
pLJC5-Tmem192-3xHA	Addgene	102930
mAG-GAL3	Addgene	62734
Software and Algorithms		
Primer3	Untergasser et al., 2012	http://primer3.ut.ee/
ImageJ	NIH	https://imagej.nih.gov/ij/download.html
FlowJo 10.2	FlowJo	https://www.flowjo.com/solutions/flowjo.
GraphPad Prism	GraphPad Software	https://www.graphpad.com.
Gene set enrichment Analysis	Broad Institute	http://www.broadinstitute.org/gsea.
DAVID analysis	NIAID/NIH	http://david.abcc.ncifcrf.gov

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Matthew K Waldor (mwaldor@research.bwh.harvard.edu)

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Bacterial strains, plasmids, and antibodies

Strains, plasmids, oligonucleotides and antibodies used in this study are listed in key resources

table and table S3, 4. *Escherichia coli* K-12 DH5α λ pir was used for cloning procedures and

plasmid propagation. *S. typhimurium* strain SL1344 and its ΔSPI-1 and ΔSPI-2 derivatives were

cultured in Luria-Bertani (LB) medium or on LB agar plates at 37°C supplemented with

streptomycin (100μg/ml). The SPI-1 (*prgH*) and SPI-2 (*ssaV*) genes were deleted from

wild-type (WT) SL1344 with the lambda red recombination system (Datsenko and Wanner,

2000). This approach was also used to introduce the GFP and mCherry-coding sequence with a

constitutive promoter (P_{rpsM}) into the *putP-putA* locus (Hautefort et al., 2003).

Cell lines

HeLa (ATCC, Cat No. CRM-CCL-2, female) and HEK293T (ATCC, Cat No. CRL-3216,

female) cells were cultured in Dulbecco's modified Eagle's medium (DMEM; ThermoFisher,

Cat No. 11965) supplemented with 10% fetal bovine serum (FBS; Gibco, Cat No. 16140-071).

HT29 (ATCC, Cat No. HTB-38, female) cells were cultured in McCoy's 5A modified medium

(Thermo Fisher, Cat No. 30-2007) supplemented with 10% FBS. THP-1 (ATCC, Cat No.

TIB-202, male) cells were cultured in RPMI-1640 medium (Lonza, Cat No. 12-167F) with 10%

non-heat inactivated FBS (GeminiBio, Cat NO. 100-500) and supplemented with HEPES

(Lonza, Cat No. 17-737E), 2-Mercaptoethanol (Invitrogen, Cat No. 21985023) and

L-Glutamine (Lonza, Cat No. 17-605E). All cells were cultured at 37°C in a 5% CO_2 incubator.

Infection of organoids derived from human small intestine

Primary human small intestine organoids (enteroids) were kindly provided by Dr. David Breault

at the Harvard Digestive Diseases Center (HDDC) Organoid Core. Enteroids were cultured in

the following medium: advanced DMEM/F12 (Gibco, Cat No.12634-028) supplemented with

L-WRN conditioned medium (ATCC CRL-3276; HDDC Organoids Core), HEPES (10mM, pH

7.4), GlutaMax (Gibco, Cat No.35050-061), B_{27} (Gibco, Cat No.12587010), N2 (Gibco, Cat

No.17502-048), 1mM N-acetyl-L-cysteine (Sigma, Cat No.A8199), 10mM nicotinamide

(Sigma, Cat No.N0636), 5µM A83-01 (Sigma, Cat No.SML0788), 10µM SB202190 (Sigma,

Cat No.S7067), 50ng/ml murine EGF (Peprotech, Cat No.315-09), 10nM gastrin (Sigma, Cat

No.G9145), and 10µM Y-27632 (Sigma, Cat No.Y0503). For Stm infection, approximately 100

enteroids were seeded in 50µl Matrigel (Corning, Cat No.356231) in each well of a 24-well

plate. Three to four days after seeding, enteroids were either mock treated, or primed with either

10ng/ml IFNβ or 0.5 µM pyridine-6 for 16 hours. Enteroids were then released from Matrigel

by incubation in 500µl Cell Recovery Solution (Corning, Cat No.354253) for 30 mins on ice.

Resuspended enteroids were pipetted up and down 50 times with a P1000 pipette and then

transferred to a new 24-well plate. Each well was infected with approximately 3×10^7 Stm. The

plate was centrifuged for 5 mins at 300× g before it was placed in a 37°Cincubator for 30 mins.

After infection, enteroids were transferred to microcentrifuge tubes, spundown, mixed with

50µl Matrigel per tube/sample, and seeded into a new 24-well plate. After Matrigel

solidification at 37°C, 500µl full enteroid media containing 50µg/ml gentamicin was added to

each well at 10ng/ml IFNβ or 0.5µM pyridine-6 was added to the corresponding primed

samples. At 20 hpi, propidium iodide (Invitrogen, Cat No.V13241) was used to stain the

enteroids before imaging. To quantitatively measure cell death, the media from each

well/sample was also assayed for LDH activity as described above. LDH release values of

mock treated samples were set at 1 for normalization.

Bone marrow chimera mice

C57BL/6 and *Ifnar1*[-/-] mice were purchased from The Jackson Laboratory (Bar Harbor, ME,

USA) and were maintained on a 12-hour light/dark cycle and a standard chow diet at the

Harvard Institute of Medicine specific pathogen-free (SPF) animal facility (Boston, MA, USA).

Animal experiments were performed according to guidelines from the Center for Animal

Resources and Comparative Medicine at Harvard Medical School. All protocols and

experimental plans were approved by the Brigham and Women's Hospital Institutional Animal

Care and Use Committee (Protocol #2016N000416). Littermate control male and female mice

were randomly assigned to each group and experiments were performed blinded with respect to

treatment. For bone marrow chimeras, recipient mice were irradiated two times with 600 rad

1□day before injection of bone marrow from WT or *Ifnar1*[-/-] mice. Bone marrow was extracted

from femurs of donor mice by flushing with PBS and then washed once in PBS; 1×10^6 cells

were injected into the tail vein of recipient mice. Mice were monitored for 4 weeks, at which

point engraftment was evaluated by flow cytometry.

Infection of chimeric mice

20μg poly (I:C) (Sigma, Cat No. P1530) was given intraperitoneally to chimeric mice one day

before Stm infection and every other day for a total of 3 doses to stimulate IFN production.

Food was withdrawn for 4 hours before infection. Stm inocula were prepared as described

above. Mice were infected orogastrically with 5×10^8 Stm in 100μl PBS. Food was returned

to the cages 2hpi. Infected mice were sacrificed 5 days after infection. Tissue samples of the

small intestine, spleen and liver were collected for histological analysis and enumeration of

colony-forming units (CFU). CFU were quantified by serial-dilution plating of homogenized

tissue samples on LB plates containing 100μg/ml streptomycin.

METHOD DETAILS

Pharmacologic inhibitors and IFNβ priming

JAK inhibitors pyridine-6 (BioVision, Cat No. 2534) and ruxolitinib (NCB018424)

(Selleckchem, Cat No.S1378) were used at 0.5μM. IFNβ (Peprotech, Cat No. 300-02BC) was

used at 10ng/ml for cell priming. Drug-treated cells were primed for 16 hours (unless otherwise

indicated) before Stm infection.

Stm infections

All tissue culture infections were done according to the following procedure unless otherwise

indicated. WT and mutant Stm were grown for ~16 hours at 37°C with shaking and then

sub-cultured (1:33) in LB without antibiotics for 3 hours until the cultures reached an OD_{600} of

0.8.To prepare the inoculum, cultures were first pelleted at 5,000× g for 5 min. The pellets were

resuspended in DMEM without FBS, and an appropriate volume of bacterial solution was

added to cells to reach a multiplicity of infection (MOI) of 100 bacteria per eukaryotic cell. The

cells were then incubated with bacteria for 30 min at 37°C with 5% CO_2. Extracellular bacteria

were removed by extensive washing with phosphate-buffered saline (PBS; Gibco, Cat No.

14190250) and addition of 50µg/ml gentamicin to the medium. At 2 hours post infection (hpi),

the gentamicin concentration was decreased to 5µg/ml.

CRISPR/Cas9 Stm infection screen

HT29-Cas9 CRISPR libraries were constructed as described previously (Blondel et al., 2016)

using the Avana sgRNA library, which contains four different sgRNAs targeting each human

protein-coding gene (Doench et al., 2016). For each library, two sets of four T225 flasks

(Corning, Cat No. 14-826-80) were seeded with 15×10^6 cells per flask and then incubated for

48 hours. At the time of the screen, there were ~150×10^6 cells per experimental condition,

corresponding to ~2,000× coverage per sgRNA. Cells were at ~70% confluence at the time of

infection. The infection was done as described above with minor modifications. Briefly, HT29

libraries were infected with WT Stm at an MOI of 300 for 30 min. After infection, the libraries

were expanded in McCoy's 5A + FBS containing 5µg/ml gentamicin, to both permit

intracellular bacterial cytotoxicity and minimizethe intracellular gentamicin concentration to

allow Stm invasion during the next round of infection. Flasks were checked daily to monitor

recovery of survivor cells; when 70% confluency was achieved, cells were trypsinized, pooled,

and reseeded for the next round of infection. In total, four rounds of infection were conducted.

Surviving cells from the last round of infection were used for preparation of genomic DNA.

Genomic DNA preparation, sequencing, and analyses of screen results

Genomic DNA was obtained from 75×10^6 cells after positive selection, as well as from input

cells, using the Blood and Cell Culture DNA MaxiKit (QIAGEN, Cat No. 13362). sgRNA

sequences was amplified by PCR as described (Doench et al., 2016). The read counts were first

normalized to reads per million within each condition by the following formula: reads per

sgRNA / total reads per condition $\times 10^6$. Reads per million were then $\log_2$-transformedby first

adding 1 to all values, in order calculate the log of sgRNAs with zero reads. The $\log_2$

fold-change of each sgRNA was then determined relative to the input sample for each

biological replicate. MAGeCK analysis for genome-scale CRISPR-Cas9 knockout screens was

used to evaluate the rank and statistical significance of perturbations from the ranked list (Li et

al., 2014) and enriched pathways were determined using ClueGo (Bindea et al., 2009).

Lentivirus preparation and transductions

The Galectin 3, Rab5, Rab7, LAMP1, and LC3B lentiviral expression plasmids used in the

study are listed in Table S4. Lentiviral packaging plasmids psPAX2 and pVSV-G, and the

corresponding cargo plasmid were transfected into 293T cells with the TransIT-LT1 transfection

reagent (Mirus, Cat No. MIR230). 48 hours following transfection, 293T culture supernatants

were harvested, passed through a 0.45µm pore filter, and added to target cells that were grown

to 70-80% confluency in 6-well plates. Polybrene (Sigma, Cat No. TR-1003-G) (8µg/ml) was

added and the 6-well plates were spun at 1000×g for 2 hours at 30°C, after which cells were

returned to 37°C. The infections were repeated the next day with supernatants from 72

hour-transfected 293T cultures.

Construction of cell lines with targeted gene disruptions

The sgRNA sequences used for construction of targeted HT29-Cas9 and HeLa-Cas9 mutant cell

lines are listed in Table S4. All sgRNA oligonucleotides were obtained from Integrated DNA

Technologies (IDT) and cloned into the pLentiGuide-Puro plasmid. Briefly, 5µg of plasmid

pLentiGuide-Puro was digested with BmsBI (Fermentas, Cat No. ER0451) and purified using

the QIAquick Gel Extraction Kit (Qiagen, Cat No. 28704). Each pair of oligos was annealed

and phosphorylated with T4 PNK (NEB, Cat No. M0201S) in the presence of 10× T4 DNA

ligase buffer in a thermocycler with the following parameters: i) incubation for 30 minutes at

37°C, ii) incubation at 95°C for 5 min with a ramp down to 25°C at 5°C per minute. Oligos

were then diluted 1:200, and 1µl of the diluted oligo mixture was ligated with 50ng of BsmBI

digested plasmid. Ligations were transformed into the STBL3 *Escherichia coli* strain (Thermo

Fisher, Cat No. C7373-03) and positive clones were identified by Sanger sequencing (Genewiz).

Lentiviral transduction of sgRNAs cloned into pLentiGuide-Puro into HT29-Cas9 and

HeLa-Cas9 cells was performed as described above. Targeted gene KO cell lines were selected

by puromycin (1µg/ml) for 10 days. HT29 KO cells were isolated as single clones while HeLa

cells were CRISPR KO pools after drug selection.

Cell survival assays

For cell survival assays, 5×10^4 HT29 cells were seeded into 96-well plates and primed with or without drugs in McCoy's 5Amedium supplemented with 10% FBS.HT29 cells were infected with Stms trains at an MOI of 100 as described above. Cell survival analysis was performed using an LDH assay (Promega, Cat No. G1780) according to the manufacturer's protocol at 4 and 20 hpi.

Stm invasion assays

mCherry-or GFP-tagged Stm were used in all flow cytometry and immunofluorescence experiments. Stm infections were performed as described above with varying MOIs. At 4hpi, suspended and attached cells were collected, resuspended in PBS, and immediately analyzed with a LSR II (BD Bioscience) or SH800 (Sony) flow cytometer. Data were processed with FlowJo software (v10.6.1).

Annexin V staining and FACS analysis

Cell death was detected with the FITC Annexin V Apoptosis Detection kit (BioLegend, Cat No. 640922). Infections were performed as described above with mCherry-Stm at an MOI of 100. 20 hpi suspended and attached cells were collected, resuspended in 100μl of Annexin V binding buffer at 1×10^7 cells/ml and mixed with 5μl of FITC-conjugated Annexin V. After incubation at room temperature (RT) for 15□min in the dark, 400μl of Annexin V binding buffer was added and stained cells were immediately analyzed by flow cytometry as described above.

Immunofluorescence microscopy of tissue cultured cells

HeLa cells were seeded in 12-well plates on 18 mm glass coverslips or 6-well chambers (Mat-TEK, Cat No.P06G-1.5-10-F). Cells were transiently transfected with LAMP1-GFP expressing plasmid mixed with Lipofectamine 3000 (ThermoFisher, Cat No. L3000008) according to the manufacturer's instructions. 24 hours post-transfection cells were primed with or without 10ng/ml IFNβ for 16 hours.The cells were then stained with 75nM Lysotracker (ThermoFisher, Cat No. L7528) for 15 min and then fixed with 2% PFA for 20 min at RT. The samples were washed with PBS three times, and stained with fluorescent phalloidin (1:1000) and 4,6-diamidino-2-phenylindole (DAPI, 1 μg /ml) to label actin filaments and nuclei, respectively. For experiments with LAMP1-RFP, Rab5-RFP, Rab7-RFP, Gal3-GFP, and eGFP-LC3B cell lines, cells were seeded in 6-well chambers and primed with 10ng/ml IFNβ for 16 hours before infection with fluorescently-labeled Stm at an MOI 50. Live cells were analyzed at 2 hpi by confocal microscopy to detect localization of Gal3 and Stm.

Quantification of lysosome distribution

Lysosome distribution was analyzed as described (Li et al., 2016); the area occupied by nuclei was excluded from analyses. Average LAMP1 intensities were measured for the area within 5μm of the nucleus ($I_{\text{perinuclear}}$), and the area >10μm from the nucleus ($I_{\text{peripheral}}$). The average intensities were calculated and normalized to cell areas. The perinuclear index was defined as $I_{\text{perinuclear}}$ /$I_{\text{peripheral}}$. Quantifications were carried out on10 cells per group with ImageJ.

Measurement of lysosome acidity

Cells with no treatment or with either 10ng/ml IFNβ or 5nM BfA1 (SantaCruz, Cat No. sc-201550) treatment for 16 hours were stained with 75nM Lysotracker or Lysosensor (Thermofisher, Cat No. L7545) for 15 min and washed with PBS. The fluorescence intensity of the stained cells was determined by flow cytometry.

Cathepsin D activity assay

HeLa cells were seeded in 96-well plates with or without 10ng/ml IFNβ priming for 16 hours. A fluorogenic peptide substrate of cathepsin D, Mca-P-L-G-L-Dpa-A-R-NH2 (Biovision, Cat No. K143), was added to the cells to a final concentration of 200μM for 2 hours. The fluorescence intensity of each well was measured with a fluorescence plate reader. Each sample was assayed in triplicate and normalized to a standard curve.

Endocytosis and lysosome function assays

HeLa cells were seeded in 24-well plates with or without 10ng/ml IFNβ priming. Cells were treated with either 50μg/ml Dextran 568 (ThermoFisher, Cat No. D22912) or 25μg/ml DQ-Red BSA (ThermoFisher, Cat No. D12050) for 2 hours in growth medium. Then, cells were washed with PBS and trypsinized for fluorescence quantification by flow cytometry.

Lysosome immunopurification (LysoIP)

LysoIP was performed largely as described (Abu-Remaileh et al., 2017). Briefly,

pLJC5-3×HA-TMEM192 was used to introduce a lysosomal tag protein in WT and *Ifnar2* KO

HeLa cells. 15 million cells were used for each replicate. Cells were rinsed twice with

pre-chilled PBS and then scraped in 1ml of PBS containing protease and phosphatase inhibitors

and pelleted at 100×g for 2 min at 4°C. Cells were resuspended in 950µl of the same buffer, and

25µl (equivalent to 2.5% of the total number cells) was reserved for further processing to

generate the whole-cell sample. The remaining cells were gently homogenized with 25 strokes

of a 2ml Dounce-type homogenizer. The homogenate was then centrifuged at 100×g for 2 min

at 4°C to pellet the cell debris and intact cells, while cellular organelles including lysosomes

remained in the supernatant. The supernatant was incubated with 150µl of anti-HA magnetic

beads preequilibrated with PBS on a rotator shaker for 3 min. Immunoprecipitates were then

gently washed three times with PBS on a DynaMag Spin Magnet. Beads with bound lysosomes

were resuspended in 100µl pre-chilled 1% Triton-X lysis buffer to extract proteins. After 10

min incubation on ice, the beads were removed with the magnet. 5µl of each sample were

subjected to 12.5%-acrylamide SDS-PAGE and immunodetected using antibody listed in Table

S6, while the remainder was submitted to the Thermo Fisher Center for Multiplexed Proteomics

of Harvard Medical School (Boston, MA, USA) for Isobaric Tandem Mass Tag (TMT)-based

quantitative proteomics.

Immunoblot analyses

Mammalian cell lysates were prepared in radioimmuno-precipitation assay (RIPA) buffer

supplemented with 1 tablet of EDTA-free protease inhibitor (Roche, Cat No. C762Q77) per

25ml buffer. Lysates were kept at 4°C for 30 min and then clarified by centrifugation in a

microcentrifuge at 13,000 rpm at 4°C for 10 min. Proteins were denatured by the addition of

SDS sample buffer and boiling for 5 min. Proteins were separated by electrophoresis in 10%

Tris-Glycine gels (ThermoFisher, Cat No. XP00102BOX), and then transferred onto

nitrocellulose membranes (Invitrogen, Cat No. IB23002). The antibodies and dilutions used are

listed in Table S6. Blots were developed with the SuperSignal West Pico Enhanced

Chemiluminescence kit (ThermoFisher, Cat No. 34577), and imaged with a Chemidoc

(Bio-Rad).

qRT-PCR quantification of Stm virulence gene expression

Hela cells were seeded at 1.5×10^6 cells per 6-well plates. After drug-treatment for 16 hours,

cells were infected with Stm at an MOI of 50 as described above. Cells were washed with PBS

and lysed in Trizol (Invitrogen, Cat No. 15596018) at 1and 4 hpi. RNA was purified with the

PureLink Micro-to-Midi total RNA purification system (Invitrogen, Cat No. 12183) according

to the manufacturer's instructions. RNA samples were treated for residual DNA contamination

using Ambion Turbo DNA-free DNase (Invitrogen, Cat No. AM1907). Purified RNA was

quantified on a Nanodrop 1000 (Thermo Scientific). RNA was reverse transcribed for

quantitative RT-PCR (qRT-PCR) experiments by adding 10µg of total RNA to a mixture

containing random hexamers (Life Technologies), 0.01M dithiothreitol, 25 mM dNTPs

(Thermo Scientific, Cat No. R0191), reaction buffer and 200 units of SuperScript III reverse

transcriptase (Invitrogen, Cat No. 18080085). cDNA was diluted 1:50 in dH$_2$O and mixed with

an equal volume of target-specific primers and Roche 2×SYBR master mix (Roche, Cat

No.04707516001). Plates were centrifuged at 1000 rpm for 1 min and stored at 4°C in the dark

until ready for use. Primer pairs were designed to minimize secondary structures, a length of

~20-nucelotides and a melting temperature of 60°C using the primer design software Primer 3.

Primer sequences are listed in Table S3. For data normalization, quadruplicate C$_t$ values for

each sample were averaged and normalized to C$_t$ values of the control gene *rpoB*. The relative

gene expression level of Stm in infection conditions was normalized to LB-cultured Stm.

Flow cytometric analysis of Stm virulence gene expression

HeLa cells were infected with mCherry-and *sifB*-GFP-expressing-Stm as described above. Cell

lysis was performed 4 hpi by washing three times with PBS and subsequent incubation for 10

min with PBS containing 0.1% Triton X-100. Cell lysates were then analyzed by flow

cytometry. Stm were first identified by gating on the mCherry signal and *sifB* expression was

quantified by gating on the mCherry+/GFP+ population. LB cultured Stm served as negative

control.

Gentamicin protection assay

Gentamicin protection assays were carried out as described (Knodler et al., 2014). Briefly,

HeLa cells in 96-well plates were infected in triplicate with Stm at an MOI of 50. Cells were

washed three times with PBS and incubated in medium containing 100µg/ml gentamicin for 30

min to eliminate extracellular bacteria. Then, media with either 10, 100 or 400µg/ml gentamicin

was applied to the cells. Cells were lysed 2hpi by washing three times with PBS and subsequent

incubation for 10 min with PBS containing 0.1 % Triton X-100. Colony forming units (CFUs)

were enumerated by plating serial dilutions of the lysates onto LB plates with 100µg/ml

streptomycin. Data was normalized to the CFU of WT HeLa cells at gentamicin 10.

Histology and tissue immunofluorescence

Formalin-fixed, paraffin-embedded distal small intestinal samples sections of 4µm thickness

were mounted on glass slides and stained with hematoxylin and eosin. Histology score was

evaluated as described (Erben et al., 2014). For immunofluorescence analysis, distal small

intestinal samples were collected and flushed with PBS and fixed in 4% paraformaldehyde

(PFA) overnight at 4°C followed by washing with PBS. Tissues were embedded in Optimal

Cutting Temperature Compound (Tissue-Tek) and stored at -80°C before sectioning on a

CM1860 UV cryostat (Leica). 6µm-thick slides were stained with TUNEL (ThermoFisher, Cat

No. A23210) according to the manufacturer's instructions and then incubated with

anti-E-cadherin antibodies at 4°C overnight at a 1:200 in PBS. The next day, AF568-conjugated

secondary antibody, diluted at 1:500, was applied to the slides for 1 hour. Nuclei were stained

with DAPI at RT for 5 min in the dark. Samples were imaged with an Eclipse Ti confocal

microscope with a 20×□objective (Nikon).

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analyses were carried out using the two-tailed Student's *t* test or one-way analysis of

variance (ANOVA) with Dunnet's post-correction on GraphPad Prism5.

DATA AND CODE AVAILABILITY

Original data for results of CRISPR screening is in Table S1, and original data for mass

spectrometry of lysosome proteomic is in Table S2.

Supplementary items

Table S1: CRISPR/Cas9 screening results, related to figure 1

Table S2: Mass spectrometry of lysosome proteomic, related to figure 3

Table S3: qPCR primers

Table S4: CRISPR KO primers

REFERENCES

Abu-Remaileh, M., Wyant, G.A., Kim, C., Laqtom, N.N., Abbasi, M., Chan, S.H., Freinkman, E., and Sabatini, D.M. (2017). Lysosomal metabolomics reveals V-ATPase- and mTOR-dependent regulation of amino acid efflux from lysosomes. Science *358*, 807-813.

Bifulco, M., Laezza, C., Stingo, S., and Wolff, J. (2002). 2',3'-Cyclic nucleotide 3'-phosphodiesterase: a membrane-bound, microtubule-associated protein and membrane anchor for tubulin. Proc Natl Acad Sci U S A *99*, 1807-1812.

Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., Fridman, W.H., Pages, F., Trajanoski, Z., and Galon, J. (2009). ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. Bioinformatics *25*, 1091-1093.

Blondel, C.J., Park, J.S., Hubbard, T.P., Pacheco, A.R., Kuehl, C.J., Walsh, M.J., Davis, B.M., Gewurz, B.E., Doench, J.G., and Waldor, M.K. (2016). CRISPR/Cas9 Screens Reveal Requirements for Host Cell Sulfation and Fucosylation in Bacterial Type III Secretion System-Mediated Cytotoxicity. Cell host & microbe *20*, 226-237.

Boxx, G.M., and Cheng, G. (2016). The Roles of Type I Interferon in Bacterial Infection. Cell host & microbe *19*, 760-769.

Butor, C., Griffiths, G., Aronson, N.N., Jr., and Varki, A. (1995). Co-localization of hydrolytic enzymes with widely disparate pH optima: implications for the regulation of lysosomal pH. J Cell Sci *108 (Pt 6)*, 2213-2219.

Chakraborty, S., Mizusaki, H., and Kenney, L.J. (2015). A FRET-based DNA biosensor tracks OmpR-dependent acidification of Salmonella during macrophage infection. PLoS biology *13*, e1002116.

Datsenko, K.A., and Wanner, B.L. (2000). One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A *97*, 6640-6645.

Desjardins, M., Celis, J.E., van Meer, G., Dieplinger, H., Jahraus, A., Griffiths, G., and Huber, L.A. (1994). Molecular characterization of phagosomes. J Biol Chem *269*, 32194-32200.

Doench, J.G., Fusi, N., Sullender, M., Hegde, M., Vaimberg, E.W., Donovan, K.F., Smith, I., Tothova, Z., Wilen, C., Orchard, R.*, et al.* (2016). Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. Nat Biotechnol *34*, 184-191.

Erben, U., Loddenkemper, C., Doerfel, K., Spieckermann, S., Haller, D., Heimesaat, M.M., Zeitz, M., Siegmund, B., and Kuhl, A.A. (2014). A guide to histomorphological evaluation of intestinal inflammation in mouse models. Int J Clin Exp Pathol *7*, 4557-4576.

Galan, J.E., Lara-Tejero, M., Marlovits, T.C., and Wagner, S. (2014). Bacterial type III secretion systems: specialized nanomachines for protein delivery into target cells. Annual review of microbiology *68*, 415-438.

Garmendia, J., Beuzon, C.R., Ruiz-Albert, J., and Holden, D.W. (2003). The roles of SsrA-SsrB and OmpR-EnvZ in the regulation of genes encoding the Salmonella typhimurium SPI-2 type III secretion system. Microbiology *149*, 2385-2396.

Geoghegan, J.L., and Holmes, E.C. (2018). The phylogenomics of evolving virus virulence. Nature reviews Genetics *19*, 756-769.

Gonzalez-Navajas, J.M., Lee, J., David, M., and Raz, E. (2012). Immunomodulatory functions of type I interferons. Nature reviews Immunology *12*, 125-135.

Gunn, J.S., Alpuche-Aranda, C.M., Loomis, W.P., Belden, W.J., and Miller, S.I. (1995). Characterization of the Salmonella typhimurium pagC/pagD chromosomal region. J Bacteriol *177*, 5040-5047.

Hautefort, I., Proenca, M.J., and Hinton, J.C. (2003). Single-copy green fluorescent protein gene fusions allow
accurate measurement of Salmonella gene expression in vitro and during infection of mammalian cells. Appl
Environ Microbiol *69*, 7480-7491.

Hess, C.B., Niesel, D.W., and Klimpel, G.R. (1989). The induction of interferon production in fibroblasts by invasive
bacteria: a comparison of Salmonella and Shigella species. Microb Pathog *7*, 111-120.

Hos, N.J., Ganesan, R., Gutierrez, S., Hos, D., Klimek, J., Abdullah, Z., Kronke, M., and Robinson, N. (2017). Type I
interferon enhances necroptosis of Salmonella Typhimurium-infected macrophages by impairing antioxidative
stress responses. J Cell Biol *216*, 4107-4121.

Hubel, P., Urban, C., Bergant, V., Schneider, W.M., Knauer, B., Stukalov, A., Scaturro, P., Mann, A., Brunotte, L.,
Hoffmann, H.H.*, et al.* (2019). A protein-interaction network of interferon-stimulated genes extends the innate
immune system landscape. Nature immunology *20*, 493-502.

Hurley, D., McCusker, M.P., Fanning, S., and Martins, M. (2014). Salmonella-host interactions - modulation of the
host innate immune system. Front Immunol *5*, 481.

Hybiske, K., and Stephens, R.S. (2008). Exit strategies of intracellular pathogens. Nat Rev Microbiol *6*, 99-110.

Jo, E.K. (2019). Interplay between host and pathogen: immune defense and beyond. Experimental & molecular
medicine *51*, 1-3.

Kernbauer, E., Maier, V., Rauch, I., Muller, M., and Decker, T. (2013). Route of Infection Determines the Impact of
Type I Interferons on Innate Immunity to Listeria monocytogenes. PLoS One *8*, e65007.

Knodler, L.A., Nair, V., and Steele-Mortimer, O. (2014). Quantitative assessment of cytosolic Salmonella in
epithelial cells. PLoS One *9*, e84681.

Kovarik, P., Castiglia, V., Ivin, M., and Ebner, F. (2016). Type I Interferons in Bacterial Infections: A Balancing Act.
Front Immunol *7*, 652.

Kuhnl, A., Musiol, A., Heitzig, N., Johnson, D.E., Ehrhardt, C., Grewal, T., Gerke, V., Ludwig, S., and Rescher, U.
(2018). Late Endosomal/Lysosomal Cholesterol Accumulation Is a Host Cell-Protective Mechanism Inhibiting
Endosomal Escape of Influenza A Virus. mBio *9*.

Lauterbach, H., Bathke, B., Gilles, S., Traidl-Hoffmann, C., Luber, C.A., Fejer, G., Freudenberg, M.A., Davey, G.M.,
Vremec, D., Kallies, A.*, et al.* (2010). Mouse CD8alpha+ DCs and human BDCA3+ DCs are major producers of
IFN-lambda in response to poly IC. J Exp Med *207*, 2703-2717.

Lee, H.C., Chathuranga, K., and Lee, J.S. (2019). Intracellular sensing of viral genomes and viral evasion.
Experimental & molecular medicine *51*, 1-13.

Li, W., Xu, H., Xiao, T., Cong, L., Love, M.I., Zhang, F., Irizarry, R.A., Liu, J.S., Brown, M., and Liu, X.S. (2014).
MAGeCK enables robust identification of essential genes from genome-scale CRISPR/Cas9 knockout screens.
Genome Biol *15*, 554.

Li, X., Rydzewski, N., Hider, A., Zhang, X., Yang, J., Wang, W., Gao, Q., Cheng, X., and Xu, H. (2016). A molecular
mechanism to regulate lysosome motility for lysosome positioning and tubulation. Nat Cell Biol *18*, 404-417.

McFerran, B., and Burgoyne, R. (1997). 2',3'-Cyclic nucleotide 3'-phosphodiesterase is associated with
mitochondria in diverse adrenal cell types. J Cell Sci *110 (Pt 23)*, 2979-2985.

Myrdal, S.E., Johnson, K.C., and Steyger, P.S. (2005). Cytoplasmic and intra-nuclear binding of gentamicin does not
require endocytosis. Hear Res *204*, 156-169.

O'Connell, R.M., Saha, S.K., Vaidya, S.A., Bruhn, K.W., Miranda, G.A., Zarnegar, B., Perry, A.K., Nguyen, B.O., Lane,
T.F., Taniguchi, T.*, et al.* (2004). Type I interferon production enhances susceptibility to Listeria monocytogenes

infection. J Exp Med *200*, 437-445.

Omotade, T.O., and Roy, C.R. (2019). Manipulation of Host Cell Organelles by Intracellular Pathogens.
Microbiology spectrum *7*.

Perkins, D.J., Rajaiah, R., Tennant, S.M., Ramachandran, G., Higginson, E.E., Dyson, T.N., and Vogel, S.N. (2015).
Salmonella Typhimurium Co-Opts the Host Type I IFN System To Restrict Macrophage Innate Immune
Transcriptional Responses Selectively. Journal of immunology *195*, 2461-2471.

Prost, L.R., Daley, M.E., Le Sage, V., Bader, M.W., Le Moual, H., Klevit, R.E., and Miller, S.I. (2007). Activation of the
bacterial sensor kinase PhoQ by acidic pH. Mol Cell *26*, 165-174.

Reis, R.C., Sorgine, M.H., and Coelho-Sampaio, T. (1998). A novel methodology for the investigation of intracellular
proteolytic processing in intact cells. Eur J Cell Biol *75*, 192-197.

Ribet, D., and Cossart, P. (2015). How bacterial pathogens colonize their hosts and invade deeper tissues.
Microbes and infection *17*, 173-183.

Robinson, N., McComb, S., Mulligan, R., Dudani, R., Krishnan, L., and Sad, S. (2012). Type I interferon induces
necroptosis in macrophages during infection with Salmonella enterica serovar Typhimurium. Nature immunology
13, 954-962.

Roy, D., Liston, D.R., Idone, V.J., Di, A., Nelson, D.J., Pujol, C., Bliska, J.B., Chakrabarti, S., and Andrews, N.W. (2004).
A process for controlling intracellular bacterial infections induced by membrane injury. Science *304*, 1515-1518.

Schoggins, J.W., MacDuff, D.A., Imanaka, N., Gainey, M.D., Shrestha, B., Eitson, J.L., Mar, K.B., Richardson, R.B.,
Ratushny, A.V., Litvak, V., *et al.* (2014). Pan-viral specificity of IFN-induced genes reveals new roles for cGAS in
innate immunity. Nature *505*, 691-695.

Schoggins, J.W., and Rice, C.M. (2011). Interferon-stimulated genes and their antiviral effector functions. Current
opinion in virology *1*, 519-525.

Song, F., Yi, Y., Li, C., Hu, Y., Wang, J., Smith, D.E., and Jiang, H. (2018). Regulation and biological role of the
peptide/histidine transporter SLC15A3 in Toll-like receptor-mediated inflammatory responses in macrophage. Cell
death & disease *9*, 770.

Spence, J.S., He, R., Hoffmann, H.H., Das, T., Thinon, E., Rice, C.M., Peng, T., Chandran, K., and Hang, H.C. (2019).
IFITM3 directly engages and shuttles incoming virus particles to lysosomes. Nature chemical biology *15*, 259-268.

Steele-Mortimer, O. (2008). The Salmonella-containing vacuole: moving with the times. Curr Opin Microbiol *11*,
38-45.

Stefan, K.L., Fink, A., Surana, N.K., Kasper, D.L., and Dasgupta, S. (2017). Type I interferon signaling restrains
IL-10R+ colonic macrophages and dendritic cells and leads to more severe Salmonella colitis. PLoS One *12*,
e0188600.

Thurston, T.L., Wandel, M.P., von Muhlinen, N., Foeglein, A., and Randow, F. (2012). Galectin 8 targets damaged
vesicles for autophagy to defend cells against bacterial invasion. Nature *482*, 414-418.

Tuli, A., and Sharma, M. (2019). How to do business with lysosomes: Salmonella leads the way. Curr Opin
Microbiol *47*, 1-7.

Unsworth, K.E., Way, M., McNiven, M., Machesky, L., and Holden, D.W. (2004). Analysis of the mechanisms of
Salmonella-induced actin assembly during invasion of host cells and intracellular replication. Cell Microbiol *6*,
1041-1055.

van der Velden, A.W., Lindgren, S.W., Worley, M.J., and Heffron, F. (2000). Salmonella pathogenicity island
1-independent induction of apoptosis in infected macrophages by Salmonella enterica serotype typhimurium.

Infection and immunity *68*, 5702-5709.

Wee, Y.S., Roundy, K.M., Weis, J.J., and Weis, J.H. (2012). Interferon-inducible transmembrane proteins of the

innate immune response act as membrane organizers by influencing clathrin and v-ATPase localization and

function. Innate Immun *18*, 834-845.

Welch, M.D. (2015). Why should cell biologists study microbial pathogens? Molecular biology of the cell *26*,

4295-4301.

Wilson, R.P., Tursi, S.A., Rapsinski, G.J., Medeiros, N.J., Le, L.S., Kotredes, K.P., Patel, S., Liverani, E., Sun, S., Zhu, W.,

et al. (2019). STAT2 dependent Type I Interferon response promotes dysbiosis and luminal expansion of the

enteric pathogen Salmonella Typhimurium. PLoS Pathog *15*, e1007745.

Xu, Y., Zhou, P., Cheng, S., Lu, Q., Nowak, K., Hopp, A.K., Li, L., Shi, X., Zhou, Z., Gao, W., *et al.* (2019). A Bacterial

Effector Reveals the V-ATPase-ATG16L1 Axis that Initiates Xenophagy. Cell *178*, 552-566 e520.

Yeung, A.T.Y., Choi, Y.H., Lee, A.H.Y., Hale, C., Ponstingl, H., Pickard, D., Goulding, D., Thomas, M., Gill, E., Kim, J.K.,

et al. (2019). A Genome-Wide Knockout Screen in Human Macrophages Identified Host Factors Modulating

Salmonella Infection. mBio *10*.

Yoshimori, T., Yamamoto, A., Moriyama, Y., Futai, M., and Tashiro, Y. (1991). Bafilomycin A1, a specific inhibitor of

vacuolar-type H(+)-ATPase, inhibits acidification and protein degradation in lysosomes of cultured cells. J Biol

Chem *266*, 17707-17712.

Figure. 1

Figure. 2

Figure. 3

Figure. 4

Figure. 5

Figure. 6

Figure. 7