Article type: Research article

Title: Analysis of healthy reference ranges for clinically relevant gut microbial indicators: A large web-based cohort study in Japan

Authors

Satoshi Watanabe[1*], Shoichiro Kameoka[1], Natsuko O. Shinozaki[1], Ryuichi Kubo[1], Akifumi Nishida[1], Minoru Kuriyama[1], Aya K. Takeda[1]

Affiliation

[1]Cykinso Inc. Tokyo, Japan.

Corresponding author:

*Satoshi Watanabe, PhD

Cykinso Inc. Tokyo, Japan.

1-36-1 Yoyogi, Shibuya, Tokyo 151-0053, Japan

Tel. & FAX: +81-3-5309-2522

E-mail: s.watanabe@cykinso.co.jp

Short title: Japanese reference ranges for gut microbiome indices

Word count: 6534 words

Conflicts of interest: The authors declare that no competing interests exist.

Grant support: None

Financial Disclosure: The authors disclose no financial arrangement.

Author contributions to manuscript: SW and AKT: study concept and design, analysis and interpretation of data, drafting of the manuscript, critical revision of the manuscript for important intellectual content, and statistical analysis. SK, NOS, and RK: interpretation of data and drafting of the manuscript. AN and MK: statistical analysis, technical support. All authors read and approved the final manuscript.

ABSTRACT

Background: Although several large cohort studies have focused on the human gut microbiome, large cohort study of the Japanese gut microbiome is scarce, especially that of healthy or non-diseased individuals. The purpose of this study was to establish a reference range for gut microbial indices by collecting Japanese real-world microbiome data from the web-based cohort.

Methods: We collected stool samples and original survey on lifestyle from 5,843 Japanese people through web-based services. From the obtained 16S rRNA sequence data derived from the stool sample, the ratio and distribution of each taxa were analyzed. The relationship between different epidemiological attributes and gut microbial indicators were statistically tested.

Results: In total, 100 taxa (genus level) of gut microbiota were shared by more than 10% of Japanese people, and 20 taxa (genus level) by 90% or more. Bacteroides-dominant type was the most popular entero-type. The qualitative and quantitative indicators of these common gut microbiota were confirmed to be strongly correlated with age, constipation / diarrhea, and a history of lifestyle-related diseases. Therefore, we set up the healthy sub-cohort that controlled for these attribute factors, and defined a reference range from the distribution of gut microbial index in that population.

Conclusions: The gut microbiota of Japanese people had high beta diversity, and there was no single "typical" gut microbiota type. We believe that the reference range of the gut microbial index obtained in this study can be a new reference value for determining the balance and health of the gut microbiota of an individual. In the future, it is necessary to clarify the clinical validity of this reference value by comparing with the clinical disease cohort.

Word count: 272 words

Keywords: Gut microbiota, 16S rRNA, Japanese, reference range, microbial index

INTRODUCTION

Since the establishment of next-generation 16S rRNA metagenome sequencing analysis, multiple large cohort studies focused on the human gut microbiome have been conducted, such as the US Human Microbiome Project [1] and MetaHIT in Europe [2]. An integrated catalog of human fecal microbial metagenomes from 1200 people in the United States, China, and Europe has identified 9.9 million microbial genes across fecal microbiota [3]. However, Japanese gut microbiome study is scarce, especially that of healthy or non-diseased individuals. Further, in recent studies comparing gut microbiomes by race/nationality, clear impacts of dietary habit were demonstrated [4], suggesting that, due to Japanese unique food culture, the gut of Japanese individuals could harbor different flora from individuals in western countries. Therefore, to advance research on the gut microbiome and various diseases in Japan, it is critical to characterize the healthy gut microbiome in the Japanese population.

Host parameters such as age, gender, and body mass index (BMI) have been reported to be related to individual differences in gut microbiota composition [5][6][7][8][9]. Further, differences in dietary habits have been shown to affect the bacterial diversity and enterotype of human gut microbiota [10][11], which may partially explain why differences in residential areas/country are strongly associated with differences in gut microbiota composition [12][13].

Recently, several studies have revealed gender differences in gut microbiota [14][15][16][17]. Min et al [18] conducted an association study to identify bacterial compositions associated with men and women and showed similar microbiota characteristics, including overall abundance and diversity, between men and women. However, they also showed gender differences at the species level between microbial taxa related to fat distribution, suggesting the existence of a gender-specific microbiome signature corresponding to gender-specific fat distribution, which may also contribute to the observed sex-specific immunity differences [19]. Thus, several immune pathophysiology may be involved in gender differences in gut bacterial composition.

Age is also an important factor affecting the gut microbiota [8][20][21][22]. Recent reports have described differences in gut microbiota between children and adults, and an adult-like composition of bacterial communities is established around 3–4 years old or more [7] [8][23][24][25]. In addition, intestinal microbiota have been shown to change with aging, although the definition of old people differs by report, such as over 60, 65, 70, or 100 [14][26][27][28]. However, the mechanism of change with aging is still unknown. Yatsunenko et al. [8] conducted a large study of subjects aged 0–83 years and showed continuous changes that occurred with age. Their report provided important insights. First, the period required to form an adult-like gut microbiota was the 3-year period following birth. Second, interpersonal variation was significantly greater between children than between adults. Third, the dominance of Bifidobacterium in the baby microbiota continued throughout the first year of life although this dominance composition diminished with age. Nevertheless, due to the limited number of subjects over 60 years of age, the continuous changes that occur in older people remain unknown. Recently, Odamaki et al. [7] reported age-related compositional differences from infants to centenarians in a Japanese cross-sectional study. They found that Bifidobacterium decreased and Enterobacteriaceae increased with age, as observed in some previous studies [8][20][21][22].

Relationships have also been shown between gut microbiota and diarrhea/constipation. Vandeputte et al. [29] described the association between stool consistency and gut microbiota composition in 53 healthy female subjects. Tigchelaar et al. [30] also reported an association between stool consistency and the structure of gut microbiota. Hadizadeh et al. [31] demonstrated a correlation between the number of bowel movements and gut microbiota. In a Japanese cohort, Takagi et al. [32] reported significant differences in microbial structure between individuals with different stool consistency (Bristol stool scale type). Therefore, investigating the relationship between bowel habits and intestinal bacterial composition can provide important information on gastrointestinal motility function.

However, Japan has its own food culture and customs compared to western countries, and the intestinal flora of Japanese individuals contain more genes for polysaccharide-degrading enzymes derived from water-soluble dietary fiber than Americans [33]. This feature may be related to the long life expectancy of Japanese people and the low body mass index (BMI) [34][35]. Nishijima et al. [36] clearly showed significant differences in the gut microbiota of the Japanese population compared to other countries, which cannot be explained by meals alone. Therefore, the structure of the intestinal flora may be highly dependent on an individual's country/region and lifestyle [37]. In this study, we investigated the relative abundance ranges of these taxa in stool samples from a large healthy human cohort. Further, we analyze the relationship between the above focused genera or the gut microbiota composition and the Japanese demographic features, life style, and bowel habits. Finally, we developed a reference range using a large healthy Japanese cohort, and considered the effect of age, gender, diarrhea, and constipation.

METHODS

Study Design and Participants

From November 2015 through June 2019, an online cohort of 5,843 individuals who had submitted fecal samples (one sample per subject) were selected from the ongoing Mykinso science microbiome research study. All participants provided written informed consent to participate after verbal and written information about the study. All procedures complied with the principles of the Declaration of Helsinki and were approved by the Institutional Review Board at our institution, registered as UMIN000028887 and UMIN000028888 in the UMIN Clinical Trials Registry System.

Symptoms of bowel habits and **Disease and medication data**

Self-reported disease data were collated from original surveys about health status completed by Mykinso online users at various times between 2016 and 2018 (Table S1). Individuals were scored positive for a disease if they replied yes to any original survey question, negative if they replied no, and unknown if data were unavailable across all original surveys. Diseases found in at least 1% of the wider cohort were considered common and retained in analyses. Self-reported prescription antibiotic use was scored from a single original survey. These data were cleaned to resolve spelling errors, followed by manual classification of entries into drug classes and subclasses by a health professional. Individuals were assumed not to be taking a medication if they had completed the original survey without listing it.

Fecal sampling, DNA extraction, and sequencing

Fecal samples were collected using brush-type collection kits containing guanidine thiocyanate solution (Techno Suruga Laboratory, Shizuoka, Japan) and stored at 4□. DNA extraction from the fecal samples was performed using an automated DNA extraction machine (GENE PREP STAR PI-480, Kurabo Industries Ltd, Osaka, Japan) according to the manufacturer's protocol. The V1–V2

region of the 16S rRNA gene was amplified using forward primer (16S_27Fmod: TCG TCG GCA

GCG TCA GAT GTG TAT AAG AGA CAG AGR GTT TGA TYM TGG CTC AG) and reverse

primer (16S_338R: GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA GTG CTG CCT

CCC GTA GGA GT) with KAPA HiFi Hot Start Ready Mix (Roche). To sequence 16S amplicons

by Illumina MiSeq platform, dual index adapters were attached using the Nextera XT Index kit.

Each library was diluted to 5 ng/µL, and equal volumes of the libraries were mixed to 4 nM. The

DNA concentration of the mixed libraries was quantified by qPCR with KAPA SYBR FAST qPCR

Master mix (KK4601, KAPA Biosystems) using primer 1 (AAT GAT ACG GCG ACC ACC) and

primer 2 (CAA GCA GAA GAC GGC ATA CGA). The library preparations were carried out

according to 16S library preparation protocol of Illumina (Illumina, San Diego, CA, USA).

Libraries were sequenced using the MiSeq Reagent Kit v2 (500 Cycles), 250 bp paired-end.

Taxonomy assignment based on 16S rRNA gene sequences

The paired-end reads of the partial 16S rRNA gene sequences were clustered by 97% nucleotide

identity, and then assigned taxonomic information using Greengenes database (v13.8)[38] through

QIIME pipeline (v1.8.0)[39]. The steps for data processing and assignment based on the QIIME

pipeline were as follows: (i) joining paired-end reads; (ii) quality filtering with an accuracy of Q30

(>99.9%) and a read length >300 bp; (iii)10,000 reads per sample were randomly extracted for the

subsequent analysis; (iv) clustering operational taxonomic units (OTUs) with 97% identity by

UCLUST (v1.2.22q) [40]; (v) assigning taxonomic information to each OTU using RDP classifier

[41] with the full-length 16S gene data of Greengenes (v13.8) to determine the identity and

composition of the bacterial genera.

Transformation of compositional microbiome data for hypothesis testing

The centered log-ratio (clr) transformed values were used as inputs for multivariate hypothesis

testing [42] to manage 0 count values as both point estimates using zCompositions R package [43]

and as a probability distribution using ALDEx2 [44] available on Bioconductor.

Group differences in beta-diversity

Aitchison distance, which is the Euclidean distance between samples after clr transformation, and

the distances between samples are the same as the phylogenetic ilr [42]. The replacement for β-

diversity exploration of microbiome data is the variance-based compositional principal component

(PCA) biplot [45], in which the relationship between inter-OTU variance and sample distance can

be observed [46]. Compositional PCA biplots display the relationships between OTUs and the dis

tances between samples on a common plot to clean substantial qualitative information regarding the

quality of the dataset and the relationships between groups [46].

Group differences in alpha-diversity

Microbiota diversity was assessed by Shannon index based on 97% nucleotide sequence identity.

These values were calculated by QIIME [[39]] with a depth of 10,000. To test two-group

differences between male and female groups, we calculated p-values by the two-sided unpaired

Welch's t-test. To test group differences among age-class group in diversity index, we calculated p-

values by the one-way analysis of variance (ANOVA).

Group differences in taxonomic abundance

To compare the taxonomic abundance between the groups , we conducted the univariate statistical

test using the ALDEx2 tool [44]. The false discovery rate (FDR) control was performed based on

the Benjamini-Hochberg procedure to correct for multiple testing, i.e., 'p.adjust' in R. Analysis was

confined to taxa with a prevalence greater than 10% and a maximum proportion (relative

abundance) greater than 0.005. An FDR-adjusted P value (or Q-value) less than 5% was considered

to be significant.

RESULTS

Cohort characteristics

Participants primarily resided in Japan (n = 5843) (Table S2), with a greater range in age, stool type,

and lifestyle than other Japanese large-scale microbiome projects [7][32][47]. Using an original

survey, participants (n = 4479) reported general health status, disease history, and lifestyle data

(participants missing any of these data were excluded, Table S3). In accordance with our

institutional review board (IRB), all survey questions were optional (question response rate,

76.65%). Eligible volunteers included male and female subjects who were considered to not have

common diseases. Disease status for individuals within the Mykinso online cohort was collected

from self-reported original surveys, and 12 common diseases (those reported in >1% of the total

cohort) were selected for consideration (Table S4). Eligible criteria included no self-reported

history of the 12 common diseases, including significant gastrointestinal inflammatory disease such

as inflammatory bowel disease or functional gastrointestinal disorders such as irritable bowel

syndrome (IBS). The primary investigation focused on a "non-disease adult" subset of individuals;

no self-reported history of inflammatory bowel disease (IBD) and diabetes. Finally, 2865

individuals were included for subsequent analysis (Table 1, Supplementary Figure 1).

Sex-related gut microbiota

Taxonomic differences in the microbial community were evaluated at the genus level. As shown in

Figure 1, the comparison of microbial composition between male and female subjects showed a

significant increase in the abundance of 12 genera in male subjects and in 13 genera in female

subjects (blue and red pch in Figure 1, respectively). These were characterized by an increase in

representative genera Prevotella, Megamonas, Collinsella, Dorea, Megasphaera, and Fusobacterium

(all p < 0.001) in male subjects and an increase in representative genera Oscillospira, Coprobacillus,

Ruminococcus, Bacteroides, Eggerthella, Anaerotruncus, Trabulsiella, and Akkermansia (all p <

0.001) in female subjects. Subsequently, we evaluated the diversity of gut microbiota using the

alpha-diversity index [Shannon index (OTU evenness estimation)]. The alpha-diversity index

showed no statistically significant differences between male (mean = 6.013) and female (mean =

6.008) subjects (Welch Two Sample t-test; t(2352.5) = -0.198, p = 0.843, 95%CI = -0.049–0.060).

Next, the overall structure of the gut microbiome for male and female subjects using beta-diversity

indices was calculated for Aitchison distance (Figure 2). PCA revealed that there were microbial

structural differences between male and female subjects (PERMANOVA, R^2 = 0.060, p = 0.001).

Age-related gut microbiota

Further, differences in the gut microbial structure in each age group were taxonomically evaluated

at the phylum level (Figure 3). In agreement with previous results, the microbiota composition

included four predominant phyla (Firmicutes, Bacteroidetes, Actinobacteria, Proteobacteria). Of

which, Actinobacteria showed significant decreases in 60 years subjects (p = 0.040, p.adjust =

0.640) and 70 years over subjects (p = 0.048, p.adjust = 0.700) compared with 19 years under

subjects.

The alpha-diversity index showed significant differences across age groups in our cohort (ANOVA;

F(6,2851) = 5.045, p < 0.001) (Figure 4). We also tested each pair difference with the Benjamini &

Hochberg method, and statistically significant differences were found between 20s group and 60s

group (p < 0.001), between 30s group and 60s group (p = 0.008), between 20s group and 40s group

(p = 0.006), and between 50s group and 60s group (p = 0.043) (Table 2). Additionally, we created

three age groups (0-19, 20–59, and 60 years or older), and the overall structure of the gut

microbiome using beta-diversity indices was calculated for Aitchison distance (Figure 5). PCA

revealed that there were microbial structural differences among the three age groups

(PERMANOVA, R^2 = 0.034, P < 0.001).

Bowel habits-related gut microbiota

Considering the heterogeneity and varying generations of samples in this dataset, we excluded samples from children (0–19 years) and aged adults (over 60 year), which might cause bias in the subsequent analysis [14][26][27][28][7]. As a result, 2675 samples were included in the resulting dataset (Table 3, Supplementary Figure 1).

The bowel habits (stool shape and defecation frequency) of all participants enrolled in this study were recorded and classified using the self-reported original survey. Additionally, perceived symptoms of diarrhea/constipation were recorded and classified. According to the stool shape, bowel frequency, and perceived symptom scores, participants were classified as normal bowel habit type, diarrhea type, constipation type, or mixed type. For the constipation type, when stool types 1 (hard stool), defecation frequency type 4 (less one time per week), or perceived constipation symptoms frequented within 1 month, they were classified as constipation. For the diarrhea type, when stool types 7 (liquid stools), defecation frequency type 1 (more three time per day), or perceived diarrhea symptoms frequented within 1 month, they were classified as diarrhea. And when the patient had both the constipation type and the diarrhea type, they were classified as mixed type. Constipation group (female, n = 337, 20.87%; male, n=37, 3.49%), diarrhea group (female, n =357, 22.11%; male, n = 457, 43.11%), mixed group (female, n =139, 8.61%; male, n = 29, 2.74%), and normal group (female, n = 782, 48.42%; male, n = 537, 50.66%) were observed (Table 3). However, in this study, to evaluate the association between bowel habits and gut microbiota, we created three bowel habit groups.

Importantly, the alpha-diversity index for each bowel habit group showed significant difference among groups in our cohort (ANOVA; $F(3,2667) = 1.761$, $p < 0.001$) (Figure 6). We also tested each pair difference with the Benjamini & Hochberg method, which showed statistically significant differences between the normal and diarrhea groups ($p < 0.001$), constipation and diarrhea groups ($p < 0.001$), and mixed and diarrhea groups ($p = 0.001$). These differences are visualized in Figure 6.

Additionally, the overall structure of the gut microbiome among the three bowel habit groups using beta-diversity indices was calculated for Aitchison distance and visualized by PCA according to Aitchison distance (Figure 7). An additional PERMANOVA analysis based on categorical variables of their abundance showed that the bowel habit type was a significant factor contributing to variation of the gut microbiota (p < 0.001). Approximately 0.7% of the variance in beta diversity was explained by the bowel habit type (PERMANOVA; $F(3,2667) = 6.714$, $R^2 = 0.007$, $P < 0.001$), which was competitive with many microbiome covariates.

Subsequently, at the genus level, we identified several altered bacteria among the three bowel habit groups. Interestingly, significantly higher relative abundance of Fusobacterium in the diarrhea group and Oscillospira (p < 0.001) in the constipation group was observed. In addition, the relative abundance of Ruminococcus, Anaerotruncus, Alistipes, and Akkermansia (all p < 0.001) were significantly higher in the constipation group, while Dorea (p < 0.01) was higher in the diarrhea group.

Reference ranges from Japanese healthy cohort

Considering the heterogeneity and bowel habits of the sample dataset, we excluded samples from individuals with diarrhea or constipation (Supplementary Figure 1), which might cause bias in the reference range [29][30][31][32]. Finally, 1319 samples were selected as the healthy reference dataset (Supplementary Figure 1). We identified 453 genera and 20 phyla of Bacteria and Archaea in the gut microbiomes of the 1319 individuals from self-reported healthy individuals. The genera with an average relative abundance of ≥ 0.5% in the Japanese healthy reference are listed in Supplementary Figure 2. At the genus level, the Japanese healthy reference was characterized by the highest abundance of Bacteroides, Faecalibacterium, Prevotella, Blautia, Bidfidobacterium, Coprococcus, and Parabacteroides (Supplementary Fig 2).

Many microorganisms that are clinically related to health and disease are present to some extent in the intestines of healthy people. The present reference ranges obtained from Japanese healthy populations could be new microbiome indices for determining and/or monitoring the state of health and/or balance of the intestinal microbiota of an individual. In this study, health-related microbiome indices were selected based on peer-reviewed studies in academic journals and in-house data analyses (Table 4). To determine the reference range of 11 target microbiome indices, a cohort of 1319 individuals from healthy individuals self-reported in the Mykinso's microbiota research study was established. Microbiome data from this cohort were analyzed to determine the empirical reference ranges for two indices of overall community structure, two complex genus indices, one class, and six genera. For each of the 1319 samples, we determined the relative abundance of each target within the microbial population, revealing the distribution of the relative abundance of each target in the cohort (Table 4). These data were used to define a central 80% healthy range with confidence intervals for each target. Many of the targets show significant spread, highlighting the importance of defining a reference range for health-related species.

DISCUSSION

We developed a reference range using a large healthy Japanese cohort. The reference range considers the effect of age, gender, diarrhea, and constipation to aid physicians in accurate diagnosis of the intestinal bacterial composition ratio using a standard value derived from a healthy population. Eighteen intestinal bacterial indicators suggested to be associated with health status were selected. Using intestinal bacterial composition test panels, detection of intestinal bacterial indicators outside the healthy range can be useful to reinforce the medical plan.

Gut microbiota and sex/gender

Some characteristics of gender-specific immune differences are induced by gut microbiota. Fransen et al [48] investigated significant differences in bacterial groups at family or genus levels. Females had higher abundance of Desulfovibrionaceae, Lactobacillaceae (Lactobacillus at the genus level), and Verrucomicrobiaceae (Akkermansia at the genus level), whereas males had higher abundance of Ruminococcaceae and Rikenellaceae (Alistipes at the genus level). In this study several characteristic differences were observed between male and female subjects regarding the genus abundance of gut microbiota. The genera Prevotella, Megagamonas, Fusobacterium, and Megasphaera were significantly increased in male subjects, while Bifidobacterium, Ruminococcus, and Akkermansia were significantly increased in female subjects. These results are consistent with the results of previous Japanese studies [7][32], and may be considered as the characteristic of gender differences of composition of intestinal microbiota in the Japanese population.

Gut microbiota and age/generation

Recent reports have shown a clear difference in the composition of intestinal flora of infants, adults, and the elderly [7][8][32]. The microbiota composition initially shifts after birth, followed by significant shifts during childhood and in later years [7][8]. In this study, we segmented by age

group (young group, middle-aged group, elderly group). The diversity index showed slight

differences between each age group from 20 to 89 years old (Figure 4), which are consistent with

previous reports [7]. Further, our results are in agreement with studies indicating clear differences in

gut microbiota composition among infants, adults, and the elderly [7][49][50]. Moreover, this study

indicated that Actinobacteria abundance and alpha-diversity index were gut microbiome indices

related to aging.

Gut microbiota and bowel habits (diarrhea/constipation)

In this study, we evaluated the association between bowel habits and gut microbiota. Similar to

Vandeputte et al. [29], we found a significant association between the bowel habits (stool shape and

defecation frequency) and gut microbiota diversity. Furthermore, deviation of the gut microbiota

composition in several genera, including Oscillospira, Ruminococcus, Anaerotruncus, Alistipes, and

Akkermansia, in constipation subjects and Fusobacterium and Dorea in diarrhea subjects were

confirmed, which are consistent with a previous report [32]. Although the role of these genera in

stool consistency remains unclear, the results illustrate the effect of gut microbiota on stool

consistency in Japanese healthy subjects.

Gut microbiota and racial/regional differences

Our results showed that Japanese adults (20–59 years old) had a greater abundance of genera

Bacteroides and Faecalibacterium (interquartile ranges (IQR) of 27.43% (19.03–35.26) and 6.83%

(3.39–10.06), respectively) and a relatively lower abundance of genera Clostridium (IQR 0.20

(0.04–0.44) %), compared to previous studies in other Japanese cohorts [36]. However, the

estimated abundance of Bifidobacterium and Blautia was greater (interquartile ranges (IQR) of

2.47% (0.86–5.78) and 5.31% (2.94-7.85), than a previous study in other nations, which were <

0.5% and > 5%, in the US and China, respectively [36]. These bacterial compositions may be

characteristic of the intestinal microbiota of the Japanese population, but may also reflect

differences in DNA extraction methods [51][52] and the amplified region of the 16S rRNA [53].

The high abundance of Bifidobacterium has been also observed in the gut microbiome of Japanese

children based on the 16S rRNA gene analysis [12], indicating its high prevalence throughout the

Japanese population. Bifidobacterium is thought to be a beneficial microbe that contains more

glycoside hydrolases for degrading starch than other intestinal microbes [54][55]. Therefore, the

high abundance of Bifidobacterium may be the consequence of the intake of various saccharides in

traditional and unique Japanese foods. However, it is unknown which foods or nutrients contribute

to the high abundance of Bifidobacterium. As future prospects, it is essential to create a reference

microbiota for the Japanese population by age group, increasing the number of subjects in the

young and the elderly age groups. Additionally, investigations of geographical differences within

Japan are of interest.

Clinical relevance and Reference ranges

All 11 microbiome indices successfully identified using 16S rRNA gene sequencing were

associated with specific health conditions.

The alpha-diversity, including Shannon index, and observed genera number appeared to be

associated with better health [56]. A recent meta-analysis has proposed reduced alpha diversity as a

reliable indicator of diarrhea-associated dysbiosis [57]. These microbiota diversity indices

(Shannon index and observed OTUs) showed significant differences in the healthy aging group,

indicating that a healthy, diverse diet promotes greater diversity in gut microbiota [56].

Previous studies have proposed that Bifidobacterium is inversely associated with IBD and diarrhea-

associated dysbiosis [58], and the consumption of probiotics, inulin, and oligofructoses promotes an

increase in Bifidobacterium abundance [59]. Additionally, Faecalibacterium has been proposed as a

dominant member of human intestinal microbiota in healthy adults, and especially as a health

sensor for active Crohn's disease patients [60]. A recent meta-analysis showed a reduction in butyrate-producing Clostridiales, including Coprococcus, Roseburia, Butyricicoccus, Faecalibacterium, Anaerostipes, and Butyricimonas, which have been associated with a healthy gut [57].

While not all genera within the order Lactobacillales are verified lactic acid producers, the dominant genera within this order (including Lactobacillus, Pediococcus, Leuconostoc, Lactococcus, Weissella) are known to harbor genes for lactic acid production and are often enriched in case patients across multiple diseases [57]. Lactobacillales genera have been shown to adapt to the lower pH of the upper gastrointestinal tract [61]. Thus, the shared disease-associated taxa may be indicators of shorter stool transit times and disruptions in the redox state and/or pH of the lower intestine, rather than specific pathogens. Genera within Lactobacillaceae and Streptococcaceae families are dominant in the upper gastrointestinal tract and are present in the stool of many individuals at low frequency [57]. These taxa likely become enriched with faster stool transit time (i.e., diarrhea signatures) [57].

Previous studies have proposed Fusobacterium to be associated with various human diseases [62]. Dysbiosis associated with colorectal cancer is generally characterized by increased prevalence of known pathogenic or pathogen-associated Fusobacterium and Enterobacter genera, which were higher in colorectal cancer patients in two or more studies [57]. Furthermore, other oral community genera, such as members of Porphyromonas, Peptostreptococcus, and Parvimonas, were found with Fusobacterium on colonic tumors [63].

This reference database study characterized the diversity of the Japanese industrialized human gut microbiome, revealing novel relationships with health, lifestyle, and dietary factors and establishing the Mykinso resource and infrastructure as a living platform for discovery.

Acknowledgements

We thank Dr Shota Nakamura (Osaka University), iMet Laboratory team (Osaka University) and

BikenBiomics team for sequencing and technical support, and Dr Takeaki Uno (National Institute

of Informatics) for support with data analysis. We also would like to thank Editage

(www.editage.com) for English language editing.

REFERENCES

[1]   The Human Microbiome Project Consortium, "A framework for human microbiome research," *Nature*, vol. 486, no. 7402, pp. 215–221, Jun. 2012, doi: 10.1038/nature11209.

[2]   MetaHIT Consortium *et al.*, "A human gut microbial gene catalogue established by metagenomic sequencing," *Nature*, vol. 464, no. 7285, pp. 59–65, Mar. 2010, doi: 10.1038/nature08821.

[3]   J. Li *et al.*, "An integrated catalog of reference genes in the human gut microbiome," *Nature Biotechnology*, vol. 32, p. 834, Jul. 2014.

[4]   S. V. Lynch and O. Pedersen, "The Human Intestinal Microbiome in Health and Disease," *New England Journal of Medicine*, vol. 375, no. 24, pp. 2369–2379, Dec. 2016, doi: 10.1056/NEJMra1600266.

[5]   M. B. Flak, J. F. Neves, and R. S. Blumberg, "Welcome to the Microgenderome," *Science*, vol. 339, no. 6123, pp. 1044–1045, Mar. 2013, doi: 10.1126/science.1236226.

[6]   J. K. Goodrich *et al.*, "Human Genetics Shape the Gut Microbiome," *Cell*, vol. 159, no. 4, pp. 789–799, Nov. 2014, doi: 10.1016/j.cell.2014.09.053.

[7]   T. Odamaki *et al.*, "Age-related changes in gut microbiota composition from newborn to centenarian: a cross-sectional study," *BMC Microbiol*, vol. 16, no. 1, p. 90, Dec. 2016, doi: 10.1186/s12866-016-0708-5.

[8]   T. Yatsunenko *et al.*, "Human gut microbiome viewed across age and geography," *Nature*, vol. 486, no. 7402, pp. 222–227, Jun. 2012, doi: 10.1038/nature11053.

[9]   J. Zhang *et al.*, "Mongolians core gut microbiota and its correlation with seasonal dietary changes," *Sci Rep*, vol. 4, no. 1, p. 5001, May 2015, doi: 10.1038/srep05001.

[10]   G. D. Wu *et al.*, "Linking Long-Term Dietary Patterns with Gut Microbial Enterotypes," *Science*, vol. 334, no. 6052, pp. 105–108, Oct. 2011, doi: 10.1126/science.1208344.

[11]   C. De Filippo *et al.*, "Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa," *Proceedings of the National Academy of Sciences*, vol. 107, no. 33, pp. 14691–14696, Aug. 2010, doi: 10.1073/pnas.1005963107.

[12]   J. Nakayama *et al.*, "Diversity in gut bacterial community of school-age children in Asia," *Scientific Reports*, vol. 5, no. 1, Jul. 2015, doi: 10.1038/srep08397.

[13]   O. La-ongkham, M. Nakphaichit, V. Leelavatcharamas, S. Keawsompong, and S. Nitisinprasert, "Distinct gut microbiota of healthy children from two different geographic regions of Thailand," *Archives of Microbiology*, vol. 197, no. 4, pp. 561–573, May 2015, doi: 10.1007/s00203-015-1089-0.

[14]   S. Mueller *et al.*, "Differences in Fecal Microbiota in Different European Study Populations in Relation to Age, Gender, and Country: a Cross-Sectional Study," *Applied and Environmental Microbiology*, vol. 72, no. 2, pp. 1027–1033, Feb. 2006, doi: 10.1128/AEM.72.2.1027-1033.2006.

[15]   C. Haro *et al.*, "Intestinal Microbiota Is Influenced by Gender and Body Mass Index," *PLoS ONE*, vol. 11, no. 5, p. e0154090, May 2016, doi: 10.1371/journal.pone.0154090.

[16]   L. Yurkovetskiy *et al.*, "Gender Bias in Autoimmunity Is Influenced by Microbiota," *Immunity*, vol. 39, no. 2, pp. 400–412, Aug. 2013, doi: 10.1016/j.immuni.2013.08.013.

[17]   J. G. M. Markle *et al.*, "Sex Differences in the Gut Microbiome Drive Hormone-Dependent Regulation of Autoimmunity," *Science*, vol. 339, no. 6123, pp. 1084–1088, Mar. 2013, doi: 10.1126/science.1233521.

[18]   Y. Min *et al.*, "Sex-specific association between gut microbiome and fat distribution," *Nat Commun*, vol. 10, no. 1, p. 2408, Dec. 2019, doi: 10.1038/s41467-019-10440-5.

[19]   J. L. Round and S. K. Mazmanian, "The gut microbiota shapes intestinal immune responses during health and disease," *Nat Rev Immunol*, vol. 9, no. 5, pp. 313–323, May 2009, doi: 10.1038/nri2515.

[20]   A. Zhernakova *et al.*, "Population-based metagenomics analysis reveals markers for gut microbiome composition and diversity," *Science*, vol. 352, no. 6285, pp. 565–569, Apr. 2016, doi: 10.1126/science.aad3369.

[21]   G. Falony *et al.*, "Population-level analysis of gut microbiome variation," *Science*, vol. 352, no. 6285, pp. 560–564, Apr. 2016, doi: 10.1126/science.aad3503.

[22]   C. F. Maurice, H. J. Haiser, and P. J. Turnbaugh, "Xenobiotics Shape the Physiology and Gene Expression of the Active Human Gut Microbiome," *Cell*, vol. 152, no. 1–2, pp. 39–50, Jan. 2013, doi: 10.1016/j.cell.2012.10.052.

[23]   E. Avershina, O. Storrø, T. Øien, R. Johnsen, P. Pope, and K. Rudi, "Major faecal microbiota shifts in composition and diversity with age in a geographically restricted cohort

of mothers and their children," *FEMS Microbiol Ecol*, vol. 87, no. 1, pp. 280–290, Jan. 2014, doi: 10.1111/1574-6941.12223.

[24]   A. Bergstrom *et al.*, "Establishment of Intestinal Microbiota during Early Life: a Longitudinal, Explorative Study of a Large Cohort of Danish Infants," *Applied and Environmental Microbiology*, vol. 80, no. 9, pp. 2889–2900, May 2014, doi: 10.1128/AEM.00342-14.

[25]   J. Cheng *et al.*, "Discordant temporal development of bacterial phyla and the emergence of core in the fecal microbiota of young children," *ISME J*, vol. 10, no. 4, pp. 1002–1014, Apr. 2016, doi: 10.1038/ismej.2015.177.

[26]   M. J. Claesson *et al.*, "Composition, variability, and temporal stability of the intestinal microbiota of the elderly," *Proceedings of the National Academy of Sciences*, vol. 108, no. Supplement_1, pp. 4586–4591, Mar. 2011, doi: 10.1073/pnas.1000097107.

[27]   D. Mariat *et al.*, "The Firmicutes/Bacteroidetes ratio of the human microbiota changes with age," *BMC Microbiol*, vol. 9, no. 1, p. 123, 2009, doi: 10.1186/1471-2180-9-123.

[28]   E. Biagi *et al.*, "Through ageing, and beyond: gut microbiota and inflammatory status in seniors and centenarians," *PLoS ONE*, vol. 5, no. 5, p. e10667, May 2010, doi: 10.1371/journal.pone.0010667.

[29]   D. Vandeputte, G. Falony, S. Vieira-Silva, R. Y. Tito, M. Joossens, and J. Raes, "Stool consistency is strongly associated with gut microbiota richness and composition, enterotypes and bacterial growth rates," *Gut*, vol. 65, no. 1, pp. 57–62, Jan. 2016, doi: 10.1136/gutjnl-2015-309618.

[30]   E. F. Tigchelaar, M. J. Bonder, S. A. Jankipersadsing, J. Fu, C. Wijmenga, and A. Zhernakova, "Gut microbiota composition associated with stool consistency," *Gut*, vol. 65, no. 3, p. 540, Mar. 2016, doi: 10.1136/gutjnl-2015-310328.

[31]   F. Hadizadeh *et al.*, "Stool frequency is associated with gut microbiota composition," *Gut*, vol. 66, no. 3, p. 559, Mar. 2017, doi: 10.1136/gutjnl-2016-311935.

[32]   T. Takagi *et al.*, "Differences in gut microbiota associated with age, sex, and stool consistency in healthy Japanese subjects," *J Gastroenterol*, vol. 54, no. 1, pp. 53–63, Jan. 2019, doi: 10.1007/s00535-018-1488-5.

[33]   J.-H. Hehemann, G. Correc, T. Barbeyron, W. Helbert, M. Czjzek, and G. Michel, "Transfer of carbohydrate-active enzymes from marine bacteria to Japanese gut

microbiota," *Nature*, vol. 464, no. 7290, pp. 908–912, Apr. 2010, doi: 10.1038/nature08937.

[34]   World Health Organization, "Global Health Observatory (GHO) data: overweight and obesity," 2015.

[35]   K. Kurokawa *et al.*, "Comparative Metagenomics Revealed Commonly Enriched Gene Sets in Human Gut Microbiomes," *DNA Research*, vol. 14, no. 4, pp. 169–181, 2007, doi: 10.1093/dnares/dsm018.

[36]   S. Nishijima *et al.*, "The gut microbiome of healthy Japanese and its microbial and functional uniqueness," *DNA Research*, vol. 23, no. 2, pp. 125–133, Apr. 2016, doi: 10.1093/dnares/dsw002.

[37]   M. Arumugam *et al.*, "Enterotypes of the human gut microbiome," *Nature*, vol. 473, no. 7346, pp. 174–180, May 2011, doi: 10.1038/nature09944.

[38]   T. Z. DeSantis *et al.*, "Greengenes, a Chimera-Checked 16S rRNA Gene Database and Workbench Compatible with ARB," *Applied and Environmental Microbiology*, vol. 72, no. 7, pp. 5069–5072, Jul. 2006, doi: 10.1128/AEM.03006-05.

[39]   J. G. Caporaso *et al.*, "QIIME allows analysis of high-throughput community sequencing data," *Nature Methods*, vol. 7, no. 5, pp. 335–336, May 2010, doi: 10.1038/nmeth.f.303.

[40]   R. C. Edgar, "Search and clustering orders of magnitude faster than BLAST," *Bioinformatics*, vol. 26, no. 19, pp. 2460–2461, Oct. 2010, doi: 10.1093/bioinformatics/btq461.

[41]   J. R. Cole *et al.*, "The Ribosomal Database Project: improved alignments and new tools for rRNA analysis," *Nucleic Acids Research*, vol. 37, no. Database, pp. D141–D145, Jan. 2009, doi: 10.1093/nar/gkn879.

[42]   J. Aitchison, C. Barcelo-Vidal, J. A. Martın-Fernandez, and V. Pawlowsky-Glahn, "Logratio Analysis and Compositional Distance," *Mathematical Geology*, p. 2, 2000.

[43]   J. Palarea-Albaladejo and J. A. Martín-Fernández, "zCompositions — R package for multivariate imputation of left-censored data under a compositional approach," *Chemometrics and Intelligent Laboratory Systems*, vol. 143, pp. 85–96, Apr. 2015, doi: 10.1016/j.chemolab.2015.02.019.

[44]   G. B. Gloor and G. Reid, "Compositional analysis: a valid approach to analyze microbiome high-throughput sequencing data," *Can. J. Microbiol.*, vol. 62, no. 8, pp. 692–703, Aug. 2016, doi: 10.1139/cjm-2015-0821.

[45]   J. Aitchison and M. Greenacre, "Biplots of compositional data," *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, vol. 51, no. 4, pp. 375–392, 2002, doi: 10.1111/1467-9876.00275.

[46]   G. B. Gloor, J. M. Macklaim, V. Pawlowsky-Glahn, and J. J. Egozcue, "Microbiome Datasets Are Compositional: And This Is Not Optional," *Front. Microbiol.*, vol. 8, p. 2224, Nov. 2017, doi: 10.3389/fmicb.2017.02224.

[47]   S. Nishijima *et al.*, "The gut microbiome of healthy Japanese and its microbial and functional uniqueness," *DNA Res*, vol. 23, no. 2, pp. 125–133, Apr. 2016, doi: 10.1093/dnares/dsw002.

[48]   F. Fransen *et al.*, "The Impact of Gut Microbiota on Gender-Specific Differences in Immunity," *Front. Immunol.*, vol. 8, p. 754, Jun. 2017, doi: 10.3389/fimmu.2017.00754.

[49]   M. Conlon and A. Bird, "The Impact of Diet and Lifestyle on Gut Microbiota and Human Health," *Nutrients*, vol. 7, no. 1, pp. 17–44, Dec. 2014, doi: 10.3390/nu7010017.

[50]   P. Vernocchi, F. Del Chierico, and L. Putignani, "Gut Microbiota Profiling: Metabolomics Based Approach to Unravel Compounds Affecting Human Health," *Front. Microbiol.*, vol. 7, Jul. 2016, doi: 10.3389/fmicb.2016.01144.

[51]   A. Wesolowska-Andersen *et al.*, "Choice of bacterial DNA extraction method from fecal material influences community structure as evaluated by metagenomic analysis," *Microbiome*, vol. 2, no. 1, p. 19, 2014, doi: 10.1186/2049-2618-2-19.

[52]   B. Wagner Mackenzie, D. W. Waite, and M. W. Taylor, "Evaluating variation in human gut microbiota profiles due to DNA extraction method and inter-subject differences," *Front. Microbiol.*, vol. 6, Feb. 2015, doi: 10.3389/fmicb.2015.00130.

[53]   J. H. Campbell *et al.*, "Host genetic and environmental effects on mouse intestinal microbiota," *ISME J*, vol. 6, no. 11, pp. 2033–2044, Nov. 2012, doi: 10.1038/ismej.2012.54.

[54]   J.-H. Lee and D. J. O'Sullivan, "Genomic Insights into Bifidobacteria," *Microbiology and Molecular Biology Reviews*, vol. 74, no. 3, pp. 378–416, Sep. 2010, doi: 10.1128/MMBR.00004-10.

[55]   S. Liu *et al.*, "Starch and starch hydrolysates are favorable carbon sources for Bifidobacteria in the human gut," *BMC Microbiol*, vol. 15, no. 1, p. 54, 2015, doi: 10.1186/s12866-015-0362-3.

[56]   M. J. Claesson *et al.*, "Gut microbiota composition correlates with diet and health in the elderly," *Nature*, vol. 488, no. 7410, pp. 178–184, Aug. 2012, doi: 10.1038/nature11319.

[57]   C. Duvallet, S. M. Gibbons, T. Gurry, R. A. Irizarry, and E. J. Alm, "Meta-analysis of gut microbiome studies identifies disease-specific and shared responses," *Nat Commun*, vol. 8, no. 1, p. 1784, Dec. 2017, doi: 10.1038/s41467-017-01973-8.

[58]   M. Joossens *et al.*, "Dysbiosis of the faecal microbiota in patients with Crohn's disease and their unaffected relatives," *Gut*, vol. 60, no. 5, p. 631, May 2011, doi: 10.1136/gut.2010.223263.

[59]   D. Meyer and M. Stasse-Wolthuis, "The bifidogenic effect of inulin and oligofructose and its consequences for gut health," *Eur J Clin Nutr*, vol. 63, no. 11, pp. 1277–1289, Nov. 2009, doi: 10.1038/ejcn.2009.64.

[60]   S. Miquel *et al.*, "Faecalibacterium prausnitzii and human intestinal health," *Current Opinion in Microbiology*, vol. 16, no. 3, pp. 255–261, Jun. 2013, doi: 10.1016/j.mib.2013.06.003.

[61]   G. P. Donaldson, S. M. Lee, and S. K. Mazmanian, "Gut biogeography of the bacterial microbiota," *Nat Rev Microbiol*, vol. 14, no. 1, pp. 20–32, Jan. 2016, doi: 10.1038/nrmicro3552.

[62]   Y. W. Han, "Fusobacterium nucleatum: a commensal-turned pathogen," *Current Opinion in Microbiology*, vol. 23, pp. 141–147, Feb. 2015, doi: 10.1016/j.mib.2014.11.013.

[63]   K. J. Flynn, N. T. Baxter, and P. D. Schloss, "Metabolic and Community Synergy of Oral Bacteria in Colorectal Cancer," *mSphere*, vol. 1, no. 3, pp. e00102-16, /msph/1/3/e00102-16.atom, Jun. 2016, doi: 10.1128/mSphere.00102-16.

Confidential

TABLES

Table 1 Distribution of primary eligible subjects

Table 2 Pairwise comparisons of Shannon-index between age groups

Table 3 Distribution of bowel habits by sex and age-class (20–50)

Table 4 Reference ranges from healthy Japanese subjects for 11 clinically relevant indices

Table S1 Original survey questions

Table S2 Distribution of all subjects

Table S3 Distribution of question non-responding subjects

Table S4 Distribution of subjects excluded from reference population due to reporting at least one

of 12 common diseases

FIGURE LEGENDS

Figure 1 Relative abundance of gut microbiota in male and female subjects. Genera were significantly different between male and female subjects. diff_btw = median difference in clr values between female and male groups. magenta: positive diff_btw in female group; cyan: negative diff_btw in female group

Figure 2 Plot of individual samples from PCA output (magenta: female samples, cyan: male samples). The distance between points is proportional to the Euclidian distance of CLR vectors of the samples (Aitchison distance). The multivariate distance between samples was estimated using the Aitchison distance, which showed significantly different composition between female and male samples (ANOSIM, $R^2 = 0.060$, $p = 0.001$)

Figure 3 Comparative analyses of taxonomic composition of microbial community at the phylum level for each age group. Each component of the cumulative bar chart indicates a phylum

Figure 4 Age-related difference in alpha-diversities of gut microbiota

Figure 5 (a) Plot of individual samples from PCA output (red: senior samples, blue: junior samples, and gray: adult samples). The distance between points is proportional to the Euclidian distance of CLR vectors of the samples (Aitchison distance). (b) The multivariate distance between samples was estimated using the Aitchison distance, which showed significantly different composition in the junior, adult and senior samples (PERMANOVA, $R^2 = 0.034$, $P < 0.001$)

Figure 6 Alteration of alpha-diversity of gut microbiota associated with stool consistency. Comparison of α-diversity indices: Shannon-index (OTU evenness estimation). Bowel habit was categorized into three groups: normal, diarrhea, and constipation

Figure 7 (a) Plot of individual samples from PCA output (gray: normal samples, red: diarrhea samples, green: constipation samples, and blue: mixed-type samples). The distance between points is proportional to the Euclidian distance of CLR vectors of the samples (Aitchison distance). The multivariate distance between samples was estimated using the Aitchison distance, which showed

significantly different composition in the junior, adult and senior samples (PERMANOVA, R^2 =

0.007, P < 0.001). (b) RDA triplot of CLR vectors of the samples constrained by bowel habit group

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

• Figure 6.

- Figure 7

Table 1.

Years	Female	Male	Number of samples
19 under	20	10	30
20-29	322	135	457
30-39	559	408	967
40-49	485	358	843
50-59	249	159	408
60-69	73	52	125
70 over	14	21	35
Sum	1722	1143	2865

Table 2.

pair	diff	lwr	upr	p.adj
20-19under	0.001	-0.398	0.400	1.000
30-19under	0.109	-0.284	0.502	0.983
30-20	0.108	-0.012	0.228	0.112
40-19under	0.151	-0.242	0.545	0.917
40-20	0.151	0.027	0.274	0.006
40-30	0.043	-0.057	0.142	0.870
50-19under	0.128	-0.272	0.529	0.965
50-20	0.127	-0.017	0.272	0.125
50-30	0.019	-0.106	0.145	0.999
50-40	-0.023	-0.151	0.105	0.998
60-19under	0.349	-0.082	0.779	0.203
60-20	0.348	0.134	0.562	<0.001
60-30	0.240	0.039	0.441	0.008
60-40	0.197	-0.006	0.400	0.063
60-50	0.220	0.004	0.437	0.043
70over-19under	0.303	-0.224	0.830	0.618
70over-20	0.302	-0.069	0.674	0.199
70over-30	0.194	-0.170	0.559	0.701
70over-40	0.152	-0.214	0.517	0.885
70over-50	0.175	-0.198	0.548	0.812
70over-60	-0.046	-0.451	0.359	1.000

diff: Differences in mean levels
lwr: 95% confidence lower level
upr: 95% confidence upper level
p.adj: p-value adjusted by Benjamin & Hochberg medhods

Table 3.

Years	Normal stool	Diarrhea	Constipation	Mixed	Number of samples
20-29: male	59	69	2	5	135
20-29: female	149	76	57	40	322
30-39: male	196	185	14	13	408
30-39: female	243	134	128	54	559
40-49: male	185	150	15	8	358
40-49: female	233	112	111	29	485
50-59: male	97	53	6	3	159
50-59: female	157	35	41	16	249
Sum	1319	814	374	168	2675

Table 4.

Microbiome Index	unit	group	lower limit [10%]	Median [50%]	upper limit [90%]	Reference
Shannon	value	all	5.08	6.01	6.88	[53, 54]
		male	5.15	6.03	6.93	
		female	5.05	6	6.85	
Observed genera	genus	all	54	65	80	[53, 54]
		male	54	65	81	
		female	54	66	79	
Bifidobacterium	%	all	0.18	2.47	9.6	[55, 56]
		male	0.12	2.18	8.45	
		female	0.21	2.79	10.41	
Faecalibacterium	%	all	0.55	6.83	12.87	[57]
		male	0.37	6.36	12.09	
		female	0.57	7.23	13.27	
butyrate-producing genus	%	all	4.25	12.16	20.47	[54]
		male	3.64	11.53	20.03	
		female	4.66	12.88	20.89	
Clostridium	%	all	0	0.19	0.79	[54]
		male	0	0.19	0.7	
		female	0	0.2	0.89	
lactic acid-producing genus	%	all	0	0.01	0.22	[54, 58]
		male	0	0.01	0.24	
		female	0	0.01	0.21	
Streptococcus	%	all	0.05	0.38	2.58	[54, 58]
		male	0.03	0.34	2.46	
		female	0.05	0.41	2.65	
Oral bacteria related genus	%	all	0.18	1.4	9.74	[59, 60]
		male	0.15	1.31	9.71	
		female	0.2	1.48	9.9	
Fusobacterium	%	all	0	0	1.37	[59, 60]
		male	0	0	2.92	
		female	0	0	0.83	
Observed genera within class gamma proteobacteria	class	all	2	5	8	[60]
		male	2	5	8	
		female	2	5	8	