# Suppression of Inflammation Delays Hair Cell Regeneration and Functional Recovery Following Lateral Line Damage in Zebrafish Larvae

Ru Zhang[1,2,3], Xiao-Peng Liu[3,4], Ya-Juan Li[2], Ming Wang[1,3], Lin Chen[1,3*], Bing Hu[1,2*]

1   CAS Key Laboratory of Brain Function and Diseases, School of Life Sciences, University of Science and Technology of China, Hefei 230027, China.

2   Lab of Neurodevelopment and Repair, University of Science and Technology of China, Hefei 230027, China.

3   Auditory Research Laboratory, University of Science and Technology of China, Hefei 230027, China.

4   Center for Hearing and Deafness, State University of New York at Buffalo, NY14214, USA.

*Corresponding authors.

Bing Hu, PhD, Lab of Neurodevelopment and Repair, University of Science and Technology of China, Hefei 230027, China.

Tel.: +86 (551) 6360-2489

E-mail: bhu@ustc.edu.cn

Lin Chen, PhD, Auditory Research Laboratory, University of Science and Technology of China, Hefei 230027, China.

Tel.: +86 (551) 6360-7623

E-mail: linchen@ustc.edu.cn

## Abstract:

**Background:** Human cochlear hair cells cannot spontaneously regenerate after loss. In contrast, those in fish and amphibians have a remarkable ability to regenerate after damaged. Previous studies focus on signaling mechanisms of hair cell regeneration, such as Wnt and Notch signals but seldom on the fact that the beginning of regeneration is accompanied by a large number of inflammatory responses. The detailed role of this inflammation in hair cell regeneration is still unknown. In addition, there is no appropriate behavioral method to quantitatively evaluate the functional recovery of lateral line hair cells after regeneration.

**Results:** In this study, we found that when inflammation was suppressed, the regeneration of lateral line hair cells and the recovery of the rheotaxis of the larvae were significantly delayed. Calcium imaging showed that the function of the neuromasts in the inflammation-inhibited group was weaker than that in the non-inflammation-inhibited group at the Early Stage of regeneration, and returned to normal at the Late Stage. Calcium imaging also revealed the cause of the mismatch between the function and quantity during regeneration.

**Conclusions:** Our results, meanwhile, suggest that suppressing inflammation delays hair cell regeneration and functional recovery when hair cells are damaged. This study may provide a new knowledge for how to promote hair cell regeneration and functional recovery in adult mammals.

**Keywords:** inflammation, hair cell regeneration, neuromast, lateral line, zebrafish larva, rheotaxis, calcium imaging.

## Background:

Deafness and hearing defects are usually caused by loss of sensory hair cells or defect of auditory function. The loss of hair cells is result of aging, infection, genetic factors, hypoxia, autoimmune disorder, ototoxic drugs or noise exposure. Unfortunately, including humans, hair cells cannot regenerate in mammals (Oesterle and Stone, 2008; Yorgason. et al., 2006). In contrast, hair cells in some non-mammalian vertebrates have a remarkable ability to regenerate, such as birds, reptiles, amphibians and fish (Matsui. and Cotanchea., 2004; Popper and Hoxter, 1984; Stone. and Rubel., 2000). It could suggest that if we figure out the mechanism of hair cell regeneration in these species, we probably can promote hair cell regeneration in mammals.

When hair cells are damaged, support cells proliferate into both hair cells and support cells, or convert into hair cells directly (Baird et al., 1996; Lopez-Schier and Hudspeth, 2006; Raphael, 1992; Roberson et al., 2004). Hair cell regeneration is finely regulated by the interaction of multiple signaling pathways, such as Notch signaling (Ma et al., 2008; Mizutari et al., 2013), Wnt/b-catenin signaling (Aman and Piotrowski, 2008; Chai et al., 2012; Shimizu et al., 2012), Fgf signaling (Aman and Piotrowski, 2008; Nechiporuk and Raible, 2008), retinoic acid (Rubbini et al., 2015) and so on. In the process of hair cell damaged, it is accompanied by a lot of inflammatory reaction, which has been found to play a role in tissue regeneration in recent years (Mescher, 2017). For example, macrophages are considered having main function in the inflammatory resolution stage and being required for fin regeneration (Li et al., 2012) and hair cell regeneration in zebrafish (Carrillo et al., 2016). In addition, it has been confirmed that neutrophils in mice play a central role in inflammation-induced optic nerve regeneration (Kurimoto et al., 2013).

In recent years, zebrafish (Danio rerio) has become an ideal model for studying inflammation and hair cell regeneration because it has conservative

innate immunity (Renshaw and Trede, 2012) and strong regeneration ability in

lateral line system (Lush and Piotrowski, 2014) which makes zebrafish larvae

to perceive the change of surrounding flow, detect their prey and avoid

predators (Coombs. et al., 2014; Dijkgraaf, 1962). The lateral system of a larva

is composed of neuromasts which located on the surface of the body.

The neuromasts on the head consist of the anterior lateral line system (aLL)

and the ones along the body comprise the posterior lateral line system

(pLL)(Thomas et al., 2015). The center of the neuromast is composed of hair

cells and they are surrounded by support cells and mantle cells. At the top of

the hair cells, rows of short stereocilia and a long kinocilium extend out of the

body called the hair bundle and are covered in a gelatinous cupula. The

arrangement of stereocilia and kinocilium determines the polarity of hair cells

and the polarity of the hair cells is planar cell polarity (PCP), which is arranged

symmetrically (Flock and Wersall, 1962), half in each direction.

When hair bundles are deflected, hair cells release transmitters and cause

exciting spikes in afferent neurons (Dijkgraaf, 1962). And then, larvae show a

robust behavior called rheotaxis (Olszewski et al., 2012). This behavior can be

applied to evaluate the function of hair cells (Suli et al., 2012).

In recent years, calcium imaging has become a popular method to

measure the function of neural cells in detail and quantitatively (Zhang et al.,

2016). When the mechanical hair bundle deflected, calcium and other cations

enter into cytoplasm through mechanotransduction channels. It changes the

membrane potential and activates voltage-gated calcium channels which allow

rapid calcium inflow to trigger synaptic transmission. GCaMPs, a

genetically-encoded calcium indicator(GECIs), are single fluorescent proteins,

which can bind calcium directly and alter conformation to respond the change

of calcium concentration (Tian et al., 2012). These significant,

activity-dependent signals can reflect the function of hair cells in a single

neuromast (Zhang et al., 2018; Zhang et al., 2016).

Previous research has found that the deletion of macrophages by

morpholino leads to the delay of hair cell regeneration (Carrillo et al., 2016).

However, does it still cause the delay of hair cell regeneration when the

macrophages are intact, and the pro-inflammatory factors are suppressed as

the hair cells are damaged? Is there any delay in the functional recovery of the

lateral line?

In order to figure out the above problems, we used an anti-inflammatory

agent, BRS-28, to suppress the inflammation when hair cells are damaged by

copper. BRS-28 is a derivative of 5α-cholestan-6-one, which was confirmed to

be a remarkably suppressor of the production of pro-inflammatory factors,

such as NO, TNF-α、IL-1β、iNOS and cox-2 (Yang et al., 2014). We count the

number of neutrophils and macrophages in Tg(corola-eGFP; lyz-Dsred)

transgenic line. Then, AB/WT zebrafish larvae were used to count the number

of regenerated hair cells. Since there is no appropriate behavioral method to

quantitatively evaluate the function of lateral line hair cells, we designed and

built devices to test rheotaxis behavior in AB/WT larvae. A behavioral analysis

software was applied for quantitative evaluation of rheotaxis, so as to reflect

the holistic functional recovery of the posterior lateral line. Finally, the function

of the regenerated hair cells in a single neuromast was evaluated by the

method of calcium imaging in Huc:h2b-gcamp6f transgenic line.

## Results

**$CuSO_4$ damaged hair cells in lateral line of zebrafish.**

Sensory hair cells in a 6-day post fertilization (dpf) AB/WT zebrafish larva

were labeled with 0.05% DASPEI clearly (**Fig. 1A**). L2、LII3、L3 neuromasts

(circles in **Fig.1 A**) were three of the posterior lateral neuromasts, which

located along the flat truck body and easily to be observed. A lateral view of the

neuromasts showed the elongated kinocilia extending from the body (**Fig. 1B**).

The neuromasts are consisted of hair cells surrounded by support cells, which

are surrounded by mantle cells (**Fig. 1C**). In order to study the effects of

inflammation on hair cell regeneration, we established a hair-cell-damaged

model. Hair cells were damaged completely, when treated with 5 µM $CuSO_4$

for 1 h (**Fig. 1D**). Labeled with 0.05% DASPEI, hair cells displayed close

arrangement and clear boundary. Only treated with $CuSO_4$ solution for 20 min,

hair cells became loose and unclear which suggested that they were already

injured. The number of hair cells decreased with weaker fluorescence intensity

and obscure cell boundary at 40 min. Hair cells were completely disappeared

at 60 min, indicating that they had been completely damaged. TUNEL assay

revealed the missing hair cells underwent apoptosis (**Supplementary Fig. 1**).

After being transferred to embryo medium (EM), the number of hair cells

quickly returned to normal (**Fig. 1E**).

**BRS-28 reduced the number of neutrophils and macrophages migrating**

**to the injured neuromasts.**

Neutrophils (**Fig. 2B, C**, blue arrows) and macrophages (**Fig. 2B, C**, white

arrows) could be marked and distinguished in larvae of Tg(corola-eGFP;

lyz-Dsred) transgenic line (**Supplementary Fig. 2**). Normally, neutrophils and

macrophages were almost absent from the neuromasts (example, **Fig. 2A**).

When treated with $CuSO_4$ solution, hair cells were damaged. Neutrophils and

macrophages migrated to the neuromasts within 1 hours (example, **Fig. 2B**).

When larvae were immerged in BRS-28, an anti-inflammatory agent, before

treated with $CuSO_4$ solution, less neutrophils and macrophages migrated to

the damaged neuromasts (example, **Fig. 2C**). When the inflammation

suppressed, the numbers of neutrophils appeared around the damaged

neuromasts were lower at 0.5,1,3 and 4 h after adding the $CuSO_4$ solution in

BRS+CuSO$_4$ group than in CuSO$_4$ group (**Fig. 2D**). In addition, we observed

BRS+CuSO$_4$ group had fewer macrophages at 0.5, 1, 2 and 3 h than CuSO$_4$

group (**Fig. 2E**). Collectively, the data strongly suggested that BRS-28 reduced

the number of neutrophils and macrophages migrating to the injured

neuromasts. It was worth noting that compared with control, there was no

significant difference in the numbers of neutrophils and macrophages between

CuSO$_4$ group and BRS+CuSO$_4$ group at 5 and 6 h, indicating that the

inflammation was almost resolved.

**Suppressing inflammation delayed hair cell regeneration.**

In order to investigate whether the regeneration of hair cells were delayed

after suppressing inflammation, we observed hair cells in the L2, LII3 and L3

neuromasts. We found that the regeneration of hair cells was delayed after the

inflammation was suppressed by the inflammatory inhibitor, BRS-28. Live

imaging showed regenerated hair cells in CuSO$_4$, BRS+CuSO$_4$ group at 24, 48

and 96 hours post injured (hpi) by CuSO$_4$(**Fig. 3A**). Control group was showed

at the same time point. Further analysis revealed that the numbers of

regenerated hair cells were significantly decreased in BRS+CuSO$_4$ group than

that in CuSO$_4$ group at 16 hpi (P=0.0061), 24 hpi (P=0.0021) and 48 hpi

(P<0.0001) (**Fig. 3B**, n = 30 neuromasts). These results indicated that the

regeneration of hair cells was delayed in BRS+CuSO$_4$ group within 48 hpi.

Compared with Control group, there was no difference in the number of hair

cells between CuSO$_4$ group and BRS+CuSO$_4$ group at 96 hpi, suggesting that

hair cells were regenerated to the normal level at 96 hpi. We also analyzed the

number of hair cells when only treated with BRS-28 (BRS group) without hair

cell damage. As expected, BRS group had no difference compared with

Control group at any time point, excluding the effect of BRS-28 on hair cells.

Since hair cells did not regenerate at a uniform rate, we defined the time

of regeneration into two periods: the Early Stage which includes the time from

to 48 hpi and the Late Stage which includes the time after 48 hpi. The regeneration of hair cells was fast in the Early Stage and slow in the Late Stage. Linear analysis was conducted on the number of hair cell regeneration in the Early Stage. The slope in $CuSO_4$ group (0.1879) was higher than that in BRS+$CuSO_4$ group (0.148) , meanwhile, x-intercept in $CuSO_4$ group (4.16) was higher than that in BRS+$CuSO_4$ group (8.287) (**Fig. 3C, D**). These implied that the hair cell regeneration in BRS+$CuSO_4$ group may begin later and slower than that in $CuSO_4$ group.

To explore whether the time window of inflammatory suppression had contribute to delayed regeneration, we changed the start time of BRS-28 treatment. We found that compared with the $CuSO_4$ group, whether BRS-28 was added at the same time as $CuSO_4$ ($CuSO_4$+BRS 0 h group), or 30 minutes after the addition of $CuSO_4$ ($CuSO_4$+BRS 0.5 h group), or 1 hour after the addition of $CuSO_4$ ($CuSO_4$+BRS 1 h group) (**Fig. 3E**), there was no statistical difference on the number of regenerated hair cells.

To sum up, the regeneration of hair cells in lateral line was delayed after the inflammation was suppressed by the inflammatory inhibitor BRS-28.

**The functional recovery of the lateral line system was delayed when inflammation was suppressed.**

Since the rheotaxis could be a suitable functional readout of the lateral line, we designed a behavioral device to test the rheotaxis of zebrafish (**Fig. 4A**, see details in Materials and Methods). Larvae were placed from the right platform, and they sense the water flow direction from the right to the left. **Figure 4B, C** were two examples of the larval rheotaxis processed by behavioral analysis software: the former was a larva with excellent rheotaxis (**Fig. 4B**) while the latter was a larva performed failure in the rheotaxis test (**Fig. 4C**). The left panels in these two examples showed the swimming track of this larva. The behavioral analysis software mapped its movement path of larvae

by line segment. The color of the line segment represented the direction of movement of the larvae. All the movements from right to left were represented by purplish or red segments, where purple indicated that the velocity along the flow direction was greater than or equal to the flow velocity, and red indicated that the velocity along the flow direction was less than the flow velocity. All the movements from left to right were represented by green segments, and the higher the brightness was, the faster the speed was. The right panels displayed the motion vector. The lengths of the blue segments represented the distance of each movement, and the direction of the blue segment represented the direction of that movement. The length of the red line segment was the ratio of motion vectors sum to the motion arithmetic sum and the direction was the direction of the sum of the vectors.

When the red segment was long and had a small angle of 0 degree, it indicated that the motion of the larva was consistent with the opposite direction of flow. It represented that the larva had a good rheotaxis, indicating its lateral line system executed its function very well. Therefore, the software reported a high score. On the contrary, when the red segment was short or had a small angle of 180 degree, it indicated that the larva moved randomly and had a poor rheotaxis, indicating its lateral line system had poor function. In this case, the software reported a low score. The scores reported by the software were plotted into bar charts and showed in **Figure 4D**. After the hair cells were damaged by $CuSO_4$, there was poor rheotaxis in both $CuSO_4$ group and BRS+$CuSO_4$ group. At 24 and 48 hpi, the rheotaxis of BRS+$CuSO_4$ group was significantly lower than that of Control group. On the contrary, the rheotaxis of $CuSO_4$ group was not significantly different from that of Control group within 24 hpi. Therefore, it indicated that the functional recovery of lateral line system was delayed in BRS+$CuSO_4$ group. The rheotaxis of BRS group at each time point was not different from that of Control group, suggesting that BRS-28

alone had no significant effect on the rheotaxis. In addition, we noted that the speed and distance of each movements were consistent within different times and between different groups: both were stable at around 22 mm/s (**Fig. 4E, F**), which indicated that BRS-28 or $CuSO_4$ did not affect the movement of zebrafish.

We concluded that the regenerated hair cells still had the ability to sense water flow, but the functional recovery of lateral line system was delayed when inflammation was suppressed.

**Calcium imaging revealed the function of a single neuromast after hair cell regeneration**

Since we found a mismatch between the function of the lateral line and the amount of hair cell regeneration, that is, after the zebrafish lateral line was damaged by copper sulfate, it took 96 h for the hair cells to return to normal, while the flow ability returned to normal at 24 h. The function of a single neuromast can be evaluated by observing its calcium activity (Zhang et al., 2016). The L3 neuromast, located in flat trunk, was stimulated by water flow from an electrode (**Fig. 5A**). Since hair cells had polarities, the yellow and green hair cells represented opposite polarities. Chou et al. reported that the polarity of the L3 neuromast is parallel to the anterior-posterior body axis (Chou et al., 2017). Thus, by adjusting the direction of the electrode, water was controlled to flow in two directions: anterior to posterior (A-P) direction or posterior to anterior (P-A) direction. We found that not all hair cells responded to the water flow, and only some hair cells were active (example, **Fig. 5B**, circled cells). These active cells only responded to stimulus in one direction: P-A direction (**Fig. 5C**, yellow ones and yellow circles in **Fig. 5B**) or A–P direction (**Fig. 5D**, green ones and green circles in **Fig. 5B**). Because the neuromasts were stereoscopic, some of the active hair cells were far from this focal plane (dashed circles in **Fig. 5B**) and were not included in subsequent

fluorescence intensity analysis.

Similar to the results of the rheotaxis, the fluorescence intensity ($\Delta F/F$) of

the regenerated hair cells were reduced significantly when inflammation was

suppressed at the Early Stage of regeneration (within 48 hpi)(**Fig. 5E**). It was

noteworthy that compared to Control group, the fluorescence intensity in

$CuSO_4$ group did not decrease significantly in the Early Stage of regeneration.

This might explain that why the rheotaxis in $CuSO_4$ group had been recovered

at 24 hpi. The fluorescence intensity of BRS+$CuSO_4$ group was not

significantly different from that of Control group and $CuSO_4$ group in the Late

Stage of regeneration (72-96 hpi) (**Fig. 5F**). Additionally, the fluorescence

intensity showed no differences between the BRS group and Control group

(**Fig. 5G**), indicating that BRS-28 had no effect on the fluorescence intensity.

Normally, only a part of the hair cells in the neuromast responds to the

stimulation of water flow. Is it the same for regenerated hair cells? We found

that only a few regenerated hair cells in $CuSO_4$ group and BRS+$CuSO_4$ group

responded to flow stimuli. The number of active cells in each neuromast in

these two group were approximately the same at 24 to 96 hpi, and were

consistent with that in Control group (**Fig. 5H**).

Furthermore, we noticed that most hair cells that responded to the flow in

the opposite direction came in pairs (**Supplementary Fig. 3A**). Although the

numbers of hair cells responding to flow in P-A direction were similar to that in

A-P direction, the fluorescence intensity of hair cells responding to P-A

direction was significantly higher than that of hair cells responding to A-P

direction (**Supplementary Fig. 3B**). It indicated that L3 neuromast was more

sensitive to the flow from the P-A direction.

The results altogether demonstrated that the recovery of hair cell function

was delayed at the Early Stage of regeneration when inflammation was

suppressed.

## Discussion

**BRS-28 suppresses inflammation and delays the initiation of hair cell regeneration.**

Although the downregulation of Notch signal during lateral line regeneration induces the proliferation of support cells by activating Wnt/b-Catenin signal (Romero-Carvajal et al., 2015), it is still unknown how the downregulation of Notch signal is triggered after hair cell death. Kniss et al. propos a hypothesis of triggering hair cell regeneration (Kniss et al., 2016). Studies in Drosophila wing disc and eye have found that JNK, Shh, EGF, and TNF signaling pathways are required during apoptosis-induced compensatory proliferation (Fan et al., 2014; Perez-Garijo et al., 2009; Ryoo et al., 2004). Kniss et al. assume that a similar process may be involved in the regeneration of hair cells. On the basis of this hypothesis, we speculate that when hair cells are damaged by $CuSO_4$, it cause apoptosis in lateral line hair cells, trigger the rise of reactive oxygen species (ROS) and reactive nitrogen species (RNS), and induce the oxidative stress. This process may improve AP-1,HIF-1α and NF-κB activity, and thus increase pro-inflammatory cytokines and chemokines, such as NO, IL-1β, TNF-α, cox-2,iNOS and so on (Pereira et al., 2016). BRS-28, suppress the production of NO、IL-1β、TNF-α、cox-2、iNOS (Yang et al., 2014), reducing the number of neutrophils and macrophages migrating to the damage of neuromasts. Besides that, the decrease of pro-inflammatory factors may reduce the activation of macrophages. These processes would decrease the production of TNF ligands and inhibit the JNK signal, which contributes to initiating cells regeneration, and eventually leads to delay initiation of compensatory proliferation and delay regeneration of hair cells.

We found that when the initiate time of inflammatory inhibitors was changed, there was no delay in hair cell regeneration (**Fig. 3E**). This also

suggests that the timing of inflammation suppression is important: when

inflammation occurs, compensatory proliferation of the support cells is

triggered and hair cells begin to regenerate. If inflammation suppression does

not take effect, regeneration seems to be unaffected.

In addition, neutrophils can also remove dead cell debris, and

macrophages can phagocyte apoptotic neutrophils or fragments of dead cells.

We believe that when the number and activity of neutrophils and macrophages

decrease, the clearance of damaged tissue areas slows down, and hair cells

cannot obtain a good regeneration environment. Since damaged neuromasts

need more time to clean up these cell fragments, this may also delay the

regeneration of hair cells.

**Suppression of inflammation delays functional recovery of regenerated**

**hair cells.**

In this study, we found that when inflammation was suppressed, hair cell

regeneration was delayed, as was the recovery of function. Finally, the quantity

and the function of hair cells returned to normal level at the Late Stage of

regeneration. Therefore, although the suppression of inflammation delayed the

regeneration of hair cells, it did not affect the overall process of hair cell

regeneration, and the function of regenerated hair cells eventually tended to be

intact. However, the effect of inflammation on the regeneration of lateral hair

cells seems to be different from that of the fin. Li et al. found that when

zebrafish larvae lack macrophages, vacuoles appear in the regenerated fin,

suggesting that macrophages may also be involved in fin regeneration (Li et al.,

2012). In our research, although the suppression of inflammation delay

regeneration of hair cells and their functional recovery at the Early Stage of

regeneration, they eventually return to the normal status at the Late Stage of

regeneration. This is not because inflammation is not suppressed sufficiently,

as Carrillo et al. found that the number of hair cells finally completed

regeneration even when macrophages is knockout (Carrillo et al., 2016).

However, this may be because the injured organs are different, and the intact

function of lateral hair cells is crucial for the survival of zebrafish. It is

suggested that the hair cells in lateral line may have more complex regulation

mechanisms during the regeneration process.

**The functional recovery of hair cells is much faster than its quantity.**

Previous studies have focused on the morphological and quantitative

recovery of regenerated hair cells in zebrafish (Carrillo et al., 2016;

Romero-Carvajal et al., 2015). Since the regeneration takes 3-4 days post

injured, it is easy to assume that the recovery of the function of the neuromasts

may be proportional to the number of regenerated hair cells. In this study, for

the first time, we performed a method to evaluate the function of regenerated

hair cells. We found that the $CuSO_4$ group already performed excellent

rheotaxis at 24 hpi (**Fig. 4C**), even though the average number of hair cells

was only 3.667 at that time (**Fig. 3B**). Therefore, hair cells recover the function

much more quickly than their numbers. In other words, although it takes 72-96

h to complete regeneration, the function of hair cells can be recovered within

24 hours which is critical for the survival of zebrafish. When BRS-28 is used to

suppress the inflammation, the amplitude of calcium activity of hair cells is

significantly lower than that of Control and $CuSO_4$ group at the Early Stage of

regeneration, and the rheotaxis of larvae is poor during this period. Therefore,

the suppression of inflammation not only delays the hair cell regeneration, but

also delays the functional recovery.

There is a mismatch between the function and quantity during

regeneration. Calcium image reveal that only a part of regenerated hair cells in

one neuromast respond to the flow. This finding is consistent with previous

study (Zhang et al., 2018). In our research, we found that this phenomenon

also exists in regeneration group ($CuSO_4$ and BRS+$CuSO_4$ group).

Regardless of the number of regenerated hair cells, the number of hair cells

that respond to water flow remain stable during the regeneration process,

which is not different from Control group (**Fig. 5H**). Besides that, in the Early

Stage of regeneration, the magnitude of fluorescence intensity and reaction

time of $CuSO_4$ group are also consistent with that of the controls. This explains

why the number of regeneration in the $CuSO_4$ group at 24 h is only 3.667 on

average, but the function of the lateral line has been restored to a level very

close to that of Control group.

In this study, we only performed calcium imaging on the L3 neuromast,

which was confirmed as the polarity of the A-P body axis in the study of Chou

et al (Chou et al., 2017). Consistent with their results, this neuromast is indeed

insensitive to the flow in the dorsal-ventral (D-V) body axis (data not shown).

Therefore, this study only focuses on the stimulus response in the A-P body

axis direction, and does not further analyze the stimulus data in the D-V body

axis direction. Compared with hair cells with polarity in the A-P direction, hair

cells with polarity in the P-A direction have greater $\Delta F/F0$ when stimulated by

water flow (**Supplementary Fig. 3B**; sample, **Fig. 5 C, D**). It indicated that L3

neuromast is more sensitive to the flow from the P-A direction. This finding is

consistent with the results measured by Chou et al. using microphonic

potentials evoked by sinusoidal stimuli (Chou et al., 2017). Most active hair

cells that responded to the opposite flow come in pairs (**Supplementary Fig.**

**3A**), suggesting that it appears to be pre-arranged rather than random.

In summary, our research suggests that suppression of inflammation

delays functional regeneration of lateral hair cells in zebrafish larvae. The

inflammation plays positive and permissive roles in hair cell regeneration.

## Materials and Methods

**Zebrafish strains and maintenance**

AB/Wild-type strain, Tg(corola-eGFP;lyz-Dsred) and Huc:h2b-gcamp6f

transgenic line were used in this study. Embryos were generated by paired

mating and maintained at 28.5℃ in EM and on a 14/10 h light/dark cycle

according to the standard protocols.

All animal manipulations were conducted strictly in accordance with the

guidelines and regulations set forth by the University of Science and

Technology of China (USTC) Animal Resources Center and the University

Animal Care and Use Committee. The protocol was approved by the

Committee on the Ethics of Animal Experiments of the USTC (Permit Number:

USTCACUC1103013).

**Hair cell damage and inflammation inhibition**

In order to damage hair cells in lateral line, 4 dpf Larvae were treated with

5 µM $CuSO_4$ (Sangon, China) diluted in embro medium (EM) for 1 h. Then,

they were washed three times and recovered in EM.

To suppress inflammation, 4 dpf larvae were immersed in 0.1% BRS-28,

an anti-inflammatory agent, for 3 h before being moved into $CuSO_4$ to damage

hair cells.

**Live imaging**

AB/Wild-type larvae were used to count the number of regenerated hair

cells in L2、LII3、L3 neuromasts (**Fig. 1A**). Hair cells were marked by

0.01 %DAPI for 5 minutes. Larvae were anesthetized in 0.02% MS-222

(Tricaine mesylate, Sigma, USA) and imaged under a fluorescence

microscope (Olympus BX-60, Japan).

In order to exhibit the damage of hair cells in copper sulfate solution and

the regeneration of hair cells in different phases, hair cells were labeled by

0.05 % DASPEI (Sigma, USA), and larvae were anesthetized in MS-222 and

imaged under a confocal microscopy (Zeiss LSM 880 +Airyscan, Germany).

Tg(corola-eGFP; lyz-Dsred) transgenic line was used to observe the

number of neutrophils and macrophages migrating to the injured neuromasts

*in vivo*. In this transgenic line, neutrophils co-expressed *lyz-Dsred* and

*coro1a-GFP* and show yellow fluorescence after these two channels are

merged, while macrophages only express *coro1a-GFP* and show green

fluorescence (Li et al., 2012). To show the neutrophils and macrophages

migrating to damaged neuromasts, larvae were anesthetized in MS-222 and

imaged under a confocal microscopy (Zeiss LSM 880 +Airyscan). In order to

count neutrophils and macrophages, we set the area around the L2 LII3 L3

neuromasts with a diameter of 100 μm as the region of interest (ROI).

Zebrafish larvae were anesthetized and imaged by the fluorescence

microscope (Olympus BX-60) with a green and a red channel.

**Rheotaxis behavior experiments**

A U-shaped tank was designed to test the rheotaxis behavior of larvae

(**Fig. 4A**). The bottom of the two cubic tanks (7 cm length *8 cm width*8 cm

height) were connected by a platform (10 cm length *8 cm width*0.5 cm height).

A peristaltic pump (Longer Pump YZ1515x, China) was used to move EM

solution from the left tank to the right tank, so that the platform formed a steady

water flow from right to left (v=10 mm/s). AB/WT zebrafish larvae were applied

to detect the ability of rheotaxis. Larvae were released at the right side of the

platform with an initial velocity almost equals 0. To avoid visual cues,

experiments were operated in the dark and rheotaxis performs were recorded

by an infrared CCD (IR850, weixinshijie, China).

Rheotaxis data were analyzed by our own rheotaxis software edited in

Matlab (2015a, MathWorks, USA). This software can plot the movement track

of zebrafish larvae in the platform, measure the direction and distance of each

swimming and calculate the speed. Finally, it reports scores based on the

magnitude in the horizontal direction of the ratio of motion vectors sum to the

motion arithmetic sum.

### Calcium imaging and data analyses

Huc: h2b-GCamp6f transgenic line was used in calcium imaging which

expressed pan-neuronal nucleus-labelled GCamp6f. Larvae were

anesthetized and fixed by a net pressure. The one-step pulled micropipette

had a long, wispy tip which must be trimmed by rubbing it against with another

pulled micropipette to generate a tip with an outer diameter of approximately

40 µm. The micropipette was filled with 0.02% MS-222 and fixed to the holder

of a micromanipulator (MX7500, Scientific Design Company, USA). The tip of

the micropipette was positioned at a distance of approximately 100 µm from

the top of the kinocilia (**Fig. 5A**). The duration of flow was controlled by

three direct links which were linked with a syringe.

Calcium imaging was collected by a confocal microscopy (FV 1000,

Olympus, Japan). To make as many hair cells as possible in the observation

area at the same time, a single z-axis was adjusted. ROI was set to 110*108.

We took 100 time-lapse images for each neuromast, and the total capture time

was 29.7 s (0.297 s per slice). Flow stimulation occurred from 10.098 to 19.899

s.

Since the neuromasts are three-dimensional, different hair cells have

different levels of fluorescence intensity. Namely, they have different levels of F

prime. The relative fluorescence intensity change ($\Delta F/F_0$) is more commonly

used. For each hair cell, the average fluorescence intensity before flow stimuli

(0-10 s) was set as $F_0$. The data would be excluded when $F_0<95$, which means

these hair cells were too far from the focal plane. When more than two hair

cells in the neuromast respond to flow stimulation, two hair cells with the

strongest fluorescence were selected and included in the statistics of

fluorescence intensity curve.

### Statistical analysis

500     All data were shown as mean ± S.E.M. or as relative proportions of 100 %

as indicated in the appropriate legends. The data were analyzed in either one-way ANOVA with Tukey's multiple comparisons test or two-way ANOVA with Tukey's multiple comparisons test by GraphPad Prism version 7.0 (Prism, San Diego, CA, USA). The level of significance was set to P < 0.05. *, **and ***represent P < 0.05, P < 0.01 and P < 0.001, respectively.

## Acknowledgments

The authors thank Drs. Wen Zilong for providing the Tg(corola-eGFP; lyz-Dsred) transgenic fish line, Drs. Wen Quan for providing the Huc:h2b-gcamp6f transgenic fish line. The authors thank Drs. Zhen Xuechu for providing BRS-28 and the compound-26 in their study is the BRS-28 mentioned in this study.

## References

Aman, A., and Piotrowski, T. (2008). Wnt/beta-catenin and Fgf signaling control collective cell migration by restricting chemokine receptor expression. Developmental cell *15*, 749-761.

Baird, R.A., Steyger, P.S., and Schuff, N.R. (1996). Mitotic and nonmitotic hair cell regeneration in the bullfrog vestibular otolith organs. Annals of the New York Academy of Sciences *781*, 59-70.

Carrillo, S.A., Anguita-Salinas, C., Pena, O.A., Morales, R.A., Munoz-Sanchez, S., Munoz-Montecinos, C., Paredes-Zuniga, S., Tapia, K., and Allende, M.L. (2016). Macrophage Recruitment Contributes to Regeneration of Mechanosensory Hair Cells in the Zebrafish Lateral Line. Journal of cellular biochemistry *117*, 1880-1889.

Chai, R., Kuo, B., Wang, T., Liaw, E.J., Xia, A., Jan, T.A., Liu, Z., Taketo, M.M., Oghalai, J.S., Nusse, R.*, et al.* (2012). Wnt signaling induces proliferation of sensory precursors in the postnatal mouse cochlea. Proceedings of the National Academy of Sciences of the United States of America *109*, 8167-8172.

Chou, S.W., Chen, Z., Zhu, S., Davis, R.W., Hu, J., Liu, L., Fernando, C.A., Kindig, K., Brown, W.C., Stepanyan, R.*, et al.* (2017). A molecular basis for water motion detection by the mechanosensory lateral line of zebrafish. Nature communications *8*, 2234.

Coombs., S., Bleckmann., H., Fay., R.R., and Popper, A.N. (2014). The Lateral Line System (New York: Springer).

Dijkgraaf, S. (1962). The functioning and significance of the lateral-line organs. Biological Reviews of the Cambridge Philosophical Society *38: 51-105*.

Fan, Y., Wang, S., Hernandez, J., Yenigun, V.B., Hertlein, G., Fogarty, C.E., Lindblad, J.L., and

Bergmann, A. (2014). Genetic models of apoptosis-induced proliferation decipher activation of JNK
and identify a requirement of EGFR signaling for tissue regenerative responses in Drosophila. PLoS
genetics *10*, e1004131.

Flock, A., and Wersall, J. (1962). A study of the orientation of the sensory hairs of the receptor
cells in the lateral line organ of fish, with special reference to the function of the receptors. The
Journal of cell biology *15*, 19-27.

Kniss, J.S., Jiang, L., and Piotrowski, T. (2016). Insights into sensory hair cell regeneration from
the zebrafish lateral line. Current opinion in genetics & development *40*, 32-40.

Kurimoto, T., Yin, Y., Habboub, G., Gilbert, H.Y., Li, Y., Nakao, S., Hafezi-Moghadam, A., and
Benowitz, L.I. (2013). Neutrophils express oncomodulin and promote optic nerve regeneration. The
Journal of neuroscience : the official journal of the Society for Neuroscience *33*, 14816-14824.

Li, L., Yan, B., Shi, Y.Q., Zhang, W.Q., and Wen, Z.L. (2012). Live imaging reveals differing roles of
macrophages and neutrophils during zebrafish tail fin regeneration. The Journal of biological
chemistry *287*, 25353-25360.

Lopez-Schier, H., and Hudspeth, A.J. (2006). A two-step mechanism underlies the planar
polarization of regenerating sensory hair cells. Proceedings of the National Academy of Sciences of
the United States of America *103*, 18615-18620.

Lush, M.E., and Piotrowski, T. (2014). Sensory hair cell regeneration in the zebrafish lateral line.
Developmental dynamics : an official publication of the American Association of Anatomists *243*,
1187-1202.

Ma, E.Y., Rubel, E.W., and Raible, D.W. (2008). Notch signaling regulates the extent of hair cell
regeneration in the zebrafish lateral line. The Journal of neuroscience : the official journal of the
Society for Neuroscience *28*, 2261-2273.

Matsui., J.I., and Cotanchea., D.A. (2004). Sensory hair cell death and regeneration two halves
of the same equation. Hearing Science.

Mescher, A.L. (2017). Macrophages and fibroblasts during inflammation and tissue repair in
models of organ regeneration. Regeneration *4*, 39-53.

Mizutari, K., Fujioka, M., Hosoya, M., Bramhall, N., Okano, H.J., Okano, H., and Edge, A.S.
(2013). Notch inhibition induces cochlear hair cell regeneration and recovery of hearing after
acoustic trauma. Neuron *77*, 58-69.

Nechiporuk, A., and Raible, D.W. (2008). FGF-Dependent Mechanosensory Organ Patterning in
Zebrafish. Science *320*, 1774-1777.

Oesterle, E.C., and Stone, J.S. (2008). Hair Cell Regeneration: Mechanisms Guiding Cellular
Proliferation and Differentiation. In Hair Cell Regeneration, Repair and Protection, RJ Salvi , A.
Popper, and R. Fay, eds. (New York: Springer), pp. 141-197.

Olszewski, J., Haehnel, M., Taguchi, M., and Liao, J.C. (2012). Zebrafish Larvae Exhibit Rheotaxis
and Can Escape a Continuous Suction Source Using Their Lateral Line. PloS one *7*, e36661.

Pereira, T.C., Campos, M.M., and Bogo, M.R. (2016). Copper toxicology, oxidative stress and
inflammation using zebrafish as experimental model. Journal of applied toxicology : JAT *36*, 876-885.

Perez-Garijo, A., Shlevkov, E., and Morata, G. (2009). The role of Dpp and Wg in compensatory
proliferation and in the formation of hyperplastic overgrowths caused by apoptotic cells in the
Drosophila wing disc. Development *136*, 1169-1177.

Popper, A.N., and Hoxter, B. (1984). Growth of a fish ear: 1. Quantitative analysis of hair cell
and ganglion cell proliferation. Hearing research *15*, 133-142.

Raphael, Y. (1992). Evidence for supporting cell mitosis in response to acoustic trauma in the
avian inner ear. J Neurocytol *21*, 663-671.

Renshaw, S.A., and Trede, N.S. (2012). A model 450 million years in the making: zebrafish and
vertebrate immunity. Disease models & mechanisms *5*, 38-47.

Roberson, D.W., Alosi, J.A., and Cotanche, D.A. (2004). Direct transdifferentiation gives rise to
the earliest new hair cells in regenerating avian auditory epithelium. Journal of Neuroscience
Research *78*, 461-471.

Romero-Carvajal, A., Navajas Acedo, J., Jiang, L., Kozlovskaja-Gumbriene, A., Alexander, R., Li,
H., and Piotrowski, T. (2015). Regeneration of Sensory Hair Cells Requires Localized Interactions
between the Notch and Wnt Pathways. Developmental cell *34*, 267-282.

Rubbini, D., Robert-Moreno, A., Hoijman, E., and Alsina, B. (2015). Retinoic Acid Signaling
Mediates Hair Cell Regeneration by Repressing p27kip and sox2 in Supporting Cells. The Journal of
neuroscience : the official journal of the Society for Neuroscience *35*, 15752-15766.

Ryoo, H.D., Gorenc, T., and Steller, H. (2004). Apoptotic cells can induce compensatory cell
proliferation through the JNK and the Wingless signaling pathways. Developmental cell *7*, 491-501.

Shimizu, N., Kawakami, K., and Ishitani, T. (2012). Visualization and exploration of Tcf/Lef
function using a highly responsive Wnt/β-catenin signaling-reporter transgenic zebrafish.
Developmental Biology *370*, 71-85.

Stone., J.S., and Rubel., E.W. (2000). Cellular studies of auditory hair cell regeneration in birds.
PNAS.

Suli, A., Watson, G.M., Rubel, E.W., and Raible, D.W. (2012). Rheotaxis in Larval Zebrafish Is
Mediated by Lateral Line Mechanosensory Hair Cells. PloS one *7*, e29727.

Thomas, E.D., Cruz, I.A., Hailey, D.W., and Raible, D.W. (2015). There and back again:
development and regeneration of the zebrafish lateral line system. Wiley Interdiscip Rev Dev Biol *4*,
1-16.

Tian, L., Hires, S.A., and Looger, L.L. (2012). Imaging neuronal activity with genetically encoded
calcium indicators. Cold Spring Harbor protocols *2012*, 647-656.

Yang, Y.X., Zheng, L.T., Shi, J.J., Gao, B., Chen, Y.K., Yang, H.C., Chen, H.L., Li, Y.C., and Zhen, X.C.
(2014). Synthesis of 5alpha-cholestan-6-one derivatives and their inhibitory activities of NO
production in activated microglia: discovery of a novel neuroinflammation inhibitor. Bioorganic &
medicinal chemistry letters *24*, 1222-1227.

Yorgason., J.G., Fayad., J.N., and Kalinec., F. (2006). Understanding drug ototoxicity: molecular
insights for preventionand clinical management. Biological Reviews.

Zhang, Q., Li, S., Wong, H.C., He, X.J., Beirl, A., Petralia, R.S., Wang, Y.X., and Kindt, K.S. (2018).
Synaptically silent sensory hair cells in zebrafish are recruited after damage. Nature
communications *9*, 1388.

Zhang, Q.X., He, X.J., Wong, H.C., and Kindt, K.S. (2016). Functional calcium imaging in zebrafish
lateral-line hair cells. Methods in cell biology *133*, 229-252.

**Fig. 1 CuSO₄ damaged hair cells in lateral line of zebrafish.**

**(A)** Lateral line hair cells in a 6 day post fertilization (dpf) AB/WT zebrafish larvae are labeled with 0.05% DASPEI. L2, LII3 and L3 neuromasts are marked with circles. Scale bar represents 500 µm.

**(B)** Lateral view of a neuromast shows sensory hair cells in the center labeled with DASPEI and a bundle of kinocilia (arrow) extend out of the periderm. Scale bar represents 50 µm.

**(C)** A cartoon illustrating the structure of the neuromast.

**(D)** Time lapse imaging shows that when merged in 5 µM CuSO₄ solution, hair cells were gradually injured and damaged within 60 min. Scale bar represents 10 µm.

**(E)** DASPEI staining displays that hair cells regenerate completely within 96 hours post injured (hpi). Scale bar represents 10 µm.

**Fig. 2 BRS-28 reduces the number of neutrophils and macrophages migrating to the injured neuromasts.**

(A-C) Live imaging (×40) displays the regions of L3 neuromasts of larvae at GFP channel, Dsred channel, and bright field (BF) channel and superimposed image in different group. Neutrophils (show both green and yollow fluorescence, indicated by white arrows) and macrophages (show olny green fluorescence, indicated by blue arrows) around the neuromasts can be observed in Tg(corola-eGFP; lyz-Dsred) larvae. They are almost absent from the neuromasts in Control group (A). Many neutrophils and macrophages migrate to injured neuromasts in CuSO4 group (B) while fewer neutrophils and macrophages migrate to injured neuromasts in BRS+CuSO4 group (C). The image is captured after adding CuSO4 solution for 1 h. Scale bar represents 50

μm.

**(D-E)** Line charts reveal decreased numbers of neutrophils **(D)** and

macrophages **(E)** within a radius of 50 μm from the center of neuromasts at

different time points after adding $CuSO_4$ in BRS+$CuSO_4$ group (n≥16) than

$CuSO_4$ group (n≥15). Control group (n≥11) is observed at the same time

points.

To **(D)** and **(E)**, comparisons were performed by using two-way ANOVA,

with Tukey's multiple comparisons test. All Error bars show mean ± S.E.M., ***

$P < 0.001$, **$P < 0.01$,*$P < 0.05$.

**Fig. 3  Suppressing inflammation delays hair cell regeneration.**

(A) Real-time imaging (×40) displays regenerated hair cells in the $CuSO_4$ and $BRS+CuSO_4$ group at 24, 48 and 96 hpi. Control group is taken at the same time point. Scale bar represents 10 μm.

(B) The numbers of regenerated hair cells were significantly decreased in $BRS+CuSO_4$ group than that in $CuSO_4$ group at 16 (P=0.0061), 24 (P=0.0021) and 48(P<0.0001) hpi. At 96 hpi, hair cells in both $CuSO_4$ group and $BRS+CuSO_4$ group regenerated to normal levels.

Linear analysis in $CuSO_4$ group (C) and $BRS+CuSO_4$ group (D) were conducted on the number of regeneration within 48 hours. The slope in $CuSO_4$ group (0.1879) is higher than that in $BRS+CuSO_4$ group (0.148) and x-intercept in $CuSO_4$ group (4.16) is higher than that in BRS+CuSO4 group (8.287).

(E) When delay the time window of inflammatory suppression, there is no delay in the regeneration of hair cells. BRS-28 was added at the same time as $CuSO_4$ ($CuSO_4$+BRS 0 h group), or 30 minutes after the addition of $CuSO_4$

678     ($CuSO_4$+BRS 0.5 h group), or 1 hour after the addition of $CuSO_4$ ($CuSO_4$+BRS

679     1 h group)(n≥27 neuromasts in each time point of each group).

680        To **(B)** and **(E)**, comparisons were performed by using two-way ANOVA,

681     with Tukey's multiple comparisons test. All Error bars show mean ± S.E.M., ***

682     $P < 0.001$, **$P < 0.01$,*$P < 0.05$.

**Fig.4 The recovery of the functional of lateral line system was delayed when inflammation was suppressed.**

(A) A U-shaped tank was designed to test the rheotaxis behavior of larvae. A peristaltic pump was used to form flow at the bottom of the tank. Larvae were placed from the right platform, and they sense the water flow from right to left. Rheotaxis perform was recorded by an infrared CCD.

A larva with excellent rheotaxis (B) and a larva with poor rheotaxis (C) were analyzed by behavioral analysis software. Moving traces were plotted in left panels and the motion vector were displayed in right panels. The lengths of the blue segments represented the distance of each movement, and the direction of the blue segment represented the direction of that movement. The length of the red line segment was the ratio of motion vectors sum to the motion arithmetic sum and the direction was the direction of the sum of the vectors.

(D) Rheotaxis score revealed that at 24 and 48 hpi, the rheotaxis of

BRS+CuSO$_4$ group was significantly lower than that of Control group. On the

contrary, the rheotaxis of CuSO$_4$ group was not significantly different from that

of Control group within 24 hpi.

The speed **(E)** and distance **(F)** of larvae swimming at each time were

consistent within different times and between different groups.

To **(D-F)**, comparisons were performed by using two-way ANOVA, with

Tukey's multiple comparisons test. All Error bars show mean ± S.E.M., $**P <$

$0.01, *P < 0.05$.

**Fig.5 Calcium imaging revealed the function of a single neuromast after hair cell regeneration**

(A) Schematic diagram shows an electrode filled with fluid is located about 100 μm away from the top of kinocilia to stimulate the neuromast. The yellow and green hair cells represent different polarities.

(B) When stimulated by the flow, only a part of hair cells respond in this focal plane (circled cells), and some are far from this focal plane (dashed circled cells). The No. 2, 4, and 6 active hair cells (yellow circles) only respond to the flow in P-A direction (C). At the same time, the No. 1, 3, and 5 active hair cells   (green circles) only respond to the flow in A-P direction (D). Scale bar in (B) represents 10 μm.

(E) The fluorescence intensity (ΔF/F) of the BRS + $CuSO_4$ group is significantly lower than that of the $CuSO_4$ group in the Early Stage of regeneration (within 48 hpi)( $P < 0.001$).

(F) The ΔF/F of BRS+$CuSO_4$ group is not significantly different from that of Control group and $CuSO_4$ group in the Late Stage of regeneration (72-96 hpi)

(G) There is no difference in ΔF/F between the BRS group and the Control group.

(H) During the regeneration process, the number of active hair cells in $CuSO_4$ and BRS+$CuSO_4$ group is basically the same, and did not increase

732  with the total number of regenerated hair cells.

733      To **(E-H)**, comparisons were performed by using one-way ANOVA, with

734  Tukey's multiple comparisons test.

**Supplementary Fig. 1 CuSO$_4$ caused apoptosis in hair cells.**

TUNEL assay revealed hair cells occurred apoptosis when treated with CuSO$_4$. Nuclei were stained with DAPI. BF: Bright Field. Scale bar represents 20 μm.

**Supplementary Fig. 2 Tg(corola-eGFP; lyz-Dsred) transgenic line could mark both neutrophils and macrophages.**

In Tg(corola-eGFP; lyz-Dsred) transgenic line, neutrophils co-expressed *lyz-Dsred* and *coro1a-GFP* and show yellow fluorescence after these two channels are merged, while macrophages only express *coro1a-GFP* and show green fluorescence. Scale bar represents 200 μm.

752
753
754

755
756 **Supplementary Fig. 3 Most active hair cells are polar in pairs and are**
757
758 **sensitive to flow in the P-A direction.**

759 **(A)** Most hair cells that responded to the flow in the opposite direction
760 come in pairs.

761 **(B)** The fluorescence intensity of hair cells responding to P-A direction is
significantly higher than that of hair cells responding to A-P direction.

762 To **(A,B)**, comparisons were performed by using one-way ANOVA, with

763 Tukey's multiple comparisons test. All Error bars show mean ± S.E.M.