Structure-based drug design, virtual screening and high-throughput screening rapidly identify antiviral leads targeting COVID-19

Zhenming Jin[1,2,9], Xiaoyu Du[2,9], Yechun Xu[3,9], Yongqiang Deng[4,9], Meiqin Liu[5,9], Yao Zhao[1], Bing Zhang[1], Xiaofeng Li[4], Leike Zhang[5], Yinkai Duan[1], Jing Yu[1], Lin Wang[1], Kailin Yang[6], Fengjiang Liu[1], Tian You[1], Xiaoce Liu[1], Xiuna Yang[1], Fang Bai[1], Hong Liu[3], Xiang Liu[7], Luke W. Guddat[8], Gengfu Xiao[5], Chengfeng Qin[4], Zhengli Shi[5], Hualiang Jiang[1,3]*, Zihe Rao[1,2,7]* & Haitao Yang[1]*

[1]Shanghai Institute for Advanced Immunochemical Studies and School of Life Science and Technology, ShanghaiTech University, Shanghai, China.

[2]School of Life Sciences, Tsinghua University, Beijing, China.

[3]Drug Discovery and Design Center, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai, China.

[4]Department of Virology, State Key Laboratory of Pathogen and Biosecurity, Beijing Institute of Microbiology and Epidemiology, Academy of Military Medical Sciences, Beijing, China.

[5]CAS Key Laboratory of Special Pathogens, Wuhan Institute of Virology, Center for Biosafety Mega-Science, Chinese Academy of Sciences, Wuhan, China.

[6]Taussig Cancer Center, Cleveland Clinic, Cleveland, USA.

[7]State Key Laboratory of Medicinal Chemical Biology, Frontiers Science Center for Cell Response, College of Life Sciences, College of Pharmacy, Nankai University, Tianjin, China.

[8]School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, QLD 4072, Australia.

[9]These authors contributed equally: Zhenming Jin, Xiaoyu Du, Yechun Xu, Yongqiang Deng, Meiqin Liu.

*e-mail: hljiang@simm.ac.cn; raozh@tsinghua.edu.cn; yanght@shanghaitech.edu.cn

Abstract

A coronavirus identified as 2019 novel coronavirus (COVID-19) is the etiological agent responsible for the 2019-2020 viral pneumonia outbreak that commenced in Wuhan[1-4]. Currently there is no targeted therapeutics and effective treatment options remain very limited. In order to rapidly discover lead compounds for clinical use, we initiated a program of combined structure-assisted drug design, virtual drug screening, and high-throughput screening to identify new drug leads that target the COVID-19 main protease (M^{pro}). M^{pro} is a key coronavirus enzyme, which plays a pivotal role in mediating viral replication and transcription, making it an attractive drug target for this virus[5,6]. Here, we identified a mechanism-based inhibitor, N3, by computer-aided drug design and subsequently determined the crystal structure of COVID-19 M^{pro} in complex with this compound. Next, through a combination of structure-based virtual and high-throughput screening, we assayed over 10,000 compounds including approved drugs, drug candidates in clinical trials, and other pharmacologically active compounds as inhibitors of M^{pro}. Seven of these inhibit M^{pro} with IC_{50} values ranging from 0.48 to 16.62 μM. Ebselen, thiadiazolidinone-8 (TDZD-8) and N3 also exhibited strong antiviral activity in cell-based assays. Our results demonstrate the efficacy of this screening strategy, and establishes a new paradigm for the rapid discovery of drug leads with clinical potential in response to new infectious diseases where no specific drugs or vaccines are available.

An epidemic of contagious pneumonia has been spreading rapidly from its likely origin in

Central China since December 2019[1,2]. The infectious pneumonia is now found in all

provinces of China, with cases also confirmed in other 35 countries. As of February 25,

2020, around 80,000 people were infected and over 2,500 have succumbed to the

epidemic[7,8]. Symptoms associated with this disease include fever, cough, myalgia or

fatigue and dyspnea[9,10]. The World Health Organization announced that the outbreak of this

deadly and fast-spreading infectious pneumonia constitutes a global health emergency[8]. A

coronavirus named 2019 novel coronavirus (COVID-19), has been identified as the

etiological agent responsible for the outbreak[3,4]. At present, there are no targeted

therapeutics and effective treatment options remain very limited.

CoVs infect humans and other animal species, causing a variety of highly prevalent and

severe diseases, including Severe Acute Respiratory Syndrome (SARS) and Middle East

Respiratory Syndrome (MERS)[11]. CoVs are positive-sense, single-stranded RNA viruses

featuring the largest viral RNA genomes known to date[12]. The COVID-19 genome is

comprised of ~30,000 nucleotides; its replicase gene encodes two overlapping polyproteins,

pp1a and pp1ab, required for viral replication and transcription[3,4]. The functional

polypeptides are released from the polyproteins by extensive proteolytic processing,

predominantly by a 33.8-kDa main protease (M^{pro}), also referred to as the 3C-like protease.

M^{pro} digests the polyprotein at no less than 11 conserved sites, starting with the autolytic

cleavage of this enzyme itself from pp1a and pp1ab[5]. The functional importance of M^{pro} in

the viral life cycle, together with the absence of closely related homologues in humans,

identify the M^{pro} as an attractive target for antiviral drug design[6].

To facilitate the rapid discovery of antiviral compounds with clinical potential, we

developed a strategy combining structure-assisted drug design, virtual drug screening and high-throughput screening to repurpose existing drugs to target M^{pro}, which is a key enzyme for the viral life cycle. Our strategy has successfully identified several new compounds that showed both strong inhibition of the target enzyme and antiviral activity in cell-based assays. These studies pave the way for the further development of novel antiviral drugs against this CoV.

Establishing a high-throughput activity assay

Recombinant COVID-19 M^{pro} with native N and C termini was expressed in *Escherichia coli* and subsequently purified (Extended Data Fig. 1a, b). The molecular weight of COVID-19 M^{pro} as determined by mass spectroscopy is 33797.0 Da, consistent with its theoretical molecular weight 33796.8 Da. In order to characterize its enzymatic activity and to carry out high throughput screening of inhibitors, we developed a fluorescence resonance energy transfer (FRET) assay. To do this, a fluorescently labeled substrate, MCA-AVLQ↓SGFR-Lys(Dnp)-Lys-NH$_2$, derived from the N-terminal auto-cleavage sequence of the viral protease was designed and synthesized for time-dependent kinetic analysis (Extended Data Fig. 1e). The catalytic efficiency (k_{cat}/K_m) for COVID-19 M^{pro} was measured to be 28,500 M^{-1}s^{-1} which is slightly higher than that for SARS-CoV M^{pro} (k_{cat}/K_m=26,500 M^{-1}s^{-1})[13], but over 30-fold higher than that of human rhinovirus 3C protease (k_{cat}/K_m=920 M^{-1}s^{-1})[14].

N3 is a potent irreversible inhibitor of COVID-19 M^{pro}

In a previous study, we designed a Michael acceptor inhibitor N3 using computer-aided drug design (CADD) (Extended Data Fig. 1c), which can specifically inhibit multiple CoV M^{pro}s, including those from SARS-CoV and MERS-CoV[15-18]. It also has displayed potent

antiviral activity against infectious bronchitis virus in an animal model[16]. Next, we

constructed a homology model for COVID-19 M^{pro} and used molecular docking to see if

N3 could target this new CoV M^{pro}. A docking pose showed that it could fit inside the

substrate-binding pocket. To assess the efficacy of N3 for COVID-19 M^{pro} , kinetic analysis

was performed[15]. A progress curve showed that it is time-dependent irreversible inhibitor

of this enzyme. Further, the shape of this curve supports the mechanism of two-step

irreversible inactivation. The inhibitor first associates with COVID-19 M^{pro} (EI) with a

dissociation constant K_i; then, a stable covalent bond is formed between N3 and M^{pro} (E−I).

The evaluation of this time-dependent inhibition requires both the equilibrium-binding

constant K_i (designated as k_2/k_1) and the inactivation rate constant for covalent bond

formation k_3 (Extended Data Fig. 1d). However, N3 exhibits very potent inhibition of

COVID-19 M^{pro} (Extended Data Fig. 1e), such that measurement of K_i and k_3 proved

difficult. When very rapid inactivation occurs, $k_{obs}/[I]$ was utilized to evaluate the inhibition

as an approximation of the pseudo second-order rate constant (k_3/K_i)[15]. The value of $k_{obs}/[I]$

of N3 for COVID-19 M^{pro} was determined to be $11,300 \pm 880$ M^{-1}s^{-1}, suggesting this

Michael acceptor has potent inhibition.

110 The crystal structure of COVID-19 M^{pro} in complex with N3

In order to elucidate the inhibitory mechanism of this compound, we determined the crystal

structure of COVID-19 M^{pro} in complex with N3 to 2.1-Å resolution. The asymmetric unit

contains only one polypeptide (Extended Data Table 1). However, two of these associate

to form a dimer by a crystallographic 2-fold symmetry axis (the two molecules are

designated protomer A and B) (Fig. 1b). All residues (residues 1–306) are visible in electron

density maps. Each protomer is composed of three domains (Fig. 1a). Domains I (residues

8–101) and II (residues 102–184) have an antiparallel β-barrel structure. Domain III

(residues 201–303) contains five α-helices arranged into a largely antiparallel globular

cluster, and is connected to domain II by means of a long loop region (residues 185–200).

COVID-19 M^{pro} has a Cys–His catalytic dyad, and the substrate-binding site is located in

a cleft between Domain I and II. These features are similar to those of other M^{pro}s reported

previously[5,16,18-20]. The electron density map shows that N3 binds in the substrate-binding

pocket in an extended conformation (Fig. 1c, Extended Data Fig. 2), with the inhibitor

backbone atoms forming an antiparallel sheet with residues 164–168 of the long strand$_{155-}$

$_{168}$ on one side, and with residues 189–191 of the loop linking domains II and III.

Here we detail the specific interactions of N3 with M^{pro} (Fig. 1c). The electron density

shows that the Sγ atom of C145-A forms a covalent bond (1.8-Å) with the Cβ of the vinyl

group, confirming that the Michael addition has occurred (Fig. 1c, d). The S1 subsite has

an absolute requirement for Gln at the P1 position. The side chains of F140-A, N142-A,

E166-A, H163-A, H172-A, S1-B (from protomer B), and main chains of F140-A and L141-

A are involved in S1 subsite formation, which also includes two ordered water molecules

(named W1 and W2). The lactam at P1 inserts into the S1 subsite and forms a hydrogen

bond with H163-A. The side chain of Leu at P2 site deeply inserts into the hydrophobic S2

subsite, which consists of the side chains of H41-A, M49-A, Y54-A, M165-A, and the alkyl

portion of the side chain of D187-A. The side chain of Val at P3 is solvent-exposed,

indicating that this site can tolerate a wide range of functional groups. The side chain of

Ala at P4 side is surrounded by the side chains of M165-A, L167-A, F185-A, Q192-A and

the main chain of Q189-A, all of which form a small hydrophobic pocket. P5 makes van

der Waals contacts with P168-A and the backbone of residues 190–191. The bulky benzyl

group extends into the S1′ site, possibly forming van der Waals interactions with T24-A

and T25-A. In addition, N3 forms multiple hydrogen bonds with the main chain of the

residues in the substrate-binding pocket, which also helps lock the inhibitor inside the

substrate-binding pocket (Fig. 1c, Extended Data Fig. 2).

Virtual screening

The structure of COVID-19 M^{pro} in complex with N3 provides a model for rapidly

identifying lead inhibitors to target COVID-19 M^{pro} through *in silico* screening. To achieve

this, an in-house database of potential binding compounds was docked using Glide (version

8.2) of Schrödinger 2019-1. The results show that cinanserin fits snugly into the substrate-

binding pocket, by interacting with H41 and E166 of M^{pro} by cation-π. Subsequently we

determined this compound has an IC_{50} value of 124.93 μM for M^{pro}, which validated our

virtual screening approach (Extended Data Fig. 3). Moreover, cinanserin is a well-

characterized serotonin antagonist, which has undergone preliminary clinical testing in

humans in the 1960s[21] and has previously been shown to inhibit SARS-CoV M^{pro} [22]. Thus,

it has potential for further optimization as an anti-viral drug lead.

High-throughput screening

Next, we used our FRET assay, to screen a library of ~10,000 compounds consisting of

approved drugs, clinical trial drug candidates and natural products. Primary hits included

seven compounds that are either FDA-approved drugs or clinical trial/preclinical drug

candidates. We then determined their IC_{50} values, which are in the range from 0.48 to 16.62

μM (Fig. 2). Amongst them, disulfiram and carmofur are FDA-approved drugs, whilst

ebselen, TDZD-8, shikonin, tideglusib, and PX-12 are currently in clinical trials or

preclinical studies. Ebselen has the strongest inhibition of M^{pro} activity with an IC_{50} of 0.48

µM. Since our structural data is only for N3, we investigated if molecular docking could

predict how these inhibitors bind to this protein. In all cases, reasonable docking poses

were found, demonstrating that they could fit inside the substrate-binding pocket (Extended

Data Fig. 4).

Antiviral activity assay

To further substantiate the enzymatic inhibition results *in vitro*, we then evaluated the

ability of these compounds to prevent cells from being infected by COVID-19. As shown

in Fig. 3, ebselen, TDZD-8 and N3 showed the most potent antiviral effects at a

concentration of 10 µM, entirely preventing the cytopathic effect (CPE) in COVID-19-

infected Vero cells. Quantitative real-time RT-PCR (qRT-PCR) demonstrated that treatment

of ebselen reduced the amounts of COVID-19 20.3-fold compared with the treatment in

the absence of inhibitor, and TDZD-8 and N3 exhibited 10.9-fold and 8.4-fold reduction in

COVID-19 growth, respectively. Ebselen is an organoselenium compound with anti-

inflammatory, anti-oxidant and cytoprotective properties. This compound has been

investigated for the treatment of multiple diseases, such as bipolar disorders and hearing

loss. The toxicity of ebselen in humans has been evaluated in a number of clinical trials[23-

25]. These data strongly suggest the clinical potential of ebselen for CoV treatment. TDZD-

8 was originally designed as a lead compound for the treatment of Alzheimer's disease,

which has been shown to be effective in animal models[26]. It is also interesting that

cinanserin displayed moderate inhibition against COVID-19 with an IC_{50} value of 20.61

µM (Extended Data Fig. 3), which is superior to that in the enzymatic inhibition assay,

suggesting that cinanserin might have multi-drug targets in preventing viral infection.

Discussion

In the last twenty years, new infectious agents have emerged to cause epidemics, such as SARS and MERS[11]. The timely development of effective antiviral agents for clinical use is extremely challenging because conventional drug development approaches normally take years of investigations and cost billions of dollars [27]. The repurposing of approved pharmaceutical drugs and drug candidates provides an alternative approach to rapidly identify potential drug leads to manage rapidly emerging viral infections[28]. Cell-based phenotypic screening has proven to be valuable[29], but the corresponding methodological complexity is not readily compatible with high-throughput pipelines, and fails to inform the identity of the molecular target or mechanism of action[30]. In this regard, we combined structure-based *ab initio* drug design, virtual screening and high-throughput screening to discover new drugs for treating COVID-19 infection. The convergence of these three approaches proved to be an efficient strategy in finding three lead compounds with good antiviral activity. Ebselen is of particular interest for further investigation since its safety in humans has been supported by different clinical trials[23-25]. Our methodology establishes a paradigm of rapid discovering drug leads with clinical potential in response to new emerging infectious diseases lacking specific drugs and vaccines.

The crystal structural and docking data have shown that these drug leads can bind to the substrate-binding pocket of COVID-19 M$^{\text{pro}}$. In our previous study, we proposed that all the CoV M$^{\text{pro}}$s share a highly conservative substrate-recognition pocket, which serves as a drug target for broad-spectrum inhibitor design[15]. Recent discovery of new CoVs and accumulation of structural data for CoV M$^{\text{pro}}$s from various species provided the opportunity to further examine our previous hypothesis. Superposition of the 12 crystal

structures of M^{pro}s[15,16,18,20,31-35] have shown that the most variable regions were the helical

domain III and surface loops, but the substrate-binding pockets located in a cleft between

domains I and II are still highly conserved among all CoV M^{pro}s, suggesting the antiviral

inhibitors targeting at this site should have wide-spectrum anti-CoV activity (Fig. 4). This

strongly supports our hypothesis that development of a single antiviral agent could provide

an effective first line of defense against all CoV-associated diseases.

References

1   Zhu, N. *et al.* A novel coronavirus from patients with pneumonia in China, 2019. *New England*
Journal of Medicine (2020).
2   Qun Li *et al.* Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected
Pneumonia. *New England Journal of Medicine* (2020).
3   Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin.
Nature, doi:10.1038/s41586-020-2012-7 (2020).
4   Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature*,
doi:10.1038/s41586-020-2008-3 (2020).
5   Anand, K., Ziebuhr, J., Wadhwani, P., Mesters, J. R. & Hilgenfeld, R. Coronavirus main proteinase
(3CL(pro)) structure: Basis for design of anti-SARS drugs. *Science* **300**, 1763-1767,
doi:10.1126/science.1085658 (2003).
6   Pillaiyar, T., Manickam, M., Namasivayam, V., Hayashi, Y. & Jung, S. H. An Overview of Severe
Acute Respiratory Syndrome-Coronavirus (SARS-CoV) 3CL Protease Inhibitors: Peptidomimetics
and Small Molecule Chemotherapy. *Journal of Medicinal Chemistry* **59**, 6595-6628,
doi:10.1021/acs.jmedchem.5b01461 (2016).
7   National Health Commission, PRC. *Novel Coronavirus(2019-nCoV) Situation Report of China*,
<http://www.nhc.gov.cn/xcs/yqtb/list_gzbd.shtml> (2020).
8   WHO. Novel Coronavirus(2019-nCoV) Situation Report. (2020).
9   Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China.
The Lancet, doi:10.1016/s0140-6736(20)30183-5 (2020).
10   Michelle L. Holshue, C. D., Scott Lindquist, Kathy H. Lofy, John Wiesman, Hollianne Bruce,
Christopher Spitters, for the Washington State 2019-nCoV Case Investigation Team. First Case of
2019 Novel Coronavirus in the United States.   (2020).
11   de Wit, E., van Doremalen, N., Falzarano, D. & Munster, V. J. SARS and MERS: recent insights
into emerging coronaviruses. *Nature Reviews Microbiology* **14**, 523-534,
doi:10.1038/nrmicro.2016.81 (2016).
12   Perlman, S. & Netland, J. Coronaviruses post-SARS: update on replication and pathogenesis. *Nature*
Reviews Microbiology **7**, 439-450, doi:10.1038/nrmicro2147 (2009).
13   Xue, X. Y. *et al.* Production of authentic SARS-CoV M-pro with enhanced activity: Application as
a novel tag-cleavage endopeptidase for protein overproduction. *Journal of Molecular Biology* **366**,
965-975, doi:10.1016/j.jmb.2006.11.073 (2007).
14   Wang, Q. M., Johnson, R. B., Cox, G. A., Villarreal, E. C. & Loncharich, R. J. A Continuous
Colorimetric Assay for Rhinovirus-14 3C Protease Using Peptidep-Nitroanilides as Substrates.
Analytical biochemistry **252**, 238-245 (1997).
15   Yang, H. T. *et al.* Design of wide-spectrum inhibitors targeting coronavirus main proteases. *Plos*
Biology **3**, 2044-2044, doi:10.1371/journal.pbio.0030428 (2005).
16   Xue, X. Y. *et al.* Structures of two coronavirus main proteases: Implications for substrate binding

and antiviral drug design. *Journal of Virology* **82**, 2515-2527, doi:10.1128/jvi.02114-07 (2008).

17   Ren, Z. L. *et al.* The newly emerged SARS-Like coronavirus HCoV-EMC also has an "Achilles'
heel": current effective inhibitor targeting a 3C-like protease. *Protein & Cell* **4**, 248-250,
doi:10.1007/s13238-013-2841-3 (2013).

18   Wang, F. H. *et al.* Michael Acceptor-Based Peptidomimetic Inhibitor of Main Protease from Porcine
Epidemic Diarrhea Virus. *Journal of Medicinal Chemistry* **60**, 3212-3216,
doi:10.1021/acs.jmedchem.7b00103 (2017).

19   Yang, H. T. *et al.* The crystal structures of severe acute respiratory syndrome virus main protease
and its complex with an inhibitor. *Proceedings of the National Academy of Sciences of the United*
States of America **100**, 13190-13195, doi:10.1073/pnas.1835675100 (2003).

20   Zhao, Q. *et al.* Structure of the main protease from a global infectious human coronavirus, HCoV-
HKU1. *Journal of virology* **82**, 8647-8655, doi:10.1128/JVI.00298-08 (2008).

21   Rubin, B. & Waugh, M. H. Antiphlogistic effects of antiserotonin (SQ 10,643) and aminopyrine in
rats versus endotoxin and other agents. *Proceedings of the Society for Experimental Biology and*
Medicine **119**, 438-443 (1965).

22   Chen, L. *et al.* Cinanserin is an inhibitor of the 3C-like proteinase of severe acute respiratory
syndrome coronavirus and strongly reduces virus replication in vitro. *Journal of virology* **79**, 7095-
7103 (2005).

23   Lynch, E. & Kil, J. in *Seminars in Hearing.* 047-055 (© Thieme Medical Publishers).

24   Masaki, C. *et al.* Effects of the potential lithium-mimetic, ebselen, on impulsivity and emotional
processing. **233**, 2655-2661, doi:10.1007/s00213-016-4319-5 (2016).

25   Kil, J. *et al.* Safety and efficacy of ebselen for the prevention of noise-induced hearing loss: a
randomised, double-blind, placebo-controlled, phase 2 trial. *The Lancet* **390**, 969-979 (2017).

26   Koehler, D., Shah, Z. A. & Williams, F. E. The GSK3β inhibitor, TDZD-8, rescues cognition in a
zebrafish model of okadaic acid-induced Alzheimer's disease. *Neurochemistry international* **122**,
31-37 (2019).

27   Bedford, J. *et al.* A new twenty-first century science for effective epidemic response. *Nature* **575**,
130-136, doi:10.1038/s41586-019-1717-y (2019).

28   Farha, M. A. & Brown, E. D. Drug repurposing for antimicrobial discovery. *Nature Microbiology* **4**,
565-577, doi:10.1038/s41564-019-0357-1 (2019).

29   Xu, M. *et al.* Identification of small-molecule inhibitors of Zika virus infection and induced neural
cell death via a drug repurposing screen. *Nature Medicine* **22**, 1101-1107, doi:10.1038/nm.4184
(2016).

30   Aulner, N., Danckaert, A., Ihm, J., Shum, D. & Shorte, S. L. Next-Generation Phenotypic Screening
in Early Drug Discovery for Infectious Diseases. *Trends in parasitology* (2019).

31   Wang, F., Chen, C., Tan, W., Yang, K. & Yang, H. Structure of Main Protease from Human
Coronavirus NL63: Insights for Wide Spectrum Anti-Coronavirus Drug Design. *Sci Rep* **6**, 22677-
22677, doi:10.1038/srep22677 (2016).

32   Lee, C.-C. *et al.* Structural basis of inhibition specificities of 3C and 3C-like proteases by zinc-
coordinating and peptidomimetic compounds. *J Biol Chem* **284**, 7646-7655,
doi:10.1074/jbc.M807947200 (2009).

33   Wang, F. H. *et al.* Crystal Structure of Feline Infectious Peritonitis Virus Main Protease in Complex
with Synergetic Dual Inhibitors. *Journal of Virology* **90**, 1910-1917, doi:10.1128/jvi.02685-15
(2016).

34   Cui, W. *et al.* The crystal structure of main protease from mouse hepatitis virus A59 in complex with
an inhibitor. *Biochemical and Biophysical Research Communications* **511**, 794-799,
doi:10.1016/j.bbrc.2019.02.105 (2019).

35   Needle, D., Lountos, G. T. & Waugh, D. S. Structures of the Middle East respiratory syndrome
coronavirus 3C-like protease reveal insights into substrate specificity. *Acta Crystallographica*
Section D-Structural Biology **71**, 1102-1111, doi:10.1107/s1399004715003521 (2015).

Fig. 1 | The crystal structure of COVID-19 M^pro in complex with N3.

a, Cartoon representation of one protomer of the dimeric M^pro-inhibitor complex. N3 is shown in green

sticks. **b,** Surface representation of the homodimer of M^pro. Protomer A is in blue while Protomer B in

salmon; N3 is presented as green sticks. **c,** A zoomed view of the substrate-binding pocket. The key

residues forming the binding pocket are shown in sticks; the two waters, assigned as W1 and W2, are

shown as red spheres; the $2F_o$-F_c density map contoured at 1.2 σ is shown around N3 molecule (blue

mesh), C145-A (yellow mesh), and the two waters (blue mesh). **d,** The C-S covalent bond.

Fig. 2 | Plots showing the inhibitory activity for the seven drug leads. The hydrolytic activity of

COVID-19 M^{pro} is measured in the presence of varying concentrations of compounds. Curves represent

best fits for calculating the IC$_{50}$ values, error bars represent mean ± SEM based on three independent

measurements.

Fig. 3 | **Antiviral activity assay.** Virus-induced cytopathic effect in Vero cells in the presence of M[Pro]
inhibitors. **a-c,** Treated with Ebselen, TDZD-8 and N3 respectively. **d,** Cells infected with COVID-19.
e, Mock-infection with DMSO (0.1%). **f,** The relative concentration of vRNA present in the supernatant
at 72 h post infection (p.i.) determined by qRT-PCR analysis.

Fig. 4 | The substrate-binding pocket of M^pros is conserved among 12 different CoVs.

a, Superposition of crystal structures of M^pros (Cα 1-300) from 12 CoVs, including COVID-19, SARS-

CoV, MERS-CoV, HCoV-HKU1, BtCoV-HKU4, MHV-A59, PEDV, FIPV, TGEV, HCoV-NL63,

HCoV-229E and IBV. The color spectrum represents the root-mean-square deviation (RMSD) of the

aligned Cα atoms. **b,** Surface presentation of conserved substrate-binding pockets of 12 CoV M^pros. N3

is shown as green sticks; the background (blue) is the surface of COVID-19 M^pro. Red: residues are

entirely identical among all 12 M^pros; violet: conserved substitution in one CoV M^pro; orange: conserved

substitution in more than one CoV M^pros. S1, S2, S4, and S1′ subsites are indicated.

Methods

Cloning, protein expression and purification

The full-length gene encoding COVID-19 M^{pro} was optimized and synthesized for *Escherichia coli* expression (GENEWIZ). Cloning strategy for producing authentic viral M^{pro} has been reported previously[13]. The expression plasmid was transformed into *Escherichia coli* BL21 (DE3) cells and then cultured in Luria broth (LB) media containing 100 μg/mL ampicillin at 37°C, 220 rpm. When the cells were grown to OD$_{600}$ of 0.6-0.8, 0.5 mM IPTG was added to the cell culture to induce the expression of the recombinant COVID-19 M^{pro} at 16°C, 180 rpm. After 10 hours, the cells were harvested by centrifugation at 3,000×g for 20 min.

The cell pellets were resuspended in lysis buffer (20 mM Tris-HCl, pH 8.0, 300 mM NaCl), lysed by high-pressure homogenization, and then centrifuged at 25,000×g for 40 min. The supernatant was loaded onto Ni-NTA affinity column (QIAGEN), and washed by the resuspension buffer containing 20 mM imidazole. The His tagged M^{pro} was eluted by cleavage buffer (50 mM Tris-HCl, pH 7.0, 150 mM NaCl) including 300 mM imidazole. Human rhinovirus 3C protease was added to remove the C-terminal His tag. The COVID-19 M^{pro} was further purified by anion-exchange chromatography and size-exclusion chromatography (Superdex 200, GE Healthcare). The purified COVID-19 M^{pro} was stored in 50 mM Tris-HCl, pH 7.3, 1 mM EDTA.

Crystallization

COVID-19 M^{pro} was incubated with 10 mM N3 for 30 min and the complex (5 mg/ml) was crystallized by hanging drop vapor diffusion method at 20°C. The best crystals were grown with well buffer containing 2% polyethylene glycol (PEG) 6000, 3% DMSO, 1 mM DTT,

0.1 M MES (pH 6.0). The cryo-protectant solution contained 30% PEG 400, 0.1 M MES

(pH 6.0).

Data collection and structure determination

X-ray data were collected on beamline BL17U1 at Shanghai Synchrotron Radiation

Facility (SSRF) at 100 K and at a wavelength of 1.07180 Å using an Eiger X 16M image

plate detector. Data integration and scaling were performed using the program Xia2. The

structure was determined by molecular replacement (MR) with the Phaser module[36] in

CCP4[37] using the SARS-CoV M^{pro} (PDB ID: 2H2Z) as a search template. The output model

from MR was subsequently subjected to iterative cycles of manual model adjustment with

Coot[38] and refinement was finished with Phenix[39]. The inhibitor N3 was built according to

the omit map. The phasing and refinement statistics are summarized in Extended Data

Table 1. The R_{work}/R_{free} values are 0.202/0.235, respectively. 97.3% residues are in most

favored regions of the Ramachandran plot, and no residues are found in disallowed regions.

Coordinates and structure factors for COVID-19 M^{pro} in complex with the inhibitor N3

have been deposited in Protein Data Bank with accession number 6LU7.

Enzymatic activity and inhibition assays

The enzyme activity assays have been described previously[13]. Briefly, the activity of

COVID-19 M^{pro} was measured by continuous kinetic assay, with the substrate MCA-

AVLQSGFR-Lys(Dnp)-Lys-NH2 (GL Biochem Shanghai Ltd), using wavelengths of 320

and 405 nm for excitation and emission, respectively. The assays started by immediately

mixing 0.2 μM COVID-19 M^{pro} with different concentrations of substrate (2.5-100 μM).

Fluorescence intensity was monitored with EnVision multimode plate reader

(PerkinElmer). Initial rates were obtained by fitting the linear portion of the curves (usually

the first 3 min of the assay curves) to a straight line. The kinetic parameters K_m and k_{cat}

were calculated from a double-reciprocal plot. As N3 is a mechanism-based irreversible

inhibitor for COVID-19 M^{pro}, k_{obs}/[I] was used as an approximation of the pseudo second-

order rate constant to evaluate the inhibition effect of the inhibitor N3[15]. In this case, the

measurement was carried out with 0.2 μM of enzyme, 20 μM of substrate and inhibitor at

6 different concentrations (0-1 μM).

Virtual screening

The virtual screening was performed using our in-house database via a workflow

application of Glide (v8.2)[40], which is a module of molecular simulation software package,

Maestro (Schrodinger 2019-1a). All compounds in the database were considered to be at

pH 7.4 ± 0.2 to estimate their protonation state using the program EpiK[41]. Their three

dimensional conformations were generated by the ligPrep module of Maestro. The newly

obtained complex structure of COVID-19 M^{pro} (PDB ID: 6LU7) was used to generate

receptor grid for docking simulations. The center of active site of the grid was determined

according to the N3 in the structure. The flexibility of the receptor hydroxyl and thiol

groups in side chains of C145, S46 and Y54 were considered. At the very beginning, a

relatively fast but raw screening was performed by using the glide standard precision model,

and the top 20% of compounds were kept. Then covalent docking simulations were

performed by choosing Michael Addition as the reaction type for these top 20% compounds

that contained Michael acceptors, and C145 was set as the source of Michael donor. Finally,

the candidate molecules were picked out by analysing the predicted binding modes and

their scores.

High-throughput drug screen and IC$_{50}$ measurement

Potential inhibitors against COVID-19 M^{pro} were screened by enzymatic inhibition assay.

When different compounds were added into the enzymatic reaction, the change of initial

rates was calculated to evaluate their inhibitory effect. Five drug libraries, Approved Drug

Library (TargetMol, USA), Clinic Compound Library (TargetMol, USA), FDA-approved

Drug Library (Selleck, USA), Natural Product Library (Selleck, USA), and Anti-virus drug

library (Shanghai Institute for Advanced Immunochemical Studies, SIAIS), which includes

~ 10,000 compounds, were used. The preliminary screening reaction mixture included 0.2

μM protein, 20 μM substrate and 50 μM compounds. The compounds of interest were

defined as those with a percentage of inhibition over 60% compared with the reaction in

the absence of inhibitor. IC_{50} values were measured at 12 concentrations and three

independent experiments were performed. All experimental data was analyzed using

GraphPad Prism software.

Molecular docking

To understand the binding interaction of these molecules with COVID-19 M^{pro}, two

different molecular docking methods, i.e., Glide (v8.2)[40] and iFitDock[42] were used to

predict their binding poses. Then a 3D molecular similarity calculation method, SHAFTs[43],

was used for molecular alignment poses enumeration by matching the critical

pharmacophore and volumetric overlay between the N3 molecule within the M^{pro} structure

and the seven drug candidates. However, the selenium atom of ebselen could not be treated

by any of these above methods, so sulfur was used to replace it in the calculations. Then

the obtained optimal superposition of these molecules were used to assist the reasonability

of the predicted binding poses from the two docking methods, and only the binding

orientations which were consistent among different methods were kept for constructing the

initial complexes. Finally, these complexes were further optimized (refined binding poses

are illustrated in Extended Data Fig. 4) and re-scored (shown in Extended Data Table 2) by

using MM-GBSA module[44] of Schrödinger, and the residues within 5 Å around the ligand

were refined.

Antiviral assay for compounds identified from high-throughput screening

Vero (African green monkey kidney) cells were incubated at 37°C in Dulbecco's modified

Eagle's medium (DMEM; Invitrogen) with 10% fetal bovine serum (FBS), 100 U/mL

penicillin, and 100 μg/mL streptomycin.

The COVID-19 was originally isolated from a throat swab specimen from a pneumonia

patient and amplified in Vero cells. The *in vitro* antiviral effects of the selected compounds

in Vero cells were assayed as previously described[45]. Briefly, cultured Vero cells were

infected with COVID-19 in the presence of each compounds at the concentration of 10 μM.

Cells were then cultured for 72 hours, and the CPE was observed daily and viral RNA copy

numbers in supernatants were quantified by in the COVID-19-specific qRT-PCR.

Antiviral assay for cinanserin

Vero E6 cell line was obtained from American Type Culture Collection (ATCC) and

maintained in Dulbecco's Modified Eagle Medium (DMEM; Gibco Invitrogen)

supplemented with 10% fetal bovine serum (FBS; Gibco Invitrogen) at 37 °C in a

humidified atmosphere of 5% CO_2. A clinical isolate COVID-19 [3] was propagated in Vero

E6 cells, and viral titer was determined as described previously[45]. All the infection

experiments were performed at biosafety level-3 (BLS-3).

Pre-seeded Vero E6 cells (5×10^4 cells/well) were pre-treated with the different

concentration of the indicated compounds for 1 hour and the virus was subsequently added

(MOI of 0.05) to allow infection for 2 hours. Then, the virus-drug mixture was removed

and cells were further cultured with fresh drug containing medium. At 24 h p.i., the cell

supernatant was collected and viral RNA in supernatant was subjected to qRT-PCR analysis.

McCoy, A. J. *et al.* Phasercrystallographic software. *Journal of Applied Crystallography* **40**, 658-674, doi:10.1107/s0021889807021206 (2007).

Potterton, L. *et al.* CCP4i2: the new graphical user interface to the CCP4 program suite. *Acta Crystallographica Section D-Structural Biology* **74**, 68-84, doi:10.1107/s2059798317016035 (2018).

Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta Crystallographica Section D: Biological Crystallography* **66**, 486-501 (2010).

Afonine, P. V. *et al.* Towards automated crystallographic structure refinement with phenix.refine. *Acta Crystallographica Section D-Structural Biology* **68**, 352-367, doi:10.1107/s0907444912001308 (2012).

Friesner, R. A. *et al.* Glide: A new approach for rapid, accurate docking and scoring. 1. Method and assessment of docking accuracy. *Journal of Medicinal Chemistry* **47**, 1739-1749, doi:10.1021/jm0306430 (2004).

Greenwood, J. R., Calkins, D., Sullivan, A. P. & Shelley, J. C. Towards the comprehensive, rapid, and accurate prediction of the favorable tautomeric states of drug-like molecules in aqueous solution. *Journal of computer-aided molecular design* **24**, 591-604 (2010).

Bai, F. *et al.* Free energy landscape for the binding process of Huperzine A to acetylcholinesterase. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 4273-4278, doi:10.1073/pnas.1301814110 (2013).

Liu, X. F., Jiang, H. L. & Li, H. L. SHAFTS: A Hybrid Approach for 3D Molecular Similarity Calculation. 1. Method and Assessment of Virtual Screening. *Journal of Chemical Information and Modeling* **51**, 2372-2385, doi:10.1021/ci200060s (2011).

Guimaraes, C. R. W. & Cardozo, M. MM-GB/SA rescoring of docking poses in structure-based lead optimization. *Journal of Chemical Information and Modeling* **48**, 958-970, doi:10.1021/ci800004w (2008).

Wang, M. *et al.* Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Research*, doi:10.1038/s41422-020-0282-0 (2020).

Acknowledgments We would like to thank Ying Lei and Juan Kong from High Throughput

Platform, staff from Analytical Chemistry Platform at Shanghai Institute for Advanced

Immunochemical Studies, for their technical support. We are grateful to National Centre

for Protein Science Shanghai and The Molecular and Cell Biology Core Facility of the

School of Life Science and Technology, ShanghaiTech University for use of their

instrumentation and technical assistance. We thank Prof. Wenqing Xu, Prof. Zhijie Liu, and

Haixia Su for discussion. We also thank the staff from beamlines BL17U1, BL18U1 and

BL19U1 at Shanghai Synchrotron Radiation Facility (China). This work was supported by

grants from National Key R&D Program of China (grant No. 2017YFC0840300) and

Project of International Cooperation and Exchanges NSFC (grant No. 81520108019) to

Z.R..

Author contributions Z.R. and H.Y. conceived the project. Z.J., H.J., Z.R., and H.Y.

designed the experiments; Z.J., X.D., Y.X., Y.D., M.L., Y.Z., B.Z., X. Li, L.Z., Y.D., J.Y.,

F.L., T.Y., Xiaoce Liu and X.Y. performed the experiments. Z.J., B.Z. and F.L. collected

the diffraction data. B.Z. and Xiang Liu solved the crystal structure. Z.J., X.D., Y.X., K.Y.,

F.B., H.L., Xiang Liu, L.G., G.X., C.Q., Z.S., H.J., Z.R. and H.Y. analyzed and discussed

the data. Z.J., X.D., F.B., Xiang Liu, L.G., G.X., C.Q., Z.S., H.J., Z.R. and H.Y wrote the

manuscript.

Competing interests The authors declare no competing interests.

Data and materials availability: The PDB accession No. for the coordinates of COVID-

19 M^{pro} in complex with N3 is 6LU7 (Deposited: 2020-01-26; Released: 2020-02-05).

505
506 **Extended Data Fig. 1 | The purification of COVID-19 M^pro and the inhibitory assay of N3**
507 **compound. a,** the SDS-PAGE gel of COVID-19 M^pro. The first lane: marker; the second lane: M^pro
508 before treating with rhinovirus 3C protease; third lane: the M^pro after the cleavage of C-terminal tag. **b,**
509 Size-exclusion chromatography profile of M^pro. **c,** The chemical structure of N3 inhibitor. **d,** Inhibition
510 mechanism for N3. **e,** Typical inhibition curves for N3.
511

512
Extended Data Fig. 2 | The interactions between M^pro and N3. a, The F_o-F_c omit map (contour level
= 3 σ, shown as the blue mesh). **b,** Detailed view of the interactions between the N3 and COVID-19
M^pro. The hydrogen bonds are shown as black dashed lines. The covalent bond between N3 and C145-
A is in purple.

Extended Data Fig. 3 | Cinanserin is an inhibitor for M^{pro}. a, The docking result of cinanserin. The

structure of COVID-19 M^{pro} is shown as a white cartoon; cinanserin is shown as cyan sticks and balls;

residues predicted to be interacting with cinanserin are shown as sticks. **b,** Inhibiton of cinanserin on

M^{pro}. **c,** Antiviral activity of cinanserin.

Extended Data Fig. 4 | Docking Poses of different COVID-19 M^{pro} inhibitors. The docking results
show that the inhibitors identified through the high-throughput screening are likely to occupy the same
pocket as N3.

Extended Data Fig. 5 | Virus-induced CPE in Vero cells treated with M^{pro} inhibitors. a-d, Treated
with disulfiram, tideglusib, carmofur and PX-12 respectively; **e,** Cells infected with COVID-19; **f,**
Mock-infection with DMSO (0.1%). **g,** The relative concentration of vRNA present in the supernatant
at 72h p.i. determined by qRT-PCR analysis. Because shikonin showed cellular toxicity at the test
concentration, its antiviral activity assay did not further proceed.

536 **Extended Data Table 1 | Data collection and refinement statistics**

	PDB code: 6LU7
Data collection	
Space group	$C2$
Cell dimensions	
a, b, c (Å)	97.931, 79.477, 51.803
α, β, γ (°)	90, 114.55, 90
Resolution (Å)	50.00-2.16 (2.22-2.16)
R_{merge}	18.9 (147.2)
$I / \sigma I$	6.3 (3.2)
Completeness (%)	100.0 (100.0)
Redundancy	6.6 (6.1)
Refinement	
Resolution (Å)	50.00-2.16
No. reflections	19455 (1431)
R_{work} / R_{free}	0.2020/0.2350
No. atoms	
Protein	2367
Ligand/ion	49
Water	84
B-factors	
Protein	42.7
Ligand/ion	46.3
Water	44.2
R.m.s. deviations	
Bond lengths (Å)	0.002
Bond angles (°)	0.474

538 *Values in parentheses are for highest-resolution shell.

540 **Extended Data Table 2 | Predicted binding affinities for the drug leads to COVID-19 M^{pro} by**

541 **using MM-GBSA module integrated in Schrödinger**

Drug	Predicted ΔG_{bind} (kcal/mol)
PX-12	-50.84
Shikonin	-67.95
TDZD-8	-43.77
Tideglusib	-61.79
Carmofur	-47.12
Ebselen	-53.83
Disulfiram	-46.16

543