The vulnerability of working memory to distraction is rhythmic

Malte Wöstmann, Troby Ka-Yan Lui, Kai-Hendrik Friese, Jens Kreitewolf, Malte Naujokat &

Jonas Obleser

Department of Psychology, University of Lübeck, Lübeck, Germany

Corresponding authors:

Malte Wöstmann (malte.woestmann@uni-luebeck.de); Jonas Obleser (jonas.obleser@uni-

luebeck.de)

Keywords: working memory, distraction, oscillation, attention, suppression, rhythm

Acknowledgements

We thank Molly Henry for fruitful discussions about the analysis pipeline used in the present

study. The authors declare no conflict of interest.

Abstract

Recent research posits that the cognitive system samples target stimuli in a rhythmic fashion, characterized by target detection fluctuating at frequencies of ~3–8 Hz. Besides prioritized encoding of targets, a key cognitive function is the protection of working memory from distractor intrusion. Here, we test to which degree the vulnerability of working memory to distraction is rhythmic. In an Irrelevant-Speech Task, $N = 23$ human participants had to retain the serial order of nine numbers in working memory while being distracted by task-irrelevant speech with variable temporal onsets. The magnitude of the distractor-evoked N1 component in the event-related potential as well as behavioural recall accuracy, both measures of memory distraction, were periodically modulated by distractor onset time in approximately 2–4 cycles per second (Hz). Critically, an underlying 2.5-Hz rhythm explained variation in both measures of distraction such that stronger phasic distractor encoding mediated lower phasic memory recall accuracy. In a behavioural follow-up experiment, we tested whether these results would replicate in a task design without rhythmic presentation of target items. Participants ($N = 6$ with on average >2,500 trials, each) retained two line-figures in memory while being distracted by acoustic noise of varying onset across trials. In agreement with the main experiment, the temporal onset of the distractor periodically modulated memory performance. Together, these results suggest that during working memory retention, the human cognitive system implements distractor suppression in a temporally dynamic fashion, reflected in ~400-ms long cycles of high versus low distractibility.

Introduction

Since our environments are rich in sensory stimuli, our attentional system faces two closely intertwined, yet conceptually dissociable tasks: First, relevant target stimuli need to be prioritized by attention in order to encode them into working memory. Second, irrelevant distractors need to be suppressed to avoid working memory interference. The cognitive system has to orchestrate these two processes dynamically in time (for review, see Nobre & van Ede, 2018), in order to prioritize targets and to suppress distractors from the ongoing stream of sensory inputs. While recent research suggests that the attentional sampling of target stimuli is not constant throughout time but periodically modulated at frequencies in the range of 3–8 Hz (for review, see Fiebelkorn & Kastner, 2019), it is at present unclear whether the vulnerability to distraction is rhythmic as well.

It is long known that rhythmic sensory stimulation has the potency to entrain so-called "attending rhythms" (Large & Jones, 1999), which eventually results in enhanced target stimulus detection at time points in-phase versus out-of-phase with the entrained rhythm (in vision: de Graaf et al., 2013; in audition: M. R. Jones, Johnston, & Puente, 2006). Studies have also revealed the neural basis of this phenomenon: Entrainment of brain oscillations by rhythmic stimulation relates to phasic modulation of target detection (Henry & Obleser, 2012; Spaak, de Lange, & Jensen, 2014). Furthermore, there is some evidence that direct entrainment of neural oscillations by transcranial alternating current stimulation (tACS) induces phasic modulation of target stimulus detection (Gundlach, Muller, Nierhaus, Villringer, & Sehm, 2016; Helfrich et al., 2014). An important interim summary of this research is that the sensory gain for processing to-be-attended target stimuli fluctuates periodically, likely so as a result of neural entrainment (for recent review articles, see Lakatos, Gross, & Thut, 2019; Obleser & Kayser, 2019).

Interestingly, several studies have demonstrated that target detection fluctuates rhythmically even in the absence of entraining sensory input or rhythmic brain stimulation (e.g. Fiebelkorn, Saalmann, & Kastner, 2013; Landau & Fries, 2012). In essence, these studies found that the time interval between an alerting stimulus and subsequent target presentation periodically modulated target detection at frequencies in the theta range of 3–8 Hz. On the neural level, rhythmic attentional sampling was accompanied by theta-rhythmic modulation of gamma responses (Landau, Schreyer, van Pelt, & Fries, 2015) and frontoparietal theta activity in humans (Helfrich et al., 2018) and non-human primates

(Fiebelkorn, Pinsk, & Kastner, 2018). These results suggest that attention samples target
stimuli at a theta rhythm.

Attention and working memory are tightly coupled neuro-cognitive faculties (Awh,
Vogel, & Oh, 2006). This is evidenced also by retro-cueing paradigms that require the flexible
allocation of attention to items in memory (e.g. Lim, Wöstmann, & Obleser, 2015; Oberauer
& Hein, 2012; Schneider, Mertes, & Wascher, 2016). It is thus straight-forward to assume that
sampling of target objects in memory might be theta-rhythmic as well. Speaking directly to
this, Peters and colleagues (2018) varied the time-interval between a retro-cue (which cued
one of two spatially separated visual objects in memory) and a subsequent memory probe.
They found rhythmic modulation of response times to the probe at a frequency of 6 Hz.
Interestingly, rhythmic trajectories of response times across probe times for the cued versus
non-cued object were in anti-phase, suggesting that target items in working memory were
sampled in alternation. From a more general perspective, theta oscillations have been
implicated in working memory function as they increase during temporal order
maintenance (Hsieh, Ekstrom, & Ranganath, 2011) and induce increased working memory
performance (for a recent review, see Hanslmayr, Axmacher, & Inman, 2019) if entrained by
rhythmic transcranial magnetic stimulation (Albouy, Weiss, Baillet, & Zatorre, 2017) or by
rhythmic visual stimulation (Köster, Martens, & Gruber, 2019).

Here, we test whether the temporal onset of task-irrelevant auditory stimuli would prove
disruptive in a slow oscillatory fashion. During working memory retention, distractor onset
periodically modulated both, the distractor-evoked N1 component in the event-related
potential (at ~3.5 Hz) and memory recall accuracy (at ~2 Hz). Critically, these two measures
of distraction were co-modulated by an underlying ~2.5-Hz rhythm. In a behavioural follow-
up experiment, we conceptually replicated the rhythmic vulnerability of working memory
to distraction in a visual match-to-sample paradigm.

Materials and methods

The main study poses a re-analysis of a previously published dataset. Below, we describe essential methodological aspects but we refer to the original article for details (Wöstmann, Lim, & Obleser, 2017).

Participants. In the main study and in the follow-up experiment, we tested $N = 23$ (12 females & 11 males; mean age: 24.5 years) and $N = 6$ (5 females & 1 male; mean age: 22 years) participants, respectively. Participants provided informed consent and were compensated financially or by course credit. Experimental procedures were approved by the ethics committee of the University of Lübeck.

Task design and stimuli in the main study. For the main study, to-be-remembered auditory materials were the German numbers from 1 to 9, spoken by a female voice (average number duration 0.6 s). To-be-ignored distractors were short German sentences (5–8 words, average duration: 2.1 s), taken from a German version of speech-in-noise sentences (Erb, Henry, Eisner, & Obleser, 2012), spoken by the same female voice as the numbers.

We used an adapted version of the Irrelevant-Speech Task (Colle & Welsh, 1976; D. M. Jones & Morris, 1992). On each trial, participants had to retain nine target numbers in memory, which were presented in random order with onset-to-onset delay of 0.75 s (presentation rate of 1.33 Hz). The most important manipulation for the present study was the temporal onset of the speech distractor. During the ensuing 5-s memory retention period, a task-irrelevant distractor sentence was presented, which had to be ignored. On each trial, the distractor onset was drawn from a uniform distribution including all millisecond values between 1.035 and 1.835 s relative to the offset of the last target number. On average, the distractor sentence was centred in the middle of the retention period, which corresponded to a distractor onset of 1.435 s.

At the end of each trial, participants were presented with a number pad on a computer screen. They had to select the numbers in their order of presentation using a computer mouse. In order to prevent participants' motor preparation for a particular behavioural response, the number pad was randomly arranged on each trial.

The experiment was implemented in Presentation software (Neurobehavioral Systems). All auditory materials were presented via Sennheiser HD-25 headphones at a comfortable level of ~65 dB A.

Additional manipulations of no interest in the main study. Two additional manipulations of the speech distractor were implemented in the main study but these have been analysed previously (Wöstmann et al., 2017) and were thus not of primary interest here. First, acoustic detail of distractor sentences was manipulated using noise-vocoding. In brief, noise-vocoding parametrically degrades the spectral content of the speech signal but leaves the temporal information largely intact (Shannon, Zeng, Kamath, Wygonski, & Ekelid, 1995). The fewer spectral bands are used for noise-vocoding, the stronger the degradation and the lower the resulting intelligibility of speech materials. In the present study, 1, 4, and 32 spectral bands were used for vocoding. As we know from previous studies that a logarithmic increase in spectral bands of noise-vocoded distractors linearly increases the degree of distraction (Wöstmann et al., 2017; Wöstmann & Obleser, 2016), we regressed out noise-vocoding (coded 1, 2, and 3 for 1, 4, and 32 spectral bands, respectively) in the statistical analysis (see below).

Second, half of the distractor sentences ended in a final word that was highly predictable (e.g. "She covers the bed with fresh sheets"; translated from German), whereas the final word was unpredictable in the other half (e.g. "We are very happy about the sheets"). Since we found in a previous study (Wöstmann & Obleser, 2016) and in the current dataset (Wöstmann et al., 2017) that final word predictability of distractor sentences had no effect on performance in the Irrelevant-Speech Task, predictability was not further considered in the present study.

EEG recording and preprocessing in the main study. The EEG was recorded at 64 active scalp electrodes (Ag/Ag-Cl; ActiChamp, Brain Products, München, Germany) at a sampling rate of 1,000 Hz, with a DC–280 Hz bandwidth, against a left-mastoid reference (channel TP9). All electrode impedances were kept below 5 kOhm. Offline, the continuous EEG data were filtered (0.3-Hz high-pass; 180-Hz low-pass) and segmented into epochs relative to the onset of the first target number (–2 to +16 s). As described in detail in (Wöstmann et al., 2017), an independent component analysis (ICA) and subsequent automatic artefact rejection were used to clean the EEG data.

For the purpose of the present study, the single-trial epoched EEG data were aligned to the onset of the task-irrelevant speech distractor (–2 to +4 s), re-referenced to the average of all electrodes, low-pass filtered at 10 Hz in order to obtain robust single-trial distractor-

evoked ERPs, and baseline corrected by subtraction of mean voltage in the time window

–0.1 to 0 s relative to distractor onset. For the EEG data analyses we used the Fieldtrip

toolbox (version 2018-06-14; Oostenveld, Fries, Maris, & Schoffelen, 2011) for Matlab

(R2018a) and customized Matlab scripts.

Analysis of periodic modulation of recall behaviour and distractor-evoked N1

amplitude in the main study. Each participant performed 180 experimental trials. After

removal of artefactual EEG epochs, an average of 153 trials (SD = 23.4) per participant

remained for the analyses of behavioural and EEG data.

The two major outcome variables of interest were memory recall accuracy and neural

distractor encoding. The former was quantified as the proportion of numbers recalled at

their respective positions of presentation; the latter was quantified as the mean EEG

amplitude at electrode Cz in the time interval 0.09–0.13 s relative to distractor onset. This

time interval was determined by inspection of the grand-average ERP waveform (see Fig.

1D). To quantify and visualize memory recall and N1 amplitude as a function of distractor

onset time, both of these single-trial measures were averaged for a 0.1-s long sliding

window, which was moved in 0.01-s steps over distractor onset time for each participant

(resulting in 71 time bins; see Fig. 1D).

To quantify rhythmic modulation of memory recall and distractor-evoked N1 across

distractor onset time, linear mixed-effects models were used to regress single-trial outcome

measures on circular sine and cosine predictors (a method that also proved superior in a

recent article that compared different approaches to analyse phasic modulation of neural

and behavioural responses; Zoefel, Davis, Valente, & Riecke, 2019).

Before the analysis of rhythmic memory recall modulation, single-trial accuracy (ranging

from 0 to 9 out of 9 correctly recalled numbers) was logit transformed. Since the logit for

proportions 0 and 1 is undefined, the following adjusted equation was used to logit-

transform single-trial proportion correct (PC) values (Fox & Weisberg, 2019):

$$logit_{adj}(PC) = ln\left(\frac{0.5+0.95\cdot(PC-0.5)}{1-(0.5+0.95\cdot(PC-0.5))}\right) \quad \text{(Eq. 1)}$$

Prior to mixed-effects model analyses, predictor and outcome variables were z-

transformed.

We first regressed out potential effects of auditory distractor vocoding, by regression of

single-trial accuracy and N1 amplitude on vocoding level plus random subject intercept,

using the *fitlme* function in Matlab and the model formulae 'Accuracy ~ Vocoding +

(1|Subject ID)' and 'N1 amplitude ~ Vocoding + (1|Subject ID)', respectively. Next, we

regressed the residuals of these regression models on sine- and cosine transformed

distractor onset times, using the model formulae 'Accuracy residuals ~ Sine predictor +

Cosine predictor + (1|Subject ID)' and 'N1 residuals ~ Sine predictor + Cosine predictor +

(1|Subject ID)'. Sine- and cosine-transformed distractor onsets for a given frequency *freq*

were determined using the formulae:

$Sine\ predictor = \sin(2 \cdot \pi \cdot freq \cdot Distractor\ onset)$ (Eq. 2)

$Cosine\ predictor = \cos(2 \cdot \pi \cdot freq \cdot Distractor\ onset)$ (Eq. 3)

This step was iteratively repeated for frequencies 0.5 to 8 Hz in steps of 0.25 Hz. For each

frequency, spectral magnitude was computed from the resulting coefficients (coef) for the

Sine and Cosine predictor using the formula:

$Spectral\ Magnitude = \sqrt{Sine\ coef^2 + Cosine\ coef^2}$ (Eq. 4)

Significance of rhythmic modulation of measures of distraction (accuracy & N1

amplitude) was derived using a permutation approach. For each frequency, we compared

the empirical value of spectral magnitude to 5,000 permutations of the data, generated by

randomly shuffling single-trial sine- and cosine-transformed distractor onset time within

individual participants. Spectral magnitude was considered significantly larger than

expected under the null hypothesis if the empirical value of spectral magnitude for a given

frequency exceeded the 95th percentile of permuted data (i.e. one-sided testing).

Analysis of joint modulation of behaviour and N1 amplitude in the main study. We

computed the cross-correlation (using the *xcorr* function in Matlab) of average binned

accuracy and N1 amplitude as a function of distractor onset time. We also calculated the

cross power spectral density of these two measures (using the *cpsd* function in Matlab;

rectangular window length = 0.3 s, window overlap = 0.1 s). Empirical cross-correlation and

cross power spectral density were compared to surrogate data, generated by 5,000

permutations of single-trial distractor onset time within single participants before binning
the data.

In order to test whether a joint 2.5-Hz rhythm, which turned out most prominent in the
cross power spectral density analysis, modulates memory recall accuracy directly or
indirectly (mediated via N1 amplitude), we performed a mediation analysis. To this end, we
extracted the 2.5-Hz phase ($\varphi_{2.5\,Hz}$) of the mixed-effects model to regress single trial accuracy
on sine-and cosine-transformed distractor onset time (using the *atan* function in Matlab to
calculate the inverse tangent) using the formula:

$$\varphi_{2.5Hz} = \text{atan}\left(\frac{Cosine\ coef}{Sine\ coef}\right) \quad \text{(Eq. 5)}$$

To estimate individual path coefficients between the 2.5-Hz rhythm, average accuracy,
and N1 amplitude binned for distractor onset time, linear regression analyses were used
(using the *fitlm* function in Matlab). Significance of the *indirect path*, which would link the
2.5-Hz rhythm and memory recall accuracy via the mediator distractor-evoked N1
amplitude, was calculated in the *medmod* package in Jamovi (version 1.1.9).

Task design, stimuli and statistical analysis in the follow-up experiment. In order to test
whether the results of the main experiment would conceptually replicate in a task that does
not contain rhythmic presentation of to-be-remembered target stimuli, we used a visual
match-to-sample task in a behavioural follow-up experiment. On each trial, participants
were presented with a central black fixation cross on a screen with grey background, flanked
by two line-figures, each one of which was generated by connecting 20 randomly drawn
points with black lines. Edge length of line-figures was ~4.05 * 4.05 cm and each line figure
was displaced from the central fixation cross by ~0.405 cm. In order that the entire visual
display did not exceed 5° visual angle, participants were seated at a distance of ~102 cm
from the screen.

Participants had the task to encode both line-figures into visual working memory while
remaining central fixation. Line-figures were presented for a duration of 0.3 s. In order to
reset the phase of a potential rhythm that samples the two line-figures in memory, the
corners of one of the two line-figures were highlighted in white within the last 0.05 s of the
encoding period. Note that such a "flash event" was also used in previous studies that tested

the rhythmic attentional sampling of visual target objects (e.g., Fiebelkorn et al., 2013;
Landau & Fries, 2012).

During the ensuing 3-s retention period, participants kept the line-figures in memory
while being distracted by a broad-band noise distractor (0.1–5 kHz) of 1-s duration, with
0.01-s linear onset and offset ramps. The distractor onset varied across 24 linearly spaced
time points in the time interval 0.5–1.5 s relative to the end of the encoding period. In the
end of a trial, one line-figure was presented on the left or right side of the fixation cross.
Participants had to indicate via button press whether the line-figure matched the one
presented on the same side during encoding. Note that the follow-up experiment also
contained trials without a distractor stimulus, as well as trials with the flash event on the side
opposite to the probed line-figure. These trials were excluded from the present analysis. The
experiment was implemented in Psychtoolbox (Brainard, 1997) for Matlab. Auditory
materials were presented via Sennheiser HD-280 Pro headphones.

The follow-up experiment was designed to increase the signal-to-noise ratio in the data
by testing relatively few participants on a huge number of experimental trials. Participants
($N = 6$) were tested in two data recording sessions of ~3.5–4 h duration, each (on two
different days). For one participant, data of the first session were excluded due to a
misunderstanding of task instructions. One other participant did not finish the experiment,
so that 77 trials were missing. On average, 2,537 trials (SD: 566) of each participant were
included in the present analysis, corresponding to an average of 105.7 trials per participant
and distractor onset time. Note that the approach to collect rich behavioural data of few
individuals has been adopted also in previous studies that investigated rhythmic sampling
of target stimuli in monkeys (Fiebelkorn et al., 2018) and in humans (Tomassini, Spinelli,
Jacono, Sandini, & Morrone, 2015).

For the statistical analysis, we used the same procedure as for the main study with the
following exceptions. First, no factors were regressed out prior to regression of accuracy on
sine- and cosine- transformed distractor onset time. Second, since single-trial accuracy was
binary (1 = correct response; 0 = incorrect response), we used logistic regression (using the
fitglme function in Matlab for a *binomial* distribution of the response variable and a *logit* link
function).

For visualization purposes, average accuracy was calculated for each participant and
distractor onset time, followed by smoothing via replacement of accuracy for bin_b by: 0.25 *

accuracy bin_{b-1} + 0.25 * accuracy bin_{b+1} + 0.5 * accuracy bin_b. Then, smoothed accuracy data

were averaged across participants, detrended, and z-transformed.

Results and Discussion

In the main study, participants (*N* = 23) retained the serial order of nine numbers in working memory, while being distracted by a task-irrelevant spoken sentence of varying temporal onset across trials (Fig. 1A&B). As it is typical for an Irrelevant-Speech Task (e.g. D. M. Jones, Macken, & Murray, 1993; Kreitewolf, Wöstmann, Tune, Plöchl, & Obleser, 2019; Wöstmann & Obleser, 2016), primacy and recency effects in working memory (Henson, 1998) were reflected in relatively high recall accuracy for early number positions (1–4) and for the last number position, respectively (Fig. 1C).

Figure 1. (**A**) Irrelevant-Speech Task. During retention of numbers in memory, participants were distracted by a task-irrelevant spoken sentence. At the end of each trial, participants had the task to select the numbers from a visual display in the order of presentation. (**B**) Histogram of distractor onset times across all (N = 23) participants. (**C**) Heatmap shows average recall accuracy as a function of number position (x-axis) and binned distractor onset time (y-axis). Blue lines show marginal means. Shaded areas show ±1 between-subject standard error of the mean (SEM). (**D**) Top: Grand-average distractor-evoked event-related potential (ERP) at electrode Cz. The topographic map shows the N1 component in the time window .09–.13 s after distractor onset (red shaded area). Bottom: Lines show z-transformed grand-average proportion correct (blue) and N1 amplitude (red) as a function of binned distractor onset time. Note that negative N1 amplitude values (referring to stronger distractor encoding) are plotted upwards.

Distractor onset periodically modulates target recall and distractor encoding. In order to test whether the vulnerability of working memory to distraction is rhythmic, we investigated two outcome measures of interest. First, on the behavioural level, serial recall

accuracy averaged across number positions was thought to be inversely related to the
degree of distraction. That is, the more a participant is distracted by irrelevant speech, the
lower the recall accuracy on an individual trial. Second, on the neural level, the amplitude of
the N1 component of the distractor-evoked ERP was utilized as a measure of neural
distractor encoding.

Both average recall accuracy and N1 amplitude fluctuated periodically across distractor
onset times (Fig. 1D), with such distractor onsets associated with stronger distractor
encoding (more negative N1 amplitude) coinciding with lower recall accuracy. As to be
expected for two measures of the same underlying construct (i.e. distraction), average serial
recall accuracy and N1 amplitude binned for distractor onset time were significantly
correlated ($r = 0.536$; $p < 0.001$).

To statistically analyse the periodic modulation of both measures of distraction, we used
linear mixed-effects models to regress single-trial accuracy and N1 amplitude on sine- and
cosine-transformed distractor onset time (Fig. 2A–C; see Materials and Methods for details).
Empirical spectra from mixed-effects models were contrasted against surrogate spectra,
derived from 5,000 permutations of single-subject distractor onset time. We found
significant periodic modulation of both measures by distractor onset time, for memory recall
accuracy at frequencies ~1.25–2.5 Hz and for N1 amplitude at frequencies ~3–3.75 Hz (Fig.
2D&E).

While previous research has shown that the attentional sampling of exogenous target
stimuli (Fiebelkorn et al., 2013; Landau & Fries, 2012) and target stimuli retained in working
memory is rhythmic (Peters et al., 2018), we demonstrate here that the vulnerability of
working memory to distraction is rhythmic as well. Mechanistically, a straight-forward
interpretation of the present results is that the suppression of distractors fluctuates
rhythmically, which results in alternating periods of higher versus lower vulnerability to
distraction. If the onset of a distractor stimulus falls into a period of high distractibility (i.e.
low distractor suppression), neural encoding of the distractor is enhanced, resulting in
stronger memory disruption. Such an interpretation implies that distractor encoding and
memory recall are co-modulated by a joint rhythm, which we tested next.

Figure 2. (**A–C**) Schematic depiction of the analysis procedure for periodic modulation of single-trial measures of distraction. (**A**) Dots show single-trial accuracy as function of distractor onset for one exemplary participant. For visualization, the shaded line shows accuracy averaged across twenty neighbouring distractor onsets. (**B**) For statistical analysis, sine- and cosine-transforms of distractor onset time were used as predictors in a linear regression to model single-trial accuracy. Spectral magnitude was calculated as the square root of the sum of squared sine and cosine estimates. Variables with uppercase names denote vectors that contain multiple entries; lowercase variables contain a single entry. (**C**) Spectral magnitudes derived from multiple linear models for frequencies 0.5–8 Hz. The unit circle shows sine/cosine estimates and spectral magnitude (length of pink arrow) for a frequency of 2.5 Hz. (**D**) Solid line shows spectral magnitude of periodic modulation of single-trial accuracy by distractor onset (derived from linear mixed-effects models). Shaded area shows the 95th percentile of surrogate spectra (derived from 5,000 permutations of single-trial distractor onset time within single participants). (**E**) Same as D but for periodic modulation of N1 amplitude.

A joint rhythm modulates distractor encoding and memory recall. We tested to what degree memory recall accuracy and the distractor-evoked N1 amplitude were periodically modulated by a joint underlying rhythm. To this end, we employed average accuracy and N1 amplitude binned for distractor onset time (shown in Fig. 1D) and computed their cross-correlation and cross power spectral density, which significantly exceeded surrogate data (derived from 5,000 permutations of single-subject distractor onset time) at frequencies of ~1.5–4 Hz and peaking at 2.5 Hz (Fig. 3). These results are in line with the notion that a common slow-oscillatory rhythm (here, 2.5 Hz) modulates memory recall accuracy and the distractor-evoked N1 amplitude.

Figure 3. (A) Lines show average N1 amplitude (red), accuracy (blue), and the 2.5-Hz rhythm (orange) extracted from the mixed-effects model to regress single-trial accuracy on 2.5-Hz sine- and cosine-transformed distractor onset time. (B) Line shows cross-correlation of average recall accuracy and N1 amplitude binned for distractor onset time. Shaded area shows 95th percentile of surrogate cross-correlations, computed on surrogate data (derived from 5,000 permutations of single-subject distractor onset times). (C) Line shows cross power spectral density of average recall accuracy and N1 amplitude binned for distractor onset time. Shaded area shows the 95th percentile of surrogate data (computed for the same permutations as in B). (D) Depiction of mediation analysis. Numbers at arrows show beta-coefficients of linear regression models. The total effect of the 2.5-Hz rhythm on memory recall accuracy ($\beta = 0.610$) was weakened when N1 amplitude was controlled for (*direct effect*; $\beta = 0.472$), resulting in a significant partial mediation via N1 amplitude (*indirect effect*; $\beta = 0.139$). *** $p < 0.001$.

If the rhythmic modulation of distractor encoding drives synchronized modulation of memory recall accuracy, the 2.5-Hz periodic modulation of memory recall should be mediated by the distractor-evoked N1 amplitude. To test this, we performed a mediation analysis. We extracted the phase of 2.5-Hz rhythm from the mixed-effects model to regress single-trial accuracy on 2.5-Hz sine- and cosine-transformed distractor onset time ($\varphi_{2.5\,Hz} = 2.79$). The 2.5-Hz rhythm significantly predicted N1 amplitude ($\beta = 0.399$; $p < 0.001$) and memory recall (*total effect*; $\beta = 0.610$; $p < 0.001$). Importantly, the latter effect was significantly reduced when the regression model included N1 amplitude as an additional predictor (*direct effect*; $\beta = 0.472$; $p < 0.001$). A significant indirect effect (*total effect − direct*

effect; β = 0.139; p = 0.009) revealed that the 2.5-Hz rhythmic modulation of memory recall

accuracy was partially mediated by the distractor-evoked N1 amplitude (22.7 % mediation).

This supports the view that periodic fluctuation of distractibility induces synchronized states

of increased versus decreased neural distractor encoding, which in turn result in increased

versus decreased working memory disruption, respectively.

These results provide converging behavioural and neural evidence that a 2.5-Hz rhythm

governs the degree to which task-irrelevant stimuli disrupt working memory (see also next

section). Notably, however, rhythmic vulnerability to memory distraction found in the

present study appears to be somewhat slower than rhythmic sampling of target stimuli,

which is typically observed in the theta range of 3–8 Hz (Fiebelkorn & Kastner, 2019). Why

would that be?

Recent neuroscience work suggests that distractor suppression is not the mere inverse of

target enhancement, but an independent neuro-cognitive mechanism that operates on

distractors in the sensory environment (Noonan et al., 2016; Wöstmann, Alavash, & Obleser,

2019), as well as on distractors in working memory (Schneider, Goddertz, Haase, Hickey, &

Wascher, 2019). In this view, it is well conceivable that rhythms of different frequency

orchestrate the dynamic sampling of target stimuli (at 3–8 Hz) versus suppression of

distractors (at ~2.5 Hz).

Alternatively, it might be that the nature of the present experimental paradigm induced

rhythmic distractibility at 2.5 Hz. In contrast to previous studies that investigated rhythmic

sampling of visual targets, the present study used a serial recall working memory task in the

auditory modality. Furthermore, the rhythmic presentation of target stimuli in the present

study might have entrained distractibility to a certain extent. To test whether these

alternative explanations can be ruled out, we next analysed the data of a behavioural follow-

up experiment.

The origin of rhythmic distractibility. An obvious question is whether the observed

rhythmic modulation of memory distractibility is entrained by external sensory stimulation

or whether it reflects a spontaneous internal rhythm underlying memory function.

Although the present study cannot provide a definite answer to this question, we address

both options below.

In theory, it is possible that the rhythmic presentation of target numbers entrained an

attending rhythm at 1.33 Hz, which periodically modulated also the distractibility during

memory retention. Consistent with this notion, distractor onset time modulated recall
accuracy at frequencies including this attending rhythm (~1.25–2.5 Hz). Furthermore, the
time point of highest memory distractibility (at ~1.26 s following target offset; see Fig. 1D)
falls precisely in-between two peaks of the hypothetical 1.33-Hz attending rhythm (i.e., in-
between the onsets of putative number positions 11 and 12). However, the joint periodic
modulation of accuracy and distractor-evoked N1 amplitude was strongest at a frequency
of 2.5 Hz rather than 1.33 Hz, which speaks against the view of entrained distractibility.

Alternatively, the observed ~2.5-Hz rhythm might be a spontaneous internal rhythm
underlying memory function (see Fig 3D). If this is the case, a similar rhythm should be
observed also in the absence of a potentially entraining stimulus sequence. In a behavioural
follow-up experiment, we used a visual match-to-sample task to test this hypothesis. While
participants retained two line-figures in working memory, they were distracted by an
auditory noise distractor of varying onset time (Fig. 4). Distractor onset time periodically
modulated task accuracy at a frequency of 2.75 Hz, which supports the claim that memory
distractibility fluctuates rhythmically even when no rhythmic sensory stimulation precedes
working memory retention. Note that in addition to a frequency of 2.75 Hz, accuracy in the
follow-up experiment was modulated at lower frequencies (< 1 Hz), most likely because
distractor onset time linearly modulated task accuracy. Later distractors were more
detrimental to accuracy (linear effect of absolute distractor onset time; $\beta = -.062$; $p = 0.001$
in a logistic regression of single-trial accuracy).

The results of this follow-up experiment constitute an analogy of distractor suppression
to what has previously been shown for the attentional sampling of target stimuli (Fiebelkorn
et al., 2013; Landau & Fries, 2012): there is an inherent (~2.5-Hz) rhythm that regulates the
temporally dynamic suppression of task-irrelevant distraction during working memory
maintenance. Relatively slow oscillatory effects in the delta frequency band (< 4 Hz) have
also been related to the sequential prioritization in working memory (de Vries, van Driel, &
Olivers, 2019). One candidate neural structure where the inherent rhythm that modulates
distractibility might originate is the hippocampus, where endogenous delta oscillations
have been found to orchestrate synchronized states of working memory activation
(Leszczynski, Fell, & Axmacher, 2015). In this sense, the ~2.5-Hz rhythm of distractor
suppression might be a temporally dynamic mechanism specific to working memory. That
is, during memory retention of target items, suppression of distraction is not a constant

process but instead modulated at ~2.5 Hz, which allows for (partial) memory intrusion of

currently task-irrelevant stimuli approximately every 400 ms.

The fact that the present study found periodic modulation of distractor suppression at a

frequency of ~2.5-Hz does not preclude that the rate of distractor suppression might adapt

to task demands. For instance, we found rhythmic modulation of neural indices of

attentional filtering during stimulus encoding (i.e. target selection and distractor

suppression) in previous studies at 0.67 Hz (Wöstmann, Herrmann, Maess, & Obleser, 2016)

and 0.375 Hz (Wöstmann, Schmitt, & Obleser, 2020), corresponding to the respective rates

of stimulus presentation.

Figure 4. (A) Design of behavioural follow-up experiment. Participants (N = 6) encoded two line-figures, before
one of them was highlighted by a flash event. During the ensuing 3-s memory retention period, an auditory
distractor was presented at one of 24 linearly spaced delays between 0.5 and 1.5 s. In the end of a trial,
participants had to indicate whether the probe figure matched the one presented on the same side during
encoding (response timeout: 2 s). **(B)** Line shows spectral magnitude of periodic modulation of single-trial
accuracy by distractor onset. Shaded area shows the 95th percentile of surrogate spectra (derived from 5,000
permutations of single-trial distractor onset time within single participants). **(C)** The blue line shows average
accuracy (which was temporally smoothed, detrended, and z-transformed for purpose of visualization). The
orange line shows the 2.75-Hz rhythm derived from the mixed model to regress accuracy on sine- and cosine-
transformed distractor onset time.

Limitations. The concept of rhythmicity implies that the mechanism of interest, e.g.

attentional sampling of targets or vulnerability to distraction, is rhythmically modulated

over an extended period of time. However, for practical reasons, the present study varied

the distractor onset within relatively short time windows of 0.8 s in the main study and 1 s

in the follow-up experiment. In fact, short time windows of investigation are a common

limitation also for studies exploring the rhythmic attentional sampling of targets (1.3 s in

Fiebelkorn et al., 2018; 0.8 s in Fiebelkorn et al., 2013; 1.2 s in Helfrich et al., 2018; 1 s in Landau

& Fries, 2012; 0.8 s in Peters et al., 2018). Thus, although the present results suggest memory
distractibility to be a rhythmic process, it is at present an open question to what extent this
rhythmicity extends over longer time intervals.

Furthermore, it must be noted that the present results suggest the effect of rhythmic
working memory distraction to be relatively small. In the main experiment, the maximum
and minimum of average proportion correct as a function of distractor onset differed only
by 0.052 (0.069 in the follow-up experiment). For comparison, the range of average recall
accuracy as a function of number position, which is known to be a large effect in the
Irrelevant-Speech Task, was 0.518. While we consider it likely that the former effect could be
increased in future studies that employ task-irrelevant stimuli of higher distractibility, it must
also be considered that the rhythmic modulation of vulnerability to distraction might be a
small effect of limited practical significance.

Conclusion. The primary goal the present study was to enhance our understanding of how
the human neuro-cognitive system orchestrates the suppression of task-irrelevant
distraction dynamically in time. Convergingly, the results of two experiments indicate that
during retention of target information in working memory, the suppression of distractors
fluctuates at a rate of ~2.5 Hz. Neurally, this effect was in part mediated by the strength of
distractor encoding, which occurred in sync with behavioural indices of distraction.

The present study sets the stage for future work to test (i) whether the observed rhythmic
distractor suppression can be entrained by external stimulation, (ii) whether top-down
executive control has the potency to modulate rhythmic distractor suppression to optimise
task goals, and (iii) to what degree rhythmic distractor suppression is independent of
rhythmic target sampling and how these two interact.

References

Albouy, P., Weiss, A., Baillet, S., & Zatorre, R. J. (2017). Selective Entrainment of Theta Oscillations in the Dorsal Stream Causally Enhances Auditory Working Memory Performance. *Neuron, 94*(1), 193-206 e195. doi:10.1016/j.neuron.2017.03.015

Awh, E., Vogel, E. K., & Oh, S. H. (2006). Interactions between attention and working memory. *Neuroscience, 139*(1), 201-208. doi:10.1016/j.neuroscience.2005.08.023

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spat Vis, 10*(4), 433-436.

Colle, H. A., & Welsh, A. (1976). Acoustic Masking in Primary Memory. *Journal of Verbal Learning and Verbal Behavior, 15*(1), 17-31.

de Graaf, T. A., Gross, J., Paterson, G., Rusch, T., Sack, A. T., & Thut, G. (2013). Alpha-band rhythms in visual task performance: phase-locking by rhythmic sensory stimulation. *PLoS One, 8*(3), e60035. doi:10.1371/journal.pone.0060035

de Vries, I. E. J., van Driel, J., & Olivers, C. N. L. (2019). Decoding the status of working memory representations in preparation of visual selection. *Neuroimage, 191*, 549-559. doi:10.1016/j.neuroimage.2019.02.069

Erb, J., Henry, M. J., Eisner, F., & Obleser, J. (2012). Auditory skills and brain morphology predict individual differences in adaptation to degraded speech. *Neuropsychologia, 50*(9), 2154-2164. doi:10.1016/j.neuropsychologia.2012.05.013

Fiebelkorn, I. C., & Kastner, S. (2019). A Rhythmic Theory of Attention. *Trends Cogn Sci, 23*(2), 87-101. doi:10.1016/j.tics.2018.11.009

Fiebelkorn, I. C., Pinsk, M. A., & Kastner, S. (2018). A Dynamic Interplay within the Frontoparietal Network Underlies Rhythmic Spatial Attention. *Neuron, 99*(4), 842-853 e848. doi:10.1016/j.neuron.2018.07.038

Fiebelkorn, I. C., Saalmann, Y. B., & Kastner, S. (2013). Rhythmic sampling within and between objects despite sustained attention at a cued location. *Curr Biol, 23*(24), 2553-2558. doi:10.1016/j.cub.2013.10.063

Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression* (3rd Edition ed.): Sage.

Gundlach, C., Muller, M. M., Nierhaus, T., Villringer, A., & Sehm, B. (2016). Phasic Modulation of Human Somatosensory Perception by Transcranially Applied Oscillating Currents. *Brain Stimul, 9*(5), 712-719. doi:10.1016/j.brs.2016.04.014

Hanslmayr, S., Axmacher, N., & Inman, C. S. (2019). Modulating Human Memory via Entrainment of Brain Oscillations. *Trends Neurosci, 42*(7), 485-499. doi:10.1016/j.tins.2019.04.004

Helfrich, R. F., Fiebelkorn, I. C., Szczepanski, S. M., Lin, J. J., Parvizi, J., Knight, R. T., & Kastner,
S. (2018). Neural Mechanisms of Sustained Attention Are Rhythmic. *Neuron, 99*(4),
854-865 e855. doi:10.1016/j.neuron.2018.07.032

Helfrich, R. F., Schneider, T. R., Rach, S., Trautmann-Lengsfeld, S. A., Engel, A. K., & Herrmann,
C. S. (2014). Entrainment of brain oscillations by transcranial alternating current
stimulation. *Curr Biol, 24*(3), 333-339. doi:10.1016/j.cub.2013.12.041

Henry, M. J., & Obleser, J. (2012). Frequency modulation entrains slow neural oscillations and
optimizes human listening behavior. *Proc Natl Acad Sci U S A, 109*(49), 20095-20100.
doi:10.1073/pnas.1213390109

Henson, R. N. (1998). Short-term memory for serial order: the Start-End Model. *Cogn Psychol,
36*(2), 73-137. doi:10.1006/cogp.1998.0685

Hsieh, L. T., Ekstrom, A. D., & Ranganath, C. (2011). Neural oscillations associated with item
and temporal order maintenance in working memory. *J Neurosci, 31*(30), 10803-
10810. doi:10.1523/JNEUROSCI.0828-11.2011

Jones, D. M., Macken, W. J., & Murray, A. C. (1993). Disruption of visual short-term memory
by changing-state auditory stimuli: the role of segmentation. *Mem Cognit, 21*(3), 318-
328.

Jones, D. M., & Morris, N. (1992). Irrelevant speech and serial recall: implications for theories
of attention and working memory. *Scand J Psychol, 33*(3), 212-229.

Jones, M. R., Johnston, H. M., & Puente, J. (2006). Effects of auditory pattern structure on
anticipatory and reactive attending. *Cogn Psychol, 53*(1), 59-96.
doi:10.1016/j.cogpsych.2006.01.003

Kreitewolf, J., Wöstmann, M., Tune, S., Plöchl, M., & Obleser, J. (2019). Working-memory
disruption by task-irrelevant talkers depends on degree of talker familiarity. *Atten
Percept Psychophys, 81*(4), 1108-1118. doi:10.3758/s13414-019-01727-2

Köster, M., Martens, U., & Gruber, T. (2019). Memory entrainment by visually evoked theta-
gamma coupling. *Neuroimage, 188*, 181-187. doi:10.1016/j.neuroimage.2018.12.002

Lakatos, P., Gross, J., & Thut, G. (2019). A New Unifying Account of the Roles of Neuronal
Entrainment. *Curr Biol, 29*(18), R890-R905. doi:10.1016/j.cub.2019.07.075

Landau, A. N., & Fries, P. (2012). Attention samples stimuli rhythmically. *Curr Biol, 22*(11),
1000-1004. doi:10.1016/j.cub.2012.03.054

Landau, A. N., Schreyer, H. M., van Pelt, S., & Fries, P. (2015). Distributed Attention Is
Implemented through Theta-Rhythmic Gamma Modulation. *Curr Biol, 25*(17), 2332-
2337. doi:10.1016/j.cub.2015.07.048

Large, E. W., & Jones, M. R. (1999). The dynamics of attending: How people track time-varying
events. *Psychological Review, 106*, 119-159.

Leszczynski, M., Fell, J., & Axmacher, N. (2015). Rhythmic Working Memory Activation in the
Human Hippocampus. *Cell Rep, 13*(6), 1272-1282. doi:10.1016/j.celrep.2015.09.081

Lim, S. J., Wöstmann, M., & Obleser, J. (2015). Selective Attention to Auditory Memory
Neurally Enhances Perceptual Precision. *J Neurosci, 35*(49), 16094-16104.
doi:10.1523/JNEUROSCI.2674-15.2015

Nobre, A. C., & van Ede, F. (2018). Anticipated moments: temporal structure in attention. *Nat
Rev Neurosci, 19*(1), 34-48. doi:10.1038/nrn.2017.141

Noonan, M. P., Adamian, N., Pike, A., Printzlau, F., Crittenden, B. M., & Stokes, M. G. (2016).
Distinct Mechanisms for Distractor Suppression and Target Facilitation. *J Neurosci,
36*(6), 1797-1807. doi:10.1523/JNEUROSCI.2133-15.2016

Oberauer, K., & Hein, L. (2012). Attention to Information in Working Memory. *Current
Directions in Psychological Science, 21*(3), 164-169. doi:10.1177/0963721412444727

Obleser, J., & Kayser, C. (2019). Neural Entrainment and Attentional Selection in the Listening
Brain. *Trends Cogn Sci, 23*(11), 913-926. doi:10.1016/j.tics.2019.08.004

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software
for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput
Intell Neurosci, 2011*, 156869. doi:10.1155/2011/156869

Peters, B., Rahm, B., Kaiser, J., & Bledowski, C. (2018). Attention fluctuates rhythmically
between objects in working memory. *Journal of Vision, 18*(186).
doi:https://doi.org/10.1167/18.10.186

Schneider, D., Goddertz, A., Haase, H., Hickey, C., & Wascher, E. (2019). Hemispheric
asymmetries in EEG alpha oscillations indicate active inhibition during attentional
orienting within working memory. *Behav Brain Res, 359*, 38-46.
doi:10.1016/j.bbr.2018.10.020

Schneider, D., Mertes, C., & Wascher, E. (2016). The time course of visuo-spatial working
memory updating revealed by a retro-cuing paradigm. *Sci Rep, 6*, 21442.
doi:10.1038/srep21442

Shannon, R. V., Zeng, F. G., Kamath, V., Wygonski, J., & Ekelid, M. (1995). Speech recognition
609   with primarily temporal cues. *Science, 270*(5234), 303-304.

610   Spaak, E., de Lange, F. P., & Jensen, O. (2014). Local entrainment of alpha oscillations by visual
611   stimuli causes cyclic modulation of perception. *J Neurosci, 34*(10), 3536-3544.
612   doi:10.1523/JNEUROSCI.4385-13.2014

613   Tomassini, A., Spinelli, D., Jacono, M., Sandini, G., & Morrone, M. C. (2015). Rhythmic
614   oscillations of visual contrast sensitivity synchronized with action. *J Neurosci, 35*(18),
615   7019-7029. doi:10.1523/JNEUROSCI.4568-14.2015

616   Wöstmann, M., Alavash, M., & Obleser, J. (2019). Alpha oscillations in the human brain
617   implement distractor suppression independent of target selection. *J Neurosci, 39*(49),
618   9797-9805. doi:10.1523/JNEUROSCI.1954-19.2019

619   Wöstmann, M., Herrmann, B., Maess, B., & Obleser, J. (2016). Spatiotemporal dynamics of
620   auditory attention synchronize with speech. *Proc Natl Acad Sci U S A, 113*(14), 3873-
621   3878. doi:10.1073/pnas.1523357113

622   Wöstmann, M., Lim, S. J., & Obleser, J. (2017). The Human Neural Alpha Response to Speech
623   is a Proxy of Attentional Control. *Cereb Cortex, 27*(6), 3307-3317.
624   doi:10.1093/cercor/bhx074

625   Wöstmann, M., & Obleser, J. (2016). Acoustic detail but not predictability of task-irrelevant
626   speech disrupts working memory. *Front Hum Neurosci, 10*(538).
627   doi:10.3389/fnhum.2016.00538

628   Wöstmann, M., Schmitt, L. M., & Obleser, J. (2020). Does Closing the Eyes Enhance Auditory
629   Attention? Eye Closure Increases Attentional Alpha-Power Modulation but Not
630   Listening Performance. *J Cogn Neurosci, 32*(2), 212-225. doi:10.1162/jocn_a_01403

631   Zoefel, B., Davis, M. H., Valente, G., & Riecke, L. (2019). How to test for phasic modulation of
632   neural and behavioural responses. *Neuroimage, 202*, 116175.
633   doi:10.1016/j.neuroimage.2019.116175

634