1 **SRRF 'n' TIRF - FCS: Simultaneous spatiotemporal super-resolution microscopy**
2

3 **Jagadish Sankaran[1],*, Harikrushnan Balasubramanian[1],*, Wai Hoh Tang[2], Xue Wen Ng[1],**
4 **Adrian Röllin[2], Thorsten Wohland[1,3]**
5

[1]Department of Biological Sciences and NUS Centre for Bio-Imaging Sciences, National
University of Singapore, 14 Science Drive 4, 117557 Singapore

[2]Department of Statistics and Applied Probability, National University of Singapore, 6 Science
Drive 2, Singapore 117546

[3]Department of Chemistry, National University of Singapore, 3 Science Drive 3, 117543
Singapore

*contributed equally to the work

13
14
15 **Abstract**
16

Super-resolution microscopy and single molecule fluorescence spectroscopy require mutually
exclusive experimental strategies optimizing either time or spatial resolution. To achieve both, we
implement a GPU-supported, camera-based measurement strategy that highly resolves spatial
structures (~60 nm), temporal dynamics (≤ 2 ms), and molecular brightness from the exact same
data set. Using epidermal growth factor receptor (EGFR), we demonstrate that pixel-wise
correlation of parameters obtained from different techniques provides new biological knowledge.

23
24 **Key Words**
25

Full knowledge of a biological system requires not only information on its spatial structure but
also its temporal dynamics. However, the acquisition of structure and dynamics requires
complementary, often mutually exclusive optimization strategies[1]. Spatial resolution depends on
the number of photons collected and thus sets a lower limit on acquisition time. Molecular
dynamics requires acquisition times shorter than the dynamics of interest and thus sets an upper
limit. Since these two limits, in general, do not lead to an overlap region, the combination of
spatiotemporal super-resolution microscopy has remained a challenge. Attempts in the past either
restrict time resolution[2,3], require specialized instrumentation[4,5] or sample labeling[6,7].

While simultaneous multi-parametric fluorescence detection (MFD) has been established for
point-measurements[8,9], such attempts in an imaging mode have been hampered by lack of
strategies that can bridge the limitations imposed by spatial and temporal resolution requirements
and by the computationally expensive data evaluation procedures required to treat the large data
sets. Here, we overcome these problems by acquiring images with high sensitivity and high-speed
using commercially available cameras and GPU-based data processing. The resulting datasets are
evaluated by a combination of spectroscopy and super-resolution techniques that include: imaging
fluorescence correlation spectroscopy (Imaging FCS)[10,11] to measure dynamics; FCS diffusion law
analysis to obtain information about the dynamic sub-resolution molecular organization[12]; Number
and Brightness analysis (N&B)[13] to determine oligomerization or aggregation states; and super-
resolution radial fluctuation microscopy (SRRF)[14] to obtain super-resolved images (Fig. S1).
Using a recently published GPU-based algorithm for SRRF[14] and a newly developed GPU
accelerated version of Imaging FCS (Fig. S2), an ImageJ plugin developed in our group, we
achieve measurements with high spatial information (~60 nm) and temporal dynamics ($\leq$ 2 ms)
from the exact same data within ~5 min, including measurement time (Figs. S3-S5, Tables 1, S1-
S2).

As a proof-of-principle, we analyzed EGFR receptor dynamics and organization and the
cytoskeletal structure on CHO-K1 cell membranes using a commercially available total internal
reflection fluorescence (TIRF) microscope. We recorded image stacks of mApple labelled EGFR
(EGFR-mApple) and LifeAct[15], an actin binding protein labelled with EGFP, on whole cells using
EMCCD and sCMOS cameras for detection (50,000 frames at 2 ms time resolution covering areas
as large as 128×128 pixels). For the simultaneous acquisition of the mApple and EGFP signals on
two halves of one camera, we used a wavelength-based image splitter.

A comparison of the structure and the dynamics of LifeAct in the green channel alone shows clear
correlations (Fig S6). The diffusion coefficient on the fibres was found to be 0.77 ± 0.53 µm^2/s.
The region between fibres contain only few lifeact particles[16]. Using 4×4 pixel binning to improve
the signal-to-noise ratio (SNR), the diffusion coefficient was 2.89 ± 3.20 µm^2/s, indicating free
diffusion of LifeAct or LifeAct bound to globular actin[15] (Fig. S7). The measured optical
resolution of SRRF was in the range of 78 - 120 nm (Table S3). Using SRRF to localize the
cytoskeleton more precisely, we can select only those pixels in the FCS diffusion coefficient map
that lie directly on top of the cytoskeleton fibres and are thus less influenced by freely diffusing
LifeAct. After using the SRRF image as a selection mask, the diffusion coefficient on the fibre
was found to be 0.80 ± 0.39 µm^2/s whereas in neighbouring pixels it was found to be 0.75 ± 0.66
µm^2/s. (Fig. S8). SRRF selection thus improves the estimated diffusion coefficient precision by
about a factor 1.4. Furthermore, the FCS results corroborate the SRRF resolved positions of the

fibres as there is a correlation between the SRRF actin fibre position and the SNR of the FCS data
(Fig. S8, Table S4).
86
The full extent of the information however is gained when analyzing both wavelength channels.
To investigate the interrelation between the actin cytoskeleton structure, which is relatively static
on the time scale of our measurements, and EGFR mobility and organization, which is highly
dynamic, we optimized the signals by spatiotemporal binning for either FCS and FCS diffusion
law, N&B (red channel, Figs. 1 a-d, Fig. S9) or SRRF (green channel, Figs. 1 e-g) analysis. SRRF
analysis led to an improvement of the FWHM from 465 nm to 97 nm of actin fibres. In the whole
cell, N&B analysis revealed that 63 ± 3 % of EGFR existed as apparent dimers (supplement). The
diffusion coefficient was found to be 0.19 ± 0.15 µm^2/s with a FCS diffusion law intercept of 2.58
s, indicating intermittent trapping of EGFR. The large error associated with the diffusion
coefficient is attributed to the inherent heterogeneity in the cell membrane. The various factors
contributing to the heterogeneity are investigated using a multi-parametric analysis.

EGFR-mApple expressing cells show both regions of uniform intensity and regions exhibiting
visible clustering (Fig. 1a). Analysis of the regions of homogeneous brightness in the mApple
channel by Imaging FCS and the FCS diffusion law resulted in a diffusion coefficient of $D = 0.24$
± 0.16 µm^2/s and a positive diffusion law intercept of 1.81 s (Fig. S10), indicating transiently
trapped diffusion and possibly EGFR oligomerization or clustering[17]. The existence of oligomers
is corroborated by N&B analysis of the same regions. EGFR-mApple showed a brightness
intermediate between monomers and dimers (Fig. S9), suggesting that 18 ± 6% of receptors are
found in dimers (Fig. S10), although we cannot exclude the existence of small amounts of higher
oligomers[18]. When analyzing the regions with visible clusters present, we obtain a $D = 0.11 \pm 0.08$
µm^2/s (Fig. S10) via FCS. In this case, a simple dimer model is not reasonable anymore. A
brightness value of 5.87 (average value of the cluster containing areas in Fig. 1 and S10) indicates
that at least trimers, if not higher oligomers, are present.

The existence of preformed oligomers is consistent with literature with up to ~65% dimerization
previously detected[18-20] in the same cell line. Diffusion law intercepts did not show a statistically
significant difference when comparing areas where visible clusters or actin is present with areas
where they were absent (Table S5, p > 0.14). This indicates that the domain partitioning of EGFR
is primarily determined by EGFR-membrane interactions, possibly sub-resolution clustering and
inter-EGFR interactions but independent of EGFR-actin interactions.

Since all evaluations stem from the exact same data set we overlap the parameter maps and
pairwise analyze their scatter plots to determine the interdependence of protein location with its
dynamics. We performed a logarithmic transformation of the entire dataset to identify power law
scaling between parameters (Fig. 1h-m).
120
Using whole cell data, we find a negative correlation between particle brightness and diffusion
coefficient (Pearson's correlation coefficient (R) = -0.34, Table S6 and Fig. 1h), indicating that
brighter particles diffuse slower than dimmer particles. This is unexpected since membrane
diffusion is only weakly dependent on size[21]. This implies either that EGFR oligomers are located
in a more viscous membrane environment, e.g. lipid domains[22] or clathrin-coated pits, or that we
detect multiple EGFR molecules in larger complexes[18,23,]. A negative correlation (R= -0.83) is also

seen between diffusion coefficient and the positive diffusion law intercept indicating that trapping
is responsible for lower EGFR mobility as expected (Table S6).
129
Brightness and SRRF do not show any correlation ($R = 0.00$, Fig. 1 l), implying that clustering is
not directly linked to the cytoskeleton. Similarly, we do not see any correlation between EGFR
diffusion and cytoskeletal structures ($R = -0.05$, Fig. 1 m), unlike in the case of LifeAct. This also
verifies our earlier observation that diffusion law intercepts are independent of the presence or
absence of actin. As it has been shown previously that EGFR dynamics changes with cytoskeleton
disruption[24], our observations suggest that it is not the static cytoskeleton itself that changes EGFR
dynamics but that the influence is indirect and cytoskeleton coupling to the plasma membrane and
resulting membrane changes are responsible for changes in EGFR dynamics[25]. Similar results were
obtained by performing the experiment using sCMOS camera as a detector (Figs. S11-S13).

Living systems are highly dynamic with processes happening on spatial scales well below the
optical diffraction limit and on time scales on the millisecond scale or faster. Therefore, the
extraction of the maximum information available on biological processes requires the
simultaneous acquisition of data with high spatiotemporal resolution. This poses particular
problems to data recording and data evaluation strategies that allow data treatment in an acceptable
time frame, ideally in real-time. We used GPU-based data evaluation and modern camera
technology to optimize data evaluation of fluorescence microscopy images by selected spatial and
temporal binning to extract multiple physicochemical parameters in parallel from a single data set,
almost in real time, using standard and widely available instrumentation. Parameter correlations
and the lack thereof provide information not available on sequentially acquired data due to the
dynamical nature of biological samples as shown on the example of EGFR. This approach is easily
extendable to other fluorescence parameters, does not require specialized instrumentation, and thus
is immediately applicable to a wide range of situations.

152
1   Weigert, M. *et al.* Content-aware image restoration: pushing the limits of fluorescence
microscopy. *Nature Methods* **15**, 1090-1097, doi:10.1038/s41592-018-0216-7 (2018).
2   Ashdown, G. W., Cope, A., Wiseman, P. W. & Owen, D. M. Molecular flow quantified
beyond the diffraction limit by spatiotemporal image correlation of structured
illumination microscopy data. *Biophysical journal* **107**, L21-23,
doi:10.1016/j.bpj.2014.09.018 (2014).
3   Kisley, L. *et al.* Characterization of Porous Materials by Fluorescence Correlation
Spectroscopy Super-resolution Optical Fluctuation Imaging. *ACS Nano* **9**, 9158-9166,
doi:10.1021/acsnano.5b03430 (2015).
4   Leutenegger, M., Ringemann, C., Lasser, T., Hell, S. W. & Eggeling, C. Fluorescence
correlation spectroscopy with a total internal reflection fluorescence STED microscope
(TIRF-STED-FCS). *Optics express* **20**, 5243-5263, doi:10.1364/OE.20.005243 (2012).
5   Hedde, P. N. *et al.* Stimulated emission depletion-based raster image correlation
spectroscopy reveals biomolecular dynamics in live cells. *Nature Communications* **4**,
2093, doi:10.1038/ncomms3093 (2013).
6   Manley, S. *et al.* High-density mapping of single-molecule trajectories with
photoactivated localization microscopy. *Nature Methods* **5**, 155-157,
doi:10.1038/nmeth.1176 (2008).

7   Strauß, S. *et al.* Modified aptamers enable quantitative sub-10-nm cellular DNA-PAINT
imaging. *Nature Methods* **15**, doi:10.1038/s41592-018-0105-0 (2018).
8   Weidtkamp-Peters, S. *et al.* Multiparameter fluorescence image spectroscopy to study
molecular interactions. *Photochemical & Photobiological Sciences* **8**, 470-480,
doi:10.1039/B903245M (2009).
9   Esposito, A. & Venkitaraman, A. R. Enhancing Biochemical Resolution by
Hyperdimensional Imaging Microscopy. *Biophysical journal* **116**, 1815-1822,
doi:https://doi.org/10.1016/j.bpj.2019.04.015 (2019).
10   Kannan, B. *et al.* Spatially resolved total internal reflection fluorescence correlation
microscopy using an electron multiplying charge-coupled device camera. *Anal. Chem.*
79, 4463-4470, doi:10.1021/ac0624546 (2007).
11   Wohland, T., Shi, X., Sankaran, J. & Stelzer, E. H. K. Single Plane Illumination
Fluorescence Correlation Spectroscopy (SPIM-FCS) probes inhomogeneous three-
dimensional environments. *Opt. Express* **18**, 10627-10641 (2010).
12   Wawrezinieck, L., Rigneault, H., Marguet, D. & Lenne, P. F. Fluorescence correlation
spectroscopy diffusion laws to probe the submicron cell membrane organization.
Biophysical journal **89**, 4029-4042, doi:10.1529/biophysj.105.067959 (2005).
13   Unruh, J. R. & Gratton, E. Analysis of molecular concentration and brightness from
fluorescence fluctuation data with an electron multiplied CCD camera. *Biophysical
journal* **95**, 5385-5398, doi:10.1529/biophysj.108.130310 (2008).
14   Gustafsson, N. *et al.* Fast live-cell conventional fluorophore nanoscopy with ImageJ
through super-resolution radial fluctuations. *Nature Communications* **7**, 12471,
doi:10.1038/ncomms12471 (2016).
15   Riedl, J. *et al.* Lifeact: a versatile marker to visualize F-actin. *Nature Methods* **5**, 605-
607, doi:10.1038/nmeth.1220 (2008).
16   Sankaran, J. *et al.* Single microcolony diffusion analysis in Pseudomonas aeruginosa
biofilms. *npj Biofilms and Microbiomes* **5**, 35, doi:10.1038/s41522-019-0107-4 (2019).
17   Abulrob, A. *et al.* Nanoscale imaging of epidermal growth factor receptor clustering:
effects of inhibitors. *The Journal of biological chemistry* **285**, 3145-3156,
doi:10.1074/jbc.M109.073338 (2010).
18   Saffarian, S., Li, Y., Elson, E. L. & Pike, L. J. Oligomerization of the EGF receptor
investigated by live cell fluorescence intensity distribution analysis. *Biophysical journal*
93, 1021-1031, doi:10.1529/biophysj.107.105494 (2007).
19   Liu, P. *et al.* Investigation of the dimerization of proteins from the epidermal growth
factor receptor family by single wavelength fluorescence cross-correlation spectroscopy.
Biophysical journal **93**, 684-698, doi:10.1529/biophysj.106.102087 (2007).
20   Zanetti-Domingues, L. C. *et al.* The architecture of EGFR's basal complexes reveals
autoinhibition mechanisms in dimers and oligomers. *Nature Communications* **9**, 4325,
doi:10.1038/s41467-018-06632-0 (2018).
21   Saffman, P. G. & Delbrück, M. Brownian motion in biological membranes. *Proceedings
of the National Academy of Sciences* **72**, 3111-3113, doi:10.1073/pnas.72.8.3111 (1975).
22   Sergé, A., Bertaux, N., Rigneault, H. & Marguet, D. Dynamic multiple-target tracing to
probe spatiotemporal cartography of cell membranes. *Nature Methods* **5**, 687-694,
doi:10.1038/nmeth.1233 (2008).

23   Nagy, P. *et al.* Activation-dependent clustering of the erbB2 receptor tyrosine kinase
detected by scanning near-field optical microscopy. *Journal of Cell Science* **112**, 1733
(1999).
24   Bag, N., Huang, S. & Wohland, T. Plasma Membrane Organization of Epidermal Growth
Factor Receptor in Resting and Ligand-Bound States. *Biophysical journal* **109**, 1925-
1936, doi:10.1016/j.bpj.2015.09.007 (2015).
25   Vogel, S. K., Greiss, F., Khmelinskaia, A. & Schwille, P. Control of lipid domain
organization by a biomimetic contractile actomyosin cortex. *eLife* **6**, e24350,
doi:10.7554/eLife.24350 (2017).
224
225
226

a Red channel TIRF image
(EGFR-mApple)

b D = 0.19 ± 0.15 µm²/s
(EGFR-mApple)

c B = 4.93 ± 0.03
Dimer fraction = 0.63 ± 0.03
(EGFR-mApple)

d $\tau_0 = 2.58$ s
Diffusion law
analysis
(EGFR-mApple)

e Green channel TIRF image
(Lifeact-EGFP)

f SRRF map (200 ms binning)
(Lifeact-EGFP)

g 465 nm 97 nm

h B-D maps merge

i SRRF-B maps merge

j SRRF-D maps merge

k R = -0.34

l R = 0.00

m R = -0.05

Fig. 1: Multi-parametric analysis from a single dual-channel fluorescence data set.
(a) TIRF image of EGFR-mApple. **(b)** Diffusion map of EGFR-mApple ($D = 0.19 \pm 0.15$ $\mu m^2/s$).
(c) Brightness map of EGFR-mApple ($B = 4.93 \pm 0.03$, dimer fraction = 0.63 ± 0.03; mean $\pm$
SEM). **(d)** Diffusion law analysis of EGFR-mApple (refer Fig. S10 for more details); intercept =
2.58 s. **(e)** TIRF image of Lifeact-EGFP. **(f)** SRRF map (200 ms binning) of Lifeact-EGFP. **(g)**
Normalized intensity profile across an actin fibre before and after SRRF microscopy. Thickness
(FWHM of the Gaussian fit) of the actin fibre before SRRF (blue curve) = 465 nm; measurement
point marked with blue line on image (e). Thickness of the actin fibre after SRRF (red curve) = 97
nm; measurement point marked with red line on image (f). **(h)** Merge of B and D maps of EGFR-
mApple. The B pixels are coloured yellow, and the D pixels are coloured magenta. Correlated
pixels between the two maps are coloured white. **(i)** Merge of SRRF (Lifeact-EGFP) and B (EGFR-
mApple) maps. SRRF pixels are coloured cyan, and B pixels are coloured yellow. Correlated pixels
between the two maps are coloured white. The SRRF map was spatially binned to the same
dimensions as the B map. **(j)** Merge of SRRF (Lifeact-EGFP) and D (EGFR-mApple) maps of
EGFR-mApple. SRRF pixels are coloured cyan, and D pixels are coloured magenta. Correlated
pixels between the two maps are coloured white. The SRRF map was spatially binned to the same
dimensions as the D map. **(k)** 2D frequency plot of log B vs log D values from image (h). **(l)** 2D
frequency plot of log SRRF vs log B values from image (i). **(m)** 2D frequency plot of log SRRF
vs log D values from image (j). The scale bars in white measure 5 µm in images (a)-(e), (h)-(j).
This figure uses cell 1 in Table S6. The reported values are averages obtained from analysis of an
entire cell.
249
250

Online methods

251
252
253 **Sample Preparation**
254
CHO-K1 (Chinese Hamster Ovary) cells (CCL-61) were obtained from ATCC (Manassas,
Virginia, USA) and were cultured in Dulbecco's Modified Eagle Medium (DMEM/High glucose
with L-glutamine, without sodium pyruvate – SH30022.FS; HyClone, GE Healthcare Life
Sciences, Utah, USA) supplemented with 1% penicillin-streptomycin (Gibco, Thermo Fisher
Scientific, Massachusetts, USA), and 10% fetal bovine serum (FBS; Gibco, Thermo Fisher
Scientific, Massachusetts, USA) at 37 °C in a 5% (v/v) CO_2 environment (Forma Steri-Cycle CO_2
incubator, Thermo Fisher Scientific, Massachusetts, USA).

The construction of the plasmids PMT-mApple, PMT-mApple$_2$ and EGFR-mApple are detailed
in the supplement. Lifeact-EGFP plasmid was a gift from Prof. Wu Min (NUS, Singapore). All
the plasmids were amplified using ZymoPURE II Plasmid Midiprep Kit (D4201; Zymo Research,
California, USA) and their concentration and purity were confirmed by a UV-Vis
spectrophotometer (NanoDrop 2000, Thermo Fisher Scientific, Massachusetts, USA). The
plasmid amounts used for transfection were 100 ng for PMT-mApple, PMT-mApple$_2$ and Lifeact-
EGFP, and 1 µg for EGFR-mApple.
269
For transfection, cell cultures that were ~90% confluent were used. The spent media removed from
the culture flask was discarded. The flask was washed twice with 5 ml PBS (phosphate buffered
saline; without Ca^{2+} and Mg^{2+}). 2 ml Trypsin-EDTA (0.25%; Gibco, Thermo Fisher Scientific,
Massachusetts, USA) was added and the flask was incubated at 37°C for 2-3 minutes to detach the
cells. 5 ml culture media was added to the flask to inhibit trypsin. The media containing the
detached cells was collected in a falcon tube and centrifuged at 1000 rpm for 3 minutes. The
supernatant was discarded and the cell pellet was resuspended in 5 ml PBS. Cell counting was
done using a cell counter (Bio-Rad, Singapore). The required number of cells was aliquoted into
falcon tubes and centrifuged at 1000 rpm for 3 minutes. The supernatant was discarded and the
cells were resuspended in R buffer (Neon Transfection Kit, Thermo Fisher Scientific,
Massachusetts, USA). Suitable amounts of Lifeact-EGFP and EGFR-mApple (or PMT-mApple,
or PMT-mApple$_2$) (100 ng for Lifeact-EGFP; 1 µg for EGFR-mApple) plasmids were mixed with
the cells for co-transfection. The cells were electroporated using Neon Transfection system
(Thermo Fisher Scientific, Massachusetts, USA) according to the manufacturer's protocol
(electroporation settings: pulse voltage = 1000 V, pulse width = 30 ms, and pulse no. = 2). After
transfection, the cells were seeded onto culture dishes (MatTek, Massachusetts, USA) containing
DMEM (supplemented with FBS). The cells were incubated at 37°C and 5% CO_2 for 36-48 hours
before measurements.
288
Before EGFR measurements, the cells were washed with HBSS (Hank's Balanced Salt Solution,
with Ca^{2+} and Mg^{2+}; #14025134; Gibco, Thermo Fisher Scientific, Massachusetts, USA) and
starved in DMEM not containing phenol red (#21063029; Gibco, Thermo Fisher Scientific,
Massachusetts, USA) for at least 4 hours. To avoid internalization of EGFR, internalization
inhibitors were added to the cells 30 minutes before the measurements. The internalization
inhibitors used were 2 mM NaF, 10 mM NaN_3 and 5 mM 2-deoxy-D-glucose (Sigma-Aldrich,
Singapore).

296
297 **Instrumentation**
298
The TIRF microscopy set-up included an inverted epi-fluorescence microscope (IX83, Olympus,
Singapore), a motorized TIRF illumination combiner (IX3-MITICO, Olympus, Singapore), and a
dual-emission image splitter (OptoSplit II; Cairn Research, Faversham, UK). We used either an
electron multiplying charge-coupled device (EMCCD; iXon^{EM}+ 860, 128×128 pixels, Andor,
Oxford Instruments, UK) camera or a scientific complementary metal oxide semiconductor
(sCMOS; Sona 4.2B-11, Andor, Oxford Instruments, UK) for detection. 488 nm (LAS/488/100,
Olympus, Singapore) and 561 nm (LAS/561/100, Olympus, Singapore) lasers were connected to
the TIRF illumination combiner. We used a 100X, NA 1.49 oil-immersion objective (Apo N,
Olympus) and a magnification changer slider (IX3-MITICO, Olympus, Singapore) to increase
magnification two-fold to 200X where required. For the cell measurements, 37°C temperature and
5% CO_2 atmosphere were maintained using an on-stage incubator (Chamlide TC, Live Cell
Instrument). The laser power used was 100 µW for the 488 nm laser and 900 µW (EMCCD) for
the 561 nm laser (as measured at the back aperture of the objective).
312
For the dual-channel measurements, the fluorescence light was passed through a dichroic (Di01-
R488/561; Semrock) and split by the image splitter on two halves of the camera chip. The image
splitter was fitted with an emission dichroic (FF560-FDi01; Semrock) and band-pass filters
(510AF23 and 685ALP, respectively; Omega Optical). A bright-field image of a stage micrometer
was used to align the image splitter. This was done in µManager (https://micro-manager.org). The
image was aligned in both the channels following the manual instructions. To check how good the
alignment was, a self-written program in µManager was used to find the similarity in both
channels. We considered the channels sufficiently aligned if the similarity was ≥ 95%.In the case
of EMCCD, the measurements were done by recording a stack of 50,000 frames of 128×128 pixels
at 500 frames per second (fps) (for cell measurements)/1000 fps (for bilayer measurements). Andor
Solis was used for image acquisition. The kinetic mode of image acquisition was used and the
'baseline clamp' was always used to minimize the baseline fluctuation. The camera was operated
using 10 MHz pixel readout speed. Maximum analog-to-digital gain was set to 4.7 and 0.45 µs
vertical shift speed were used. The EM gain used was 300.
327
328 **Table 1: Acquisition parameters for the various experimental configurations**
329

Camera	Analysis	Laser wavelength [nm]	Time per frame [ms]*	No. of frames	Image pixel size [nm]	Map dimensions [Pixels]
EMCCD	FCS	561	2	50000	240	120×56
Fig.1	Diffusion law	561	2	50000	240	114×56
	N&B	561	20	5000	240	120×56
	SRRF	488	200	500	48	600×280

330 *: acquisition time was 2 ms; longer times are reached by time binning of frames.
331
DATA ANALYSES

The data analyses was performed on a computer with the following configuration – Windows 10
Home 64-bit operating system, Intel® Core™ i7-7800X CPU @ 3.50 GHz processor, 32 GB
RAM, NVIDIA Titan Xp GPU with 3840 CUDA cores and 12.3 GB memory.

FCS

The image stacks from the cell measurements were loaded in Imaging FCS[27] 1.52 plugin for Fiji
and the ACFs calculated. The values used for the parameters were: frame time = 0.002 s, correlator
(p, q) = (16, 12), pixel size = 24 µm, NA = 1.49, λ1 = 565 nm (red channel). The EMCCD data
analysis was performed at 1×1 binning for 100X magnification. Bleach correction[26] was
performed with a polynomial of order 8. ACFs were fitted with a one-component diffusion model.

$$G(t) = \frac{1}{N} \frac{g(t)}{g(0)} \qquad \text{Eq. 1}$$

$$g(\tau) = \left(\frac{\sqrt{4D\tau + \omega_{xy}^2}}{a\sqrt{\pi}} \left(e^{-\frac{a^2}{4D\tau + \omega_{xy}^2}} - 1 \right) + \text{erf}\left(\frac{a}{\sqrt{4D\tau + \omega_{xy}^2}} \right) \right)^2 + G_\infty \qquad \text{Eq. 2}$$

342
where a is the pixel size, τ is the lag time, N is the number of particles, D is the diffusion coefficient,
ω_{xy} is the PSF (xy) ($1/e^2$ radius) direction. To exclude outliers or non-converged fits, only data
with $0.01 < D < 10$ µm^2/s were considered for further analysis.

DIFFUSION LAW

The image stack was analyzed using the Imaging FCS 1.52 plugin for Fiji. The values used for the
parameters were: frame time = 0.002 s, correlator (p, q) = (16, 12), pixel size = 24 µm (EMCCD),
magnification = 100, NA = 1.49, λ_1 = 565 nm (red channel), PSF (xy) = 0.96 (EMCCD red channel)
and bleach correction = polynomial of order 8. In the "Diff. Law" tab, the diffusion law plot was
generated and fit with a straight line for square binning 1 to 5 (EMCCD).

N&B

The 2 ms, 50000 frames image stack was temporally binned (sum binning of 10 frames each) to a
20 ms, 5000 frames stack in Fiji. The images were analyzed using the Imaging FCS 1.52 plugin.
The values used for the parameters were: frame time = 0.02 s, binning = 1×1 (EMCCD), and bleach
correction = polynomial of order 8. An intensity filter with a suitable range was set, as defined by
the background intensity in an image, to exclude the background pixels and include only the pixels
containing the cell. A dark image (image taken with the camera shutter closed) with the same
spatiotemporal dimensions as the measurement image was loaded for background correction in the
"Bgr NUM" tab. Then in the "N&B" tab, "G1" was selected in the "NB mode" and the "N&B"
button in "N&B analysis" was pressed to generate the N&B maps. The N&B equations are defined
as[13]:

$$N = \frac{(\langle I \rangle - \text{offset})^2}{\sigma^2 - \sigma_0{}^2} \qquad \text{Eq. 3}$$

$$B = \frac{\sigma^2 - \sigma_0{}^2}{\langle I \rangle - \text{offset}} \qquad \qquad Eq.\ 4$$

363
364 where, N is the apparent number of particles in the observation volume, $\langle I \rangle$ is the average intensity,
365 *offset* is the intensity offset of EMCCD, σ^2 is the variance of the signal, B is the apparent brightness
366 of a particle, and $\sigma_0{}^2$ is the variance of the readout noise in the EMCCD. The offset and $\sigma_0{}^2$ can be
367 obtained from a dark image captured by the EMCCD. The variance is typically affected by shot
368 noise, and hence we used the covariance. This is referred to as G1 analysis.
369

$$\text{Covariance}(F(t), F(t + \Delta t)) = \langle F(t)F(t + \Delta t) \rangle \qquad \qquad Eq.\ 5$$

370 where, $F(t)$ is the fluorescence signal at time t and $F(t+\Delta t)$ is the fluorescence signal at time $t+\Delta t$.
371
372 The dimer fraction is determined using the equation below:
373

$$\text{Dimer fraction} = \frac{r - 1}{2p - r + 1} \qquad \qquad Eq.\ 6$$

374
where r is the ratio of the brightness of EGFR to the brightness of monomer, p is the proportion of
the molecules which are fluorescent. The procedure to determine p based on the brightness of
monomer and dimer is provided in supplement. The error propagation to estimate the error
associated with the dimer fraction is also provided in supplement.

SRRF

The 2 ms, 50000 frames image stack was temporally binned (average binning of 100 frames each)
to a 200 ms, 500 frames stack (for Fig. S3 other temporal binnings were also used) in Fiji using
the Image → Transform → Bin command sequence. The NanoJ-SRRF[14] plugin was used. The
"SRRF analysis" option in the plugin was chosen. The default settings were used, with an addition
– the "Do Gradient Smoothing" ("Show Advanced Settings" → "Radiality" → "Do Gradient
Smoothing") was also activated. To determine the fibre thickness, the straight line tool in Fiji was
used to draw a line segment across a fibre in the generated SRRF maps. The "Plot Profile" function
was used to generate an intensity histogram, which was fitted using the "Curve Fitting" tool in
Fiji. The curve fitting function used was a Gaussian given by the equation below.

$$y = a + (b - a)e^{\frac{-(x-c)^2}{2d^2}} \qquad \qquad Eq.\ 7$$

Where a is the offset, b-a is the height at the centre c, and d is the standard deviation of the
Gaussian. The SRRF maps had black borders so the image was cropped to get rid of them. As a
result, the original acquired image was cropped before performing FCS, diffusion law and N&B
analyses to maintain the same cell area in all the maps (for the image dimensions, refer table 1).

Statistical Analysis

The frequency plots in Fig. 1 were generated using the "*Bivariate histogram*" command in
IgorPro© (Wavemetrics Inc, Oregon, USA). A linear fit was performed to estimate the correlation
coefficient (*R*). Graphpad Quickcalcs was used to perform t-tests. Only those pixels were included
in the scatter plots that had valid values for both parameters.

Code availability

The   Imaging   FCS   1.52.   ImageJ   plugin   is   available   at
http://www.dbs.nus.edu.sg/lab/BFL/imfcs_image_j_plugin.html or alternatively is included in the
ImageJ update site.
402
403 **Data availability**
404
The datasets generated during the current study are available from the corresponding author on
reasonable request.

Competing Interests

The authors declare that there are no competing interests.

References

26   Macháň, R., Foo, Y. H. & Wohland, T. On the Equivalence of FCS and FRAP:
Simultaneous Lipid Membrane Measurements. *Biophysical journal* **111**, 152-161,
doi:10.1016/j.bpj.2016.06.001 (2016).
27   Sankaran, J., Shi, X., Ho, L. Y., Stelzer, E. H. & Wohland, T. ImFCS: a software for
imaging FCS data analysis and visualization. *Optics express* **18**, 25468-25481,
doi:10.1364/oe.18.025468 (2010).
416