Stimulus transformation into motor action: Dynamic graph analysis reveals aging effects on brain network communication

Nils Rosjat[1,2*], Bin A. Wang[1,a], Liqing Liu[1,2,b], Gereon R. Fink[1,3], Silvia Daun[1,2]

1 Cognitive Neuroscience, Institute of Neuroscience and Medicine (INM-3), Forschungszentrum Jülich, Jülich, Germany

2 Institute of Zoology, University of Cologne, Cologne, Germany

3 Department of Neurology, Faculty of Medicine and University Hospital Cologne, University of Cologne, Cologne, Germany

¤a Current address: Department of Neurology, BG University Hospital Bergmannsheil, Ruhr-University Bochum, Bochum, Germany.

¤b Current address: Faculty of Psychology, Key Research Base of Humanities and Social Sciences of Ministry of Education, Tianjin Normal University, Tianjin, China

* n.rosjat@fz-juelich.de

Abstract

The vast majority of our actions result from a complex interplay of various brain regions. To date our knowledge of aging-associated functional changes in the brain networks that are involved in the transformation of visual information into motor action remains sketchy. To investigate this topic, we employed dynamical graphs based on phase-locking of Electroencephalography signals recorded from healthy younger (YS) and older subjects (OS) while performing a simple visually-guided finger-tapping task. The network analysis yielded specific age-related dynamic network structures in the low frequencies (2-7 Hz), which seem to be a necessary prerequisite for stimulus processing, movement initiation and execution. The networks in OS, however, contained several additional, particularly interhemispheric, connections and showed an overall increased coupling density. Cluster analyses revealed reduced variability of the subnetworks in OS, particularly during movement preparation. In YS, occipital, parietal, sensorimotor and central regions were - temporally arranged in this order - heavily involved in hub nodes. Whereas in OS, a hub in frontal regions preceded the noticeably delayed occurrence of sensorimotor hubs, indicating different neural information processing in OS.

The increased interhemispheric connectivity and additional inclusion of frontal regions further extend previous fMRI data, which report an overactivation, especially in the prefrontal and pre-motor areas, associated with a loss of hemispheric lateralization in OS. All observed changes in network organization are compatible with a compensatory mechanism to maintain motor function in OS.

Keywords: Compensation, HAROLD, low frequencies, PASA, Phase locking, information processing

Introduction

Aging leads to a slowing down of cognitive abilities. This includes cognitive processes that are essential for motor performance. For example, increased deficits in the execution and planning of movements, as well as a slow down in response to external stimuli [1–3]. A study by Niessen et al. [4] recently showed that older subjects have deficits primarily in the accuracy of processing visual stimuli into motor actions and less in the pure execution of the motor act itself. Even though several studies investigated the effects of non-pathological aging on human motor performance, to date, our understanding of the neurobiological underpinnings of these changes, remains sketchy.

Several fMRI studies reported an association between aging and changes in movement-related neural activity. In particular, a generally enhanced activation was reported, especially in pre-frontal and pre-motor areas [5–7]. This effect is often referred to as the so-called PASA (posterior to anterior shift in aging) phenomenon [8]. The HAROLD (hemispheric asymmetry reduction in older subjects) model proposes another effect often reported by fMRI studies, namely a reduced hemispheric asymmetry during cognitive processing, which is thought to be rather a general aging-associated than task-specific phenomenon (for a summary see [9]). It is assumed that these effects (PASA and HAROLD) reflect either a compensatory mechanism, i.e., a recruitment of additional brain areas to support weakened functionality in core brain regions [10–12], or difficulties in recruiting specialized task-specific subnetworks in older subjects [9, 13, 14].

Up to now, the neural mechanisms underlying these changes in healthy aging remain elusive. One hypothesis states that they result from changes in inter-regional neural synchronization since this is a crucial mechanism underlying motor and cognitive tasks [15, 16]. Synchronization of distinct neural populations represents the organization of their temporally coordinated activities (for a review, see [17]). Several studies using data from EEG/MEG/single-unit/ECog recordings while subjects performed a variety of different motor tasks showed that remote neural populations synchronize over a short-time period [18–21], suggesting that coordinated timing constitutes a fundamental principle involved in motor and cognitive processing [20, 22–25].

In our previous study [26], we could show that during *voluntary*, i.e. internally-triggered movements, motor-

related networks built on inter-regional neural synchronization in younger as well as older subjects. In that study, we performed a phase-locking analysis between the neural signals recorded from the electrodes lying above the motor cortex with a low time resolution of 100 ms. In particular, we calculated the phase-locking value (PLV) between the electrodes of interest as it is a general characteristic for the quantification of synchronization between neural signals [27]. The results of this previous study exhibited an increase in interhemispheric connectivity in older subjects compatible with the HAROLD model. However, due to the restriction on electrodes above the motor cortex, we were unable to find evidence for the PASA phenomenon.

In the current study we were interested in age-related changes in the processing of neural information during externally-guided movements through visual information. In particular, we were interested in the transformation of visual information towards motor execution in such visually-guided movements. For the investigation of this aspect it is essential to obtain information about the *timecourse* of the network structure. As the results mentioned above originate from fMRI studies or were analyzed without or only minor time resolution, little information is available about aging-effects on the contribution of the different stages of stimulus transformation into motor action, i.e., stimulus processing, movement planning, and execution. Accordingly, we here aimed at investigating how changes related to healthy aging affect the organization of brain networks with a particular interest in the role of occipital, parietal and prefrontal and premotor regions besides the classical motor-related ones. Notably, we focused on connectivity patterns and their temporal evolution during different movement phases, as indexed by the phase relationship of distinct neural signals. For this reason, we used the high temporal resolution of the rPLV and applied it to Electroencephalography (EEG) data. The data were recorded from younger and older healthy participants who performed a simple visually-guided finger-tapping task and dynamic networks based on these phase relationships were created. A dynamic network (also dynamic graph) is a composition of temporally successive connectivity networks (or graphs) [28, 29]. In contrast to static networks, dynamic networks take into account the temporal development of the connectivity. They, therefore, serve to provide a more precise temporal account of the different coupling structures of the respective age groups, especially about the influence of aging on the routing of neural information during the different movement phases.

Materials and Methods

Participants In this study, we included EEG data of a group of twenty-one younger healthy individuals (10F/11M, age: 22-35 years) and of a group of thirty-one older healthy subjects (15F/16M, age: 60-78 years) first presented in two earlier studies [3, 16]. All participants were right-handed according to the Edinburgh Handedness Inventory [30] and had normal or corrected to normal color vision. The older participants had no history of psychiatric or neurological disease (as assessed by the Trail making test (TMT A: 41.16 ± 16.27 s,

TMT B: $78.59 \pm 28.02\,$s) [31], the Mini-mental-state-test (28.97 ± 1.29) [32], the Clock-drawing test [33], and the Beck depression inventory (4.58 ± 3.21) [34]. All participants had given their written informed consent prior to the study. The local ethics committee of the Faculty of Medicine at the University of Cologne had approved both studies.

Experimental Design   We recorded EEG data while both groups performed a simple finger-tapping task. The experimental paradigm reported here consisted of two conditions, a visually-cued condition with intention to act, i.e. a visually-cued motor condition (Fig. 1 (A)) and a visually-cued condition without intention to act, i.e. a vision-only condition (Fig. 1 (B)). In the motor condition (visually-cued tapping, Fig. 1 (A)) the subjects were presented with a right- or left-pointing arrow (2° wide and 1.2° high, expressed as visual angles) for 200 ms on a screen with inter-stimulus intervals of varying length $\geq 4\,$s. The participants had to press a button with their left or right index finger, corresponding to the direction of the arrow, as fast as possible. In the second condition (vision-only, Fig. 1 (B)) the same stimuli as in the visually-cued tapping condition were presented. However, this time the participants were instructed to pay attention to the arrows but to refrain from performing or imagining to perform the button press. Thus, no motor action was performed in this condition (for full details see [3, 16]). During the experiment, a third, self-initiated tapping condition (not reported here) was recorded. In this condition, the participants were free to choose when to move which hand without any external stimuli (for details see [35]). The overall duration of the experiment including breaks was 70 min.

Our analyses were restricted to data acquired from the visually-cued and vision-only condition (Fig. 1 (A) and (B)) and the conditions were analyzed locked to the onset of the visual stimulus.

EEG recording and preprocessing   While the subjects performed the task, we continuously collected EEG data from 64 active Ag/AgCl electrodes (Brain Products GmbH, Munich, Germany), placed according to the international 10-20 system. The reference electrode was placed at the left earlobe. Bipolar horizontal and left vertical electrooculograms (EOG) were recorded with three of the 64 scalp electrodes (previously located at FT9, FT10, and TP10 in the 10-20 system). These were placed at the bilateral outer canthi and under the left eye to monitor eye movements and blinking. Before the experiment, it was ensured that the impedances of all electrodes were below 15 kΩ. The EEG signals were amplified, bandpass filtered in the frequency range $0.87 - 500\,$Hz, and digitized at a sampling rate of $2.5\,$kHz. Index finger movement onsets were detected by acceleration sensors attached to the tip of each index finger. We also used the information from the acceleration signals to monitor the subjects' behavior, e.g., to rule out errors such as mirror movements. We defined the onset of the finger movement as the instant of time at which the numerical time derivative of the acceleration signal exceeded a predefined threshold.

The acquired raw data went through several offline pre-processing steps. First, we band-pass filtered the

Fig 1. Experimental paradigm. Organizational setup of the experiment. (A) Visually-cued finger movements and (B) vision-only condition. Each condition is represented by a unique reminder (e.g. blue square in the visually-cued tapping condition). The arrows indicate the hand (left or right) to be used (adapted from Fig. 1 in [35]).

data from 0.5 to 48 Hz to remove slow voltage drifts. We next downsampled the data to 200 Hz to reduce the file size and thus the computing time. We then removed artifacts using a semi-automated process in EEGLAB using independent component analysis (ICA) [36]. Finally, the data were epoched to intervals $[-1500, +2500]$ ms, which centered around stimulus onset.

For our analysis, we excluded trials that contained EEG artifacts or included movements during the pre-stimulus period detected by the accelerometer. After excluding trials based on these criteria, we only considered data from participants who had at least 30 correct trials per hand. Based on these quality considerations, data from three younger and seven older subjects were excluded from further analysis. Thus, EEG data from 18 younger and 24 older subjects were used for further processing.

Spatial filtering A crucial pre-processing step for the following connectivity analysis was spatial filtering. In the EEG, volume conduction is a primary concern in determining the relationship of signals. The activity of a single neural generator can influence the signal in several different electrode positions and thus lead to inaccurate phase-locking between these recording sites. To reduce the detection of false-positive connectivity, we applied spatial filtering using surface Laplacians, which demonstrably improves spatial resolution and allows the analysis of electrodes close to the region of interest [27]. Thus, our preprocessed data were

re-referenced to a small Laplacian reference to improve the spatial resolution of the signals and thus their suitability for subsequent connectivity analysis [37]. Electrodes on the outer edge, which do not have four neighbours, were excluded from further analyses, so that a total of 42 electrodes were included in the investigation.

Phase-locking analysis Following preprocessing, we transformed the epoched and cleaned data to the time-frequency domain using Morlet wavelets [38] in a $\delta - \beta$ frequency range (2-30 Hz) with a step size of 1 Hz (5 cycles). This time-frequency decomposition was performed with the Statistical Parametric Mapping toolbox (SPM12, Wellcome Trust Centre for Neuroimaging, London UK) implemented in MatLab R2018b (The MathWorks Inc., Massachusetts, USA). We then analyzed the amplitude and phase information from the time-frequency decomposition using customized MATLAB scripts.

We obtained the temporal evolution of the amplitude $A(f,t)$ and phase $\varphi(f,t)$ by applying the complex Morlet wavelet transformation for each frequency separately to our data. We quantified the communication of two different brain regions, i.e., acquisition sites, by synchronization determined by the single-frequency phase-locking value (sPLV; adapting the phase-locking value defined in [27]). For a pair of channels m and n, sPLV is defined as:

$$sPLV_{m,n}(f,t) = \frac{1}{N}\left|\sum_{k=1}^{N} e^{i(\varphi_{m_k}(f,t)-\varphi_{n_k}(f,t))}\right| \tag{1}$$

here φ_{m_k} denotes the phase of the EEG at channel m in the k-th trial. N is the total number of trials, and i is the imaginary unit. While minimum sPLV $= 0$ represents a random distribution of phase differences over all trials without any coherence, a maximal sPLV $= 1$ occurs only in the case of perfect inter-trial phase locking of the phase differences between the EEG signals at the two channels m and n over all trials.

Since we were interested in an event-related measure that represented the synchronization increases and decreases during stimulus processing, movement preparation and execution, we normalized the sPLV of the electrophysiological recordings at each pair of channels with respect to its baseline value and calculated its relative change over the entire epoch. We refer to these normalized sPLVs as *relative phase-locking values* (rPLVs):

$$rPLV_{m,n}(f,t) = \frac{sPLV_{m,n}(f,t) - \overline{sPLV_{m,n}(f)}}{\overline{sPLV_{m,n}(f)}} \tag{2}$$

here $\overline{sPLV_{m,n}(f)}$ denotes the sPLV of the baseline interval for each frequency, i.e. the mean sPLV at frequency f in the baseline interval, which was defined from $[-1300, -500]$ ms excluding the first 200 ms of the original interval as it contained edge artifacts of the Morlet transformation.

To analyze inter-regional synchronization, we considered four major frequency bands: δ (2-3 Hz), θ (4-7 Hz), α (8-12 Hz) and β (13-30 Hz). Instead of first filtering the signal to a specific frequency band, we calculated

Fig 2. Phase-locking analysis. Schematic illustration of the procedure to construct corresponding phase-locking networks from contrasted, i.e., visually-cued - vision-only, rPLVs (adapted from Fig. 1 in [26]). The black vertical lines represent movement onset.

the rPLV separately for each frequency with a resolution of 1 Hz before averaging these values over the frequencies of the frequency band. In this way, we ensured a true 1:1 frequency coupling that maximizes the contribution of each frequency in the band. The steps described above were performed for both the tapping and the vision-only control condition. The resulting mean rPLVs (averaged over a frequency band of interest) were then contrasted by subtracting the vision-only control condition from the visually-cued condition, resulting in positive values when phase-locking is stronger in visually-cued and negative values for stronger phase-locking in the control condition. (cf. Fig. 2). The contrasted mean rPLVs then underwent statistical testing (described below) to define the edges of the phase-locking network. The resulting networks were analyzed using graph theoretical metrics as implemented in the Brain Connectivity Toolbox [39].

By implementing the contrast condition we reduced the pure sensory effect of the visual stimulation, which would otherwise have to led to an all-to-all connectivity in the networks.

Statistical analysis We applied pointwise t-tests with a significance level of $p < 0.05$, corrected for multiple comparisons (false discovery rate, FDR, $q = 0.05$) [40], to test whether the mean relative amplitude changes (averaged over frequency band scaled logarithmically relative to pre-stimulus baseline) between YS and OS differed significantly in the intervals $[0, 1000]$ ms ($\delta - \theta$) and $[0, 1500]$ ms (α). To test whether the difference between the mean rPLV (averaged over the frequency band) and the baseline (averaged over the frequency band) was statistically significant, we compared the mean rPLV obtained for each pair of electrodes at each time point in the interval $[0, 1000]$ ms with its baseline value. To this end, we used a pointwise t-test with a significance level of $p < 0.05$, corrected for multiple comparisons (false discovery rate, FDR, $q = 0.05$) [40]. The corrections were performed concerning the number of time points, age groups, and electrodes. The

baseline was constructed by generating normally-distributed random values that had the same mean and standard deviation as the EEG at any time point of the baseline interval $[-1300, -500]$ ms.

For the construction of the rPLV networks, we defined a *connection*, i.e., an edge of the network, between two given nodes to exist at a given time point t if the mean rPLV between these two nodes was significantly increased with respect to its baseline value. Based on our previous results [16, 26], which showed a symmetrical behavior of phase-locking, and to reduce the number of statistical comparisons, the rPLVs during left- and right-hand finger movements were collapsed together after flipping the ones for the right-hand movements. Electrodes were defined as being either ipsilateral or contralateral to the moved hand. For convenience, results are presented resembling left-hand movements, i.e., electrodes over the left hemisphere are assumed to be ipsilateral while electrodes over the right hemisphere are assumed to be contralateral to the performed movement.

Dynamic network analysis The statistical analysis above defines a binary undirected graph $G_t = (V, E_t)$, where the graph G is defined by a set of vertices V and edges $E_t : V \times V \to \mathrm{R}$, for each point in time t for younger and older participants. The dynamic graph, also called temporal networks, is defined as the ordered set of graphs $G(t) = \{G_t | t \in [1, \ldots, T]\}$. It can also more efficiently be represented by a list of triples (i, j, t), defining the contacts of two nodes i and j and the time point t at which they occur [28]. In contrast to static graphs, dynamic graphs account for time-varying connectivity patterns that might reflect the formation of subnetworks due to task performance.

Besides the time-varying networks, we investigated the *aggregated graph* of the dynamic network, which is a graph that accounts for the number of occurrences of each edge over all graphs, i.e., edges that appear more often have a higher weight while rarely present edges have a smaller weight. We performed a cluster analysis using the Louvain community algorithm, which uses a greedy optimization method to minimize the ratio of the number of edges inside communities to edges outside communities, for each time step and the aggregated graph for both groups of participants to retrieve information on closely connected subnetworks [41–44]. To reduce the effect of randomly assigned cluster labels between time points, we always used the previous cluster as the initial condition for the following community detection. Additionally, we post-hoc assigned a label for each community, minimizing the number of label switches between time points, which prevents the same cluster from being assigned different cluster labels at consecutive time points. Furthermore, based on these clusters we tested the similarity of clusters between age groups by computing the variation of information (VI) [45]

$$VI(X, Y) = \frac{H(X) + H(Y) - 2MI(X, Y)}{\log(n)} \tag{3}$$

where H is entropy, MI is mutual information, and n is the number of nodes. This metric ranges from 0, i.e., identical clustering, to 1 (as it is normalized by $\log(n)$), i.e. maximally distinct clustering. To account for

actual age effects, we compared $VI(X, Y)$ (X representing YS and Y representing OS) with time-lagged ₂₁₉
self-distribution distances $VI(X(t), X(t-1))$ and $VI(Y(t), Y(t-1))$ to test whether the partition distance ₂₂₀
between younger and older subjects is larger than within group partition changes in time. ₂₂₁

The flexibility F of the dynamic network is defined by the average over all *node flexibilities* f_i [46], which are ₂₂₂
defined as the number of times that a node changed communities, normalized by the maximal number of ₂₂₃
times this node could have changed communities, ₂₂₄

$$f_i = \frac{m}{T-1} \tag{4}$$

with m the number of community changes and T the number of time points. ₂₂₅

Finally, we analyzed the hub nodes of the networks. Those nodes play a crucial role in the network, as they ₂₂₆
serve as the connection between different subnetworks. For the analysis of hub nodes, we calculated the ₂₂₇
betweenness centrality for each time point, which is defined as the fraction of shortest paths that pass through ₂₂₈
each node [47, 48]. We primarily focused on the first occurrence of each hub node in the network as well as on ₂₂₉
their timecourses. This gives an indication of how neural information is transmitted from stimulus processing ₂₃₀
to motor action. ₂₃₁

Results ₂₃₂

Behavioral results   We first tested whether there was a significant difference between older subjects (OS) ₂₃₃
and younger subjects (YS) in reaction time (RT), defined as the time that elapsed from stimulus presentation ₂₃₄
until movement onset, and accuracy rate, i.e., the percentage of correct responses among all trials. The ₂₃₅
reaction time was significantly longer in older subjects (497 ms $\pm$ 84 ms) than in younger subjects (426 ms $\pm$ ₂₃₆
77 ms), $t(82) = 3.9442, p < 0.0001$. The mean response accuracy in the visually-cued condition for older ₂₃₇
subjects (93.13% $\pm$ 5.69%) and for younger subjects (97.7% $\pm$ 2.22%) was greater than 90%, consistent ₂₃₈
with the instructions of the task. However, as expected, the accuracy was lower for the older participants, ₂₃₉
$t(40) = -3.2329, p = 0.0025$. We further tested the movement times, i.e. the time from movement onset ₂₄₀
(determined using the accelerometer) to the actual button press of younger (M =81.9 ms, Std = 8.6) vs. ₂₄₁
older (M = 88.3 ms, Std = 8.3) subjects for significant differences and found that the finger movements of ₂₄₂
older subjects were significantly (however on average only 6 ms) slower than those of the younger ones ₂₄₃
(t(82)=−3.4307, p=0.0004). ₂₄₄

In summary, and consistent with the current literature, we found an age-related deceleration in response to ₂₄₅
the external stimuli and a reduced accuracy in task performance. However, the performance of the motor act ₂₄₆
itself was only slightly impaired. ₂₄₇

Event-related changes across frequencies

Amplitude (de-)synchronization (ERD/ERS)   We next investigated the changes in amplitudes in response
to the visual stimulus to check whether the data show known basic motor-related effects (ERD/ERS) and to
clarify whether phase-locking effects could possibly be due to changes in amplitude in the same frequency
range. We performed separate analyses for the $\delta - \theta$, α and β frequency ranges for the visually-cued motor
task and for the contrast between the visually-cued motor task and the vision-only task (i.e. visual stimulation
without intention to move). During processing of visual stimulation and execution of movement, both age
groups show a relative increase in amplitude in $\delta - \theta$ frequencies (2-7 Hz), both in the visually-cued motor and
in the contrast condition. Consequently, there are no significant differences in relative amplitude between the
two age groups (Fig. 3 A, third column). In the α frequency range (8-12 Hz), event-related desynchronization
(ERD) can be observed in both groups in both the visually-cued and the contrast condition during motor
execution, i.e. around 500 ms after stimulus. In the visually-cued condition, OS show a significantly increased
ERD (i.e. decreased amplitude) in all examined electrodes compared to YS, $2.1202 <$ t's < 5.3578, p's < 0.05
(FDR corrected), df $= 40, 0.4202 <$ sd's < 3.2969 (Fig. 3 B top row). Note, that the ERD in OS also lasts
longer than in YS, reaching well into the post-movement phase in the central, motor and parietal electrodes
(up to 1300 ms). In the contrast condition, however, the differences in ERD are much less pronounced (all p's
> 0.05, apart from few p's < 0.05) (Fig. 3 B bottom row). In β frequencies (13-30 Hz), parts of the ERD are
also visible in most electrodes during movement. Additionally, especially in YS, we observe an event-related
synchronization (ERS) after the execution of the movement (approximately 1000 ms after stimulus). The
ERS is significantly stronger in YS than in OS in both, the visually-cued ($-3.4455 <$ t's < 6.4240, p's < 0.05
(FDR corrected), df $= 40, 0.2788 <$ sd's < 1.4721) and the contrast condition, $-3.8928 <$ t's < 5.2834, p's
< 0.05 (FDR corrected), df $= 40, 0.4394 <$ sd's < 1.4033 (Fig. 3 C, bottom row). In summary, we observe
age-related differences in ERD and ERS in the α and β frequencies and no age-related differences in the
$\delta - \theta$ frequencies consistent with the current literature.

Phase synchronization   We calculated the rPLVs separately for the $\delta - \theta$, α and β frequency ranges for the
visually-cued motor condition and for the contrast between the visually-cued motor and the control condition
(i.e. visual stimulation without intention to move). We found an increased phase-locking in the $\delta - \theta$ frequency
band (2-7 Hz) during stimulus processing and movement preparation and execution in both age groups and
both conditions (Fig. 4). In the visually-cued condition, a significant increase in rPLV was present throughout
all electrode pairs (Fig. 4 top row) in both age groups (Ys: $-12.3739 <$ ts < 5.6742, p < 0.05 (FDR corrected),
df $= 17, 0.0709 <$ sd < 2.1193; OS: $-11.0582 <$ ts < 4.6396, p < 0.05 (FDR corrected), df $= 23, 0.0942 <$
sd < 1.7203), whereas contrasting highlighted more specific connections (Fig. 4 bottom row) (Ys: $-8.8947 <$
ts < 5.8060, p < 0.05 (FDR corrected), df $= 17, 0.0987 <$ sd < 1.3434; OS: $-8.2287 <$ ts < 5.8060, p < 0.05

Fig 3. Event-related amplitude changes. Amplitude differences on a logarithmic scale in $\delta - \theta$ (A), α (B) and β frequencies (C) in visually-cued (top rows) and contrast conditions (lower rows) for YS (left column), OS (middle column) and differences YS-OS (right column) for all 42 considered electrodes. Non significant amplitude differences between YS and OS are greyed out in the right column. $t = 0$ represents the onset of the visual stimulus.

Significant delta-theta rPLV increase

Fig 4. Event-related rPLV changes Significant increase in phase-locking in the $\delta - \theta$ frequency band in the visually-cued condition (top) and in the contrast condition (bottom) for YS (left) and OS (right) for 819 electrode pairs. rPLV values are color coded from 0 (no change) to 0.5 (50% increase). Non significant changes are greyed out. $t = 0$ represents the onset of the visual stimulus.

(FDR corrected), df $= 23$, $0.1148 <$ sd < 1.1383). In addition, there were increased rPLVs in the β frequency band (13-30 Hz) in the post-movement period (approx. 500 ms after stimulus presentation). However, this phase-locking effect in the β frequency band could only be observed in few electrode pairs and therefore did not survive the FDR correction. No significantly increased rPLVs were observed in the α frequencies. We hence focused our further dynamical graph analysis on the $\delta - \theta$ frequencies and on the contrast condition in which the pure sensory effect of the visual stimulation was reduced.

Dynamic motor network analysis

Fig 5. Dynamic graph snapshots. Exemplary graph snapshots for YS (top) and OS (bottom) for five different time points representative for stimulus processing, movement preparation and execution and post movement. Edges related to the core motor network are color coded in blue (ipsilateral), green (contralateral) and orange (interhemispheric). Color codes are chosen as in [26] for better comparison.

Time-varying network analysis Phase locking in the $\delta - \theta$ frequency band was used to construct connectivity matrices for each time point of the trial, which resulted in a dynamic graph (see Materials and methods - *Dynamic network analysis*). Representative networks are shown in Fig. 5. The figure displays network connectivity which can roughly be assigned to stimulus processing (150 ms), movement preparation (250 ms), movement execution (350 & 450 ms) and the post movement phase (550 ms). The younger subjects displayed the densest networks in the stimulus processing and movement preparation phase, the older subjects additionally in the early phase of movement execution. The connections in YS networks clustered around occipital regions and central motor-related regions, while OS networks showed a dense connectivity spread over the whole brain. Both group of subjects associated graphs expressed pronounced connectivity between ipsilateral (blue) and contralateral (green) sensorimotor nodes in the movement execution and post movement phase. The older subjects' networks, however, additionally included an increased number of interhemispheric connections (orange), which were already present in the movement preparation phase. Overall, older subjects' networks displayed an increase in connectivity, leading to a denser graph.

We quantified this by analyzing the average node degree over time for YS and OS (Fig. 6). A paired t-test showed that the average node degree was significantly increased in OS compared to YS in whole network connectivity, i.e., taking all nodes of the network into account, t's < -2.9027, p's < 0.05 (FDR-corrected), df $= 60$, $0.3404 \leq$ sd's ≤ 4.7489. The same effect was observed for motor network connectivity, i.e., taking only nodes above the motor-related regions into account, t's < -3.3773, p's < 0.05 (FDR-corrected), df $= 19$,

Fig 6. Node degree dynamics. A: Timecourses of the average overall node degrees of YS (blue) and OS (red) surrounded by shaded regions representing the confidence interval. B: Timecourses of the average motor network node degrees of YS (blue) and OS (red) surrounded by shaded regions representing the confidence interval. Movement periods are marked with blue (YS) and red (OS) solid lines at the x-axes. Intervals with significant differences between age groups are marked with black dots.

$1.0954 \leq$ sd's ≤ 4.8284. In both cases, the significant differences were mainly present in the time interval from 200 to 400 ms.

In the following, we applied the Louvain clustering method (clustering parameter $q = 0.9$) which measures the density of edges inside communities to edges outside communities to separate the graphs into sets of closely interconnected subnetworks (Fig. 7 A, B). During stimulus processing, YS networks were separated in a cluster with a focus on occipital and parietal electrodes (red nodes) and a cluster containing motor, frontal, and central electrodes (light blue nodes). OS networks, on the other hand, were divided into three more widespread clusters with electrodes from almost all eight regions (Fig. 7 B (100 ms)). In the movement preparatory and early movement execution phases (200-300 ms), OS network clusters were reduced to two extended, stable clusters that integrated electrodes from all 8 regions (cf. Fig. 7 B (200 and 300 ms)). The YS networks, on the other hand, broke up into several smaller clusters, also showing no clear regional distinction. A paired t-test showed a significant reduction in node flexibility in OS, t(40) = 5.6589, p < 0.001, $\Delta_{\overline{f_y} - \overline{f_o}} = 0.1742 \pm 0.0622$, in the interval (200, 300) ms. Even when the interval was increased to (1-700) ms the node flexibility was significantly reduced in OS, t(40) = 2.7318, p $= 0.00925$, $\Delta_{\overline{f_y} - \overline{f_o}} = 0.0426 \pm 0.0315$. During movement execution the number of connections decreased (cf. Fig. 6 A), which led to a sparser network that for YS could be divided mainly into a contralateral motor - central - ipsilateral frontal community (red) and several smaller communities (Fig. 7 B, 400 ms). In OS, the motor, frontal, and central electrodes were more densely connected, which led to less distinct clusters as in YS.

We calculated the variance of information (VI), which serves as a marker of the difference of two clusters of the same nodes, between both groups for each time point (VI(YS,OS)). As a comparison and, in particular, to

Fig 7. Louvain clustering. A: Channel partitions for YS (left) and OS (right). Clusters are labeled from 1 to 6 (with arbitrary colors). Unconnected channels are displayed in blue (cluster 0). Movement periods are marked with black horizontal lines. B: Circular graphs showing representative clusterings from 100 ms, 200 ms, 300 ms, and 400 ms post stimulus. Nodes are colored corresponding to the clusters in A. Edges related to the core motor network are color-coded in blue (ipsilateral), green (contralateral), and orange (interhemispheric). C: Distribution distance for clusters computed for younger and older subjects (black) compared to self-distribution distances of younger (blue) and older subjects (red). Movement periods are marked with blue (YS) and red (OS) solid lines.

detect real effects of age, we calculated the VI for YS (VI(YS,YS)) and OS (VI(OS,OS)) for consecutive time points (Fig. 7 C). With this analysis, we could show that the variance between age groups stays at a high level in the first 700 ms of the epoch (with a value between 0.6-0.8). The variance between age groups was constantly higher compared to the self-distribution distances of YS and OS, except for the time interval [200, 300] ms. In this interval, the self variance of information for OS displayed a drop, which was related to the less variable networks, i.e., the decreased node flexibility during movement preparation described above. At the same time, YS showed an increased variance of information reflecting a high clustering variability during that time.

Aggregated networks In the following, we investigated the aggregated networks, i.e., the sum of all unweighted connectivity matrices over all time points t, for YS and OS. The resulting networks are presented in Fig. 8. YS and OS showed densely connected nodes above the occipital cortex, which were related

to the early time intervals of the dynamic graphs. In YS the motor network showed additionally dense ₃₃₇ connectivity between nodes above ipsilateral frontal and central regions (blue), diagonal interhemispheric ₃₃₈ connections between nodes above the ipsilateral frontal and contralateral motor cortex (orange), horizontal ₃₃₉ interhemispheric connections between nodes above the ipsilateral and contralateral motor cortex (orange), ₃₄₀ and between nodes above the contralateral motor and central electrodes (green) (Fig. 8 A). In contrast, ₃₄₁ OS showed a densely connected network between all nodes (cf. Fig. 6), but more strikingly an increase in ₃₄₂ interhemispheric horizontal connectivity between nodes above the frontal, frontocentral, and central regions, ₃₄₃ i.e., between nodes above the ipsilateral and contralateral frontal and above the ipsilateral and contralateral ₃₄₄ motor cortex (Fig. 8 B). ₃₄₅

Additionally, we investigated the Louvain clustering for the aggregated networks (Fig. 8 C and D). Both ₃₄₆ aggregated networks could be clustered into three communities, which are represented by different colored ₃₄₇ nodes in (blue, red, and green). The YS network clustered into an occipital-parietal cluster (red nodes), a ₃₄₈ cluster mainly involving the nodes above the ipsilateral frontal, central, and contralateral motor cortex (blue ₃₄₉ nodes) and a contralateral frontal and ispilateral motor regions one (green), whereas the OS network showed ₃₅₀ a cluster that mainly involved nodes above frontal areas and ipsi- and contralateral motor cortex (blue nodes), ₃₅₁ a cluster above parietal and central areas (green nodes), and one cluster containing the remaining nodes ₃₅₂ (red). Thereby, the clustering results of the aggregated networks supported the network differences of the ₃₅₃ dynamic networks during movement execution described above. ₃₅₄

HUB node analysis   In the last step, we analyzed the HUB nodes of the dynamical graphs using the ₃₅₅ betweenness centrality. We investigated the order of first appearance as well as the timecourse of the HUB ₃₅₆ nodes (Fig. 9). In both groups, occipital nodes appeared as the first HUB nodes in the system. In YS, HUB ₃₅₇ nodes were then shifted to parieto-central (light blue) and later to contralateral motor nodes. After most ₃₅₈ participants pressed the button (cf. Fig. 9: B (boxplot)), the HUB nodes switched back and forth between ₃₅₉ central/fronto-central (dark blue) and contralateral motor nodes. Whereas OS HUB nodes first shifted to central ₃₆₀ and fronto-central nodes, from there to contralateral frontal nodes, and only in the last step to contralateral ₃₆₁ motor nodes. A comparison of the timecourses showed that OS HUBs remained roughly 100 ms longer in ₃₆₂ occipital nodes than in YS. Thus, YS HUB nodes shifted about 100 ms earlier to parieto-central nodes and ₃₆₃ finally also to contralateral motor nodes. ₃₆₄

Because of the results presented in Fig. 9, we hypothesized that the timing of connectivity between electrodes ₃₆₅ above contralateral motor and central regions might be a crucial factor in establishing the differences in ₃₆₆ behavioral output observed between OS and YS. For this, we investigated whether the peak time of the rPLV ₃₆₇ was related to the timing of movement output, i.e., the reaction times. Since rPLV was strongest either between ₃₆₈ C4 and Cz/FCz or CP4 and Cz/FCz, we decided to create a virtual connection by averaging over all four ₃₆₉

Fig 8. Aggregated networks in aging. Top: Representation of the aggregated networks for younger (A) and older subjects (B). The thickness of the lines accounts for the number of occurrences of the corresponding edge in the respective time interval. Bottom: Circular representation of the aggregated networks for younger (C) and older subjects (D). Nodes are color-coded depending on assigned clusters. Edges related to the core motor network are color-coded in blue (ipsilateral), green (contralateral), and orange (interhemispheric).

possible combinations. The reaction time was significantly correlated to this virtual contralateral motor-central

connection, $p < 0.01, \rho = 0.42$ (see Fig. 10).

Fig 9. HUB nodes. Timepoints of first occurrences of HUB nodes (A) and timecourses of HUB nodes in the interval [0, 750] ms (B) - represented by groups of electrodes (Occipital: PO7, PO8; Central: Pz (light blue), CPz, Cz, FCz; contra-Frontal: FC2, FC4, FC6, F2, F4, F6; contra-Motor: C2, C4, C6, CP2, CP4, CP6) - as defined by betweenness centrality in the networks of younger and older subjects. The mean reaction times for each age group are depicted by boxplots on the x-axis.

Discussion

In the current study we were interested in age-related changes in the processing of neural information during externally-triggered, i.e. visually-guided movements. To this extent, we performed a dynamic graph analysis

Fig 10. Behavioral correlation Each subject's reaction time (both age groups) plotted against the rPLV peak time between virtual contralateral motor and central electrodes.

on EEG data recorded from healthy younger and older participants, while they performed a visually-cued ₃₇₅ finger-tapping task. The graphs were based on phase-locking as a measure of inter-regional synchronization, ₃₇₆ which, according to the communication through coherence (CTC) hypothesis, reflects increased functional ₃₇₇ connectivity between neural populations expressed by coherently coordinated firing patterns [22, 23]. For this ₃₇₈ purpose, we calculated the rPLV, which measures the instantaneous synchronization between two distinct ₃₇₉ recording sites. The great advantage of rPLV is that it is capable of measuring fast transient synchronizations ₃₈₀ and thus preserves the advantage of the high temporal resolution of EEG recordings, which is essential for ₃₈₁ the construction of dynamic graphs related to the motor task. In contrast, other frequently used methods like ₃₈₂ dynamic causal modeling (DCM) or Granger causality require statistically stationary signals and thus result in ₃₈₃ connectivity information with lower time resolution. Since some of the effects we observed (e.g., decreased ₃₈₄ node flexibility) lasted for only a few hundred milliseconds and, hence, could not guarantee stationary signals ₃₈₅ (as they took place close to the button press), the application of rPLV instead of DCM or Granger causality ₃₈₆ was crucial for investigating the underlying dynamic neural networks in our study. ₃₈₇

To start with, we computed the relative power changes of all considered electrodes in the $\delta - \theta, \alpha$ and β ₃₈₈ frequency range and the rPLVs between all pairs of the considered electrodes in these frequencies. Note ₃₈₉

that only rPLVs in the $\delta - \theta$ frequency range turned out to be significantly increased. The electrodes were then defined as the nodes of the dynamic graph, while the edges for each time-point represented significantly increased rPLVs. We compared the time evolution of the dynamic networks that we obtained for the younger and older subjects with a focus on stimulus processing, movement preparation and movement execution. We investigated the networks considering three different aspects: the overall connectivity, i.e., the node degree of the whole graph in general and the one of the motor-related nodes, the formation of clusters, i.e., subnetworks during the different phases of the motor act, and the evolution of HUB nodes over time.

Relative amplitude changes in oscillations The analysis of the changes in event-related amplitude differences has revealed varying effects for the individual frequency bands. In the low frequencies ($\delta - \theta$), neither in the visually-cued motor nor in the contrast condition significant differences in the strength of the amplitude changes between both groups could be seen. In the higher frequencies (α and β), well known motor effects, the so-called ERD and ERS could be observed in the oscillations [49–52]. The ERD during motor execution (approximately 500-700 ms after stimulus) differs in the mid frequency range (α). Here, the OS show a significantly stronger and longer ERD. However, this change was much less pronounced in the contrast as compared to the visually-cued motor condition. In the higher frequencies (β), a significant difference between YS and OS could be seen especially in the time after motor execution (about 1000 ms) in ERS. The YS show a clear β-rebound after movement execution, which is not present in OS. In contrast to the results in the α frequency range, this effect is visible in the visually-cued motor as well as in the contrast condition. Analogous results for aging differences in the respective frequencies, in particular in ERD and ERS have been reported [3, 53–55]

Setup of dynamic networks In the construction of the dynamic networks in the $\delta - \theta$ frequency range, we defined a connection at a given time point to be present if the respective rPLV was significantly increased compared to baseline (i.e., before the stimulus) levels. Since this involved extensive testing, we applied FDR correction (q = 0.05) to account for the considerable number of multiple comparisons. One could argue that this method is not strict enough. However, it has turned out that even with a stricter FDR correction (q = 0.01), the main differences in motor-related networks between YS and OS prevailed and the core message of our results remained unaffected (results not shown). Furthermore, we decided to analyze our data at the electrode level. At least in principle, a source localization could have provided further insights into the origin of the effects reported here. However, we refrained from doing so to avoid the ill-posed inverse problem, which only provides an assumption of the source activity and could, therefore, lead to erroneous interpretations [56–58]. Instead, we used the small Laplacian reference to substantially reduce the effect of volume conduction on our data. Although not a source localization method, through the small Laplacian, electrodes that are directly above the sources are maximally sensitive to the underlying activity. As we can see in our results, there are

many neighboring electrode pairs that do not have significant rPLVs, i.e. connections, which one would expect to see in the case of pure volume conduction (e.g. C1-Cz, C4-C6) (see Fig. 9 B). These observations suggest that our results cannot primarily be driven by volume conduction.

Dynamic graph connection density Several fMRI studies have reported an overactivation related to the aging human brain [5–8]. The results of the dynamic graphs in the $\delta - \theta$ frequency band reported here revealed a general increase in node degree in all electrodes during movement preparation and execution (Fig. 6 A). Especially in motor-related electrodes, YS showed a plateau-like behavior while OS node degree showed a peak roughly twice as high during movement preparation (Fig. 6 B). This effect might reflect *over-connectivity*, which may relate to the overactivation previously reported in fMRI studies. One might argue that this over-connectivity in OS is driven by increases in amplitude in the respective frequency band. This, however, can be ruled out due to the fact that no difference in relative change in $\delta - \theta$ amplitude could be observed between YS and OS in neither the motor and nor the contrast condition. Furthermore, significant differences in relative change in α and β amplitudes occur only after movement.

Community structure The analysis of the aggregated networks revealed a substantial increase in the number of interhemispheric connections in OS. The cluster analysis of aggregated networks in YS unraveled three clusters, a cluster mainly consisting of parietal and occipital electrodes, a cluster of ipsilateral frontal, central, and contralateral motor electrodes, and a cluster that included ipsilateral motor and contralateral frontal electrodes. In OS, on the other hand, there was no clear separation in clusters possible. This might be related to the increased number of interhemispheric connections and thus a weaker inter-regional separability, which is consistent with the HAROLD model stating reduced hemispheric asymmetry during cognitive tasks [9].

Additionally, we performed a similar cluster analysis on the dynamical graphs for each time-point. The clusters were different in both groups for almost the whole analyzed period (Fig. 7 B,C). The dynamic graph communities in YS during motor execution were, similar to the aggregated ones described above, separated in a more structured way, i.e., a dominant cluster including ipsilateral frontal, central, and contralateral motor electrodes, (cf. Fig. 7 B). The most striking difference between YS and OS was observed during movement preparation, i.e., immediately before movement onset. While YS showed a peak in VI between time-points in this epoch, i.e., a very variable network structure, OS networks exhibited a dip in VI (cf. Fig. 7 A, C in [200-300]ms). This visual inspection was confirmed by an analysis of node flexibility, which revealed a significant decrease in node flexibility in OS in that time interval. The decreased flexibility in OS networks might be related to the *over-connectivity* described above and the fact that OS expressed difficulties in recruiting task-specific subnetworks [9, 13, 14] and needed the whole network, i.e., higher effort to keep nearly the same

task performance as YS.

Network information flow   We furthermore included an analysis of HUB nodes of the dynamical graphs, i.e., nodes connecting different clusters by a large number of connections. Here, we focused on the first occurrence and the timecourses of HUB nodes as one of the key players in the network. In YS, the networks involved a direct shift of the HUB nodes from occipital via parieto-central to contralateral sensorimotor electrodes. Once YS had executed the movement, HUB nodes switched back and forth between central and contralateral motor electrodes, which might be related to the incoming sensory feedback from the button press. The HUBs of OS, in contrast, remained roughly 100ms longer in the occipital regions, then shifted to central/fronto-central electrodes and then targeted an additional HUB node above contralateral frontal electrodes, which appeared at 370 ms, i.e., approximately the same time as the shift to contralateral motor areas in YS. Finally, the connectivity to contralateral sensorimotor areas dominated, which occurred roughly 80 ms later than in YS (Fig. 9). Here, we can observe two things, on the one hand the HUB nodes in OS remain longer in occipital regions, which indicates already a delay in the processing of the visual stimulus, and on the other hand a shift to more frontal areas. This shift of the HUB nodes is compatible with the PASA (posterior-anterior shift in aging) phenomenon [8], which refers to the fact that the overall activity in OS is shifted to more frontal areas representing a stronger involvement of cognitive control. We hypothesize that this *detour* via frontal areas is accountable for longer reaction times in OS. This hypothesis is supported by the reported positive correlation between the reaction times and the peak timing of the connection between central and contralateral motor electrodes (cf. Fig. 10).

Motor network connectivity in voluntary movements   In a previous publication [26] , we reported on an invariant, i.e. present and unchanged in all movement phases, motor network in younger volunteers consisting of iPFC/iPM, mPFC/SMA, and cSM while subjects were performing voluntary, i.e. internally triggered, movements. This invariant network was significantly attenuated in OS and accompanied by increased interhemispheric connectivity. The results we have presented here reveal comparable patterns. YS established most (motor-related) connections between ipsilateral frontal, central, and contralateral motor regions (cf. aggregated networks in Fig. 8 A and dynamic networks during motor execution in Fig. 7 B). The networks of OS, on the other hand, showed additional interhemispheric connections, especially between ipsilateral frontal and contralateral frontal as well as ipsilateral motor and contralateral motor electrodes. Also, similar to the results during voluntary movements, OS exhibited higher connectivity between central and contralateral frontal electrodes. These results indicate that by investigating the contrast between visually-cued and vision-only conditions, we devised a condition alike the pure motor task (voluntary movement). Furthermore, these findings indicate that we identified a highly stable core network (iPFC/iPM - mPFC/SMA - cSM) that is important for movement execution over several experimental conditions.

Acknowledgements 488

This study was funded by the University of Cologne Emerging Groups Initiative (CONNECT) within the framework of the Institutional Strategy of the University of Cologne and the German Excellence Initiative. SD gratefully acknowledges additional support from the German Research Foundation, Germany (DA1953/5-2). GRF gratefully acknowledges additional support from the Marga and Walter Boll Foundation.

References

1. Wu T, Hallett M. The influence of normal human ageing on automatic movements. The Journal of Physiology. 2005;562(2):605–615.

2. Seidler RD, Bernard JA, Burutolu TB, Fling BW, Gordon MT, Gwin JT, et al. Motor control and aging: Links to age-related brain structural, functional, and biochemical effects. Neuroscience & Biobehavioral Reviews. 2010;34(5):721–733. doi:10.1016/J.NEUBIOREV.2009.10.005.

3. Liu L, Rosjat N, Popovych S, Wang BA, Yeldesbay A, Toth TI, et al. Age-related changes in oscillatory power affect motor action. PloS one. 2017;12(11):e0187911.

4. Niessen E, Fink GR, Hoffmann HEM, Weiss PH, Stahl J. Error detection across the adult lifespan: Electrophysiological evidence for age-related deficits. NeuroImage. 2017;152:517–529.

5. Sailer A, Dichgans J, Gerloff C. The influence of normal aging on the cortical processing of a simple motor task. Neurology. 2000;55(7):979–985.

6. Heuninckx S, Wenderoth N, Debaere F, Peeters R, Swinnen SP. Neural basis of aging: the penetration of cognition into action control. Journal of Neuroscience. 2005;25(29):6787–6796.

7. Ward NS, Swayne OBC, Newton JM. Age-dependent changes in the neural correlates of force modulation: An fMRI study. Neurobiology of Aging. 2008;29(9):1434 – 1446. doi:https://doi.org/10.1016/j.neurobiolaging.2007.04.017.

8. Dennis NA, Cabeza R. Neuroimaging of healthy cognitive aging. In: The handbook of aging and cognition. Psychology Press; 2011. p. 10–63.

9. Cabeza R. Hemispheric asymmetry reduction in older adults: the HAROLD model. Psychology and aging. 2002;17(1):85.

10. Cabeza R, Grady CL, Nyberg L, McIntosh AR, Tulving E, Kapur S, et al. Age-related differences in neural activity during memory encoding and retrieval: a positron emission tomography study. Journal of neuroscience. 1997;17(1):391–400.

11. Nolde SF, Johnson MK, Raye CL. The role of prefrontal cortex during tests of episodic memory. Trends in cognitive sciences. 1998;2(10):399–406.

12. Reuter-Lorenz PA, Jonides J, Smith EE, Hartley A, Miller A, Marshuetz C, et al. Age differences in the frontal lateralization of verbal and spatial working memory revealed by PET. Journal of cognitive neuroscience. 2000;12(1):174–187.

13. Mitrushina M, Satz P. Analysis of longitudinal covariance structures in assessment of stability of cognitive functions in elderly. Brain dysfunction. 1991;.

14. Babcock RL, Laguna KD, Roesch SC. A comparison of the factor structure of processing speed for younger and older adults: Testing the assumption of measurement equivalence across age groups. Psychology and aging. 1997;12(2):268.

15. Fell J, Dietl T, Grunwald T, Kurthen M, Klaver P, Trautner P, et al. Neural bases of cognitive ERPs: more than phase reset. Journal of cognitive neuroscience. 2004;16(9):1595–1604.

16. Popovych S, Rosjat N, Toth TI, Wang BA, Liu L, Abdollahi RO, et al. Movement-related phase locking in the delta–theta frequency band. NeuroImage. 2016;139:439–449.

17. Palva S, Palva JM. Discovering oscillatory interaction networks with M/EEG: challenges and breakthroughs. Trends in cognitive sciences. 2012;16(4):219–230.

18. Singer W. Neuronal Synchrony: A Versatile Code for the Definition of Relations? Neuron. 1999;24(1):49–65. doi:10.1016/S0896-6273(00)80821-1.

19. SINGER W. Synchrony, oscillations, and relational codes. The Visual Neurosciences. 2004; p. 1665–1681.

20. Uhlhaas PJ, Roux F, Rodriguez E, Rotarska-Jagiela A, Singer W. Neural synchrony and the development of cortical networks. Trends in Cognitive Sciences. 2010;14(2):72–80. doi:10.1016/J.TICS.2009.12.002.

21. van Wijk BCM, Beek PJ, Daffertshofer A. Neural synchrony within the motor system: what have we learned so far? Frontiers in Human Neuroscience. 2012;6:252. doi:10.3389/fnhum.2012.00252.

22. Fries P. A mechanism for cognitive dynamics: neuronal communication through neuronal coherence. Trends in cognitive sciences. 2005;9(10):474–480.

23. Fries P. Rhythms for cognition: communication through coherence. Neuron. 2015;88(1):220–235.

24. Baker S, Olivier E, Lemon R. Coherent oscillations in monkey motor cortex and hand muscle EMG show task-dependent modulation. The Journal of physiology. 1997;501(1):225–241.

25. Baker S, Spinks R, Jackson A, Lemon R. Synchronization in monkey motor cortex during a precision grip task. I. Task-dependent modulation in single-unit synchrony. Journal of Neurophysiology. 2001;85(2):869–885.

26. Rosjat N, Liu L, Wang BA, Popovych S, Tóth T, Viswanathan S, et al. Aging-associated changes of movement-related functional connectivity in the human brain. Neuropsychologia. 2018;117:520–529.

27. Lachaux JP, Rodriguez E, Martinerie J, Varela FJ. Measuring phase synchrony in brain signals. Human brain mapping. 1999;8(4):194–208.

28. Sizemore AE, Bassett DS. Dynamic graph metrics: Tutorial, toolbox, and tale. NeuroImage. 2018;180:417–427.

29. Holme P, Saramäki J. Temporal networks. Physics Reports. 2012;519(3):97 – 125. doi:https://doi.org/10.1016/j.physrep.2012.03.001.

30. Oldfield RC. The assessment and analysis of handedness: the Edinburgh inventory. Neuropsychologia. 1971;9(1):97–113.

31. Spreen O, Strauss E. A compendium of neuropsychological tests: administration, norms and commentary. Oxford University Press, New York; 1998.

32. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. Journal of psychiatric research. 1975;12(3):189–198.

33. Agrell B, Dehlin O. The clock-drawing test. Age and Ageing. 1998;27(3):399–403. doi:10.1093/ageing/27.3.399.

34. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. Archives of general psychiatry. 1961;4(6):561–571.

35. Wang BA, Viswanathan S, Abdollahi RORO, Rosjat N, Popovych S, Daun S, et al. Frequency-specific modulation of connectivity in the ipsilateral sensorimotor cortex by different forms of movement initiation. NeuroImage. 2017;159:248–260. doi:10.1016/J.NEUROIMAGE.2017.07.054.

36. Makeig S, Bell AJ, Jung TP, Sejnowski TJ. Independent component analysis of electroencephalographic data. In: Advances in neural information processing systems; 1996. p. 145–151.

37. Hjorth B. An on-line transformation of EEG scalp potentials into orthogonal source derivations. Electroencephalography and clinical neurophysiology. 1975;39(5):526–530.

38. Kronland-Martinet R, Morlet J, Grossmann A. Analysis of sound patterns through wavelet transforms. International journal of pattern recognition and artificial intelligence. 1987;1(02):273–302.

39. Rubinov M, Sporns O. Complex network measures of brain connectivity: uses and interpretations. Neuroimage. 2010;52(3):1059–1069.

40. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal statistical society: series B (Methodological). 1995;57(1):289–300.

41. Blondel VD, Guillaume JL, Lambiotte R, Lefebvre E. Fast unfolding of communities in large networks. Journal of statistical mechanics: theory and experiment. 2008;2008(10):P10008.

42. Reichardt J, Bornholdt S. Statistical mechanics of community detection. Physical Review E. 2006;74(1):016110.

43. Ronhovde P, Nussinov Z. Multiresolution community detection for megascale networks by information-based replica correlations. Physical Review E. 2009;80(1):016109.

44. Sun Y, Danila B, Josić K, Bassler KE. Improved community structure detection using a modified fine-tuning strategy. EPL (Europhysics Letters). 2009;86(2):28004.

45. Meilă M. Comparing clusterings—an information based distance. Journal of multivariate analysis. 2007;98(5):873–895.

46. Bassett DS, Wymbs NF, Porter MA, Mucha PJ, Carlson JM, Grafton ST. Dynamic reconfiguration of human brain networks during learning. Proceedings of the National Academy of Sciences. 2011;108(18):7641–7646.

47. Brandes U. A faster algorithm for betweenness centrality. Journal of mathematical sociology. 2001;25(2):163–177.

48. Kintali S. Betweenness centrality: Algorithms and lower bounds. arXiv preprint arXiv:08091906. 2008;. 595

49. Gerloff C, Richard J, Hadley J, Schulman A, Honda M, Hallett M. Functional coupling and regional 596 activation of human cortical motor areas during simple, internally paced and externally paced finger 597 movements. Brain. 1998;121:1513–1531. 598

50. Gerloff C, Uenishi N, Kunieda T, Hallett M, et al. Cortical activation during fast repetitive finger 599 movements in humans: steady-state movement-related magnetic fields and their cortical generators. 600 Electroencephalogr Clin Neurophysiol. 1998;109:444–453. 601

51. Pfurtscheller G, da Silva F. Event-related EEG/MEG synchronization and desynchronization: basic 602 principles. Clinical Neurophysiology. 1999;110:1842–1857. 603

52. Neuper C, Woertz M, Pfurtscheller G. ERD/ERS patterns reflecting sensorimotor activation and 604 deactivation. Progress in Brain Research. 2006;159:211–222. 605

53. Labyt E, Szurhaj W, Bourriez J, Cassim F, Defebvre L, Destee A, et al. Changes in oscillatory cortical 606 activity related to a visuo-motor task in young and elderly healthy subjects. Clinical Neruophysiology. 607 2003;114(6):1153–1166. 608

54. Yordonova J, Kolec V, Hohnsbein J, Falkenstein M. Sensorimotor slowing with ageing is mediated by a 609 functional dysregulation of motor-generation processes. Brain. 2004;127:351–362. 610

55. Toledo D, Manzano G, Barela J, Kohn A. Cortical correlates of response time slowing in older 611 adults: ERP and ERD/ERS analyses during passive ankle movement. Clinical Neurophysiology. 612 2016;127(1):655–663. 613

56. Bradley A, Yao J, Dewald J, Richter CP. Evaluation of electroencephalography source localization 614 algorithms with multiple cortical sources. PloS one. 2016;11(1):e0147266. 615

57. Grech R, Cassar T, Muscat J, Camilleri KP, Fabri SG, Zervakis M, et al. Review on solving the inverse 616 problem in EEG source analysis. Journal of neuroengineering and rehabilitation. 2008;5(1):25. 617

58. Wendel K, Väisänen O, Malmivuo J, Gencer NG, Vanrumste B, Durka P, et al. EEG/MEG Source 618 Imaging: Methods, Challenges, and Open Issues. Computational Intelligence and Neuroscience. 619 2009;2009:1–12. doi:10.1155/2009/656092. 620