1   Classification: Biological Sciences, Evolutionary Biology
2
3   **The generalist herbivore *Tetranychus urticae* (Koch) adapts to novel plant hosts through rapid**
4   **evolution of metabolic resistance**
5

6   Golnaz Salehipourshirazi[1,#], Kristie Bruinsma[1,#], Huzefa Ratlamwala[1,#], Sameer Dixit[1,#], Vicent
7   Arbona[3], Emilie Widemann[1], Maja Milojevic[1], Pengyu Jin[1], Nicolas Bensoussan[1], Aurelio Gómez-
8   Cadenas[3], Vladimir Zhurov[1], Miodrag Grbic[1,2,4], Vojislava Grbic[1,*]
9
10   [1]Department of Biology, The University of Western Ontario, London, Ontario, N6A 5B8, Canada.
11   [2] Instituto de Ciencias de la Vid y el Vino (CSIC, UR, Gobiernode La Rioja), 26006 Logrono, Spain.
12   [3]Departament de Ciències Agràries i del Medi Natural, Universitat Jaume I, E-12071 Castelló de
13   la Plana, Spain. [4] Department of Biology, University of Belgrade, Serbia
14
15   [#]These authors contributed equally
16
17   [*]Corresponding author: Vojislava Grbic, The University of Western Ontario, 1151 Richmond St,
18   London, Ontario, N6A 5B8, Canada; phone: 519 661-2111, e-mail: vgrbic@uwo.ca
19

Abstract

Genetic adaptation, occurring over long evolutionary time, enables host-specialized herbivores to develop novel resistance traits and to counteract the defenses of a narrow range of host plants. In contrast, physiological acclimation, leading to the suppression and/or detoxification of host defenses is hypothesized to enable generalists to shift between plant hosts. Here, we examined the long-term response of an extreme generalist, the two-spotted spider mite, *Tetranychus urticae* Koch (TSSM), to the shift to the non-preferred and novel host plant *Arabidopsis thaliana*. We identified the key requirement of two tiers of cytochrome P450 monooxygenases for TSSM adaptation to *Arabidopsis*: general xenobiotic-responsive P450s that have a limited contribution to mite adaptation to *Arabidopsis* and adaptation-associated P450s that efficiently counteract *Arabidopsis* defenses, illustrating that in about 25 generations of TSSM selection on *Arabidopsis* plants mites evolved metabolic resistances characteristic of both generalist and specialist herbivores.

Keywords: *Arabidopsis*, host shift, biochemical adaptation, plant allelochemicals, metabolic resistance, suppression of plant defenses, xenobiotic responsiveness, detoxification, *NADPH-cytochrome P450 reductase*, RNAi, plant-herbivore interaction

Introduction

During millions of years of co-evolution with their host plants, herbivores have developed two main strategies to counteract plant resistance traits. Specialist herbivores have evolved highly efficient adaptation mechanisms against a limited set of host defenses, including: modified feeding behavior (Helmus and Dussourd, 2005), suppression of plant defenses (Zhao et al., 2015), reduced xenobiotic target site sensitivity (Dobler et al., 2012), sequestration (Beran et al., 2018) and detoxification of plant toxins (Ratzka et al., 2002). In contrast, generalist herbivores evolved an innate ability to feed on a broad range of host plants that display a wide array of resistance traits (Despres et al., 2007; Barrett and Heil, 2012; Heidel-Fischer and Vogel, 2015). Attenuation of plant responses induced by herbivore feeding, consistent with the suppression of host defenses, have been described for many generalists (Zarate et al., 2007; Kant et al., 2008; Musser et al., 2012; Wu et al., 2012). Consistently, effectors that modulate plant defenses have been identified in secretions of a number of generalist herbivores belonging to different feeding guilds (Musser et al., 2002; Hogenhout and Bos, 2011; Wu et al., 2012; Bass et al., 2013; Jonckheere et al., 2016; Kaloshian and Walling, 2016; Basu et al., 2018). Their mode of action is largely unknown but, to be effective against many host plants, it is assumed that they either target conserved compounds or pathways associated with plant defense, or that they collectively have a very broad spectrum activity with only a specific subset being effective against any particular host. Another mechanism of host-adaptation is metabolic resistance whereby herbivores effectively detoxify ingested plant toxins. In specialists, metabolic resistance is based on a limited number of detoxification enzymes that have high specificity and efficiency for a given plant toxin (Ratzka et al., 2002; Li et al., 2003; Wittstock et al., 2004; Mao et al., 2006; Gloss et al., 2014; Heidel-Fischer et al., 2019). Genes encoding these enzymes are usually expressed constitutively at high level and can be further induced in the presence of plant toxin. In contrast, it is assumed that generalist herbivores rely on ubiquitous classes of detoxification enzymes (e.g. carboxyl/cholinesterases (CCEs), cytochrome P450 monooxygenases (CYP450s), glutathione S-transferases (GSTs), UDP-glycosyltransferases (UGTs), and ABC transporters (ABCs)) that were shown to accept structurally diverse substrates which they metabolize with low levels of activity (Li et al., 2004; Halon et al., 2015; Shi et al., 2018; Snoeck et al., 2019). Genes encoding general detoxification enzymes underwent extensive amplification and neofunctionalization, and are transcriptionally responsive to a wide range of xenobiotics (Govind et al., 2010; Grbic et al., 2011; Zhurov et al., 2014; Wybouw et al., 2015; Muller et al., 2017; Schweizer et al., 2017). Thus, it is hypothesized that attenuation of plant defenses combined with the expanded and functionally versatile detoxification capabilities enable generalist herbivores to cope with diverse allelochemicals and to feed on many host plants.

Tetranychus urticae Koch (TSSM) is a striking example of generalist herbivore that feeds on more than 1,100 plant species from over 100 families (Migeon and Dorkeld, 2006-2019). Such a wide host range indicates that TSSM can counteract a great diversity of plant resistance traits. However, individual TSSM populations do not perform equally well on all potential host plants. Instead, TSSM has the outstanding ability to adapt to new hosts, in just 20 to 25 generations (Gould, 1979; Fry, 1989; Magalhaes et al., 2007; Wybouw et al., 2015). The mechanism of this extremely rapid host adaptability is not known. The analysis of plant transcriptional changes following the host shift revealed that some TSSM populations can suppress plant induced responses (Kant et al., 2008; Wybouw et al., 2015). At the same time, TSSM massively reprograms its detoxification capacity (Dermauw et al., 2013; Zhurov et al., 2014; Wybouw et al., 2015) and the complement of its salivary secretions (Jonckheere et al., 2016; Villarroel et al., 2016; Jonckheere et al., 2018). However, no functional evidence explains whether these changes contribute to TSSM host adaptation or if they merely reflect stress

responses or a different feeding physiology due to the host shift. Thus, it remains unclear
whether adaptation to the new host plant by *T. urticae* requires manipulation of host plant
defenses, detoxification of host allelochemicals or both, and if the initial xenobiotic responses
are sufficient for TSSM adaptation to a new host, or if changes over a longer period of time are
required for the evolution of TSSM host-adaptation.
Arabidopsis thaliana is a non-preferred host for the TSSM London reference population
that is reared on beans – *Phaseolus vulgaris* (Zhurov et al., 2014). In an experimental
evolutionary setup, we adapted the TSSM London ancestral population to *Arabidopsis.* Taking
advantage of well characterized *Arabidopsis* induced defenses against mite herbivory (Zhurov et
al., 2014), we show that *Arabidopsis*-adapted TSSM do not suppress them. Using manipulative
pharmacological and reverse genetics experiments that independently suppressed the activities
of families of detoxification enzymes we provide, for the first time, *in vivo* functional evidence
that *T. urticae* requires two tiers of P450 enzyme activities for host adaptation. We show that
P450s responsive to the initial encounter of the new host have a small contribution to mite
adaptation to *Arabidopsis*. Instead, a second tier of P450 activity, specific to *Arabidopsis*-
adapted mites, is the major contributor to mite adaptation. Our data demonstrate that the
reprogramming of the mite detoxification system occurred over approximately 25 generations
of TSSM selection on *Arabidopsis* resulting in metabolic resistance with features characterized in
both generalist and specialist herbivores. Our results show that the detoxification mechanisms
involved in herbivore adaptation to novel host plants can evolve extremely rapidly and are key
to TSSM adaptation to *Arabidopsis*.
107
108 **Results and discussion**
109
TSSM feeding induces JA-regulated *Arabidopsis* defense compounds
Jasmonic acid (JA) and its bioactive conjugate jasmonoyl-isoleucine (JA-Ile) were shown to be
required for *Arabidopsis* defenses against TSSM herbivory (Zhurov et al., 2014). To determine if
JA-induced processes are sufficient for *Arabidopsis* resistance against TSSM, we sprayed
Arabidopsis plants with methyl jasmonate (MeJA) and determined the effect of induced JA-
responses on mite performance. Mite fecundity dramatically decreased (5-fold) on MeJA
treated Col-0 plants, Figure 1A, indicating that JA-induced defenses are major components of
Arabidopsis resistance against mite herbivory. JA responsiveness is mediated by COI1 as well as
the MYC2, MYC3 and MYC4 (MYC2,3,4) transcriptional activators that induce the expression of a
wide spectrum of JA-regulated genes, Figure 1B and (Devoto et al., 2005; Fernandez-Calvo et al.,
2011). The *myc2,3,4* mutant plants were extremely susceptible to *T. urticae*, Figure 1A. Among
MYC2,3,4-regulated genes are *CYP79B2* and *CYP79B3* that are required for the synthesis of Trp-
derived secondary metabolites, Figure 1A and (Hull et al., 2000; Mikkelsen et al., 2000;
Schweizer et al., 2013). Trp-derived secondary metabolites include several known
allelochemicals (indole glucosinolates, camalexin and cyanogenic glucosides) directly
synthesized from Trp and callose whose synthesis is indirectly regulated through this pathway
(Bednarek, 2012; Rajniak et al., 2015). Of these compounds, indole glucosinolates (IGs) were
implicated in deterrence of aphid (Kim and Jander, 2007; Kim et al., 2008), whiteflies (Elbaz et
al., 2012), TSSM (Zhurov et al., 2014) and leaf mining drosophilid (Whiteman et al., 2011;
Whiteman et al., 2012) herbivory. The modest increase in TSSM fecundity when mites fed on
cyp79b2 cyp79b3 plants, Figure 1A, indicates that Trp-dependent defenses have a relatively
minor contribution to *Arabidopsis* defenses against TSSM. These data further indicate that
MYC2,3,4 induce synthesis of additional *CYP79B2 CYP79B3*-independent *Arabidopsis* defense
compounds that prominently contribute to *Arabidopsis* resistance to TSSM. The identity of these

compounds is at present not known. Thus, upon TSSM herbivory, *Arabidopsis* plants induce
synthesis of JA-regulated defense compounds against TSSM, including Trp-derived secondary
metabolites and, *CYP79B2 CYP79B3*-independent molecules, Figure 1B.
137
Mites can adapt to a complex array of *Arabidopsis* defenses
Because of the complexity of *Arabidopsis* induced defenses against TSSM herbivory, how could a
mite population adapted to a specific host plant switch to *Arabidopsis*? In an experimental
evolutionary setup, the reference London strain, an ancestral TSSM population reared on bean
and highly susceptible to *Arabidopsis* defenses (Zhurov et al., 2014), was transferred and
continuously maintained on *Arabidopsis* plants for 18 months (≥25 generations), Figure 2A. We
initially infested *Arabidopsis* plants with approximately 1,000 fertilized female mites in a
triplicated experiment on two *Arabidopsis* genotypes: a) Col-0 wild-type plants, and b) *cyp79b2*
cyp79b3 mutant plants that lack Trp-derived defenses but express the remaining of JA-regulated
defenses, Figures 1B and 2A. The performance of the ancestral and selected mite populations
was subsequently quantified by counting the number of eggs, larvae, nymphs and adults derived
from 20 adult females in 7 days. Performance was measured in two experimental regimes: a)
direct transfer, where the initial mites were moved from their corresponding rearing plant hosts
directly to the experimental plants, and b) indirect transfer, that included mite maintenance on
bean plants for two generations prior to their transfer to the experimental plants. The
performance of the ancestral and selected mite populations on bean, or *Arabidopsis cyp79b2*
cyp79b3 and Col-0 plants had similar patterns in both direct (Supplemental Figure 1) and
indirect (Figure 2B) transfer regimes, indicative of genetic adaptation that is independent of
maternal and environmental physiological effects.
On bean plants, Figure 2B top, *Arabidopsis*-selected mite populations had slight but
significantly lower performance relative to the ancestral population, indicating that TSSMs can
evolve the ability to exploit different hosts without major reduction in their performance on the
ancestral plants. On *cyp79b2 cyp79b3* plants, Figure 2B middle, *Arabidopsis*-selected mite
populations performed significantly better than the ancestral population, suggesting that they
are able to overcome *CYP79B2 CYP79B3*-independent *Arabidopsis* defenses to a similar extent.
On Col-0 plants, Figure 2B bottom, only mites selected on Col-0 had increased performance over
the ancestral population, showing that Col-0 selected mites were able to adapt to *CYP79B2*
CYP79B3-dependent defenses. However, mites selected on *cyp79b2 cyp79b3* plants and the
ancestral mite population that were not exposed to *CYP79B2 CYP79B3*-dependent *Arabidopsis*
defenses were susceptible to these defenses and had similar and low performances when they
fed on Col-0 plants.
We have arbitrarily chosen populations #3 of *cyp79b2 cyp79b3* and Col-0 selected mites
for further studies that were performed in indirect transfer regimes. The analysis of additional
mite fitness parameters, fecundity and *Arabidopsis* leaf damage caused by mite feeding (Figure
2C and D, respectively) confirmed the adaptation status of these mite populations (from now on
referred to as *cyp*-a (for #3 *cyp79b2 cyp79b3* selected mite population), Col-a (for #3 Col-0
selected mite population) and bean-a (for the ancestral London TSSM population)). Thus, our
data demonstrate that TSSM can adapt to novel plant hosts with a complex array of defenses.
Moreover, mite adaptation to *CYP79B2 CYP79B3*-independent *Arabidopsis* defenses (present in
both types of *Arabidopsis*-selected mite populations) can be uncoupled from mite adaptation to
Trp-derived defenses (present only in mites selected on Col-0 plants).
179
180
181

182 ***Arabidopsis*-adapted mites do not suppress *Arabidopsis* defenses**
183 Suppression of induced plant defenses was proposed to be the common and most efficient
184 mechanism of TSSM adaptation to new host plants (Jonckheere et al., 2016; Villarroel et al.,
185 2016; Blaazer et al., 2018; Jonckheere et al., 2018). Indeed, interference with either JA-
186 biosynthesis or its signalling appears to be an effective way to attenuate a whole range of
187 *Arabidopsis* defense compounds that restrict TSSM herbivory (Figure 1). Furthermore, the
188 ancestral London strain previously gave rise to tomato-adapted populations that, in the process,
189 gained the ability to suppress tomato JA-regulated responses (Wybouw et al., 2015), indicating
190 that London mites are able to evolve the defense suppression trait.
191 If *Arabidopsis*-adapted mites suppress *Arabidopsis* defenses, then it is expected that
192 they elicit attenuated *Arabidopsis* responses relative to those triggered by the ancestral bean-a
193 mites. The previous transcriptome and metabolome analysis of *Arabidopsis* responses to TSSM
194 herbivory established that the expression of the *AOS, MYC2, CYP79B2* and *CYP79B3* genes
195 (labeled in red in Figure 1B) and the abundance of JA, JA-Ile and indol-3-ylmethylglucosinolate
196 (I3M) metabolites (labeled in green in Figure 1B) are reliable markers of *Arabidopsis* induced
197 defenses, at 24 h following mite infestation (Zhurov et al., 2014). Thus, to investigate whether
198 mite adaptation to *Arabidopsis* relies on the attenuation of JA-induced defenses, we determined
199 the levels of defense marker transcripts (RT-qPCR) and metabolites (liquid chromatography–
200 mass spectrometry, LC-MS) in Col-0 plants that were challenged with bean-a, *cyp*-a or Col-a
201 mites, after 24 h of mite herbivory. Contrary to what would be expected if plant defenses were
202 attenuated, *cyp*-a and Col-a mites induced all JA-regulated marker genes to similar or higher
203 levels relative to non-adapted bean-a mites, Figure 3A. Likewise, JA, JA-Ile and I3M metabolites
204 accumulated at comparable levels in bean-a, *cyp*-a and Col-a challenged Col-0 plants (Figure 3B),
205 indicating that *Arabidopsis* defenses are augmented and not attenuated by mite adaptation. In
206 addition, one of the hallmarks of defense-suppressing TSSM populations is the similarity of their
207 performance on wild-type and defense-deficient host plants (Kant et al., 2008). However,
208 *Arabidopsis*-adapted mites, like the ancestral population, performed better on *cyp79b2 cyp79b3*
209 than on Col-0 plants, Figure 2C and D, indicating that *Arabidopsis*-adapted mites do not
210 manipulate the onset of *Arabidopsis* defenses. Cumulatively, these data demonstrate that
211 adaptation of mites to *Arabidopsis* is not based on the suppression of host defenses.
212
213 ***Arabidopsis*-adapted mites are responsive to *Arabidopsis* xenobiotics**
214 Metabolic resistance is another mechanism of herbivore adaptation to xenobiotic challenges
215 previously implicated in the evolution of TSSM resistance to pesticides (Van Leeuwen and
216 Dermauw, 2016) and mite adaptation to several plant hosts (Agrawal et al., 2002; Wybouw et al.,
217 2012; Wybouw et al., 2014; Wybouw et al., 2015). These previous studies established that
218 constitutive overexpression and/or increased expression of genes encoding xenobiotic-
219 metabolizing enzymes upon exposure to a novel host (transcriptional plasticity) are correlated
220 with TSSM's increased detoxification potential and resistance to pesticides or host
221 allelochemicals. Moreover, the requirement for rapid transcriptional plasticity has been
222 demonstrated in the adaptation of the generalist *M. persicae* to a new host plant (Mathers et al.,
223 2017).
224 We have previously identified forty genes, whose expression levels correlated with
225 increasing levels of *Arabidopsis* defenses (Zhurov et al., 2014). They primarily encoded
226 cytochrome P450 monooxygenases (P450s) and UDP-glycosyltransferase (UGTs) detoxifying
227 enzymes. We have arbitrarily chosen three genes within each class *(CYP392A1, CYP392A16,*
228 *CYP392D8, UGT201A2v2, UGT204B1* and *UGT204A5)* and tracked their expression (RT-qPCR) in
229 bean-a, *cyp*-a and Col-a mites feeding on bean, *cyp79b2 cyp79b3* and Col-0 *Arabidopsis* plants.

On the ancestral bean plants, all marker transcripts were at similar low levels in both bean- and
Arabidopsis-adapted mites, Figure 4, indicating that none of the tested genes underwent
constitutive upregulation during mite adaptation to *Arabidopsis*. Consistent with the previous
report (Zhurov et al., 2014), transfer of bean-a mites to *cyp79b2 cyp79b3* or Col-0 plants
resulted in induced expression of all tested *CYP* and *UGT* genes. Their expression was also
induced in *cyp*-a and Col-a mites when they were shifted from bean to *cyp79b2 cyp79b3* or Col-0
plants. Even though there was variability in the relative expression of *CYP* and *UGT* genes in
these mites, their levels were either comparable or lower than in bean-a mites, Figure 4. Thus,
neither constitutive expression nor inducibility of tested *CYP* and *UGT* genes increased in
Arabidopsis-adapted mites, demonstrating that: a) *Arabidopsis*-adapted mites retained
responsiveness to *Arabidopsis* xenobiotics, and b) mite genes initially induced by the shift from
bean to *Arabidopsis* plants do not associate with TSSM long-term adaptation to *Arabidopsis*.
242
Metabolic resistance underlies TSSM adaptation to *Arabidopsis*
To identify changes in the detoxification potential associated with mite adaptation to
Arabidopsis, we determined global enzymatic activity for three main protein families - esterases,
glutathione-S-transferases (GSTs) and P450s - in ancestral and adapted mite populations upon
feeding on bean and *Arabidopsis* plants. Overall, activities of all three enzymatic classes were
responsive to the host plant challenge: they were lowest when mites fed on the ancestral bean
plants and they progressively increased with the complexity of *Arabidopsis* defenses, Figure 5A,
D and Supplemental Figure 2. However, of the three classes of detoxification enzymes, the
detoxification potential of P450s is the only one that was associated with TSSM adaptation to
Arabidopsis. P450 activity was consistently higher in Col-a mites across all plant hosts, indicating
that both constitutive and inducible levels of P450 activity increased in Col-a mites, Figure 5D
(three-way ANOVA; plant host: mite strain interaction F = 103.93, p = 5.451e-16, followed by
Tukey's HSD test, p < 0.01, n = 4).
To further test the requirement of esterase, GST and P450 activities for TSSM
adaptation to *Arabidopsis*, we examined whether the separate enzymatic classes may be
correlated with the mite's ability to use *Arabidopsis* as a host. If a particular class of enzymes is
required for mite adaptation to *Arabidopsis*, then the reduction of its activity is expected to
restore the susceptibility of *Arabidopsis*-adapted mites to *Arabidopsis* defense compounds. We
used S,S,S tributyl-phosphorotrithioate (DEF, an inhibitor of esterase activity), diethyl maleate
(DEM, an inhibitor of GST activity), and piperonyl butoxide (PBO) and trichlorophenylpropynyl
ether (TCPPE) (inhibitors of P450 activity) to inhibit the activity of the indicated enzymatic
classes. In preliminary experiments, concentrations of 2000 mg/L (DEF), 100 mg/L (DEM), 1000
mg/L (PBO), and 1500 mg/L (TCPPE) were identified as sublethal but nevertheless capable of
significantly reducing the matching enzymes in Col-a mites feeding on Col-0 plants, Figure 5B, E
and G, and Supplemental Figure 2. Note that inhibitors do not affect all enzymes equally within
the targeted enzymatic class (Feyereisen, 2014). Therefore, the lack of inhibitory effect on mite
performance is not a strong evidence against the involvement of a particular enzymatic class in
mite host adaptation. Conversely, the significant decrease of mite performance on a new host
plant upon the application of a given inhibitor strongly supports the requirement of enzyme(s)
within the matching class for mite host adaptation.
The application of DEF reduced esterase activities by 32% but had no effect on the
fitness of Col-a mites when feeding on either Col-0 or *cyp79b2 cyp79b3* plants, Supplemental
Figure 2, indicating that esterases may not be required for mite adaptation to *Arabidopsis*. The
application of DEM decreased the GST enzymatic activity by 43%, Figure 5B. Inhibition of GST
activity had a minor but significant effect on the fecundity of Col-a mites, suggesting that GST

activity is required for the high Col-a performance on *Arabidopsis*, Figure 5C. Isothiocyanates are
the major *Arabidopsis* anti-herbivore defense compounds that are detoxified by GSTs in many
generalist caterpillars (Wadleigh and Yu, 1988; Schramm et al., 2012; Jeschke et al., 2016) and
leaf mining drosophilids (Gloss et al., 2014). However, isothiocyanates are toxic breakdown
products of aliphatic glucosinolates, shown to be ineffective against *T. urticae* (Zhurov et al.,
2014). Isothiocyanates from Trp-derived indole glucosinolates, a glucosinolate class proposed to
be toxic to *T. urticae* (Zhurov et al., 2014), are unstable and are converted to alternative
products (Wittstock and Burow, 2010). Whether some of these alternative products confer
toxicity against *T. urticae* and are one of the potential substrates for adaptation-associated GSTs
remains to be determined.
While esterase and GST inhibitors had no or little effect on TSSM fecundity and fitness,
the application of PBO significantly reduced the P450 activity (25%) and dramatically reduced
(44%) the fecundity of Col-a mites feeding on Col-a host plants, Figure 5E and F. Since the PBO
does not interfere with all P450s equally and also inhibits esterases in some arthropods
(Feyereisen, 2014), we applied another structurally unrelated P450 inhibitor, TCPPE, to confirm
that the observed effects were due to the decreased P450 activity. TCPPE, like PBO, significantly
inhibited P450 enzyme activity in Col-a mites (13%), Figure 5G. It dramatically reduced the
fecundity of Col-a mites when they fed on Col-0 (39%) and *cyp79b2 cyp79b3* plants (33%), Figure
5H, suggesting that P450 activity is required for the ability of Col-a mites to counteract
Arabidopsis defenses.
298
P450 activity is required for TSSM adaptation to *Arabidopsis*
As P450 inhibitors were applied directly to *Arabidopsis* leaves, there is a possibility that
PBO/TCPPE perturbed *Arabidopsis* defense physiology and only secondarily affected mite
fecundity. For example, the synthesis of JA metabolites (Heitz et al., 2012; Aubert et al., 2015;
Widemann et al., 2015) and Trp-derived secondary metabolites (Hull et al., 2000) occur via
pathways that include many plant P450s. Therefore, to further dissect the requirement of mite
P450s in TSSM adaptation to *Arabidopsis*, we took advantage of recently validated RNAi gene
silencing protocol in TSSM (Suzuki et al., 2017) to silence P450 pathway.
Because the *CYP* gene family is very large in TSSM, it is difficult to characterize the
involvement of P450s in mite adaptation to *Arabidopsis* using RNAi (Grbic et al., 2011). To
circumvent this hurdle, we instead silenced the expression of *NADPH-cytochrome P450*
reductase (*Tu-CPR*), a co-enzyme required for the catalytic reactions carried out by microsomal
P450s (Masters and Okita, 1980). *Tu-CPR* is a single gene in the TSSM genome encoded by the
tetur18g03390 locus (Grbic et al., 2011). It is constitutively expressed in all tissues of adult
spider mites, including the midgut epithelial and digestive cells where digestion and
detoxification of dietary xenobiotics are expected to take place (Bensoussan et al., 2018), Figure
6A. Application of dsRNA-*Tu-CPR* to Col-a mites resulted in decreased expression of *Tu-CPR,*
Figure 6B*.* Consistent with the Tu-CPR function as an essential component of the P450 enzyme
complex, the P450 activity was reduced by 38% in the *Tu-CPR* silenced mites relative to mites
exposed to dsRNA-NC (the negative control, dsRNA-NC, is homologous to a non-transcribed
genomic region, see Supplemental Table 1 and (Suzuki et al., 2017)), Figure 6C. Strikingly, the
silencing of *Tu-CPR* reduced the fecundity of Col-a mites by almost 50% when they were
challenged on the *Arabidopsis* host, but had no effect when they infested bean, Figure 6D.
Similar phenotypes were obtained upon the application of a second, independent, dsRNA-*Tu-*
CPR-1 fragment, Supplemental Figure 3, confirming the specificity of the observed phenotypes
to a loss of *Tu-CPR* function and the requirement of P450 activity for the adaptation of Col-a
mites to *Arabidopsis*.

The RNAi reverse genetics approach combined with the application of P450 inhibitors
identified several features of P450 activity. First, the reduced P450 activity, through the
application of P450 inhibitors or dsRNA-*Tu-CPR*, did not affect TSSM fecundity when mites fed
on bean plants - an ancestral host (Figure 5H and 6D, and Supplemental Figure 3), suggesting
that P450 activity is specifically required for mites' ability to use *Arabidopsis* as a host. Second,
the experimental treatments reduced global P450 activity but did not eliminate it, Figures 5E, 5G
and 6C. Despite the incomplete inhibition of P450 activity, the fitness advantage of Col-a mites
over the ancestral population on Col-0 plants was almost completely abolished (Figures 5F, 5H
6D and Supplemental Figure 2), demonstrating that P450 activity is the main contributor to
TSSM adaptation to *Arabidopsis*. Third, P450 inhibitors had a limited but significant effect on the
fecundity of bean-a ancestral TSSM population when these mites were challenged on the
Arabidopsis host (Figures 5F and H). As bean-a mites had no prior exposure to *Arabidopsis* plants,
the effect of P450 inhibitors on their fecundity uncovers the limited contribution of initially
induced P450s to mite fitness on *Arabidopsis.* Since, *Arabidopsis*-adapted mites retained
responsiveness to shift from bean to *Arabidopsis* plants, Figure 4, it is expected that these P450s
also contribute to the resistance of *Arabidopsis*-adapted mites to *Arabidopsis* defenses. Fourth,
the reduced P450 activity profoundly affected the fecundity of Col-a mites when exposed to
Arabidopsis defenses (Figures 5F and H, and Supplemental Figure 2), indicating that Col-a mites
acquired highly efficient P450 enzymatic catalytic activity(ies) against *Arabidopsis* defenses
during the adaptation process. At least some of these P450 proteins should be encoded by *CYP*
genes with increased basal expression, as increased P450 activity was detected in bean-fed Col-a
mites, Figure 5D. Fifth, inhibition of P450 activity affected Col-a mite fecundity when they fed on
both Col-0 and *cyp79b2 cyp79b3* plants, Figures 5F and H. This indicates that P450s function to
counter CYP79B2 CYP79B3-independent defenses, shown to be major contributors to
deterrence of TSSM herbivory, Figure 1.
In summary, we provided robust *in vivo* functional data proving that P450 activity is
required for *T. urticae* adaptation to the *Arabidopsis* host plant. In an unbiased analysis, we
identified two distinct modes of P450 activities, Figures 4-6. One corresponds to the early
responsive *CYP* genes that are induced in both ancestral and *Arabidopsis*-adapted mites upon
exposure to *Arabidopsis*. These *CYPs* enable limited adaptation to *Arabidopsis* defenses ((Zhurov
et al., 2014) and this study). This is consistent with the canonical detoxification mechanism in
generalist herbivores associated with physiological acclimation whereby the exposure to
xenobiotics results in the transcriptional induction of genes encoding enzymes capable of
metabolizing these defense compounds (Li et al., 2004; Sasabe et al., 2004; Mathers et al., 2017;
Shi et al., 2018). Typically, such enzymes recognize a wide range of substrates and have low
enzymatic catalytic activity, resulting in a limited contribution to the detoxification of plant
defense compounds. The other more significant contribution of P450-mediated metabolic
resistance is realized through the action of adaptation-associated P450s (Figures 5, 6 and
Supplemental Figure 3). This pattern of metabolic counteraction is reminiscent of strategies
deployed by host-specialised herbivores, in which host-adaptation associates with mutations
that enhance the enzyme activity (Li et al., 2007; Schuler, 2011) or with the overexpression of
detoxification genes (Bass et al., 2013). Therefore, we interpret the adaptation of TSSM
populations to *Arabidopsis* as a combination of metabolic resistances characteristic of both
generalist and specialist herbivores. Remarkably, the host-shift in TSSM occurs within an
ecological timeframe, over 25 generations, enabling individual TSSM populations to expand its
host range without losing the ability of feeding on the ancestral host.
373

Conclusion

The innate ability to perceive and respond to plant defenses in a process of physiological acclimation is assumed to enable generalist herbivores to utilize a wide range of host plants. On the other hand, genetic adaptation, occurring over long evolutionary time scales, is presumed to enable specialist herbivores to evolve novel adaptation traits that efficiently counteract defenses of a narrow range of host plants (Despres et al., 2007; Barrett and Heil, 2012; Heidel-Fischer and Vogel, 2015). *T. urticae* is an extreme generalist that acclimates, like other polyphagous herbivores, to host shift by reprogramming its salivary and detoxification complements (Figure 4 and (Dermauw et al., 2013; Zhurov et al., 2014; Wybouw et al., 2015; Jonckheere et al., 2016; Villarroel et al., 2016; Jonckheere et al., 2018)). However, prolonged exposure (over generations) to initially unfavorable host plants enables *T. urticae* to dramatically increase its performance (Figure 2 and (Gould, 1979; Fry, 1989; Magalhaes et al., 2007; Wybouw et al., 2015)). As demonstrated here for the *Arabidopsis*-adapted mite populations, these mites undergo genetic adaptation to a complex mixture of defenses presented by the new host plant.

Even though suppression of plant defenses has been proposed to be the prevailing mechanism of adaptation for generalist herbivores (Jonckheere et al., 2016; Villarroel et al., 2016; Blaazer et al., 2018; Jonckheere et al., 2018), our data show that neither *cyp*-a nor Col-a mite populations attenuate *Arabidopsis* JA-induced defenses. Instead, we provide multiple independent lines of evidence indicating that *Arabidopsis*-adapted mites evolve metabolic resistances characteristic of both generalist and specialist herbivores. Notably, these adaptation mechanisms unfolded over approximately 25 generations, indicating that *T. urticae* has the ability to reset its interaction with the host plant extremely rapidly.

The genomic repertoire of detoxification genes in TSSM could be the versatile toolkit enabling that generalist mite to respond to a wide range of plant defenses. If it is so, the comparative genome and transcriptome analysis of *Arabidopsis*-adapted and ancestral TSSM populations may identify genes within the P450 family that are required for host adaptation to *Arabidopsis* and reveal if adaptation-associated P450 activity is achieved by altering gene regulation, copy number or through specific mutation(s). Whether the long-term genetic adaptation to the new host described in this study is exclusive to *T. urticae* or is widespread among generalist herbivores remains unknown. However, understanding the mechanisms involved in generalist adaptation is highly relevant in the context of anthropogenic environmental alterations characterized by habitat fragmentation and loss, climate change and invasion of alien species. These changes are reshaping ecological communities and ecological interactions, and are expected to promote the proliferation of generalist species (DeLucia et al., 2012; Stireman and Singer, 2018).

Materials and methods

Plant growing and mite rearing conditions

Columbia-0 (Col-0) seeds were obtained from the Arabidopsis Biological Resource Center (ABRC, Ohio State University), *cyp79b2 cyp79b3* from B. A. Halkier (University of Copenhagen, Denmark), *myc2 myc3 myc4*, from R. Solano (Universidad Autónoma de Madrid) and bean (*Phaseolus vulgaris*, cultivar California Red Kidney) from Stokes, Thorold, Ontario, Canada. *Arabidopsis* plants were grown under 100 to 150 µmol m^{-2} sec^{-1} cool-white fluorescent light at 24°C with a 10-h/14-h (light/dark) photoperiod in controlled growth chambers. The two spotted spider mites (*Tetranychus urticae*), London reference strain (Grbic et al., 2011), were reared on bean plants in growth chambers at 24°C, 60% relative humidity, and with a 16-h/8-h (light/dark) photoperiod for more than 100 generations.

Experimental evolution

The London strain, reared on bean plants over the last ten years (referred to as bean-a strain), is an ancestral line used in the experimental evolution. Three independent selection lines were generated by transferring approximately 1,000 randomly chosen adult bean-a females to 3-week-old *cyp79b2 cyp79b3* or Col-0 *Arabidopsis* plants that were replaced biweekly. Mite populations were allowed to propagate for at least 25 generations, generating six selected lines: three independent lines selected on *cyp79b2 cyp79b3* plants, referred to as *cyp*-a lines (#1-#3), and three independent lines selected on Col-0 plants, referred to as Col-a lines (#1-#3). Selected lines were subsequently continuously reared on their corresponding plant hosts. *cyp*-a line (#3) and Col-a line (#3) were used for the follow-up experiments.

Mite performance analysis

Twenty 3-day-old adult female mites were transferred to each of the experimental plants (bean, *cyp79b2 cyp79b3* and Col-0) to be evaluated for their performance. Mite performance was measured as total population size seven days post inoculation in direct (mites were moved directly from their rearing hosts to experimental plants) and indirect (mites were reared on bean plants for two generations prior to their transfer to the experimental plants) transfer regimes. The experiment was performed in four biological replicates for each selected line. Differences between selected line population sizes were determined using a one-way ANOVA, followed by Tukey's honestly significant difference (HSD) test. All other experiments were performed with mites that were propagated for two generations on bean plants.

Plant damage analysis

Leaf damage of *cyp79b2 cyp79b3* and Col-0 *Arabidopsis* plants upon feeding of bean-a, *cyp*-a and Col-a mite strains was performed as previously described (Zhurov et al., 2014). Briefly, ten adult female mites were placed on the rosette of *Arabidopsis* plants and allowed to feed for 3 days before plants were cut at the base of the rosette. The adaxial side of the rosette was scanned using a Canon® CanoScan 8600F model scanner (Canon U.S.A. Inc., Melville, NY, U.S.A) at a resolution of 1200 dpi and a brightness setting of +25. Scanned plants were saved as .jpg files for subsequent analysis in Adobe Photoshop 5 (Adobe Systems, San Jose, CA) as previously described. The experiment was performed using 6 biological replicates/trial, and in three experimental trials. Differences in plant damage between mite strains on different hosts were determined using a three-way ANOVA, using mite strain, plant host and experimental replication as main effects, including interaction terms, followed by Tukey's honestly significant difference (HSD) test. There was a lack of interaction between experimental trials and other main effects, so data across trials were combined for increased statistical power by use of the three-way ANOVA (Brady et al., 2015).

Mite fecundity assay

Fecundity of bean-a strain on Col-0, *cyp79b2 cyp79b3* and *myc2 myc3 myc4* as well as bean-a, *cyp*-a and Col-a strains when feeding on *cyp79b2 cyp79b3* or Col-0 *Arabidopsis* plants were performed on detached leaves. Briefly, a petiole of a detached fully-elongated adult leaf was mounted through the parafilm into a small petri plate filled with water. The detached leaf was infested with a single 3-day-old adult female mite whose fecundity was monitored over two or six days. A vented lid was placed on-top of the set up to ensure the mite containment. Inoculated leaves were kept under standard mite rearing conditions and were replaced every second day in the case of 6-day period. The experiment included five to ten biological replicates for each treatment that were repeated in three experimental trials. Differences in fecundity

between mite strains on different hosts were determined using a three-way ANOVA, using mite
strain, plant host and experimental trial as main effects, including interaction terms, followed by
Tukey's honestly significant difference (HSD) test. There was a lack of interaction between
experimental trials and other main effects, so data across trials were combined for increased
statistical power (Brady et al., 2015). For mite fecundity assay on MeJA-treated Col-0 leaves, the
rosette leaves from five-week-old Col-0 plants were sprayed 3 times within 24 h with methyl
jasmonate (MeJA) (Sigma-Aldrich, Cat # 392707, 500 µM in ethanol 0.4 % (v/v)), or with ethanol
0.4 % (v/v). Subsequently, fully-elongated adult leaves were cut and each petiole was inserted in
a PCR tube containing 340 µL of 0.5 mg/mL L-Tryptophan (Sigma-Aldrich, Cat # T0254).
Detached leaves were sprayed 3 times within 24 h with MeJA or ethanol 0.4 % solution and
were transferred in new PCR tubes containing fresh tryptophan solution. Each detached leaf was
then infested with 10 adult female mites (London strain) that were starved overnight. The
number of eggs were recorded 24 h after mite addition. The experiment included nine biological
replicates and was repeated in two experimental trials. Differences in fecundity were detected
by a two-way ANOVA, using treatment and experimental trials as main effects including an
interaction term. There was a lack of interaction between experimental trials and treatment, so
data across trials were combined for increased statistical power (Brady et al., 2015).

Real-Time Quantitative Reverse Transcription-PCR Analysis
For plant marker gene analysis, bean-a, *cyp*-a and Col-a mite strains were collected from their
rearing hosts and were propagated for two generations on bean plants after which their
corresponding 3-day-old adult female mites were transferred to Col-0 plants. Mites were
allowed to feed for 24 h, after which leaves were collected and immediately frozen in liquid
nitrogen. These samples were used for the plant marker gene/metabolite analysis. Experiment
was replicated in three biological replicates/trial and in three independent trials. For TSSM
marker gene analysis, bean-a, *cyp*-a and Col-a mite strains were collected from their rearing
hosts and were propagated for two generations on bean plants after which their corresponding
3-day-old adult female mites were transferred to bean, *cyp79b2 cyp79b3* or Col-0 plants for 24 h.
Samples of 100 mites were collected and immediately frozen in liquid nitrogen. The experiment
was replicated in three biological replicates/trial and in three independent trials. Preparation of
RNA, cDNA and the real-time quantitative reverse transcription-PCR analysis was performed as
previously described (Zhurov et al., 2014). Primer sequences and amplification efficiencies (E)
used in qPCR are shown in Supplemental Table 1. *PEROXIN4* (*AT5G25760*), was used as the
reference gene for *Arabidopsis* genes and the expression of *RP49* (*tetur18g03590*) was used as
the reference for mite genes. Ct values of three technical replicates were averaged to generate a
biological replicate Ct value. For plotting, an expression values for each target gene (T) was
normalized to the reference gene (R). Normalized relative quantity (NRQ) was calculated as NRQ
$= (1 + ER)^{CtR}/(1 + ET)^{CtT}$. NRQs were Log2-transformed for statistical analysis. Mite marker gene
data were analyzed by a two-way ANOVA with mite strain and plant host as main effects
including an interaction term, followed by Tukey's honestly significant difference (HSD) test
(Rieu and Powers, 2009). For plant marker gene analysis, the data were analysed by a two-way
ANOVA, using mite strain and experimental trial as main effects including an interaction term,
followed by Tukey's honestly significant difference (HSD) test. There was a lack of interaction
between experimental trial and mite strain, so data across trials were combined for increased
statistical power (Brady et al., 2015). For graphical representation of the expression of plant
marker genes, the NRQ data were further normalized to the 'no mite' control sample, which was
therefore set to one. Thus, these results represent fold change differences in the expression of
marker genes relative to 'no mite' control.

Metabolic Analysis

Plant material used for the RNA isolation toward the expression analysis of the plant marker genes was utilized for the quantification of plant metabolites (JA, JA-Ile and I3M) as described in Zhurov et al., 2014. Differences in metabolite levels were detected by a two-way ANOVA, using mite strain and experimental trial as main effects including an interaction term, followed by Tukey's honestly significant difference (HSD) test. There was a lack of interaction between experimental trial and mite strain, so data across trials were combined for increased statistical power (Brady et al., 2015).

Determination of cytochrome P450 monooxygenase (P450), esterase and glutathione-S-transferase (GST) activity

To perform the enzymatic activity assays, 3-day-old spider mite females of each Col-a, *cyp*-a and bean-a strains, reared for two generations on bean plants, were placed on bean, *cyp79b2 cyp79b3* or Col-0 plants. After 24 h, 200 females from each treatment were collected and immediately frozen, to be used for protein extraction and enzymatic assays. Mites were homogenized in 100 mM phosphate buffer, pH 7.6. The concentration of total protein mite extracts was determined using the Quick Start™ Bradford Protein Assay (Quick start Bradford 1x dye reagent, Bio-Rad, Cat# 500-0205), with Bovine Serum Albumin (Sigma-Aldrich, Cat # A7906) as the standard. Twenty, 10 and 10 µg of protein per reaction were used for measuring enzymatic activity of glutathione-S-transferases (GSTs), P450 monooxygenases, and esterases, respectively. The enzymatic activities were assessed spectrophotometrically using 1-chloro-2,4-dinitrobenzene (CDNB) (for GSTs) (Habig and Jakoby, 1981), 7-ethoxy-4-trifluoromethylcoumarin (7-EFC) (for P450s) (Buters et al., 1993), and 4-nitrophenyl acetate (pNPA) (for esterases) (Park et al., 1961), as substrates. Enzymatic activities were determined in four independent samples with three technical replicates per sample. Differences in enzymatic activity between mite strains on different plant hosts were detected using a two-way ANOVA, with mite strain and plant host as main effects including an interaction term, followed by Tukey's honestly significant difference (HSD) test.

Application of enzyme inhibitors

Diethyl maleate (DEM, an inhibitor of glutathione S-transferase *(GST)* activity), S,S,S tributyl-phosphorotrithioate (DEF, an inhibitor of esterase activity), and piperonyl butoxide (PBO) and trichlorophenylpropynyl ether (TCPPE) (inhibitors of P450 activities) were dissolved in acetone (1:1 by volume) and were subsequently diluted with distilled water to reach desired concentrations. A range of concentrations of DEM (100, 200, 500, 1000 and 2000 mg/L), DEF (10, 20, 100, 200 and 500 mg/L), PBO (30, 100, 500, 1000 and 2000 mg/L) and TCPPE (1500 and 2000 mg/L) were tested to identify highest sublethal inhibitor concentration (mortality below 10%). Briefly, a batch of five detached bean leaves was sprayed either with a solution containing an inhibitor or a mock solution to the point of run-off. Fifty 1-2-day-old adult bean-a female mites were transferred to the upper (adaxial) side of a treated bean leaf that was placed on wet cotton wool. After 24 h, the number of surviving mites was recorded and mortality was calculated using Abbott's formula (Abbott, 1925).

For inhibitor bioassays, detached leaves of Col-0 *Arabidopsis* plants were dipped in a solution containing one of the inhibitors (DEF, DEM, PBO, TCPPE) or water and acetone only as control. Upon drying, leaves were infested with 1-2-day-old adult female mites. For fecundity assays, one adult bean-a or Col-a female was placed on the leaf and the number of eggs laid by each female per day was recorded for six days. Treated Col-0 leaves were replaced every other day.

Fecundity was determined in three independent experimental trials with ten replicates per trial.
For enzymatic assays, each Col-0 leaf was infested with ten 1 to 2-day-old adults Col-a female
mites. After 24 h, a pool of 100 spider mites was used to measure GST, esterase and P450
activities in DEM-, DEF-, PBO- and TCPPE-treated spider mites, respectively, using the protocol
described above. Enzymatic activities were determined in five independent samples with three
technical replicates per sample. Differences between means of fecundity and enzymatic
activities of control and inhibitor-treated samples were calculated using unpaired Student's t
tests (following tests for equal variance).
574
575   *In situ* **hybridization**
576   DIG-labelled probes were produced and the whole mount *in situ* hybridization was performed
577   according to previously published methods (Dearden et al., 2002). Images were collected using a
578   Zeiss AxioCam HRc 412-312 camera mounted on a Zeiss Axioplan II microscope.
579
RNAi of *Tu-CPR*
Two non-overlapping fragments (*Tu-CPR*, 645 nt) and (*Tu-CPR-F1*, 564 nt), complementary to the
coding region of *Tu-CPR (tetur18g03390*) and dsRNA complementary to a 382 bp non-
transcribed intergenic fragment spanning the region 1690614±1690995 of the genomic scaffold
12 (a negative control dsRNA, referred to as NC) were synthesized using primers listed in
Supplementary Table 1 according to previously described protocol (Suzuki et al., 2017). A BLAST
search against the *T. urticae* genome confirmed that dsRNA sequences are unique. dsRNA
solutions at concentrations of 500 ng/µL were supplemented with 6% (v/v) blue dye
erioglaucine (McCormick, Sparks Glencoe, MD). Newly molted adult female mites were soaked
in dsRNA/dye solutions at 20°C for 24 h (Suzuki et al., 2017). Post soaking, mites with visible blue
dye in posterior midgut were selected and were transferred in batches of 10 to either detached
Col-0 leaves or bean leaf disks. Fecundity was determined in three independent experimental
trials with ten replicates/trial as number of eggs deposited by individual female mite on 3[rd] and
4[th] day post soaking. Mites used for the analysis of *CPR* expression levels were further selected
on the basis of visual phenotype (spotless). The RT-qPCR was performed in eight (for dsRNA-*CPR*)
and three (for dsRNA-*CPR-1*) experimental replicas, as described above. P450 activity in *CPR*
silenced mites was determined in five independent samples with three technical replicates per
sample, as described above. Differences in fecundity between dsRNA treated mites was
detected using a two-way ANOVA, with mite treatment and experimental trial as main effects
including an interaction term, followed by Tukey's honestly significant difference (HSD) test.
There was a lack of interaction between experimental trial and mite treatment, so data across
trials were combined for increased statistical power (Brady et al., 2015). Enzyme activity and RT-
qPCR expression levels were compared between dsRNA treated mites using unpaired Student's t
tests (following tests for equal variance).
604

Acknowledgments
The authors would like to thank Drs. Thomas Van Leeuwen and Rene Feyereisen for their
constructive comments on pharmacological experiments, and Pierre Hilson for critical
comments on the manuscript. This work was supported by the Government of Canada through
the Ontario Research Fund (RE08-067) and the Natural Sciences and Engineering Research
Council of Canada (NSERC) awarded to MG and VG.
611

References

Abbott, W.S. (1925). A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18**,** 265-267. DOI: 10.1093/jee/18.2.265a.

Agrawal, A.A., Vala, F., and Sabelis, M.W. (2002). Induction of preference and performance after acclimation to novel hosts in a phytophagous spider mite: adaptive plasticity? *American Naturalist* 159(5)**,** 553-565. DOI: 10.1086/339463.

Aubert, Y., Widemann, E., Miesch, L., Pinot, F., and Heitz, T. (2015). CYP94-mediated jasmonoyl-isoleucine hormone oxidation shapes jasmonate profiles and attenuates defence responses to *Botrytis cinerea* infection. *Journal of Experimental Botany* 66(13)**,** 3879-3892. DOI: 10.1093/jxb/erv190.

Barrett, L.G., and Heil, M. (2012). Unifying concepts and mechanisms in the specificity of plant-enemy interactions. *Trends in Plant Science* 17(5)**,** 282-292. DOI: 10.1016/j.tplants.2012.02.009.

Bass, C., Zimmer, C.T., Riveron, J.M., Wilding, C.S., Wondji, C.S., Kaussmann, M., et al. (2013). Gene amplification and microsatellite polymorphism underlie a recent insect host shift. *Proceedings of the National Academy of Sciences* 110(48)**,** 19460-19465. DOI: 10.1073/pnas.1314122110.

Basu, S., Varsani, S., and Louis, J. (2018). Altering plant defenses: Herbivore-associated molecular patterns and effector arsenal of chewing herbivores. *Molecular Plant-Microbe Interactions* 31(1)**,** 13-21. DOI: 10.1094/mpmi-07-17-0183-fi.

Bednarek, P. (2012). Chemical warfare or modulators of defence responses - the function of secondary metabolites in plant immunity. *Current Opinion in Plant Biology* 15(4)**,** 407-414. DOI: 10.1016/j.pbi.2012.03.002.

Bensoussan, N., Zhurov, V., Yamakawa, S., O'Neil, C.H., Suzuki, T., Grbic, M., et al. (2018). The digestive system of the two-spotted spider mite, *Tetranychus urticae* Koch, in the context of the mite-plant interaction. *Frontiers in Plant Science* 9**,** 1206. DOI: 10.3389/fpls.2018.01206.

Beran, F., Sporer, T., Paetz, C., Ahn, S.J., Betzin, F., Kunert, G., et al. (2018). One pathway is not enough: The cabbage stem flea beetle *Psylliodes chrysocephala* uses multiple strategies to overcome the glucosinolate-myrosinase defense in its host plants. *Frontiers in Plant Science* 9**,** 1754. DOI: 10.3389/fpls.2018.01754.

Blaazer, C.J.H., Villacis-Perez, E.A., Chafi, R., Van Leeuwen, T., Kant, M.R., and Schimmel, B.C.J. (2018). Why do herbivorous mites suppress plant defenses? *Frontiers in Plant Science* 9**,** 1057. DOI: 10.3389/fpls.2018.01057.

Brady, S.M., Burow, M., Busch, W., Carlborg, O., Denby, K.J., Glazebrook, J., et al. (2015). Reassess the *t* test: Interact with all your data via ANOVA. *Plant Cell* 27(8)**,** 2088-2094. DOI: 10.1105/tpc.15.00238.

Buters, J.T., Schiller, C.D., and Chou, R.C. (1993). A highly sensitive tool for the assay of cytochrome P450 enzyme activity in rat, dog and man. Direct fluorescence monitoring of the deethylation of 7-ethoxy-4-trifluoromethylcoumarin. *Biochemical Pharmacology* 46(9)**,** 1577-1584. DOI: 10.1016/0006-2952(93)90326-r.

Dearden, P.K., Donly, C., and Grbic, M. (2002). Expression of pair-rule gene homologues in a chelicerate: early patterning of the two-spotted spider mite *Tetranychus urticae*. *Development* 129(23)**,** 5461-5472. DOI: 10.1242/dev.00099.

DeLucia, E.H., Nabity, P.D., Zavala, J.A., and Berenbaum, M.R. (2012). Climate change: Resetting plant-insect interactions. *Plant Physiology* 160(4)**,** 1677-1685. DOI: 10.1104/pp.112.204750.

Dermauw, W., Wybouw, N., Rombauts, S., Menten, B., Vontas, J., Grbic, M., et al. (2013). A link
between host plant adaptation and pesticide resistance in the polyphagous spider mite
Tetranychus urticae. *Proceedings of the National Academy of Sciences* 110(2)**,** E113-122.
DOI: 10.1073/pnas.1213214110.
Despres, L., David, J.P., and Gallet, C. (2007). The evolutionary ecology of insect resistance to
plant chemicals. *Trends in Ecology & Evolution* 22(6)**,** 298-307. DOI:
10.1016/j.tree.2007.02.010.
Devoto, A., Ellis, C., Magusin, A., Chang, H.S., Chilcott, C., Zhu, T., et al. (2005). Expression
profiling reveals COI1 to be a key regulator of genes involved in wound- and methyl
jasmonate-induced secondary metabolism, defence, and hormone interactions. *Plant*
Molecular Biology 58(4)**,** 497-513. DOI: 10.1007/s11103-005-7306-5.
Dobler, S., Dalla, S., Wagschal, V., and Agrawal, A.A. (2012). Community-wide convergent
evolution in insect adaptation to toxic cardenolides by substitutions in the Na,K-ATPase.
Proceedings of the National Academy of Sciences 109(32)**,** 13040-13045. DOI:
10.1073/pnas.1202111109.
Elbaz, M., Halon, E., Malka, O., Malitsky, S., Blum, E., Aharoni, A., et al. (2012). Asymmetric
adaptation to indolic and aliphatic glucosinolates in the B and Q sibling species of
Bemisia tabaci (Hemiptera: Aleyrodidae). *Molecular Ecology* 21(18)**,** 4533-4546. DOI:
10.1111/j.1365-294X.2012.05713.x.
Fernandez-Calvo, P., Chini, A., Fernandez-Barbero, G., Chico, J.M., Gimenez-Ibanez, S., Geerinck,
J., et al. (2011). The *Arabidopsis* bHLH transcription factors MYC3 and MYC4 are targets
of JAZ repressors and act additively with MYC2 in the activation of jasmonate responses.
Plant Cell 23(2)**,** 701-715. DOI: 10.1105/tpc.110.080788.
Feyereisen, R. (2014). Insect P450 inhibitors and insecticides: challenges and opportunities. *Pest*
Management Science 71(6)**,** 793-800. DOI: 10.1002/ps.3895.
Fry, J.D. (1989). Evolutionary adaptation to host plants in a laboratory population of the
phytophagous mite *Tetranychus urticae* Koch. *Oecologia* 81(4)**,** 559-565. DOI:
10.1007/bf00378969.
Gloss, A.D., Vassao, D.G., Hailey, A.L., Dittrich, A.C.N., Schramm, K., Reichelt, M., et al. (2014).
Evolution in an ancient detoxification pathway is coupled with a transition to herbivory
in the Drosophilidae. *Molecular Biology and Evolution* 31(9)**,** 2441-2456. DOI:
10.1093/molbev/msu201.
Gould, F. (1979). Rapid host range evolution in a population of the phytophagous mite
Tetranychus urticae Koch. *Evolution* 33(3)**,** 791-802. DOI: 10.1111/j.1558-
5646.1979.tb04735.x.
Govind, G., Mittapalli, O., Griebel, T., Allmann, S., Böcker, S., and Baldwin, I.T. (2010). Unbiased
transcriptional comparisons of generalist and specialist herbivores feeding on
progressively defenseless *Nicotiana attenuata* plants. *Plos One* 5(1)**,** e8735. DOI:
10.1371/journal.pone.0008735.
Grbic, M., Van Leeuwen, T., Clark, R.M., Rombauts, S., Rouze, P., Grbic, V., et al. (2011). The
genome of *Tetranychus urticae* reveals herbivorous pest adaptations. *Nature* 479(7374)**,**
487-492. DOI: 10.1038/nature10640.
Habig, W.H., and Jakoby, W.B. (1981). Glutathione S-transferases (rat and human). *Methods in*
Enzymology 77**,** 218-231. DOI: 10.1016/s0076-6879(81)77029-0.
Halon, E., Eakteiman, G., Moshitzky, P., Elbaz, M., Alon, M., Pavlidi, N., et al. (2015). Only a
minority of broad-range detoxification genes respond to a variety of phytotoxins in
generalist *Bemisia tabaci* species. *Scientific Reports* 5**,** 17975. DOI: 10.1038/srep17975.

Heidel-Fischer, H.M., Kirsch, R., Reichelt, M., Ahn, S.-J., Wielsch, N., Baxter, S.W., et al. (2019). An insect counteradaptation against host plant defenses evolved through concerted neofunctionalization. *Molecular Biology and Evolution* 36(5)**,** 930-941. DOI: 10.1093/molbev/msz019.

Heidel-Fischer, H.M., and Vogel, H. (2015). Molecular mechanisms of insect adaptation to plant secondary compounds. *Current Opinion in Insect Science* 8**,** 8-14. DOI: 10.1016/j.cois.2015.02.004.

Heitz, T., Widemann, E., Lugan, R., Miesch, L., Ullmann, P., Desaubry, L., et al. (2012). Cytochromes P450 CYP94C1 and CYP94B3 catalyze two successive oxidation steps of plant pormone jasmonoyl-isoleucine for catabolic turnover. *Journal of Biological Chemistry* 287(9)**,** 6296-6306. DOI: 10.1074/jbc.M111.316364.

Helmus, M.R., and Dussourd, D.E. (2005). Glues or poisons: which triggers vein cutting by monarch caterpillars? *Chemoecology* 15(1)**,** 45-49. DOI: 10.1007/s00049-005-0291-y.

Hogenhout, S.A., and Bos, J.I. (2011). Effector proteins that modulate plant-insect interactions. *Current Opinion in Plant Biology* 14(4)**,** 422-428. DOI: 10.1016/j.pbi.2011.05.003.

Hull, A.K., Vij, R., and Celenza, J.L. (2000). *Arabidopsis* cytochrome P450s that catalyze the first step of tryptophan-dependent indole-3-acetic acid biosynthesis. *Proceedings of the National Academy of Sciences* 97(5)**,** 2379-2384. DOI: 10.1073/pnas.040569997.

Jeschke, V., Gershenzon, J., and Vassão, D.G. (2016). A mode of action of glucosinolate-derived isothiocyanates: Detoxification depletes glutathione and cysteine levels with ramifications on protein metabolism in *Spodoptera littoralis*. *Insect Biochemistry and Molecular Biology* 71**,** 37-48. DOI: 10.1016/j.ibmb.2016.02.002.

Jonckheere, W., Dermauw, W., Khalighi, M., Pavlidi, N., Reubens, W., Baggerman, G., et al. (2018). A gene family coding for salivary proteins (SHOT) of the polyphagous spider mite *Tetranychus urticae* exhibits fast host-dependent transcriptional plasticity. *Molecular Plant-Microbe Interactions* 31(1)**,** 112-124. DOI: 10.1094/mpmi-06-17-0139-r.

Jonckheere, W., Dermauw, W., Zhurov, V., Wybouw, N., Van den Bulcke, J., Villarroel, C.A., et al. (2016). The salivary protein repertoire of the polyphagous spider mite *Tetranychus urticae*: a quest for effectors. *Molecular & Cellular Proteomics* 15(12)**,** 3594-3613. DOI: 10.1074/mcp.M116.058081.

Kaloshian, I., and Walling, L.L. (2016). Hemipteran and dipteran pests: Effectors and plant host immune regulators. *Journal of Integrative Plant Biology* 58(4)**,** 350-361. DOI: 10.1111/jipb.12438.

Kant, M.R., Sabelis, M.W., Haring, M.A., and Schuurink, R.C. (2008). Intraspecific variation in a generalist herbivore accounts for differential induction and impact of host plant defences. *Proceedings of the Royal Society B-Biological Sciences* 275(1633)**,** 443-452. DOI: 10.1098/rspb.2007.1277.

Kim, J.H., and Jander, G. (2007). *Myzus persicae* (green peach aphid) feeding on Arabidopsis induces the formation of a deterrent indole glucosinolate. *Plant Journal* 49(6)**,** 1008-1019. DOI: 10.1111/j.1365-313X.2006.03019.x.

Kim, J.H., Lee, B.W., Schroeder, F.C., and Jander, G. (2008). Identification of indole glucosinolate breakdown products with antifeedant effects on *Myzus persicae* (green peach aphid). *Plant Journal* 54(6)**,** 1015-1026. DOI: 10.1111/j.1365-313X.2008.03476.x.

Li, W., Schuler, M.A., and Berenbaum, M.R. (2003). Diversification of furanocoumarin-metabolizing cytochrome P450 monooxygenases in two papilionids: Specificity and substrate encounter rate. *Proceedings of the National Academy of Sciences* 100**,** 14593-14598. DOI: 10.1073/pnas.1934643100.

Li, X., Baudry, J., Berenbaum, M.R., and Schuler, M.A. (2004). Structural and functional divergence of insect CYP6B proteins: From specialist to generalist cytochrome P450. *Proceedings of the National Academy of Sciences* 101(9)**,** 2939-2944. DOI: 10.1073/pnas.0308691101.

Li, X., Schuler, M.A., and Berenbaum, M.R. (2007). Molecular mechanisms of metabolic resistance to synthetic and natural xenobiotics. *Annual Review of Entomology* 52**,** 231-253. DOI: 10.1146/annurev.ento.51.110104.151104.

Magalhaes, S., Fayard, J., Janssen, A., Carbonell, D., and Olivieri, I. (2007). Adaptation in a spider mite population after long-term evolution on a single host plant. *Journal of Evolutionary Biology* 20(5)**,** 2016-2027. DOI: 10.1111/j.1420-9101.2007.01365.x.

Mao, W., Rupasinghe, S., Zangerl, A.R., Schuler, M.A., and Berenbaum, M.R. (2006). Remarkable substrate-specificity of CYP6AB3 in *Depressaria pastinacella*, a highly specialized caterpillar. *Insect Molecular Biology* 15(2)**,** 169-179. DOI: 10.1111/j.1365-2583.2006.00623.x.

Masters, B.S.S., and Okita, R.T. (1980). The history, properties, and function of NADPH-cytochrome P-450 reductase. *Pharmacology & Therapeutics* 9(2)**,** 227-244. DOI: 10.1016/S0163-7258(80)80020-9.

Mathers, T.C., Chen, Y., Kaithakottil, G., Legeai, F., Mugford, S.T., Baa-Puyoulet, P., et al. (2017). Rapid transcriptional plasticity of duplicated gene clusters enables a clonally reproducing aphid to colonise diverse plant species. *Genome Biology* 18(1)**,** 27. DOI: 10.1186/s13059-016-1145-3.

Migeon, A., and Dorkeld, F. (2006-2019). *Spider Mites Web: a comprehensive database for the Tetranychidae.* [Online]. Available: http://www.montpellier.inra.fr/CBGP/spmweb [Accessed August 1 2019].

Mikkelsen, M.D., Hansen, C.H., Wittstock, U., and Halkier, B.A. (2000). Cytochrome P450 CYP79B2 from *Arabidopsis* catalyzes the conversion of tryptophan to indole-3-acetaldoxime, a precursor of indole glucosinolates and indole-3-acetic acid. *Journal of Biological Chemistry* 275(43)**,** 33712-33717. DOI: 10.1074/jbc.M001667200.

Muller, C., Vogel, H., and Heckel, D.G. (2017). Transcriptional responses to short-term and long-term host plant experience and parasite load in an oligophagous beetle. *Molecular Ecology* 26(22)**,** 6370-6383. DOI: 10.1111/mec.14349.

Musser, R.O., Hum-Musser, S.M., Eichenseer, H., Peiffer, M., Ervin, G., Murphy, J.B., et al. (2002). Herbivory: caterpillar saliva beats plant defences. *Nature* 416(6881)**,** 599-600. DOI: 10.1038/416599a.

Musser, R.O., Hum-Musser, S.M., Lee, H.K., DesRochers, B.L., Williams, S.A., and Vogel, H. (2012). Caterpillar labial saliva alters tomato plant gene expression. *Journal of Chemical Ecology* 38(11)**,** 1387-1401. DOI: 10.1007/s10886-012-0198-3.

Park, J.H., Meriwether, B.P., Clodfelder, P., and Cunningham, L.W. (1961). The hydrolysis of p-nitrophenyl acetate catalyzed by 3-phosphoglyceraldehyde dehydrogenase. *Journal of Biological Chemistry* 236**,** 136-141.

Rajniak, J., Barco, B., Clay, N.K., and Sattely, E.S. (2015). A new cyanogenic metabolite in *Arabidopsis* required for inducible pathogen defence. *Nature* 525(7569)**,** 376-379. DOI: 10.1038/nature14907.

Ratzka, A., Vogel, H., Kliebenstein, D.J., Mitchell-Olds, T., and Kroymann, J. (2002). Disarming the mustard oil bomb. *Proceedings of the National Academy of Sciences* 99(17)**,** 11223-11228. DOI: 10.1073/pnas.172112899.

Rieu, I., and Powers, S.J. (2009). Real-Time Quantitative RT-PCR: Design, calculations, and statistics. *Plant Cell* 21(4)**,** 1031-1033. DOI: 10.1105/tpc.109.066001.

Sasabe, M., Wen, Z., Berenbaum, M.R., and Schuler, M.A. (2004). Molecular analysis of
CYP321A1, a novel cytochrome P450 involved in metabolism of plant allelochemicals
(furanocoumarins) and insecticides (cypermethrin) in *Helicoverpa zea*. *Gene* 338(2), 163-
175. DOI: 10.1016/j.gene.2004.04.028.
Schramm, K., Vassao, D.G., Reichelt, M., Gershenzon, J., and Wittstock, U. (2012). Metabolism of
glucosinolate-derived isothiocyanates to glutathione conjugates in generalist
lepidopteran herbivores. *Insect Biochemistry and Molecular Biology* 42(3), 174-182. DOI:
10.1016/j.ibmb.2011.12.002.
Schuler, M.A. (2011). P450s in plant–insect interactions. *Biochimica et Biophysica Acta* 1814(1),
36-45. DOI: 10.1016/j.bbapap.2010.09.012.
Schweizer, F., Fernandez-Calvo, P., Zander, M., Diez-Diaz, M., Fonseca, S., Glauser, G., et al.
(2013). *Arabidopsis* basic Helix-Loop-Helix transcription factors MYC2, MYC3, and MYC4
regulate glucosinolate biosynthesis, insect performance, and feeding behavior. *Plant Cell*
25(8), 3117-3132. DOI: 10.1105/tpc.113.115139.
Schweizer, F., Heidel-Fischer, H., Vogel, H., and Reymond, P. (2017). *Arabidopsis* glucosinolates
trigger a contrasting transcriptomic response in a generalist and a specialist herbivore.
Insect Biochemistry and Molecular Biology 85, 21-31. DOI: 10.1016/j.ibmb.2017.04.004.
Shi, Y., Wang, H., Liu, Z., Wu, S., Yang, Y., Feyereisen, R., et al. (2018). Phylogenetic and
functional characterization of ten P450 genes from the CYP6AE subfamily of *Helicoverpa
armigera* involved in xenobiotic metabolism. *Insect Biochemistry and Molecular Biology*
93, 79-91. DOI: 10.1016/j.ibmb.2017.12.006.
Snoeck, S., Pavlidi, N., Pipini, D., Vontas, J., Dermauw, W., and Van Leeuwen, T. (2019). Substrate
specificity and promiscuity of horizontally transferred UDP-glycosyltransferases in the
generalist herbivore *Tetranychus urticae*. *Insect Biochemistry and Molecular Biology* 109,
116-127. DOI: 10.1016/j.ibmb.2019.04.010.
Stireman, J.O., and Singer, M.S. (2018). Tritrophic niches of insect herbivores in an era of rapid
environmental change. *Current Opinion in Insect Science* 29, 117-125. DOI:
10.1016/j.cois.2018.07.008.
Suzuki, T., Nunes, M.A., Espana, M.U., Namin, H.H., Jin, P., Bensoussan, N., et al. (2017). RNAi-
based reverse genetics in the chelicerate model *Tetranychus urticae*: A comparative
analysis of five methods for gene silencing. *Plos One* 12(7), e0180654. DOI:
10.1371/journal.pone.0180654.
Van Leeuwen, T., and Dermauw, W. (2016). The molecular evolution of xenobiotic metabolism
and resistance in chelicerate mites. *Annual Review of Entomology* 61, 475-498. DOI:
10.1146/annurev-ento-010715-023907.
Villarroel, C.A., Jonckheere, W., Alba, J.M., Glas, J.J., Dermauw, W., Haring, M.A., et al. (2016).
Salivary proteins of spider mites suppress defenses in *Nicotiana benthamiana* and
promote mite reproduction. *Plant Journal* 86(2), 119-131. DOI: 10.1111/tpj.13152.
Wadleigh, R.W., and Yu, S.J. (1988). Detoxification of isothiocyanate allelochemicals by
glutathione transferase in three lepidopterous species. *Journal of Chemical Ecology*
14(4), 1279-1288. DOI: 10.1007/bf01019352.
Whiteman, N.K., Gloss, A.D., Sackton, T.B., Groen, S.C., Humphrey, P.T., Lapoint, R.T., et al.
(2012). Genes involved in the evolution of herbivory by a leaf-mining, Drosophilid fly.
Genome Biology and Evolution 4(9), 900-916. DOI: 10.1093/gbe/evs063.
Whiteman, N.K., Groen, S.C., Chevasco, D., Bear, A., Beckwith, N., Gregory, T.R., et al. (2011).
Mining the plant-herbivore interface with a leafmining *Drosophila* of *Arabidopsis*.
Molecular Ecology 20(5), 995-1014. DOI: 10.1111/j.1365-294X.2010.04901.x.

Widemann, E., Grausem, B., Renault, H., Pineau, E., Heinrich, C., Lugan, R., et al. (2015).
Sequential oxidation of Jasmonoyl-Phenylalanine and Jasmonoyl-Isoleucine by multiple
cytochrome P450 of the CYP94 family through newly identified aldehyde intermediates.
Phytochemistry 117**,** 388-399. DOI: 10.1016/j.phytochem.2015.06.027.
Wittstock, U., Agerbirk, N., Stauber, E.J., Olsen, C.E., Hippler, M., Mitchell-Olds, T., et al. (2004).
Successful herbivore attack due to metabolic diversion of a plant chemical defense.
Proceedings of the National Academy of Sciences 101(14)**,** 4859-4864. DOI:
10.1073/pnas.0308007101.
Wittstock, U., and Burow, M. (2010). Glucosinolate breakdown in *Arabidopsis*: Mechanism,
regulation and biological significance. *The Arabidopsis Book* 2010(8). DOI:
10.1199/tab.0134.
Wu, S., Peiffer, M., Luthe, D.S., and Felton, G.W. (2012). ATP hydrolyzing salivary enzymes of
caterpillars suppress plant defenses. *Plos One* 7(7)**,** e41947. DOI:
10.1371/journal.pone.0041947.
Wybouw, N., Balabanidou, V., Ballhorn, D.J., Dermauw, W., Grbic, M., Vontas, J., et al. (2012). A
horizontally transferred cyanase gene in the spider mite *Tetranychus urticae* is involved
in cyanate metabolism and is differentially expressed upon host plant change. *Insect*
Biochemistry and Molecular Biology 42(12)**,** 881-889. DOI: 10.1016/j.ibmb.2012.08.002.
Wybouw, N., Dermauw, W., Tirry, L., Stevens, C., Grbic, M., Feyereisen, R., et al. (2014). A gene
horizontally transferred from bacteria protects arthropods from host plant cyanide
poisoning. *Elife* 3**,** e02365. DOI: 10.7554/eLife.02365.
Wybouw, N., Zhurov, V., Martel, C., Bruinsma, K.A., Hendrickx, F., Grbic, V., et al. (2015).
Adaptation of a polyphagous herbivore to a novel host plant extensively shapes the
transcriptome of herbivore and host. *Molecular Ecology* 24(18)**,** 4647-4663. DOI:
10.1111/mec.13330.
Zarate, S.I., Kempema, L.A., and Walling, L.L. (2007). Silverleaf whitefly induces salicylic acid
defenses and suppresses effectual jasmonic acid defenses. *Plant Physiology* 143(2)**,** 866-
875. DOI: 10.1104/pp.106.090035.
Zhao, C., Escalante, L.N., Chen, H., Benatti, T.R., Qu, J., Chellapilla, S., et al. (2015). A massive
expansion of effector genes underlies gall-formation in the wheat pest *Mayetiola*
destructor. *Current Biology* 25(5)**,** 613-620. DOI: 10.1016/j.cub.2014.12.057.
Zhurov, V., Navarro, M., Bruinsma, K.A., Arbona, V., Santamaria, M.E., Cazaux, M., et al. (2014).
Reciprocal responses in the interaction between *Arabidopsis* and the cell-content-
feeding chelicerate herbivore spider mite. *Plant Physiology* 164(1)**,** 384-399. DOI:
10.1104/pp.113.231555.
883
884

 Figure Legends
Figure 1. *Arabidopsis* induced responses to feeding of the two-spotted spider mite (TSSM)
Tetranychus urticae. (A) The fecundity of bean-reared TSSM (London strain; ancestral population)
upon feeding on +/- MeJA treated Col-0 plants (left) and on Col-0 wild type, *cyp79b2 cyp79b3*
mutant plants lacking Trp-derived secondary metabolites, and *myc2 myc3 myc4* mutant plants
(myc2,3,4) that lack JA-dependent induced responses (right). For fecundity on +/- MeJA treated
Col-0 leaves, the number of eggs was counted over 24 h and is presented as mean number of
eggs laid by a female mite per day ± SEM; experiment was replicated in nine biological
replicates/trial in two independent trials. Asterisk indicates a significant difference between
treated and control samples (ANOVA: ***$P<0.001$). For fecundity on different genotypes (right),
the number of eggs was counted over two days and is presented as mean number of eggs laid
by a female mite per day ± SEM; experiment was replicated in five biological replicates/trial in
three independent trials. Different letters represent significant difference between means
(Tukey's HSD test, α = 0.05). (B) A simplified schematic of *Arabidopsis* induced responses to
TSSM feeding. Jasmonic acid (JA), its bioactive conjugate jasmonoyl-isoleucine (JA-Ile), and
MYC2, MYC3 and MYC4 (MYC2,3,4) transcriptional activators transduce the recognition of TSSM
herbivory and induce the expression of JA-regulated genes such as *ALLENE OXIDE SYNTHASE*
(*AOS), CYP79B2* and *CYP79B3. AOS* is a JA biosynthetic gene, while *CYP79B2* and *CYP79B3*
encode enzymes required for the biosynthesis of Trp-derived secondary metabolites.
904
Figure 2. Experimental evolution of *Tetranychus urticae* adaptation to *Arabidopsis*. (A) A
schematic of the experimental evolution procedure. The ancestral population was reared on
beans. Triplicated ancestral populations were selected on *cyp79b2 cyp79b3* and Col-0 plants for
at least 25 generations. (B) Performance of ancestral and selected populations on bean (top),
cyp79b2 cyp79b3 (middle) and Col-0 (bottom) plants. Mites from *Arabidopsis*-selected
populations were transferred from their respective rearing hosts, *cyp79b2 cyp79b3* and Col-0
plants, to beans for two generations prior to testing their performance (an indirect transfer
regime). The performance was measured as the size of total population derived from twenty
adult female mites and seven days post infestation. Data are represented as mean ± SEM, n=4.
Statistics were performed on total population counts. See also Figure S1. (C) Fecundity of the
ancestral (bean-a) and populations #3 of *Arabidopsis*-selected mites (*cyp*-a and Col-a) upon
feeding on Col-0 and *cyp79b2 cyp79b3* plants. Fecundity was measured over six days and data
are presented as mean number of eggs laid by a female mite per day ± SEM; experiment was
replicated in ten biological replicates/trial in tree independent trials. (D) Leaf damage of Col-0
and *cyp79b2 cyp79b3 Arabidopsis* plants upon herbivory of ancestral (bean-a), and populations
#3 of *Arabidopsis*-selected mites (*cyp*-a and Col-a). Data are represented as mean ± SEM;
experiment was replicated in six biological replicates/trial in three independent trials. Different
letters represent significant differences between means (Tukey's HSD test, α = 0.05).
923
Figure 3. Induced *Arabidopsis* responses upon feeding of bean-a, *cyp*-a and Col-a mites for 24 h.
(A) Expression of *AOS, CYP79B2, CYP79B3* and *MYC2* genes in response to bean-a, *cyp*-a and Col-
a mite feeding. Shown are means ± SEM of FCs of expression levels detected by RT-qPCR;
experiment was replicated in three biological replicates/trial in three independent trials. Primer
sequences and amplification efficiencies (E) used in qPCR are shown in Supplemental Table 1.
PEROXIN4 (*AT5G25760*), was used as the reference gene. (B) Levels of JA and JA-Ile (ng g^{-1} fresh
weight), and relative level of the indol-3-ylmethyl glucosinolate (I3M, shown as normalized peak
area), in 3-week-old Col-0 plants after herbivory of bean-a, *cyp*-a and Col-a mites for 24 h.
Values are means ± SEM; experiment was replicated in three biological replicates/trial in two

independent trials. Different letters represent significant differences between means (Tukey's
HSD test, α = 0.05).

Figure 4. The effect of mite host-adaptation state and plant hosts on the expression levels of
candidate *cytochrome P450 (CYP)* and *UDP-glycosyltransferases (UGT)* mite genes. (A) *CYP392A1*
(*tetur07g06410*), *CYP392A16* (*tetur06g04520*) and *CYP392D8* (*tetur03g05070*). (B) *UGT201A2v2*
(*tetur02g02480*), *UGT204B1* (*tetur02g09830*) and *UGT204A5* (*tetur05g00090*). Shown are
means ± SEM for relative quantity of expression normalized to *RP49* (*tetur18g03590*) reference
gene, experiment was replicated in three biological replicates/trial; additional two independent
experimental trials had similar results. Primer sequences and amplification efficiencies (E) used
in qPCR are shown in Supplemental Table 1. Different letters represent significant differences
between means (Tukey's HSD test, α = 0.05).

Figure 5. The requirements of glutathione-S-transferase and cytochrome P450 activities for
Tetranychus urticae adaptation to *Arabidopsis*. (A, D) The activities of glutathione-S-transferases
(A) and cytochrome P450s (D) in bean-a, *cyp*-a and Col-a mites (reared on beans for two
generations) feeding on bean, *cyp79b2 cyp79b3* or Col-0 plants. Data are represented as mean ±
SEM, n=4. Different letters represent significant differences between means (Tukey's HSD test, α
= 0.05). (B, E, G) The activities of glutathione-S-transferases (B) and cytochrome P450s (E, G) in
Col-a mites feeding on Col-0 plants after the application of diethyl maleate (DEM, an inhibitor of
GST activity; in B), piperonyl butoxide (PBO, an inhibitor of P450 activity; in E) and
trichlorophenylpropynyl ether (TCPPE, inhibitor of P450 activity; in G). Data are represented as
mean ± SEM, n=5. (C, F) The effects of DEM (C) and PBO (F) treatments on fecundity of bean-a
and Col-a mites upon feeding on Col-0 and *cyp79b2 cyp79b3* plants. Fecundity was measured
over six days and data are presented as mean number of eggs laid by a female mite per day ±
SEM; experiment was replicated in ten biological replicates/trial in three independent trials. (H)
The effect of TCPPE treatment on fecundity of bean-a and Col-a mites upon feeding on Col-0,
cyp79b2 cyp79b3 or bean plants. Fecundity was measured over six days and data are presented
as mean number of eggs laid by a female mite per day ± SEM; experiment was replicated in ten
biological replicates/trial in three independent trials. Asterisks in panels B, C, E-H indicate a
significant difference between treated and control samples (unpaired Student's t test: *P <0.05,
**P<0.01, ***P<0.001). See also Figure S2.

Figure 6. The requirement of CPR and P450 activity for *Tetranychus urticae* adaptation to
Arabidopsis. (A) Whole-mount *in-situ* hybridization with anti-sense probe of *Tu-CPR* in *T. urticae*
(i) adult female, (ii) adult male, (iii) background control (hybridization with sense probe),
enlarged view of panel 'i' showing (iv) epithelium (v) ovaries and (vi and vii) digestive cells of
adult female. (B-D) The effects of dsRNA-*Tu-CPR*. (B) Relative level of *Tu-CPR* transcript
normalized with *RP49* in dsRNA treated Col-a mites (mean ± SEM, n=8, unpaired Student's t test
*** P<0.001). (C) The activity of cytochrome P450s in dsRNA treated Col-a mite upon feeding on
Col-0 plants (mean ± SEM, n=5, unpaired Student's t test *** P<0.001). (D) The fecundity of
dsRNA treated Col-a mites feeding on Col-0 or bean plants. Fecundity was measured over two
days (3 and 4 dpi) and data are presented as mean number of eggs laid by a female mite per day
± SEM; the experiment was replicated in ten (for Col-0 plants) and 4 (for bean plants) biological
replicates/trial in three independent trials. Data are represented as mean ± SEM (unpaired
Student's t test *** P<0.001). Scale bars in (F): (i-iii) 20μm; (iv-vii) 100μm. See also Figure S3.

Supplemental Data

Supplemental Table 1. Gene-specific primer sequences used for real-time quantitative RT-PCR.

Supplemental Figure 1. Experimental evolution of *Tetranychus urticae* adaptation to *Arabidopsis*. Performance of ancestral and selected populations on bean (top), *cyp79b2 cyp79b3* (middle) and Col-0 (bottom) plants. Mites were transferred directly from their respective rearing hosts to experimental bean, *cyp79b2 cyp79b3* and Col-0 plants. The performance was measured as the size of total population derived from twenty adult female mites and seven days post infestation. Data are represented as mean ± SEM, n=4. Statistics were performed on total population counts.

Supplemental Figure 2. The requirement of esterase activity for *Tetranychus urticae* adaptation to *Arabidopsis*. (A) The esterase activities in bean-a, *cyp*-a and Col-a mites (reared on beans for two generations) feeding on bean, *cyp79b2 cyp79b3* or Col-0 plants. Data are represented as mean ± SEM, n=4. Different letters represent significant differences between means (Tukey's HSD test, α = 0.05). (B) Esterase activity in Col-a mites feeding on Col-0 plants after the application of S,S,S tributyl-phosphorotrithioate (DEF, an inhibitor of esterase activity). Data are represented as mean ± SEM, n=5. (C) Effects of DEF treatment on fecundity of bean-a and Col-a mites upon feeding on Col-0 and *cyp79b2 cyp79b3* plants. Fecundity was assessed as mean number of eggs laid by a female mite in six days ± SEM; the experiment was replicated three times using 10 biological replicates/trial. Asterisks in panels B, C, and E indicate a significant difference between treated and control samples (unpaired Student's t test: *P <0.05, **P<0.01).

Supplemental Figure 3. RNAi silencing of *Tu-CPR* gene. (A) A schematic of the *Tu-CPR* locus. DNA sequences used for the generation of dsRNA-*Tu-CPR* are shown in red (fragment CPR, 645 bp), and green (fragment CPR-1, 564 bp). UTR and coding sequences are shown as light and dark blue boxes, respectively. (B) The effect of dsRNA-*Tu-CPR*. Relative expression of *Tu-CPR* normalized with *RP49* in dsRNA treated Col-a mites (mean ± SEM, n=8, Student's t test, ns). (C) The fecundity of dsRNA treated Col-a mites feeding on Col-0 and bean plants. Fecundity was measured over two days (3 and 4 dpi) and data are presented as mean number of eggs laid by a female mite per day ± SEM; the experiment was replicated in ten (for Col-0 plants) and 4 (for bean plants) biological replicates/trial in three independent trials. Data are represented as mean ± SEM (Student's t test *** P<0.001).

FIGURE 1

A

Fecundity
(eggs/mite/day)

2.5
2
1.5
1
0.5

***

MeJA: -   +
        Col-0

a   b   c

Col-0   cyp79b2,b3   myc2,3,4

B

TSSM induced *Arabidopsis* responses

Linolenic acid
AOS
JA
JA-Ile
MYC2,3,4

JA biosynthesis

JA signalling

JA regulated responses

Trp-dependent
defense compounds
(e.g. **I3M**)

CYP79B2,B3 Trp

other defense
responses

FIGURE 2

A

B

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

Supplemental Table 1. Gene-specific primer sequences used for the analysis of levels of gene expression (RT-qPCR), synthesis of dsRNA (RNAi) and preparation of probe for the *in situ* transcript localization.

Locus ID and name	Application	Primers	Efficiency
tetur07g06410 CYP392A1	RT-qPCR	F: 5'-GACGCCTTCGCAAATGATGC-3' R: 5'-TAACCGTCTGTGTTCACGCC-3'	0.929
tetur06g04520 CYP392A16	RT-qPCR	F: 5'-TTGATTGGGCTTGCCCTCTT-3' R: 5'-AGCCAACAATCGGAAGACCC-3'	0.903
tetur03g05070 CYP392D8	RT-qPCR	F: 5'-ACCAGAGAGATTCCTCAGCG-3' R: 5'-AAAGCCAAAGTTGCACCAGG-3'	1.030
tetur02g02480 UGT201A2v2	RT-qPCR	F: 5'-TCGAGAACTGAGTGGAGTAGC-3' R: 5'-TAGCAGGCAAAGGTGTTCCA-3'	1.090
tetur02g09830 UGT204B1	RT-qPCR	F: 5'-GCTTCGGTTGAGAAACGTGG-3' R: 5'-AAAATCGGCATTCGCTTCGG-3'	1.000
tetur05g00090 UGT204A5	RT-qPCR	F: 5'-TGGACGGAAATCGTAGTGGA-3' R: 5'-AGCTCATCAAAGACCAGCGA-3'	1.090
tetur18g03590 RP49	RT-qPCR	F: 5'-CTTCAAGCGGCATCAGAGC-3' R: 5'-CGCATCTGACCCTTGAACTTC-3'	0.976
AT5G42650 AOS	RT-qPCR	F: 5'-AAATCCAACGGCGGAGAACT-3' R: 5'-TCGTCGCCAACGGTTGATAA-3'	0.984
AT4G39950 CYP79B2	RT-qPCR	F: 5'-GAAAAGAGGTTGTGCGGCTC-3' R: 5'-TCTCACTTCACCGTCGGGTA-3'	0.994
AT2G22330 CYP79B3	RT-qPCR	F: 5'-TCTACCGATGCTTACGGGATTG-3' R: 5'-TACAAGTTCCTTAATGGTTGGTTTG-3'	0.973
AT1G32640 MYC2	RT-qPCR	F: 5'-TCGCTTACATCAACGAGCTTAAATC-3' F: 5'- TATCTTCACTTCAATCTCCATCCCC-3'	0.900
AT5G25760 PEROXIN4	RT-qPCR	F: 5'-GCTCTTATCAAAGGACCTTCGG-3' R: 5'-CGAACTTGAGGAGGTTGCAAAG-3'	0.992
tetur18g03390 Tu-CPR	RT-qPCR	F: 5'-CCATTCTTGGCACCTATCGT-3' R: 5'-GCAAGGTGATCTCCAGCTTC-3'	0.994
tetur18g03390 Tu-CPR	RNAi (fragment 1)	F: 5'-[T7]-CCTCGACTTCAGCCACGTTA-3' R: 5'-[T7]-AACATCCCGAGCCATGTTCC-3'	not applicable
tetur18g03390 Tu-CPR	RNAi (fragment 2)	F: 5'-[T7]-ACAAACCGGTACTGCAGAGG-3' R: 5'-[T7]-TGCATACACGAACGGTCTCC-3'	not applicable
T. urticae genomic scaffold 12, position 1690614-1690995, *NC*	RNAi (negative control)	F: 5'-GCCCTCTCCTGGTTGTAAACTT-3' R: 5'-CGACCCCATCAGGCTATTGA-3'	not applicable
tetur18g03390 Tu-CPR	*in situ* (antisense)	F: 5'-[T7]-ACAAACCGGTACTGCAGAGG-3' R: 5'-TGCATACACGAACGGTCTCC-3'	not applicable
tetur18g03390 Tu-CPR	*in situ* (sense)	F: 5'-ACAAACCGGTACTGCAGAGG-3' R: 5'-[T7]-TGCATACACGAACGGTCTCC-3'	not applicable

[T7]: T7 RNA Polymerase promoter sequence, 5'-TAATACGACTCACTATAGGG-3'

SUPPLEMENTAL FIGURE 1

SUPPLEMENTAL FIGURE 2

SUPPLEMENTAL FIGURE 3

A

Tu-CPR (tetur18g03390)

B

C