Colonization with the commensal fungus *Candida albicans* perturbs the gut-brain axis through

dysregulation of endocannabinoid signaling

Laura Markey[a], Andrew Hooper[b], Laverne C. Melon[b,c], Samantha Baglot[d], Matthew N. Hill[d], Jamie

Maguire[b] and Carol A. Kumamoto[a]*

[a]Department of Molecular Biology and Microbiology, Tufts University, 150 Harrison Ave, Boston, MA,

02111

[b]Department of Neuroscience, Tufts University, 150 Harrison Ave, Boston, MA, 02111

[c]Current affiliation- Department of Biology, Wesleyan University, Middletown, CT, 06459

[d]Hotchkiss Brain Institute, University of Calgary, Calgary AB, Canada, T2N1Z4

*Corresponding Author: Carol A. Kumamoto, Tufts University, Department of Molecular Biology and

Microbiology, 136 Harrison Ave., Boston, MA 02111 USA

Abstract

Anxiety disorders are the most prevalent mental health disorder worldwide, with a lifetime prevalence of 5-7% of the human population. Although the etiology of anxiety disorders is incompletely understood, one aspect of host health that affects anxiety disorders is the gut-brain axis. We used a mouse model of gastrointestinal (GI) colonization to demonstrate that the commensal fungus *Candida albicans* affects host health via the gut-brain axis. In mice, bacterial members of the gut microbiota can influence the host gut-brain axis, affecting anxiety-like behavior and the hypothalamus-pituitary-adrenal (HPA) axis which produces the stress hormone corticosterone (CORT). Here we showed that mice colonized with *C. albicans* demonstrated increased anxiety-like behavior and increased basal production of CORT as well as dysregulation of CORT production following acute stress. The HPA axis and anxiety-like behavior are negatively regulated by the endocannabinoid anandamide (AEA). *C. albicans*-colonized mice exhibited systemic changes in the endocannabinoidome, within the GI tract and the brain, and showed a negative correlation between brain AEA levels and serum CORT. Further, increasing AEA levels using the well-characterized fatty acid amide hydrolase (FAAH) inhibitor URB597 was sufficient to reverse both neuroendocrine phenotypes in *C. albicans*-colonized mice. Thus, a commensal fungus that is a common colonizer of humans had widespread effects on the physiology of its host. To our knowledge, this is the first report of microbial manipulation of the endocannabinoid (eCB) system that resulted in neuroendocrine changes contributing to anxiety-like behavior.

Keywords

Endocannabinoid, microbiota, *Candida albicans*, anxiety, corticosterone

1. Introduction

Anxiety disorders are the most prevalent mental health disorder worldwide, with a lifetime prevalence of 5-7% of the human population (Baxter et al., 2013). The etiology of anxiety disorders is complex and incompletely understood, although many different genetic and environmental factors have been identified (Leonardo and Hen, 2008). The gut-brain axis, the bi-directional communication between

the gut and the brain through multiple host systems, has been shown to affect anxiety disorders (Luna and

Foster, 2015). Constant communication between the periphery and the brain is required for normal

homeostasis and overall health.

One component of the gut-brain axis thought to play a role in anxiety disorders is the

hypothalamus-pituitary-adrenal (HPA) axis. The endocrine output of the HPA axis, the stress hormone

corticosterone (CORT), is a broad regulator of host health including the immune, neuroendocrine,

metabolic and cardiovascular systems. Under basal conditions, CORT release is regulated by the

circadian rhythm and helps to maintain homeostasis (Dallman et al., 1993). CORT is also a key part of

the acute psychological stress response and a rapid increase in circulating CORT is essential for mounting

an effective response to stress (Sapolsky et al., 2000). Dysregulation of the HPA axis and CORT

production is correlated with mental health disorders including anxiety and depression (Kallen et al.,

2008).

The gut microbiota, the diverse assemblage of microbes that colonizes the GI tract of humans, has

been shown to affect virtually every aspect of human health, including the gut-brain axis. Researchers

using germ-free mouse models demonstrated that the gut microbiota plays an important role in the

development and regulation of the HPA axis (Sudo et al., 2004). Studies have also shown that specific

probiotic bacterial species can affect the gut-brain axis, as treatment with *Bifidobacterium sp*. and

Lactobacillus sp. decreases anxiety-like behavior and normalizes HPA axis function in mice (Messaoudi

et al., 2011). Investigation into the contribution of the gut-microbiota-brain axis to mental health could

reveal novel regulators of anxiety-like behavior and offer mechanistic insight into the etiology of the

disease.

Candida albicans is a commensal fungus that colonizes the GI tract of ~60% of the human

population (Raimondi et al., 2019). Although a major colonizer of humans, its role as a commensal

microbe is largely uncharacterized. Previous researchers have shown that GI *C. albicans* colonization

induces local and systemic immune changes which are protective against systemic *C. albicans* infection

and disease (Shao et al., 2019), and against infection with bacterial pathogens such as *C. difficile* (Markey

et al., 2018) and *S. aureus* (Shao et al., 2019). The research that follows demonstrates that *C. albicans*

affects host health beyond the GI tract. We show that *C. albicans* dysregulated HPA axis activity and

increased anxiety-like behavior in a mouse model by altering endocannabinoid signaling.

The mammalian endocannabinoid (eCB) system regulates both the HPA axis and anxiety-like

behavior. It consists of two major neuroactive lipids, N-arachidonoylethanolamide (AEA) and 2-

arachidonoylglycerol (2-AG) and their receptors CB1 (distributed throughout the nervous system) and

CB2 (present primarily on immune cells) (Devane et al., 1992; Mechoulam et al., 1995). AEA and 2-AG

are produced in response to activation in post-synaptic neurons, and act at CB1 on pre-synaptic neurons to

limit neurotransmitter release. AEA and 2-AG levels are regulated by substrate availability and the

activity of their synthetic and degradative enzymes (Placzek et al., 2008). Using genetic and

pharmacological tools in rodents, researchers have demonstrated that manipulation of CB1 signaling

affects basal and stress-induced activation of the HPA axis as well as performance in tests for anxiety-like

behavior (Barna et al., 2004; Hill et al., 2011). Here we show that treatment with the FAAH inhibitor

URB597, which increases AEA levels and thus CB1 signaling, is sufficient to reverse the effect of *C.*

albicans colonization on CORT and anxiety-like behavior. We also show that colonization with *C.*

albicans altered levels of eCBs in the gut and that brain AEA levels correlated with the increase in serum

CORT observed in the *C. albicans*-colonized mice. These results establish a mechanism by which a

specific member of the gut microbiota, *C. albicans*, affects activity in the brain.

2. Results

2.1 A single oral inoculation with *Candida albicans* is sufficient to establish gastrointestinal colonization without inflammatory disease

We used a murine colonization model to investigate the effects of *C. albicans* inoculation on the mammalian host. Co-housing of up to 23 5-week-old mice in a large cage was used to standardize gut microbiota by coprophagy and therefore only female mice were used in these studies. The mice were acclimated to our facility and handled daily while co-housed. They were then either inoculated with a single dose of *C. albicans* or mock-inoculated with buffer and transferred to small cages in groups of 3-4. *C. albicans* was measurable in fecal pellets after 24h and 3 days after inoculation (Fig. 1B) and was detected throughout the GI tract (Fig. S1). We assessed behavior two days post-inoculation and mice were sacrificed after three days of colonization (Fig. 1A). *C. albicans*-colonized mice did not exhibit symptoms of illness or lose >5% bodyweight (Fig. S2). A multiplex ELISA was used to measure inflammatory cytokines in the serum after sacrifice and low levels of circulating IL-1β, IFN-γ, IL-10 and IL-6 were measured in mock-inoculated and *C. albicans*-colonized mice (Fig. 1C). Although IL-6 was significantly increased in *C. albicans*-colonized mice (Student's t-test, p=0.041), IL-6 as well as all other cytokines were well below the levels typically observed in mice with *C. albicans* disease (Tuite et al., 2005). Thus, our model resulted in stable colonization of mice with *C. albicans* in the GI tract that was not associated with disease, modeling commensal GI colonization.

Previous work demonstrated that colonization with *C. albicans* in the absence of antibiotic treatment had a small effect on the composition of the bacterial microbiota (Erb Downward et al., 2013). To determine whether the bacterial microbiota was significantly changed upon *C. albicans* colonization, we analyzed the bacterial microbiota of the cecum of mice sacrificed three days post-inoculation using 16s rRNA DNA sequencing and the QIIME analysis pipeline (Caporaso et al., 2010). The composition and diversity of bacterial taxa were not significantly altered by the introduction of *C. albicans* (Fig. 1D-F), consistent with previous results (Erb Downward et al., 2013). Mann-Whitney U test and linear

discriminant analysis effect size ((LEFSE) analysis (Segata et al., 2011) showed that no taxa were

significantly different in relative abundance between the microbiota of mock-colonized and *C. albicans*-

colonized mice (Fig. 1D). The overall diversity of the microbiota was also not significantly different

(Fig. 1E). We analyzed beta diversity and summarized the results using principal coordinates analysis

(Fig. 1F). PERMANOVA analysis of the complete principal coordinates analysis found that there was

not significant separation of populations based on *C. albicans* colonization (pseudo-F statistic=1.7,

p=0.12, 999 permutations). To summarize, mice inoculated with a single dose of *C. albicans* were stably

colonized with neither disruption of the bacterial microbiota nor invasive disease.

2.2 *C. albicans* colonization increases anxiety-like behavior in the EPM

To assess the effect of GI *C. albicans* colonization on host emotional behavior, standard tests for

anxiety-like and stress-coping behavior were used. Two days post-inoculation (Fig. 1A), all mice were

subjected to a single behavioral test, either the elevated plus maze (EPM) test for anxiety-like behavior

(Walf and Frye, 2007) or the forced swim test (FST) for stress-coping behavior (Hascoët and Bourin,

2009). The EPM consists of two open, aversive arms and two closed arms. Mice were allowed to explore

the EPM for a 5-minute trial which was scored by a blinded observer for time spent and entries into the

closed and open arms. Anxiety-like behavior is detected as avoidance of the open arms, expressed as

percentage arm time (time spent in specified arms divided by the total time spent all arms).

C. albicans colonization significantly increased anxiety-like behavior in the EPM. *C. albicans*-

colonized mice spent significantly less time in and made fewer entries into the open arms of the EPM

compared to the mock-colonized mice (Fig. 2A-B, Mann-Whitney test, p<0.05). There was a significant

increase in the percentage of time spent in the closed arms in the *C. albicans*-colonized mice but no

difference in the number of entries into the closed arms (Fig. 2C-D). There was no effect of *C. albicans*

colonization on the duration of time spent in the center or the total number of entries into all arms (Fig.

2E-F), indicating that *C. albicans* colonization did not alter overall activity but specifically increased

anxiety-like behavior in this test.

The forced swim test (FST) for stress-coping behavior was used to examine the effect of *C.*

albicans colonization on a second aspect of emotional behavior, again two days post-inoculation. Mice

were placed in a cylinder containing 22°C water for a six-minute trial which was video-recorded and later

scored by a blinded observer for time spent immobile. A passive-coping response in this assay was

defined as an increase in the amount of time spent floating immobile rather than the active coping

response defined as swimming or actively struggling. There was no difference in the time spent immobile

between the *C. albicans*-colonized mice and the mock-colonized mice (Fig. 2G), indicating that *C.*

albicans colonization did not affect stress-coping behavior in this test.

2.3 *C. albicans* colonization alters production of the stress hormone CORT

The underlying biology of anxiety-like behavior is multifaceted (Nutt et al., 2002). One

neuroendocrine pathway that has been shown to be dysregulated in human patients with anxiety is the

hypothalamus-pituitary-adrenal (HPA) axis (Kallen et al., 2008). To determine whether *C. albicans*

colonization affected the HPA axis, we sacrificed mice three days post-inoculation and measured

circulating serum CORT under basal conditions and in response to acute psychological stress.

C. albicans-colonized and mock-colonized mice were sacrificed unstressed. *C. albicans*-

colonized mice had significantly higher unstressed basal CORT (Fig. 3A Welch's t-test, p=0.0012). When

the effect of circadian rhythm on serum CORT was examined, basal CORT in both groups was shown to

increase over time throughout the afternoon, as expected (Chung et al., 2017) (Fig. 3B). Basal CORT in

C. albicans-colonized mice, however, began to rise sooner than in mock-colonized mice and thus was

significantly higher in *C. albicans*-colonized mice sacrificed two-to-five hours prior to lights out,

consistent with a circadian advance of CORT production (Fig. 3B, ANOVA followed by Sidak's multiple

comparison test, p<0.05). These results indicate that *C. albicans* colonization dysregulates circadian

control of the HPA axis. Further studies of basal CORT were conducted only with mice sacrificed

between 14:00-17:00.

To determine whether the changes to basal CORT regulation affected the stress-responsive

function of the HPA axis, mice were subjected to 30 minutes of restraint stress and then sacrificed

immediately or after 30 minutes or 60 minutes of recovery. *C. albicans*-colonized and mock-colonized

mice had comparable peak CORT immediately after stress (Fig. 3C) but *C. albicans*-colonized mice

failed to recover from stress to the same extent as the uncolonized mice after 60 minutes of recovery time

(Fig. 3C, Student's t-test, p=0.042). These results indicated that *C. albicans* colonization advanced

circadian basal CORT production and impaired negative feedback of the HPA axis after stress and

recovery.

Activation of the HPA axis begins in the brain, through excitation of the CRH-producing neurons

of the paraventricular nucleus (PVN) of the hypothalamus. We used immunofluorescence to quantify

expression of cFos protein in the PVN as a marker for neuronal activation. Nuclear cFos protein was not

detected in the PVN of unstressed mock-colonized or *C. albicans*-colonized mice (Fig. 3D, left). In mice

subjected to 30 minutes of stress and 30 minutes of recovery, nuclear cFos protein was comparably

detected in both mock-colonized and *C. albicans*-colonized mice (Fig. 3D, right). The PVN remains

stress-responsive in the *C. albicans*-colonized mice, and was not sufficiently stimulated at baseline to be

detected by this method. These results demonstrate that *C. albicans* colonization significantly impacts the

gut-brain axis through altered basal and stress-induced HPA axis function and increased anxiety-like

behavior.

2.4 Neuroendocrine changes observed in *C. albicans*-colonized mice are mediated by the

endocannabinoid system

The neuroactive lipid endocannabinoid (eCB) N-arachidonoylethanolamide (anandamide; AEA),

through its interactions with the CB1 receptor, is a key regulator of anxiety-like behavior and both stress-

induced and basal CORT production (Moreira et al., 2008). The circadian advance of basal CORT,

reduced feedback inhibition of stress-induced CORT and increased anxiety-like behavior observed in the

C. albicans-colonized mice resemble the phenotype of the CB1 knock-out mouse (Barna et al., 2004),

suggesting a deficit of eCB-CB1 signaling as a result of *C. albicans* colonization. Therefore, we

hypothesized that *C. albicans* colonization was interacting with the gut-brain axis via the

endocannabinoid system.

To test this hypothesis, we used the well-characterized drug URB597 to increase AEA levels and

therefore amplify AEA-CB1 signaling through inhibition of the AEA degradative enzyme, fatty acid

amide hydrolase (FAAH) (Fegley et al., 2005, p. 597). Mice were given an intraperitoneal injection of

vehicle or URB597 (1mg/kg) 4 hours prior to sacrifice under unstressed conditions, to determine whether

elevating AEA levels was sufficient to alleviate the high basal CORT of the *C. albicans*-colonized mice.

C. albicans-colonized mice given vehicle injection had significantly higher basal CORT than mock-

colonized mice as expected (Fig. 4A, Welch's t-test, p=0.0043). Administration of URB597 was

sufficient to decrease basal CORT in the *C. albicans*-colonized mice (Fig. 4A, Welch's t-test, p=0.0062).

This result is consistent with an AEA deficit in the *C. albicans*-colonized mice underlying the elevated

basal CORT. URB597 did not significantly change basal CORT in the mock-colonized mice (Welch's t-

test, p=0.10).

URB597 was administered 4-6 hours prior to testing in the EPM (0.1-0.15mg/kg) to determine

whether increasing AEA-CB1 signaling was sufficient to alleviate anxiety-like behavior in the *C.*

albicans-colonized mice. Due to variability in activity level of mice, data are shown as the percentage of

time spent in the open arm normalized to the experimental mean of the mock-colonized vehicle control.

As expected, *C. albicans*-colonized mice treated with vehicle had increased anxiety-like behavior

compared to the mock-colonized mice (Fig. 4B, Mann-Whitney U-test, p=0.0058). Treatment with

URB597 significantly increased open arm time of the *C. albicans*-colonized mice (Fig. 4B, Mann-

Whitney U-test, p=0.0025), which indicates that increasing AEA-CB1 signaling was sufficient to

normalize anxiety-like behavior in the *C. albicans*-colonized mice. There was no effect of URB597

treatment on overall activity as measured by total arm entries (Fig. 4C). Together, these results

demonstrate that increasing AEA-CB1 signaling with the FAAH inhibitor URB597 reversed the effect of

C. albicans colonization on basal CORT production and anxiety-like behavior, suggesting that the

neuroendocrine phenotypes observed in *C. albicans*-colonized mice result from decreased AEA-CB1

signaling.

2.5 Basal CORT and brain AEA levels are negatively correlated in *C. albicans*-colonized mice

To test the hypothesis that AEA levels were reduced in colonized mice, AEA levels in the

forebrain were measured. Mice were sacrificed on day 3 post-inoculation and forebrain samples analyzed

by mass spectrometry. We observed only a trend towards decreased AEA in the forebrains of *C.*

albicans-colonized mice compared to mock-colonized mice (Fig. 4D, Student's t-test, p=0.30). However,

a negative correlation between basal serum CORT and AEA in the *C. albicans*-colonized mice (Fig. 4F,

R^2=0.44, F-test, p=0.05) but not in the mock-colonized mice (Fig. 4E, R^2=0.119, F-test, p=0.45) was

observed. Thus mice that responded to *C. albicans* colonization with increased basal CORT did indeed

have lower levels of AEA in their forebrain. Previous results have demonstrated a similar negative

correlation between basal CORT and AEA levels in the prefrontal cortex of mice (Hill et al., 2010).

Taken together, these results indicate that *C. albicans* colonization had variable effects on bulk AEA

levels in the mouse forebrain and reduced forebrain AEA was observed in colonized mice with greater

dysregulation of the HPA axis.

A second canonical eCB, 2-arachidonoylglycerol (2-AG), was also measured in the forebrain of

mice. There was no difference between mock-colonized and *C. albicans*-colonized mice in bulk levels or

degree of correlation between 2-AG and serum CORT (Fig. S4).

2.6 *C. albicans* colonization alters the gut endocannabinoidome

We performed untargeted metabolomic analysis of the cecum contents of mice to determine

whether *C. albicans* colonization altered the metabolite composition of the GI tract and thereby altered

eCB metabolism. Of the 735 compounds measured, 21 were significantly differentially abundant in

mock-colonized versus *C. albicans*-colonized mice (Table S2, Student's t-test, p<0.05, uncorrected).

This result is below the false discovery rate and therefore pathway enrichment analysis was performed to

investigate the relevance of the 21 hits. Of the 5 pathways that contained more than one significant

compound, two were significantly enriched for *C. albicans*-dependent changes: sterols and eCBs (Fisher's

exact test, Holm's correction for multiple comparisons, p<0.0001 and p=0.031 respectively). The

enrichment in eCB family compounds indicated that *C. albicans* colonization altered eCB metabolism in

the gut, consistent with the eCB changes observed in the brain.

Two such eCB compounds, N-acylethanolamides (NAEs), linoleoyl and linolenoyl ethanolamide,

were significantly increased in the *C. albicans*-colonized mice (Fig. 4G-H, Student's t-test, p=0.0037 and

p=0.019). An additional NAE, oleoyl ethanolamide, was increased in the *C. albicans*-colonized mice but

not significantly (Fig. 4I, Student's t-test, p=0.078). These compounds are structurally similar to

anandamide (AEA) but have different acyl groups. AEA itself is present at low levels in the GI tract and

was not detected in any of the samples. The increase in 18-C NAEs implies that *C. albicans* colonization

altered NAE metabolism. Feeding studies have shown that dietary enrichment for a specific fatty acid

(FA) results in overproduction of the NAE derived from the enriched FA and limits production of

alternate NAEs (Sihag and Jones, 2019). Membrane phospholipid acyl chains can also act as substrates

for the production of NAEs. We measured a trend towards an increase in the 18-C FA linolenate (Fig. 4J,

Student's t-test, p=0.05) as well as a significant increase in the lysophospholipid 1-linoleoyl-

glycerophosphocholine (Fig. 4K, Student's t-test, p=0.017) in the GI tract of *C. albicans*-colonized mice.

Both of these compounds could be converted into their respective 18-C NAEs and increase production of

18C NAEs linoleoyl and linolenoyl ethanolamide while limiting production of AEA. Thus *C. albicans*-

induced changes in precursor compound abundance in the GI tract could contribute to the alterations in

AEA levels observed in the *C. albicans*-colonized mice.

2.7 Altered hepatic lipid metabolism in *C. albicans*-colonized mice reflects gut metabolite changes

In addition to the specific enrichment of two lipid subpathways, sterols and eCBs, and the trend

towards increased linolenate described above, we observed an overall increase in lipid compounds

containing long-chain polyunsaturated fatty acids (PUFAs) in the *C. albicans*-colonized mice (Fig. S3).

Lipids are absorbed from the GI tract into the blood and pass through the liver, a central organ for lipid

metabolism which is highly regulated by dietary lipid levels (Xie et al., 2010). We measured liver gene

expression to determine whether the changes in lipid levels we quantified in the cecum were sufficient to

induce a physiological change in the host.

Mice were sacrificed after three days of *C. albicans* colonization and qRT-PCR was used to

measure the expression of lipid-responsive genes in the liver. Decreased expression of the mRNA (*Scd1*)

encoding the enzyme stearoyl-coA desaturase (SCD1) was detected in the *C. albicans*-colonized mice

compared to the mock-colonized mice (Fig. 5A, Mann-Whitney U-test, p=0.0086), consistent with a

transcriptional response to increased PUFA-containing lipids from the GI tract (Lee et al., 1998). Gene

expression of regulators of lipogenesis, *Fads1* (acyl-CoA 8-3 desaturase) and *Fads2* (acyl-CoA 6

desaturase), the transcription factor sterol regulatory element-binding protein 1 (SREBP1-c) encoded by

Srebf1 and fatty acid synthase (*Fasn*) (Oosterveer et al., 2009), were not significantly changed by *C.*

albicans colonization (Fig. 5B), implying that the increased lipid abundance measured in the cecum was

not a result of host lipogenesis but rather that the increased abundance of PUFAs resulted from *C.*

albicans colonization and was sufficient to induce hepatic transcriptional changes.

Also, increased mRNA expression of two enzymes that regulate rate-limiting steps of hepatic

gluconeogenesis, phosphoenoylpyruvate carboxykinase (PEPCK-C) encoded by *Pck1* and glucose-6-

phosphatase (G6Pase) encoded by *G6pc* was detected in the *C. albicans*-colonized mice compared to the

mock-colonized mice (Fig. 5C, Student's t-test, p=0.027, Fig. 5D, Mann-Whitney U-test, p=0.0134).

These enzymes can be regulated by both dietary lipid abundance (Massillon et al., 2003) and by basal

serum CORT in accordance with the circadian rhythm (Reddy et al., 2007). Thus, the increased

expression of gluconeogenic enzymes may reflect both circadian advance of CORT and increased

abundance of dietary PUFAs present in *C. albicans*-colonized mice. Regardless, these changes in hepatic

gene expression demonstrate that *C. albicans* alters host physiology beyond the GI tract and could have a

significant impact on host metabolism.

To summarize, we demonstrated that colonization of mice with the commensal fungus *C.*

albicans resulted in significant changes to the gut-brain axis including increased anxiety-like behavior

and altered activity of the HPA axis. We identified altered eCB levels in the brain and GI tract of the *C.*

albicans-colonized mice and demonstrated that increasing levels of AEA through administration of

URB597 was sufficient to reverse the *C. albicans*-induced increase in CORT production and decrease

anxiety-like behavior. These results demonstrate that the gut microbiota can alter activity in the brain

through modulation of eCB signaling.

3.1 Discussion

C. albicans is the most common fungal member of the human gut microbiota. We investigated

the effect of *C. albicans* colonization on the gut-brain axis and observed specific changes in the

endocannabinoidome related to a stress-like behavioral and neuroendocrine phenotype. Previous

investigations of the gut-microbiota-brain axis have shown that colonization with commensal bacterial

species decreased CORT produced in response to stress and decreased anxiety-like behaviors (Messaoudi

et al., 2011; Sudo et al., 2004). In contrast to these bacteria, commensal *C. albicans* increased basal

CORT and increased anxiety-like behavior, resembling the neuroendocrine phenotypes observed after

infection with the parasite *Trichuris muris* (Bercik et al., 2010) or the mouse gastrointestinal pathogen

Citrobacter rodentium (Lyte et al., 2006). Researchers demonstrated that infection with these pathogens

increased anxiety-like behavior through increased inflammation. *C. albicans* in contrast is a commensal

and did not induce significant systemic inflammation. Rather, *C. albicans* colonization changed the gut

metabolome and altered eCB signaling to produce the observed increase in basal CORT and anxiety-like

behavior. To our knowledge, this is the first communication to report that a commensal fungus, *C.*

albicans, affects neuroendocrine host phenotypes and that microbiota-induced changes to eCBs can affect

the brain and behavior.

Treatment with the FAAH inhibitor URB597 alleviated both the elevated basal CORT and

increased anxiety-like behavior observed in *C. albicans*-colonized mice, indicating a common mechanism

of insufficient AEA. Consistent with the results of the URB597 experiments, we measured a trend

towards decreased AEA in the forebrain of *C. albicans*-colonized mice, and found a correlation in *C.*

albicans-colonized mice such that those with the largest increase in basal CORT also had the lowest

AEA, similar to what has been seen in chronic stress models (Hill et al., 2010). This negative correlation

supported the model that an AEA deficit was responsible for the neuroendocrine changes observed in *C.*

albicans-colonized mice. Previous investigations found that AEA levels specifically within the prefrontal

cortex (Hill et al., 2011) and the amygdala (Hill et al., 2010) were responsible for the downregulation of

CORT during recovery from stress and for anxiety-like behavior, respectively. It is possible that within

those discrete brain regions, colonized mice exhibited a larger decrease in AEA than the decrease we

measured in the forebrain as a whole. Untargeted metabolomic analysis of cecum contents further

supported the model that *C. albicans* colonization altered eCB metabolism, as two alternate NAEs,

linoleoyl and linolenoyl ethanolamide, were increased in abundance in the *C. albicans*-colonized mice as

were compounds containing their FA precursors. We suggest that these changes in GI tract eCB levels

could reflect systemic changes in lipid availability and NAE production and therefore disruption of

normal eCB metabolism. Such disruption could contribute to the altered eCB-CB1 signaling and

consequent neuroendocrine changes observed in the *C. albicans*-colonized mice.

Adolescence in humans is a dynamic period of development during which stress and anxiety

disorders present a significant health concern (Siegel and Dickstein, 2012); therefore ongoing research

into neuroendocrine health using adolescent models is relevant to human health. The current study was

performed with mid-adolescent (5-week-old) mice, which were sensitive to HPA axis dysregulation by GI

tract colonization with *C. albicans*. Previous work investigating the gut microbiota and the HPA axis

demonstrated a window during which mice are sensitive to microbiota manipulation, as Sudo *et al* (2004)

found that hyperreactivity of the HPA axis in germ-free mice could be reversed through microbiota

transplant in adolescent (6-week-old) mice but not in adult (8-week-old or 14-week-old) mice. Lee *et al*

(2013) showed that AEA levels in the brain increased significantly between adolescence and adulthood,

and thus the adolescent brain may be especially sensitive to the degree of eCB disruption observed in the

C. albicans-colonized mice. Previous researchers have also demonstrated significant developmental

differences in HPA axis regulation and response in both rodents (Romeo et al., 2014) and humans

(Gunnar et al., 2009; Netherton et al., 2004). Women are significantly more likely to experience anxiety

disorders than men (McLean et al., 2011), and therefore this study using a female mouse model provides

valuable insight into dysregulation of the HPA axis and anxiety-like behavior in a female population.

More broadly, these results highlight the sensitivity of the HPA axis and related anxiety-like

behavior to changes in AEA levels, and provide additional evidence that the eCB system is a viable drug

target for anxiety disorders. Indeed, a recent study in healthy human volunteers demonstrated that

treatment with a different FAAH inhibitor, PF-04457845, raised AEA levels and protected against the

development of stress-induced anxiety (Mayo et al., 2019), demonstrating that FAAH inhibition and

increased AEA can reverse activation of stress responses and modulate emotional behavior in humans as

it does in mouse models. Altogether this work demonstrates the ability of the gut metabolome to have a

significant impact on the brain and behavior, and illustrates the ability of the common gut commensal

fungus *C. albicans* to affect the host beyond the GI tract.

3.2 Conclusions

We have shown that GI colonization with the most prevalent human commensal fungus, *C.*

albicans, can alter systemic host health through the gut-brain axis. Although *C. albicans* colonization

was limited to the GI tract, mice so colonized exhibited global endocannabinoidome changes that resulted

in dysregulation of basal CORT production and increased anxiety-like behavior. We suggest that *C.*

albicans is able to induce these changes through manipulation of lipid availability in the GI tract, in

agreement with previous studies that have shown the importance of GI lipid pools for eCB production

throughout the host (Sihag and Jones, 2019). This study illustrates a novel mechanism by which a

member of the microbiota can impact host health, by modulating the GI tract metabolome to change eCB

levels throughout the host and alter the gut-brain axis.

4. Materials and Methods

Detailed methods are presented in the Supplemental Information

4.1 Animals

Up to 23 five-week-old female C57BL/6 mice (Jackson Laboratory) were cohoused in a large

cage (24"x17") and given sterile food, water and bedding. After four days of acclimation, mice were

transferred to standard cages with 3-4 mice inoculated with *Candida albicans* or 3-4 mice given a mock-

inoculum in each cage. Mice were put through a behavioral test on day two and sacrificed on day three

post-inoculation without stress (Sarkar et al., 2011).

All experiments were done in compliance with NIH Guide for the Care and Use of Laboratory Animals

and Tufts University IACUC guidelines.

4.2 Strains and growth conditions

C. albicans strain CKY101 (Brown et al., 1999) was used for all experiments. For preparation of

mouse inoculum, cells were grown at 37°C in standard yeast media for 24 hours. 25µl ($5x10^7$ cells) of

cells in PBS with 2% sucrose was fed to mice for gastrointestinal colonization. 25µl of 2% sucrose in

PBS was fed to mice for mock-inoculation.

4.3 Drug treatment

URB597 (Sigma Aldrich) and URB937 (Cayman Chemical) were dissolved in 18:1:1 normal

saline:PEG400:Tween80 (Sigma Aldrich) and administered to mice via intraperitoneal injection at a

dosage of 0.1-1mg/kg bodyweight four hours prior to behavioral testing or sacrifice.

4.4 Restraint stress

Mice were placed in a 50ml conical tube with two airholes enclosed with a rubber stopper for 30

minutes. After this restraint, mice were placed back in their home cage and anesthetized and sacrificed or

allowed to recover in the home cage for 30 or 60 minutes and then anesthetized and sacrificed.

4.5 Elevated Plus Maze

Behavioral testing in the Elevated Plus Maze (EPM) was performed as described in Walf *et al*

(2007). After a 5m trial in the EPM, the mouse was placed in a fresh sterile cage. All trials were

recorded with a video-camera from above and scored by a blinded observer.

4.6 Forced Swim Test

Behavioral testing in the forced swim test was performed as described in Can *et al* (2012). A

mouse was removed from its home cage and placed in the beaker of water for a six-minute trial which

was recorded by video-camera from the side and scored after the fact by a blinded observer. After testing,

mice were placed in a fresh sterile cage.

4.7 Bacterial microbiota analysis

The cecum, including contents, was dissected from mice after sacrifice on day three and was

immediately frozen on dry ice. Microbial DNA was extracted using the QIAamp DNA Stool Mini Kit

(Qiagen) following the manufacturer's protocol. Libraries were prepared from each sample and

sequenced as described (Caporaso et al., 2012). The resulting fastq files were used as input for

downstream analysis using QIIME (1.8.0)(Caporaso et al., 2010). The resultant OTU tables contained the

relative abundance of bacterial taxa in each sample. Analysis of alpha and beta diversity were performed

using standard QIIME scripts.

4.8 Measurement of hormones and cytokines in serum

Trunk blood was collected into serum separator blood collection tubes (BD) after sacrifice by

decapitation. Tubes were then spun to separate serum. Serum was divided into aliquots and frozen at -

80°C. Serum corticosterone was measured using a Corticosterone ELISA Kit (Enzo Life Sciences)

following the manufacturer's small volume protocol. Manufacturer reported sensitivity down to 27pg/ml

of CORT. Authors observed an average intra-assay coefficient of variance of 5.7% and an average inter-

assay coefficient of variance of 8.9%. Serum cytokines were measured using a multiplex ELISA

(Quanterix) following manufacturer's protocol. Manufacturer reported sensitivity for each cytokine

individually: mIFN-γ 7.1pg/ml, mIL-1β 1.1pg/ml, mIL-6 6.4pg/ml, mIL-10 3.0pg/ml, mIL-12p70

0.37pg/ml. Due to volume of mouse serum required for assay, authors did not have sufficient sample to

calculate inter-assay coefficient of variance.

4.9 Immunohistochemistry for cFOS in hypothalamus slices

Brains were dissected and fixed in 4% (w/v) paraformaldehyde in PBS for 24h at 4°C. They were

then cryopreserved in sucrose gradient, rapidly frozen in isopentane chilled on dry ice, and stored at -

80°C. Free-floating sections were prepared using a cryostat. Sections of interest were incubated with

1:5000 dilution of rabbit anti-mouse cFos antibody (Sigma Aldrich F7799) for 72h, then stained using

anti-rabbit IgG (VectaStain Elite ABC Kit) and streptadvidin-Alexa488 (Molecular Probes). Sections

were imaged using a Zeiss microscope with Apotome attachment.

4.10 Extraction and measurement of AEA in forebrain

Lipid extraction from forebrain samples was performed as described previously(Morena et al.,

2015). AEA was measured using mass spectrometry as described previously (Qi et al., 2015).

4.11 Untargeted metabolomic analysis of cecum contents

The cecum was dissected after sacrifice on day three and contents were squeezed into a tube and

immediately frozen in dry ice/ethanol bath and stored at -80°C. Extraction of metabolites and untargeted

mass spectrometry analysis were performed by Metabolon. Statistical analysis of results was performed

using Metabolon Metabolync tools.

4.12 Real-time quantitative PCR analysis

Tissues were frozen at -80°C in RNALater (Invitrogen). RNA was purified from tissues using

QIAzol for lysis and extraction with a Qialyzer, followed by column purification using the Ambion

Purelink Mini kit (Invitrogen). cDNA was synthesized using SuperScript III (Invitrogen) with oligo-dT

priming and the manufacturer's protocol. qPCR reactions were performed using SYBR Green Master

Mix (Applied Biosystems) and a LightCycler 480 II (Roche) instrument.

4.13 Statistical analysis

Statistical anlaysis was performed using GraphPad Prism8.4.0. The majority of the data was

analyzed using a two-tailed t-test to compare the means of the two experimental groups in question.

When the standard deviation of the data was significantly different between groups, Welch's t-test was

used; otherwise Student's t-test was used. Throughout $\alpha=0.05$ was used to assess significance. When

multiple comparisons were required, the Bonferroni correction was used to adjust the value of α. Data

was checked for normality using the D'agostino-Pearson omnibus test. When data was not normally

distributed, the Mann-Whitney test was used to compare experimental groups.

5. Acknowledgments

The authors gratefully acknowledge Dr. Michael Romero for providing critical expertise at an early stage of this project and Dr. Anne Kane for technical assistance in preparation of samples for microbiota analysis. This research was supported by NIH NIAID R01 AI118898 (to C.A.K.). JM is supported by the following grants: R01AA026256, R01NS105628, R01NS105628. LM was also supported by NIH training grant T32AI07422.

Conflicts of Interest

MH is a member of the advisory board for Sophren Therapeutics and JM is a member of the advisory board for SAGE Therapeutics.

6. References

1. Barna, I., Zelena, D., Arszovszki, A.C., Ledent, C., 2004. The role of endogenous cannabinoids in the hypothalamo-pituitary-adrenal axis regulation: in vivo and in vitro studies in CB1 receptor knockout mice. Life Sciences 75, 2959–2970. https://doi.org/10.1016/j.lfs.2004.06.006
2. Baxter, A.J., Scott, K.M., Vos, T., Whiteford, H.A., 2013. Global prevalence of anxiety disorders: a systematic review and meta-regression. Psychological Medicine 43, 897–910. https://doi.org/10.1017/S003329171200147X
3. Bercik, P., Verdu, E.F., Foster, J.A., Macri, J., Potter, M., Huang, X., Malinowski, P., Jackson, W., Blennerhassett, P., Neufeld, K.A., Lu, J., Khan, W.I., Corthesy-Theulaz, I., Cherbut, C., Bergonzelli, G.E., Collins, S.M., 2010. Chronic gastrointestinal inflammation induces anxiety-like behavior and alters central nervous system biochemistry in mice. Gastroenterology 139, 2102-2112.e1. https://doi.org/10.1053/j.gastro.2010.06.063
4. Brown, D.H., Jr., Giusani, A.D., Chen, X., Kumamoto, C.A., 1999. Filamentous growth of Candida albicans in response to physical environmental cues and its regulation by the unique CZF1 gene. Molecular microbiology 34, 651–62.
5. Can, A., Dao, D.T., Arad, M., Terrillion, C.E., Piantadosi, S.C., Gould, T.D., 2012. The Mouse Forced Swim Test. J Vis Exp. https://doi.org/10.3791/3638
6. Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D., Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld, J., Knight, R., 2010. QIIME allows analysis of high-throughput community sequencing data. Nat. Methods 7, 335–336. https://doi.org/10.1038/nmeth.f.303
7. Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S.M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J.A., Smith, G., Knight, R., 2012. Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J 6, 1621–1624. https://doi.org/10.1038/ismej.2012.8

8. Chung, S., Lee, E.J., Cha, H.K., Kim, J., Kim, D., Son, G.H., Kim, K., 2017. Cooperative roles of the suprachiasmatic nucleus central clock and the adrenal clock in controlling circadian glucocorticoid rhythm. Scientific Reports 7, 46404. https://doi.org/10.1038/srep46404

9. Dallman, M.F., Strack, A.M., Akana, S.F., Bradbury, M.J., Hanson, E.S., Scribner, K.A., Smith, M., 1993. Feast and Famine: Critical Role of Glucocorticoids with Insulin in Daily Energy Flow. Frontiers in Neuroendocrinology 14, 303–347. https://doi.org/10.1006/frne.1993.1010

10. Devane, W.A., Hanus, L., Breuer, A., Pertwee, R.G., Stevenson, L.A., Griffin, G., Gibson, D., Mandelbaum, A., Etinger, A., Mechoulam, R., 1992. Isolation and structure of a brain constituent that binds to the cannabinoid receptor. Science 258, 1946–1949. https://doi.org/10.1126/science.1470919

11. Erb Downward, J.R., Falkowski, N.R., Mason, K.L., Muraglia, R., Huffnagle, G.B., 2013. Modulation of post-antibiotic bacterial community reassembly and host response by Candida albicans. Scientific reports 3, 2191. https://doi.org/10.1038/srep02191

12. Fegley, D., Gaetani, S., Duranti, A., Tontini, A., Mor, M., Tarzia, G., Piomelli, D., 2005. Characterization of the fatty acid amide hydrolase inhibitor cyclohexyl carbamic acid 3'-carbamoyl-biphenyl-3-yl ester (URB597): effects on anandamide and oleoylethanolamide deactivation. J. Pharmacol. Exp. Ther. 313, 352–358. https://doi.org/10.1124/jpet.104.078980

13. Gunnar, M.R., Wewerka, S., Frenn, K., Long, J.D., Griggs, C., 2009. Developmental changes in hypothalamus–pituitary–adrenal activity over the transition to adolescence: Normative changes and associations with puberty. Dev Psychopathol 21, 69–85. https://doi.org/10.1017/S0954579409000054

14. Hascoët, M., Bourin, M., 2009. The Forced Swimming Test in Mice: A Suitable Model to Study Antidepressants, in: Mood and Anxiety Related Phenotypes in Mice, Neuromethods. Humana Press, Totowa, NJ, pp. 85–118. https://doi.org/10.1007/978-1-60761-303-9_6

15. Hill, M.N., McLaughlin, R.J., Bingham, B., Shrestha, L., Lee, T.T.Y., Gray, J.M., Hillard, C.J., Gorzalka, B.B., Viau, V., 2010. Endogenous cannabinoid signaling is essential for stress adaptation. Proc. Natl. Acad. Sci. U.S.A. 107, 9406–9411. https://doi.org/10.1073/pnas.0914661107

16. Hill, M.N., McLaughlin, R.J., Pan, B., Fitzgerald, M.L., Roberts, C.J., Lee, T.T.-Y., Karatsoreos, I.N., Mackie, K., Viau, V., Pickel, V.M., McEwen, B.S., Liu, Q., Gorzalka, B.B., Hillard, C.J., 2011. Recruitment of prefrontal cortical endocannabinoid signaling by glucocorticoids contributes to termination of the stress response. J. Neurosci. 31, 10506–10515. https://doi.org/10.1523/JNEUROSCI.0496-11.2011

17. Kallen, V.L., Tulen, J.H.M., Utens, E.M.W.J., Treffers, P.D.A., Jong, F.H.D., Ferdinand, R.F., 2008. Associations between HPA axis functioning and level of anxiety in children and adolescents with an anxiety disorder. Depression and Anxiety 25, 131–141. https://doi.org/10.1002/da.20287

18. Lee, K.N., Pariza, M.W., Ntambi, J.M., 1998. Conjugated Linoleic Acid Decreases Hepatic Stearoyl-CoA Desaturase mRNA Expression. Biochemical and Biophysical Research Communications 248, 817–821. https://doi.org/10.1006/bbrc.1998.8994

19. Lee, T.T.-Y., Hill, M.N., Hillard, C.J., Gorzalka, B.B., 2013. Temporal changes in N-acylethanolamine content and metabolism throughout the peri-adolescent period. Synapse 67, 4–10. https://doi.org/10.1002/syn.21609

20. Leonardo, E.D., Hen, R., 2008. Anxiety as a Developmental Disorder. Neuropsychopharmacol 33, 134–140. https://doi.org/10.1038/sj.npp.1301569

21. Luna, R.A., Foster, J.A., 2015. Gut brain axis: diet microbiota interactions and implications for modulation of anxiety and depression. Curr. Opin. Biotechnol. 32, 35–41. https://doi.org/10.1016/j.copbio.2014.10.007

22. Lyte, M., Li, W., Opitz, N., Gaykema, R.P.A., Goehler, L.E., 2006. Induction of anxiety-like behavior in mice during the initial stages of infection with the agent of murine colonic

hyperplasia Citrobacter rodentium. Physiol. Behav. 89, 350–357.
https://doi.org/10.1016/j.physbeh.2006.06.019

23. Markey, L., Shaban, L., Green, E.R., Lemon, K.P., Mecsas, J., Kumamoto, C.A., 2018. Pre-
colonization with the commensal fungus Candida albicans reduces murine susceptibility to
Clostridium difficile infection. Gut Microbes 9, 497–509.
https://doi.org/10.1080/19490976.2018.1465158

24. Massillon, D., Arinze, I.J., Xu, C., Bone, F., 2003. Regulation of Glucose-6-phosphatase Gene
Expression in Cultured Hepatocytes and H4IIE Cells by Short-chain Fatty Acids ROLE OF
HEPATIC NUCLEAR FACTOR-4α. J. Biol. Chem. 278, 40694–40701.
https://doi.org/10.1074/jbc.M303182200

25. Mayo, L.M., Asratian, A., Lindé, J., Morena, M., Haataja, R., Hammar, V., Augier, G., Hill,
M.N., Heilig, M., 2019. Elevated anandamide, enhanced recall of fear extinction, and attenuated
stress responses following inhibition of fatty acid amide hydrolase (FAAH): a randomized,
controlled experimental medicine trial. Biological Psychiatry.
https://doi.org/10.1016/j.biopsych.2019.07.034

26. McLean, C.P., Asnaani, A., Litz, B.T., Hofmann, S.G., 2011. Gender differences in anxiety
disorders: Prevalence, course of illness, comorbidity and burden of illness. Journal of Psychiatric
Research 45, 1027–1035. https://doi.org/10.1016/j.jpsychires.2011.03.006

27. Mechoulam, R., Ben-Shabat, S., Hanus, L., Ligumsky, M., Kaminski, N.E., Schatz, A.R.,
Gopher, A., Almog, S., Martin, B.R., Compton, D.R., 1995. Identification of an endogenous 2-
monoglyceride, present in canine gut, that binds to cannabinoid receptors. Biochem. Pharmacol.
50, 83–90. https://doi.org/10.1016/0006-2952(95)00109-d

28. Messaoudi, M., Lalonde, R., Violle, N., Javelot, H., Desor, D., Nejdi, A., Bisson, J.-F., Rougeot,
C., Pichelin, M., Cazaubiel, M., Cazaubiel, J.-M., 2011. Assessment of psychotropic-like
properties of a probiotic formulation (Lactobacillus helveticus R0052 and Bifidobacterium
longum R0175) in rats and human subjects. Br. J. Nutr. 105, 755–764.
https://doi.org/10.1017/S0007114510004319

29. Moreira, F.A., Kaiser, N., Monory, K., Lutz, B., 2008. Reduced anxiety-like behaviour induced
by genetic and pharmacological inhibition of the endocannabinoid-degrading enzyme fatty acid
amide hydrolase (FAAH) is mediated by CB1 receptors. Neuropharmacology, Cannabinoid
Signaling in the Nervous System 54, 141–150. https://doi.org/10.1016/j.neuropharm.2007.07.005

30. Morena, M., De Castro, V., Gray, J.M., Palmery, M., Trezza, V., Roozendaal, B., Hill, M.N.,
Campolongo, P., 2015. Training-Associated Emotional Arousal Shapes Endocannabinoid
Modulation of Spatial Memory Retrieval in Rats. J. Neurosci. 35, 13962–13974.
https://doi.org/10.1523/JNEUROSCI.1983-15.2015

31. Netherton, C., Goodyer, I., Tamplin, A., Herbert, J., 2004. Salivary cortisol and
dehydroepiandrosterone in relation to puberty and gender. Psychoneuroendocrinology 29, 125–
140. https://doi.org/10.1016/s0306-4530(02)00150-6

32. Nutt, D.J., Ballenger, J.C., Sheehan, D., Wittchen, H.-U., 2002. Generalized anxiety disorder:
comorbidity, comparative biology and treatment. Int J Neuropsychopharmacol 5, 315–325.
https://doi.org/10.1017/S1461145702003048

33. Oosterveer, M.H., Dijk, T.H. van, Tietge, U.J.F., Boer, T., Havinga, R., Stellaard, F., Groen,
A.K., Kuipers, F., Reijngoud, D.-J., 2009. High Fat Feeding Induces Hepatic Fatty Acid
Elongation in Mice. PLOS ONE 4, e6066. https://doi.org/10.1371/journal.pone.0006066

34. Placzek, E.A., Okamoto, Y., Ueda, N., Barker, E.L., 2008. Mechanisms for recycling and
biosynthesis of endogenous cannabinoids anandamide and 2-arachidonylglycerol. Journal of
Neurochemistry 107, 987–1000. https://doi.org/10.1111/j.1471-4159.2008.05659.x

35. Qi, M., Morena, M., Vecchiarelli, H.A., Hill, M.N., Schriemer, D.C., 2015. A robust capillary
liquid chromatography/tandem mass spectrometry method for quantitation of neuromodulatory

endocannabinoids. Rapid Commun. Mass Spectrom. 29, 1889–1897.
https://doi.org/10.1002/rcm.7277

36. Raimondi, S., Amaretti, A., Gozzoli, C., Simone, M., Righini, L., Candeliere, F., Brun, P., Ardizzoni, A., Colombari, B., Paulone, S., Castagliuolo, I., Cavalieri, D., Blasi, E., Rossi, M., Peppoloni, S., 2019. Longitudinal Survey of Fungi in the Human Gut: ITS Profiling, Phenotyping, and Colonization. Front Microbiol 10, 1575. https://doi.org/10.3389/fmicb.2019.01575

37. Reddy, A.B., Maywood, E.S., Karp, N.A., King, V.M., Inoue, Y., Gonzalez, F.J., Lilley, K.S., Kyriacou, C.P., Hastings, M.H., 2007. Glucocorticoid signaling synchronizes the liver circadian transcriptome. Hepatology 45, 1478–1488. https://doi.org/10.1002/hep.21571

38. Romeo, R.D., Minhas, S., Svirsky, S.E., Hall, B.S., Savenkova, M., Karatsoreos, I.N., 2014. Pubertal shifts in adrenal responsiveness to stress and adrenocorticotropic hormone in male rats. Psychoneuroendocrinology 42, 146–152. https://doi.org/10.1016/j.psyneuen.2014.01.016

39. Sapolsky, R.M., Romero, L.M., Munck, A.U., 2000. How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. Endocr. Rev. 21, 55–89. https://doi.org/10.1210/edrv.21.1.0389

40. Sarkar, J., Wakefield, S., MacKenzie, G., Moss, S.J., Maguire, J., 2011. Neurosteroidogenesis Is Required for the Physiological Response to Stress: Role of Neurosteroid-Sensitive GABAA Receptors. J. Neurosci. 31, 18198–18210. https://doi.org/10.1523/JNEUROSCI.2560-11.2011

41. Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S., Huttenhower, C., 2011. Metagenomic biomarker discovery and explanation. Genome Biol 12, R60. https://doi.org/10.1186/gb-2011-12-6-r60

42. Shao, T.-Y., Ang, W.X.G., Jiang, T.T., Huang, F.S., Andersen, H., Kinder, J.M., Pham, G., Burg, A.R., Ruff, B., Gonzalez, T., Khurana Hershey, G.K., Haslam, D.B., Way, S.S., 2019. Commensal Candida albicans Positively Calibrates Systemic Th17 Immunological Responses. Cell Host & Microbe 25, 404-417.e6. https://doi.org/10.1016/j.chom.2019.02.004

43. Siegel, R.S., Dickstein, D.P., 2012. Anxiety in adolescents: Update on its diagnosis and treatment for primary care providers. Adolescent Health, Medicine and Therapeutics 3, 1. https://doi.org/10.2147/AHMT.S7597

44. Sihag, J., Jones, P.J.H., 2019. Dietary fatty acid profile influences circulating and tissue fatty acid ethanolamide concentrations in a tissue-specific manner in male Syrian hamsters. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids 1864, 1563–1579. https://doi.org/10.1016/j.bbalip.2019.07.002

45. Sudo, N., Chida, Y., Aiba, Y., Sonoda, J., Oyama, N., Yu, X.-N., Kubo, C., Koga, Y., 2004. Postnatal microbial colonization programs the hypothalamic-pituitary-adrenal system for stress response in mice. J. Physiol. (Lond.) 558, 263–275. https://doi.org/10.1113/jphysiol.2004.063388

46. Tuite, A., Elias, M., Picard, S., Mullick, A., Gros, P., 2005. Genetic control of suceptibility to Candida albicans in susceptible A/J and resistant C57BL/6J mice. Genes & Immunity 6, 672–682. https://doi.org/10.1038/sj.gene.6364254

47. Walf, A.A., Frye, C.A., 2007. The use of the elevated plus maze as an assay of anxiety-related behavior in rodents. Nature Protocols 2, 322–328. https://doi.org/10.1038/nprot.2007.44

48. Xie, Z., Li, H., Wang, K., Lin, J., Wang, Q., Zhao, G., Jia, W., Zhang, Q., 2010. Analysis of transcriptome and metabolome profiles alterations in fatty liver induced by high-fat diet in rat. Metabolism 59, 554–560. https://doi.org/10.1016/j.metabol.2009.08.022

7. Figure Legends

Figure 1: Gastrointestinal colonization with *Candida albicans* after a single inoculation without disruption of bacterial gut microbiota or invasive disease

A) Timeline for acute colonization model. 5-week-old female mice were cohoused for four days and handled daily. Mice were then moved into smaller groups in standard cages and orally inoculated with $5x10^7$CFU of *C. albicans* or given buffer. On day one post-inoculation, fecal pellets were sterilely collected from mice and plated on YPD-SA to measure colonization levels. On day two post-inoculation, all mice underwent a behavioral test. On day three post-inoculation mice were anesthetized with isoflurane and sacrificed by decapitation. Some mice were subjected to restraint stress prior to sacrifice. B) *C. albicans* CFU/g of fecal pellets was measured on day 1 post-inoculation and CFU/g of cecum contents was measured after sacrifice on day three post-inoculation. No culturable fungi were measured on either day from mock-colonized mice. Day 1: N=56, Day 3: N=50. Figure includes data from 6 cohorts. Bars indicate geometric mean. C) A multiplex ELISA was used to measure inflammatory cytokines in the serum of mice. N.D, not detectable. Student's t-test, p<0.05. Mock-colonized N=13, *C. albicans*-colonized N=14. Figure includes data from 3 cohorts. D-F) Microbiota analysis of the cecum contents of mice was performed using standard 16s rRNA DNA sequencing and QIIME analysis pipeline. Mock-colonized N=8 and *C. albicans*-colonized N=8. Figure includes data from 2 cohorts. D) The relative abundance of all bacterial families detected with a median relative abundance greater than 0 is shown. Bar shows geometric mean with SEM E) The average alpha diversity metric Chao1 was calculated to determine the diversity of the microbiota at different levels of sampling. The average Chao1 score of the experimental groups is shown. F) Beta diversity was calculated using weighted UniFrac scores and principal coordinates analysis was performed. For C-D, symbols indicate individual mice and bars indicate the average. Mock-colonized mice are represented as solid dots and bars; *C. albicans*-colonized mice are represented as open circles and bars.

Figure 2: *C. albicans* **colonization increases anxiety-like behavior**

On day two post-inoculation mice underwent either the Elevated Plus Maze (EPM) or the Forced Swim Test (FST). Trials in the behavioral tests were video-recorded and scored by a blinded observer after the fact. For the EPM data (A-F) that follows, mock-colonized N=22 and *C. albicans*-colonized N=19 and statistical analysis performed was Mann-Whitney U-test (A-C, G) or Student's t-test (D-F), $p<0.05$. Figures include data from two cohorts. A) Percentage of time spent in the open arms of the EPM. B) Total number of entries into the open arms of the EPM. C) Percentage of time spent in the closed arms of the EPM. D) Total entries into the closed arms of the EPM. E) Percentage of time spent in the neutral central square of the EPM (neither open nor closed arms). F) Total entries during the trial (sum of the open and closed arm entries), a metric for locomotor activity. G) Total time spent immobile during the six-minute FST trial. Immobility was defined as no movements beyond those required to stay afloat. For FST data: mock-colonized N=33, *C. albicans*-colonized N=24. Figure includes data from three cohorts. Throughout, symbols represent individual mice and bars indicate average. Mock-colonized mice are represented as solid dots and bars; *C. albicans*-colonized mice are represented as open circles and bars.

Figure 3: *C. albicans* **colonization increases basal CORT and alters feedback inhibition after acute stress**

A) Corticosterone (CORT) in trunk blood of mice sacrificed without stress was measured by ELISA. Mock-colonized N=15, *C. albicans*-colonized N=19. Figure includes data from four cohorts. Data was analyzed using Welch's t-test, $p < 0.05$ B) CORT in trunk blood of mice sacrificed at different times of day. In addition to the data summarized in A (mice sacrificed from 13:00-17:00), B includes mice sacrificed without stress from 17:00-20:00. Data was binned in one-hour increments. Mock-colonized: 13:00 N=5, 14:00 N=9, 15:00 N=11, 16:00 N=6, 17:00 N=9, 18:00 N=8, 19:00 N=8. *C. albicans*-colonized: 13:00 N=6, 14:00 N=13, 15:00 N=10, 16:00 N=11, 17:00 N=10, 18:00 N=6, 19:00 N=14. C) Mice were subjected to 30 minutes of restraint stress and then sacrificed immediately (0m recovered) or after 30 minutes or 60 minutes of recovery as shown on the x-axis. Mock-colonized: 0m N=14, 30m N=15, 60m N=13. *C. albicans*-colonized: 0m N=15, 30m N=13, 60m N=12. D) Immunohistochemistry was used to visualize cFos protein with Alexa488 in the paraventricular nucleus (PVN) of mice using the 20X objective. DAPI was used to visualize nuclei and Alexa488 to visualize cFOS. The PVN of mice sacrificed unstressed (left) do not exhibit cFOS staining, while PVN of mice sacrificed after 30 minutes of stress and 30 minutes of recovery (right) are positive for cFOS. Representative images are shown. For figures A and C, symbols indicate individual mice and bars indicate the average. For figure B, symbols indicate average and error bars indicate standard deviation. Throughout, mock-colonized mice are solid dots and bars and *C. albicans*-colonized mice are open circles and bars.

Figure 4: Neuroendocrine phenotypes of the *C. albicans*-colonized mice are mediated through disruption of the endocannabinoid system

        A-C) Mice were treated with the fatty acid amide hydrolase (FAAH) inhibitor URB597, a well-characterized drug which increases AEA levels by blocking degradation. Mice were given either URB597 or vehicle control by intraperitoneal injection 4-6 hours prior to sacrifice (A) or testing in the EPM (B-C). A) Mice were treated with 1mg/kg of URB597 or vehicle. ELISA was used to measure CORT in trunk blood of mice sacrificed without stress. Welch's t-test, $p < 0.05$. Mock-colonized: vehicle N=15, URB597 N=19. *C. albicans*-colonized: vehicle N=16, URB597 N=18. Figure includes data from four cohorts. B-C) Mice were treated with 0.1mg/kg (triangles) or 0.15mg/kg (circles) URB597 or vehicle, then tested in the EPM. Mock-colonized: vehicle N=17, 0.1mg/kg URB597 N=4, 0.15mg/kg URB597 N=8. *C. albicans*-colonized: vehicle N=13, 0.1mg/kg URB597 N=4 0.15mg/kg URB597 N=8. B) Percentage time spent in the open arms, normalized to the average percent open time of the mock-colonized vehicle control, is shown. Mann-Whitney U-test, $p < 0.05$. C) Total arms entries, a metric for overall locomotor activity is shown. D) AEA (pmol/g tissue) was measured in forebrain samples by mass spectrometry. E-F) Linear regression of basal serum CORT and forebrain AEA (D) in individual mice are shown. (D-F) Mock-colonized N=7, *C. albicans*-colonized N=9. Untargeted mass spectrometry was used to measure relative abundance of 735 compounds in the cecum contents of mice. N-acylethanolamides (G-I), a free fatty acid (J) and a lysophospholipids (K) containing an 18-carbon chain with varying degrees of unsaturation are shown here. Raw data was log-transformed. Mock-colonized N=8, *C. albicans*-colonized N=8. Student's t-test, $p < 0.05$. Symbols indicate individual mice and bars indicate average. Solid circles and bars are mock-colonized mice and open circles and bars are *C. albicans*-colonized mice.

Figure 5: *C. albicans* colonization affects hepatic lipid-responsive gene expression

           RNA was extracted from the liver of mice sacrificed unstressed on day three. cDNA was synthesized and qRT-PCR was used to quantify liver gene transcription. RNA gene expression is normalized to GAPDH and expressed relative to the average expression of the mock-colonized mice. A) Stearoyl-CoA desaturase 1 (*Scd1*) RNA expression in the liver is shown. Mock-colonized N=24, *C. albicans*-colonized N=27. Figure includes data from four cohorts. B) Expression of multiple lipogenic genes is shown: acyl-CoA 8-3 desaturase (*Fads1*), acyl-CoA 6 desaturase (*Fads2*), sterol regulatory element-binding protein 1 (*Srepbf1*) and fatty acid synthase (*Fasn*). Mock-colonized N=11, *C. albicans*-colonized N= 10. Figure includes data from two cohorts. C) Expression of the gluconeogenesis enzymes phosphoenolpyruvate carboxykinase (*Pck1*) (mock-colonized N=24, *C. albicans*-colonized N=23) and D) glucose-6-phosphatase (*G6Pc*) is shown (mock-colonized N=20, *C. albicans*-colonized N=18). Figures include data from four cohorts. Symbols represent individual mice and bars indicate average. Solid symbols and bars indicate mock-colonized mice and open symbols and bars indicate *C. albicans*-colonized mice. Mann-Whitney U-test was used for statistical analysis of A and D; Welch's t-test was used for analysis of B and C, $p < 0.05$.

8. Figures

Figure 1: Gastrointestinal colonization with *Candida albicans* after a single inoculation without disruption of bacterial gut microbiota or invasive disease

Figure 2: *C. albicans* **colonization increases anxiety-like behavior**

Figure 3: *C. albicans* colonization increases basal CORT and alters feedback inhibition after acute stress

Figure 4: Neuroendocrine phenotypes of the *C. albicans*-colonized mice are mediated through disruption of the endocannabinoid system

Figure 5: *C. albicans* **colonization affects hepatic lipid-responsive gene expression**