Colonization with the commensal fungus *Candida albicans* perturbs the gut-brain axis

through dysregulation of endocannabinoid signaling

Laura Markey[a], Andrew Hooper[b], Laverne C. Melon[b,c], Samantha Baglot[d], Matthew N.

Hill[d], Jamie Maguire[b] and Carol A. Kumamoto[a]*

[a]Department of Molecular Biology and Microbiology, Tufts University, 150 Harrison

Ave, Boston, MA, 02111

[b]Department of Neuroscience, Tufts University, 150 Harrison Ave, Boston, MA, 02111

[c]Current affiliation- Department of Biology, Wesleyan University, Middletown, CT,

06459

[d]Hotchkiss Brain Institute, University of Calgary, Calgary AB, Canada, T2N1Z4

***Corresponding Author:** Carol A. Kumamoto, Tufts University, Department of

Molecular Biology and Microbiology, 136 Harrison Ave., Boston, MA 02111 USA

Classification

Biological Sciences; Microbiology and Neuroscience

Abstract

Anxiety disorders are the most prevalent mental health disorder worldwide, with a lifetime prevalence of 5-7% of the human population. Although the etiology of anxiety disorders is incompletely understood, one aspect of host health that affects anxiety disorders is the gut-brain axis. We used a mouse model of gastrointestinal (GI) colonization to demonstrate that the commensal fungus *Candida albicans* affects host health via the gut-brain axis. In mice, bacterial members of the gut microbiota can influence the host gut-brain axis, affecting anxiety-like behavior and the hypothalamus-pituitary-adrenal (HPA) axis which produces the stress hormone corticosterone (CORT). Here we showed that mice colonized with *C. albicans* demonstrated increased anxiety-like behavior and increased basal production of CORT as well as dysregulation of CORT production following acute stress. The HPA axis and anxiety-like behavior are negatively regulated by the endocannabinoid anandamide (AEA). *C. albicans*-colonized mice exhibited changes in endocannabinoid levels within the GI tract and a negative correlation between brain AEA levels and serum CORT. Further, increasing AEA levels using the well-characterized fatty acid amide hydrolase (FAAH) inhibitor URB597 was sufficient to reverse both neuroendocrine phenotypes in *C. albicans*-colonized mice. Thus, a commensal fungus that is a common colonizer of humans had widespread effects on the physiology of its host. To our knowledge, this is the first report of microbial manipulation of the endocannabinoid (eCB) system that resulted in neuroendocrine changes contributing to anxiety-like behavior.

Keywords

Endocannabinoid, microbiota, *Candida albicans*, anxiety, corticosterone

Significance Statement

Anxiety disorders are the most prevalent mental health disorder worldwide yet their etiology is incompletely understood and treatment options are limited. The gut-brain axis has been shown to influence the development of anxiety disorders. Recent work has established that the bacterial gut microbiota can influence the gut-brain axis and result in changes to neuroendocrine health including anxiety-like behavior in mice. We describe a novel role for a fungal member of the gut microbiota, *Candida albicans*, that influences the neuroendocrine system through manipulation of the endocannabinoid system. Additionally, this work demonstrates that acute, limited manipulation of endocannabinoids can have a significant impact on host health, emphasizing the potential of this system as a target for the treatment of anxiety disorders.

Introduction

Anxiety disorders are the most prevalent mental health disorder worldwide, with a lifetime prevalence of 5-7% of the human population[1]. The etiology of anxiety disorders is complex and incompletely understood, although many different genetic and environmental factors have been identified[2]. The gut-brain axis, the bi-directional communication between the gut and the brain through multiple host systems, has been shown to affect anxiety disorders[3]. Constant communication between the periphery and the brain is required for normal homeostasis and overall health.

One component of the gut-brain axis thought to play a role in anxiety disorders is the hypothalamus-pituitary-adrenal (HPA) axis. The endocrine output of the HPA axis,

the stress hormone corticosterone (CORT), is a broad regulator of host health including

the immune, neuroendocrine, metabolic and cardiovascular systems. Under basal

conditions, CORT release is regulated by the circadian rhythm and helps to maintain

homeostasis[4]. CORT is also a key part of the acute psychological stress response and a

rapid increase in circulating CORT is essential for mounting an effective response to

stress[5]. Dysregulation of the HPA axis and CORT production is correlated with mental

health disorders including anxiety and depression[6].

The gut microbiota, the diverse assemblage of microbes that colonizes the GI tract

of humans, has been shown to affect virtually every aspect of human health, including the

gut-brain axis. Researchers using germ-free mouse models demonstrated that the gut

microbiota plays an important role in the development and regulation of the HPA axis[7,8].

Studies have also shown that specific probiotic bacterial species can affect the gut-brain

axis, as treatment with *Bifidobacterium sp*. and *Lactobacillus sp*. decreases anxiety-like

behavior and normalizes HPA axis function in mice[9,10]. Investigation into the

contribution of the gut-microbiota-brain axis to mental health could reveal novel

regulators of anxiety-like behavior and offer mechanistic insight into the etiology of the

disease.

Candida albicans is a commensal fungus that colonizes the GI tract of ~60% of

the human population[11]. Although a major colonizer of humans, its role as a commensal

microbe is largely uncharacterized. Previous researchers have shown that GI *C. albicans*

colonization induces local and systemic immune changes which are protective against

systemic *C. albicans* infection and disease[12], and against infection with bacterial

pathogens such as *C. difficile*[13] and *S. aureus*[12]. The research that follows demonstrates

that *C. albicans* affects host health beyond the GI tract. We show that *C. albicans*

dysregulated HPA axis activity and increased anxiety-like behavior in a mouse model by

altering endocannabinoid signaling.

The mammalian endocannabinoid (eCB) system regulates both the HPA axis and

anxiety-like behavior. It consists of two major neuroactive lipids, N-

arachidonoylethanolamide (AEA) and 2-arachidonoylglycerol (2-AG) and their receptors

CB1 (distributed throughout the nervous system) and CB2 (present primarily on immune

cells)[14–16]. AEA and 2-AG are produced in response to activation in post-synaptic

neurons, and act at CB1 on pre-synaptic neurons to limit neurotransmitter release. AEA

and 2-AG levels are regulated by substrate availability and the activity of their synthetic

and degradative enzymes[17]. Using genetic and pharmacological tools in rodents,

researchers have demonstrated that manipulation of CB1 signaling affects basal and

stress-induced activation of the HPA axis as well as performance in tests for anxiety-like

behavior[18,19]. Here we show that treatment with the FAAH inhibitor URB597, which

increases AEA levels and thus CB1 signaling, is sufficient to reverse the effect of *C.*

albicans colonization on CORT and anxiety-like behavior. We also show that

colonization with *C. albicans* altered levels of eCBs in the gut and that brain AEA levels

correlated with the increase in serum CORT observed in the *C. albicans*-colonized mice.

These results establish a mechanism by which a specific member of the gut microbiota,

C. albicans, affects activity in the brain.

Results

A single oral inoculation with *Candida albicans* is sufficient to establish gastrointestinal colonization without inflammatory disease

We used a murine colonization model to investigate the effects of *C. albicans* inoculation on the mammalian host. Co-housing of up to 23 5-week-old mice in a large cage was used to standardize gut microbiota by coprophagy and therefore only female mice were used in these studies. The mice were acclimated to our facility and handled daily while co-housed. They were then either inoculated with a single dose of *C. albicans* or mock-inoculated with buffer and transferred to small cages in groups of 3-4. *C. albicans* was measurable in fecal pellets after 24h and 3 days after inoculation (Fig. 1B) and was detected throughout the GI tract (Fig. S1). We assessed behavior two days post-inoculation and mice were sacrificed after three days of colonization (Fig. 1A). *C. albicans*-colonized mice did not exhibit symptoms of illness or lose >5% bodyweight (Fig. S2). A multiplex ELISA was used to measure inflammatory cytokines in the serum after sacrifice and low levels of circulating IL-1β, IFN-γ, IL-10 and IL-6 were measured in mock-inoculated and *C. albicans*-colonized mice (Fig. 1C). Although IL-6 was significantly increased in *C. albicans*-colonized mice (Student's t-test, p=0.04), IL-6 as well as all other cytokines were well below the levels typically observed in mice with *C. albicans* disease[21]. Thus, our model resulted in stable colonization of mice with *C. albicans* in the GI tract that was not associated with disease, modeling commensal GI colonization.

Previous work demonstrated that colonization with *C. albicans* in the absence of

antibiotic treatment had a small effect on the composition of the bacterial microbiota[22].

To determine whether the bacterial microbiota was significantly changed upon *C.*

albicans colonization, we analyzed the bacterial microbiota of the cecum of mice

sacrificed three days post-inoculation using 16s rRNA DNA sequencing and the QIIME

analysis pipeline[23]. The composition and diversity of bacterial taxa were not significantly

altered by the introduction of *C. albicans* (Fig. 1D-F), consistent with previous results[22].

Mann-Whitney U test and linear discriminant analysis effect size ((LEFSE) analysis[24]

showed that no taxa were significantly different in relative abundance between the

microbiota of mock-colonized and *C. albicans*-colonized mice (Fig. 1D). The overall

diversity of the microbiota was also not significantly different (Fig. 1E). We analyzed

beta diversity and summarized the results using principal coordinates analysis (Fig. 1F).

PERMANOVA analysis of the complete principal coordinates analysis found that there

was not significant separation of populations based on *C. albicans* colonization (pseudo-F

statistic=1.7306, p=0.117, 999 permutations).   To summarize, mice inoculated with a

single dose of *C. albicans* were stably colonized with neither disruption of the bacterial

microbiota nor invasive disease.

***C. albicans* colonization increases anxiety-like behavior in the EPM**

To assess the effect of GI *C. albicans* colonization on host emotional behavior,

standard tests for anxiety-like and stress-coping behavior were used. Two days post-

inoculation (Fig. 1A), all mice were subjected to a single behavioral test, either the

elevated plus maze (EPM) test for anxiety-like behavior[25] or the forced swim test (FST)

for stress-coping behavior[26]. The EPM consists of two open, aversive arms and two

closed arms. Mice were allowed to explore the EPM for a 5-minute trial which was

scored by a blinded observer for time spent and entries into the closed and open arms.

Anxiety-like behavior is detected as avoidance of the open arms, expressed as percentage

arm time (time spent in specified arms divided by the total time spent all arms).

C. albicans colonization significantly increased anxiety-like behavior in the EPM.

C. albicans-colonized mice spent significantly less time in and made fewer entries into

the open arms of the EPM compared to the mock-colonized mice (Fig. 2A-B, Welch's t-

test, $p < 0.05$). There was a significant increase in the percentage of time spent in the

closed arms in the *C. albicans*-colonized mice but no difference in the number of entries

into the closed arms (Fig. 2C-D). There was no effect of *C. albicans* colonization on the

duration of time spent in the center or the total number of entries into all arms (Fig. 2E-

F), indicating that *C. albicans* colonization did not alter overall activity but specifically

increased anxiety-like behavior in this test.

The forced swim test (FST) for stress-coping behavior was used to examine the

effect of *C. albicans* colonization on a second aspect of emotional behavior, again two

days post-inoculation. Mice were placed in a cylinder containing 22°C water for a six-

minute trial which was video-recorded and later scored by a blinded observer for time

spent immobile. A passive-coping response in this assay was defined as an increase in

the amount of time spent floating immobile rather than the active coping response

defined as swimming or actively struggling. There was no difference in the time spent

immobile between the *C. albicans*-colonized mice and the mock-colonized mice (Fig.

2G), indicating that *C. albicans* colonization did not affect stress-coping behavior in this

test.

C. albicans colonization alters production of the stress hormone CORT

The underlying biology of anxiety-like behavior is multifaceted[27,28]. One

neuroendocrine pathway that has been shown to be dysregulated in human patients with

anxiety is the hypothalamus-pituitary-adrenal (HPA) axis[6]. To determine whether *C.*

albicans colonization affected the HPA axis, we sacrificed mice three days post-

inoculation and measured circulating serum CORT under basal conditions and in

response to acute psychological stress.

C. albicans-colonized and mock-colonized mice were sacrificed unstressed. *C.*

albicans-colonized mice had significantly higher unstressed basal CORT (Fig. 3A

Welch's t-test, p=0.0012). When the effect of circadian rhythm on serum CORT was

examined, basal CORT in both groups was shown to increase over time throughout the

afternoon, as expected[29] (Fig. 3B). Basal CORT in *C. albicans*-colonized mice, however,

began to rise sooner than in mock-colonized mice and thus was significantly higher in *C.*

albicans-colonized mice sacrificed three-to-five hours prior to lights out, consistent with

a circadian advance of CORT production. These results indicate that *C. albicans*

colonization dysregulates circadian control of the HPA axis. Further studies of basal

CORT were conducted only with mice sacrificed between 14:00-16:00.

To determine whether the changes to basal CORT regulation affected the stress-

responsive function of the HPA axis, mice were subjected to 30 minutes of restraint stress

and then sacrificed immediately or after 30 minutes or 60 minutes of recovery. *C.*

albicans-colonized and mock-colonized mice had comparable peak CORT immediately

after stress (Fig. 3C) but *C. albicans*-colonized mice failed to recover from stress to the

same extent as the uncolonized mice after 60 minutes of recovery time (Fig. 3C,

Student's t-test, p=0.04). These results indicated that *C. albicans* colonization advanced

circadian basal CORT production and impaired negative feedback of the HPA axis after

stress and recovery.

Activation of the HPA axis begins in the brain, through excitation of the CRH-

producing neurons of the paraventricular nucleus (PVN) of the hypothalamus. We used

immunofluorescence to quantify expression of cFos protein in the PVN as a marker for

neuronal activation. Nuclear cFos protein was not detected in the PVN of unstressed

mock-colonized or *C. albicans*-colonized mice (Fig. 3D, left). In mice subjected to 30

minutes of stress and 30 minutes of recovery, nuclear cFos protein was comparably

detected in both mock-colonized and *C. albicans*-colonized mice (Fig. 3D, right). The

PVN remains stress-responsive in the *C. albicans*-colonized mice, and was not

sufficiently stimulated at baseline to be detected by this method. These results

demonstrate that *C. albicans* colonization significantly impacts the gut-brain axis through

altered basal and stress-induced HPA axis function and increased anxiety-like behavior.

Neuroendocrine changes observed in *C. albicans*-colonized mice are mediated by the

endocannabinoid system

The neuroactive lipid endocannabinoid (eCB) N-arachidonoylethanolamide

(anandamide; AEA), through its interactions with the CB1 receptor, is a key regulator of

anxiety-like behavior and both stress-induced and basal CORT production[30–32]. The

circadian advance of basal CORT, reduced feedback inhibition of stress-induced CORT

and increased anxiety-like behavior observed in the *C. albicans*-colonized mice resemble

the phenotype of the CB1 knock-out mouse[19], suggesting a deficit of eCB-CB1 signaling

as a result of *C. albicans* colonization. Therefore, we hypothesized that *C. albicans*

colonization was interacting with the gut-brain axis via the endocannabinoid system.

To test this hypothesis, we used the well-characterized drug URB597 to increase

AEA levels and therefore amplify AEA-CB1 signaling through inhibition of the AEA

degradative enzyme, fatty acid amide hydrolase (FAAH)[33,34]. Mice were given an

intraperitoneal injection of vehicle or URB597 (1mg/kg) 4 hours prior to sacrifice under

unstressed conditions, to determine whether elevating AEA levels was sufficient to

alleviate the high basal CORT of the *C. albicans*-colonized mice. *C. albicans*-colonized

mice given vehicle injection had significantly higher basal CORT than mock-colonized

mice as expected (Fig. 4A, Welch's t-test, p=0.0043). Administration of URB597 was

sufficient to decrease basal CORT in the *C. albicans*-colonized mice (Fig. 4A, Welch's t-

test, p=0.0062). This result is consistent with an AEA deficit in the *C. albicans*-

colonized mice underlying the elevated basal CORT. URB597 did not significantly

change basal CORT in the mock-colonized mice (Welch's t-test, p=0.0971).

URB597 was administered 4-6 hours prior to testing in the EPM (0.1-0.15mg/kg)

to determine whether increasing AEA-CB1 signaling was sufficient to alleviate anxiety-

like behavior in the *C. albicans*-colonized mice. Due to variability in activity level of

mice, data are shown as the percentage of time spent in the open arm normalized to the

experimental mean of the mock-colonized vehicle control. As expected, *C. albicans*-

238 colonized mice treated with vehicle had increased anxiety-like behavior compared to the

239 mock-colonized mice (Fig. 4B, Welch's t-test, p=0.0033). Treatment with URB597

240 significantly increased open arm time of the *C. albicans*-colonized mice (Fig. 4B,

241 Welch's t-test, p=0.0024), which indicates that increasing AEA-CB1 signaling was

242 sufficient to normalize anxiety-like behavior in the *C. albicans*-colonized mice. There

243 was no effect of URB597 treatment on overall activity as measured by total arm entries

244 (Fig. 4C). Together, these results demonstrate that increasing AEA-CB1 signaling with

245 the FAAH inhibitor URB597 reversed the effect of *C. albicans* colonization on basal

246 CORT production and anxiety-like behavior, suggesting that the neuroendocrine

247 phenotypes observed in *C. albicans*-colonized mice result from decreased AEA-CB1

248 signaling.

249 **Basal CORT and brain AEA levels are negatively correlated in *C. albicans*-colonized**

250 **mice**

251   To test the hypothesis that AEA levels were reduced in colonized mice, AEA

252 levels in the forebrain were measured. Mice were sacrificed on day 3 post-inoculation

253 and forebrain samples analyzed by mass spectrometry. We observed only a trend

254 towards decreased AEA in the forebrains of *C. albicans*-colonized mice compared to

255 mock-colonized mice (Fig. 4D, Student's t-test, p=0.3). However, a negative correlation

256 between basal serum CORT and AEA in the *C. albicans*-colonized mice (Fig. 4F,

257 R^2=0.4416, F-test, p=0.05) but not in the mock-colonized mice (Fig. 4E, R^2=0.119, F-

258 test, p=0.4486) was observed. Thus mice that responded to *C. albicans* colonization with

259 increased basal CORT did indeed have lower levels of AEA in their forebrain. Previous

results have demonstrated a similar negative correlation between basal CORT and AEA

levels in the prefrontal cortex of mice[35]. Taken together, these results indicate that *C.*

albicans colonization had variable effects on bulk AEA levels in the mouse forebrain and

reduced forebrain AEA was observed in colonized mice with greater dysregulation of the

HPA axis.

A second canonical eCB, 2-arachidonoylglycerol (2-AG), was also measured in

the forebrain of mice. There was no difference between mock-colonized and *C. albicans*-

colonized mice in bulk levels or degree of correlation between 2-AG and serum CORT

(Fig. S4).

***C. albicans* colonization alters the gut endocannabinoidome**

We performed untargeted metabolomic analysis of the cecum contents of mice to

determine whether *C. albicans* colonization altered the metabolite composition of the GI

tract and thereby altered eCB metabolism. Of the 735 compounds measured, 21 were

significantly differentially abundant in mock-colonized versus *C. albicans*-colonized

mice (Table S2, Student's t-test, $p < 0.05$, uncorrected). This result is below the false

discovery rate and therefore pathway enrichment analysis was performed to investigate

the relevance of the 21 hits. Of the 5 pathways that contained more than one significant

compound, two were significantly enriched for *C. albicans*-dependent changes: sterols

and eCBs (Fisher's exact test, Holm's correction for multiple comparisons, $p < 0.0001$ and

$p = 0.0305$ respectively). The enrichment in eCB family compounds indicated that *C.*

albicans colonization altered eCB metabolism in the gut, consistent with the eCB

changes observed in the brain.

Two such eCB compounds, N-acylethanolamides (NAEs), linoleoyl and

linolenoyl ethanolamide, were significantly increased in the *C. albicans*-colonized mice

(Fig. 4G-H, Student's t-test, p=0.0037 and p=0.0188). An additional NAE, oleoyl

ethanolamide, was increased in the *C. albicans*-colonized mice but not significantly (Fig.

4I, Student's t-test, p=0.08). These compounds are structurally similar to anandamide

(AEA) but have different acyl groups. AEA itself is present at low levels in the GI tract

and was not detected in any of the samples. The increase in 18-C NAEs implies that *C.*

albicans colonization altered NAE metabolism. Feeding studies have shown that dietary

enrichment for a specific fatty acid (FA) results in overproduction of the NAE derived

from the enriched FA and limits production of alternate NAEs[37,38]. Membrane

phospholipid acyl chains can also act as substrates for the production of NAEs. We

measured a trend towards an increase in the 18-C FAs linolenate and linoleate (Fig. 4J-K,

Student's t-test, p=0.05 and p=0.06) as well as a significant increase in the

lysophospholipid 1-linoleoyl-glycerophosphocholine (Fig. 4L, Student's t-test, p=0.02) in

the GI tract of *C. albicans*-colonized mice. All of these compounds could be converted

into their respective 18-C NAEs and increase production of 18C NAEs linoleoyl and

linolenoyl ethanolamide while limiting production of AEA, contributing to the alterations

in AEA levels observed in the *C. albicans*-colonized mice.

Altered hepatic lipid metabolism in *C. albicans*-colonized mice reflects gut

metabolite changes

In addition to the specific enrichment of two lipid subpathways, sterols and eCBs

and the trends towards increased 18-C FAs described above, we observed an overall

increase in lipid compounds containing long-chain polyunsaturated fatty acids (PUFAs)

in the *C. albicans*-colonized mice (Fig. S3). Lipids are absorbed from the GI tract into

the blood and pass through the liver, a central organ for lipid metabolism which is highly

regulated by dietary lipid levels[39]. We measured liver gene expression to determine

whether the changes in lipid levels we quantified in the cecum were sufficient to induce a

physiological change in the host.

Mice were sacrificed after three days of *C. albicans* colonization and qRT-PCR

was used to measure the expression of lipid-responsive genes in the liver. Decreased

expression of the mRNA (*Scd1*) encoding the enzyme stearoyl-coA desaturase (SCD1)

was detected in the *C. albicans*-colonized mice compared to the mock-colonized mice

(Fig. 5A, Student's t-test, p=0.006), consistent with a transcriptional response to

increased PUFA-containing lipids from the GI tract[40]. Gene expression of regulators of

lipogenesis, *Fads1* (acyl-CoA 8-3 desaturase) and *Fads2* (acyl-CoA 6 desaturase), the

transcription factor sterol regulatory element-binding protein 1 (SREBP1-c) encoded by

Srebf1 and fatty acid synthase (*Fasn*)[41], were not significantly changed by *C. albicans*

colonization (Fig. 5B), implying that the increased lipid abundance measured in the

cecum was not a result of host lipogenesis. Also, increased mRNA expression of two

enzymes that regulate rate-limiting steps of hepatic gluconeogenesis,

phosphoenoylpyruvate carboxykinase (PEPCK-C) encoded by *Pck1* and glucose-6-

phosphatase (G6Pase) encoded by *G6pc* was detected in the *C. albicans*-colonized mice

compared to the mock-colonized mice (Fig. 5C-D, Student's t-test, p<0.05), consistent

with a liver response to increased dietary lipids[42]. These genes are positively regulated

by endocrine mediators CORT and glucagon, as well as by dietary lipids[42,43] and so may

also be upregulated by the elevated basal CORT present in the *C. albicans*-colonized

mice. These results imply that the increased abundance of PUFAs measured in the GI

tract was sufficient to induce hepatic transcriptional changes.

To summarize, we demonstrated that colonization of mice with the commensal

fungus *C. albicans* resulted in significant changes to the gut-brain axis including

increased anxiety-like behavior and altered activity of the HPA axis. We identified

altered eCB levels in the brain and GI tract of the *C. albicans*-colonized mice and

demonstrated that increasing levels of AEA through administration of URB597 was

sufficient to reverse the *C. albicans*-induced increase in CORT production and decrease

anxiety-like behavior. These results demonstrate that the gut microbiota can alter activity

in the brain through modulation of eCB signaling.

Discussion

C. albicans is the most common fungal member of the human gut microbiota. We

investigated the effect of *C. albicans* colonization on the gut-brain axis and observed

specific changes in the endocannabinoidome related to a stress-like behavioral and

neuroendocrine phenotype. Previous investigations of the gut-microbiota-brain axis have

shown that colonization with commensal bacterial species decreased CORT produced in

response to stress and decreased anxiety-like behaviors[8,9]. In contrast to these bacteria,

commensal *C. albicans* increased basal CORT and increased anxiety-like behavior,

resembling the neuroendocrine phenotypes observed after infection with the parasite

Trichuris muris[44] or the mouse gastrointestinal pathogen *Citrobacter rodentium*[45].

Researchers demonstrated that infection with these pathogens increased anxiety-like

behavior through increased inflammation. *C. albicans* in contrast is a commensal and did

not induce significant systemic inflammation. Rather, *C. albicans* colonization changed

the gut metabolome and altered eCB signaling to produce the observed increase in basal

CORT and anxiety-like behavior. To our knowledge, this is the first communication to

report that a commensal fungus, *C. albicans,* affects neuroendocrine host phenotypes and

that microbiota-induced changes to eCBs can affect the brain and behavior.

Treatment with the FAAH inhibitor URB597 alleviated both the elevated basal

CORT and increased anxiety-like behavior observed in *C. albicans*-colonized mice,

indicating a common mechanism of insufficient AEA. Consistent with the results of the

URB597 experiments, we measured a trend towards decreased AEA in the forebrain of

C. albicans-colonized mice, and found a correlation in *C. albicans*-colonized mice such

that those with the largest increase in basal CORT also had the lowest AEA, similar to

what has been seen in chronic stress models[35]. This negative correlation supported the

model that an AEA deficit was responsible for the neuroendocrine changes observed in

C. albicans-colonized mice. Previous investigations found that AEA levels specifically

within the prefrontal cortex[18] and the amygdala[35,46] were responsible for the

downregulation of CORT during recovery from stress and for anxiety-like behavior,

respectively. It is possible that within those discrete brain regions, colonized mice

exhibited a larger decrease in AEA than the decrease we measured in the forebrain as a

whole. Untargeted metabolomic analysis of cecum contents further supported the model

that *C. albicans* colonization altered eCB metabolism, as two alternate NAEs, linoleoyl

and linolenoyl ethanolamide, were increased in abundance in the *C. albicans*-colonized

mice as were compounds containing their FA precursors. We suggest that these changes

in GI tract eCB levels could reflect systemic changes in lipid availability and NAE

production and therefore disruption of normal eCB metabolism. Such disruption could

contribute to the altered eCB-CB1 signaling and consequent neuroendocrine changes

observed in the *C. albicans*-colonized mice.

Previous work investigating the gut microbiota and the HPA axis demonstrated a

window during which mice are sensitive to microbiota manipulation, as Sudo *et al*[8] found

that hyperreactivity of the HPA axis in germ-free mice could be reversed through

microbiota transplant in adolescent (6-week-old) mice but not in adult (8-week-old or 14-

week-old) mice. The current study was performed with mid-adolescent (5-week-old)

mice, which were sensitive to HPA axis dysregulation by GI tract colonization with *C.*

albicans. Lee *et al* showed that AEA levels in the brain increased significantly between

adolescence and adulthood[47], and thus the adolescent brain may be sensitive to the degree

of eCB disruption observed in the *C. albicans*-colonized mice. Previous researchers have

also demonstrated significant developmental differences in HPA axis regulation and

response in both rodents[48] and humans[49,50]. Adolescence in humans is a dynamic period

of development during which stress and anxiety disorders present a significant health

concern[51]; therefore ongoing research into neuroendocrine health using adolescent

models is relevant to human health. Women are significantly more likely to experience

anxiety disorders than men[52], and therefore this study using a female mouse model

provides valuable insight into dysregulation of the HPA axis and anxiety-like behavior in

a female population.

More broadly, these results highlight the sensitivity of the HPA axis and related

anxiety-like behavior to changes in AEA levels, and provide additional evidence that the

eCB system is a viable drug target for anxiety disorders. Indeed, a recent study in healthy

human volunteers demonstrated that treatment with a different FAAH inhibitor, PF-

04457845, raised AEA levels and protected against the development of stress-induced

anxiety[53], demonstrating that FAAH inhibition and increased AEA can reverse activation

of stress responses and modulate emotional behavior in humans as it does in mouse

models. Altogether this work demonstrates the ability of the gut metabolome to have a

significant impact on the brain and behavior, and illustrates the ability of the common gut

commensal fungus *C. albicans* to affect the host beyond the GI tract.

Materials and Methods

Detailed methods are presented in the Supplemental Information

Animals

Up to 23 five-week-old female C57BL/6 mice (Jackson Laboratory) were

cohoused in a large cage (24"x17") and given sterile food, water and bedding. After four

days of acclimation, mice were transferred to standard cages with 3-4 mice inoculated

with *Candida albicans* or 3-4 mice given a mock-inoculum in each cage. Mice were put

through a behavioral test on day two and sacrificed on day three post-inoculation without

stress[54].

All experiments were done in compliance with Tufts University IACUC guidelines.

Strains and growth conditions

C. albicans strain CKY101[55] was used for all experiments. For preparation of

mouse inoculum, cells were grown at 37°C in standard yeast media for 24 hours. 25µl

(5×10^7 cells) of cells in PBS with 2% sucrose was fed to mice for gastrointestinal

colonization. 25µl of 2% sucrose in PBS was fed to mice for mock-inoculation.

Drug treatment

URB597 (Sigma Aldrich) and URB937 (Cayman Chemical) were dissolved in

18:1:1 normal saline:PEG400:Tween80 (Sigma Aldrich) and administered to mice via

intraperitoneal injection at a dosage of 0.1-1mg/kg bodyweight four hours prior to

behavioral testing or sacrifice.

Restraint stress

Mice were placed in a 50ml conical tube with two airholes enclosed with a rubber

stopper for 30 minutes. After this restraint, mice were placed back in their home cage

and anesthetized and sacrificed or allowed to recover in the home cage for 30 or 60

minutes and then anesthetized and sacrificed.

Elevated Plus Maze

Behavioral testing in the Elevated Plus Maze (EPM) was performed as described

in Walf *et al* 2013[25]. After a 5m trial in the EPM, the mouse was placed in a fresh sterile

cage. All trials were recorded with a video-camera from above and scored by a blinded

observer.

Forced Swim Test

Behavioral testing in the forced swim test was performed as described in Can *et al* 2012 [56]. A mouse was removed from its home cage and placed in the beaker of water for a six-minute trial which was recorded by video-camera from the side and scored after the fact by a blinded observer. After testing, mice were placed in a fresh sterile cage.

Bacterial microbiota analysis

The cecum, including contents, was dissected from mice after sacrifice on day three and was immediately frozen on dry ice. Microbial DNA was extracted using the QIAamp DNA Stool Mini Kit (Qiagen) following the manufacturer's protocol. Libraries were prepared from each sample and sequenced as described[57]. The resulting fastq files were used as input for downstream analysis using QIIME (1.8.0)[23]. The resultant OTU tables contained the relative abundance of bacterial taxa in each sample. Analysis of alpha and beta diversity were performed using standard QIIME scripts.

Measurement of hormones and cytokines in serum

Trunk blood was collected into serum separator blood collection tubes (BD) after sacrifice by decapitation. Tubes were then spun to separate serum. Serum was divided into aliquots and frozen at -80°C.

Serum corticosterone was measured using a Corticosterone ELISA Kit (Enzo Life Sciences) following the manufacturer's small volume protocol. Serum cytokines were measured using a multiplex ELISA (Quanterix) following manufacturer's protocol.

Immunohistochemistry for cFOS in hypothalamus slices

Brains were dissected and fixed in 4% (w/v) paraformaldehyde in PBS for 24h at 4°C. They were then cryopreserved in sucrose gradient, rapidly frozen in isopentane chilled on dry ice, and stored at -80°C. Free-floating sections were prepared using a cryostat. Sections of interest were incubated with 1:5000 dilution of rabbit anti-mouse cFos antibody (Sigma Aldrich F7799) for 72h, then stained using anti-rabbit IgG (VectaStain Elite ABC Kit) and streptadvidin-Alexa488 (Molecular Probes). Sections were imaged using a Zeiss microscope with Apotome attachment.

Extraction and measurement of AEA in forebrain

Lipid extraction from forebrain samples was performed as described previously[58]. AEA was measured using mass spectrometry as described previously[59].

Untargeted metabolomic analysis of cecum contents

The cecum was dissected after sacrifice on day three and contents were squeezed into a tube and immediately frozen in dry ice/ethanol bath and stored at -80°C. Extraction of metabolites and untargeted mass spectrometry analysis were performed by Metabolon. Statistical analysis of results was performed using Metabolon Metabolync tools.

Real-time quantitative PCR analysis

Tissues were frozen at -80°C in RNALater (Invitrogen). RNA was purified from tissues using QIAzol for lysis and extraction with a Qialyzer, followed by column purification using the Ambion Purelink Mini kit (Invitrogen). cDNA was synthesized using SuperScript III (Invitrogen) with oligo-dT priming and the manufacturer's protocol.

475 qPCR reactions were performed using SYBR Green Master Mix (Applied Biosystems)

476 and a LightCycler 480 II (Roche) instrument.

Acknowledgments

The authors gratefully acknowledge Dr. Michael Romero for providing critical

expertise at an early stage of this project and Dr. Anne Kane for technical assistance in

preparation of samples for microbiota analysis. This research was supported by NIH

NIAID R01 AI118898 (to Carol A. Kumamoto). JM is supported by the following grants:

R01AA026256, R01NS105628, R01NS105628. LM was also supported by NIH training

grant T32AI07422.

Conflicts of Interest

485 MH is a member of the advisory board for Sophren Therapeutics and JM is a

486 member of the advisory board for SAGE Therapeutics.

References

1.   Baxter, A. J., Scott, K. M., Vos, T. & Whiteford, H. A. Global prevalence of
anxiety disorders: a systematic review and meta-regression. *Psychological Medicine* **43**,
897–910 (2013).
2.   Leonardo, E. D. & Hen, R. Anxiety as a Developmental Disorder.
Neuropsychopharmacol **33**, 134–140 (2008).
3.   Luna, R. A. & Foster, J. A. Gut brain axis: diet microbiota interactions and
implications for modulation of anxiety and depression. *Curr. Opin. Biotechnol.* **32**, 35–41
(2015).
4.   Dallman, M. F. *et al.* Feast and Famine: Critical Role of Glucocorticoids with
Insulin in Daily Energy Flow. *Frontiers in Neuroendocrinology* **14**, 303–347 (1993).
5.   Sapolsky, R. M., Romero, L. M. & Munck, A. U. How do glucocorticoids
influence stress responses? Integrating permissive, suppressive, stimulatory, and
preparative actions. *Endocr. Rev.* **21**, 55–89 (2000).
6.   Kallen, V. L. *et al.* Associations between HPA axis functioning and level of
anxiety in children and adolescents with an anxiety disorder. *Depression and Anxiety* **25**,
131–141 (2008).

7.   Neufeld, K. M., Kang, N., Bienenstock, J. & Foster, J. A. Reduced anxiety-like behavior and central neurochemical change in germ-free mice. *Neurogastroenterol. Motil.* **23**, 255–264, e119 (2011).

8.   Sudo, N. *et al.* Postnatal microbial colonization programs the hypothalamic-pituitary-adrenal system for stress response in mice. *J. Physiol. (Lond.)* **558**, 263–275 (2004).

9.   Messaoudi, M. *et al.* Assessment of psychotropic-like properties of a probiotic formulation (Lactobacillus helveticus R0052 and Bifidobacterium longum R0175) in rats and human subjects. *Br. J. Nutr.* **105**, 755–764 (2011).

10.   Ait-Belgnaoui, A. *et al.* Prevention of gut leakiness by a probiotic treatment leads to attenuated HPA response to an acute psychological stress in rats. *Psychoneuroendocrinology* **37**, 1885–1895 (2012).

11.   Raimondi, S. *et al.* Longitudinal Survey of Fungi in the Human Gut: ITS Profiling, Phenotyping, and Colonization. *Front Microbiol* **10**, 1575 (2019).

12.   Shao, T.-Y. *et al.* Commensal Candida albicans Positively Calibrates Systemic Th17 Immunological Responses. *Cell Host & Microbe* **25**, 404-417.e6 (2019).

13.   Markey, L. *et al.* Pre-colonization with the commensal fungus Candida albicans reduces murine susceptibility to Clostridium difficile infection. *Gut Microbes* **9**, 497–509 (2018).

14.   Matsuda, L. A., Lolait, S. J., Brownstein, M. J., Young, A. C. & Bonner, T. I. Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* **346**, 561–564 (1990).

15.   Devane, W. A. *et al.* Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science* **258**, 1946–1949 (1992).

16.   Mechoulam, R. *et al.* Identification of an endogenous 2-monoglyceride, present in canine gut, that binds to cannabinoid receptors. *Biochem. Pharmacol.* **50**, 83–90 (1995).

17.   Placzek, E. A., Okamoto, Y., Ueda, N. & Barker, E. L. Mechanisms for recycling and biosynthesis of endogenous cannabinoids anandamide and 2-arachidonylglycerol. *Journal of Neurochemistry* **107**, 987–1000 (2008).

18.   Hill, M. N. *et al.* Recruitment of prefrontal cortical endocannabinoid signaling by glucocorticoids contributes to termination of the stress response. *J. Neurosci.* **31**, 10506–10515 (2011).

19.   Barna, I., Zelena, D., Arszovszki, A. C. & Ledent, C. The role of endogenous cannabinoids in the hypothalamo-pituitary-adrenal axis regulation: in vivo and in vitro studies in CB1 receptor knockout mice. *Life Sciences* **75**, 2959–2970 (2004).

20.   Tiraboschi, I. C. N. *et al.* Congenital candidiasis: confirmation of mother-neonate transmission using molecular analysis techniques. *Med Mycol* **48**, 177–181 (2010).

21.   Tuite, A., Elias, M., Picard, S., Mullick, A. & Gros, P. Genetic control of suceptibility to Candida albicans in susceptible A/J and resistant C57BL/6J mice. *Genes & Immunity* **6**, 672–682 (2005).

22.   Erb Downward, J. R., Falkowski, N. R., Mason, K. L., Muraglia, R. & Huffnagle, G. B. Modulation of post-antibiotic bacterial community reassembly and host response by Candida albicans. *Scientific reports* **3**, 2191 (2013).

23.   Caporaso, J. G. *et al.* QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods* **7**, 335–336 (2010).

24.   Segata, N. *et al.* Metagenomic biomarker discovery and explanation. *Genome Biol*
12, R60 (2011).
25.   Walf, A. A. & Frye, C. A. The use of the elevated plus maze as an assay of
anxiety-related behavior in rodents. *Nature Protocols* **2**, 322–328 (2007).
26.   Hascoët, M. & Bourin, M. The Forced Swimming Test in Mice: A Suitable Model
to Study Antidepressants. in *Mood and Anxiety Related Phenotypes in Mice* 85–118
(Humana Press, Totowa, NJ, 2009). doi:10.1007/978-1-60761-303-9_6.
27.   Kalueff, A. V., Wheaton, M. & Murphy, D. L. What's wrong with my mouse
model?: Advances and strategies in animal modeling of anxiety and depression.
Behavioural Brain Research **179**, 1–18 (2007).
28.   Nutt, D. J., Ballenger, J. C., Sheehan, D. & Wittchen, H.-U. Generalized anxiety
disorder: comorbidity, comparative biology and treatment. *Int J Neuropsychopharmacol*
5, 315–325 (2002).
29.   Chung, S. *et al.* Cooperative roles of the suprachiasmatic nucleus central clock
and the adrenal clock in controlling circadian glucocorticoid rhythm. *Scientific Reports* **7**,
46404 (2017).
30.   Patel, S., Roelke, C. T., Rademacher, D. J., Cullinan, W. E. & Hillard, C. J.
Endocannabinoid Signaling Negatively Modulates Stress-Induced Activation of the
Hypothalamic-Pituitary-Adrenal Axis. *Endocrinology* **145**, 5431–5438 (2004).
31.   Moreira, F. A., Kaiser, N., Monory, K. & Lutz, B. Reduced anxiety-like
behaviour induced by genetic and pharmacological inhibition of the endocannabinoid-
degrading enzyme fatty acid amide hydrolase (FAAH) is mediated by CB1 receptors.
Neuropharmacology **54**, 141–150 (2008).
32.   Hill, M. N. *et al.* Suppression of amygdalar endocannabinoid signaling by stress
contributes to activation of the hypothalamic-pituitary-adrenal axis.
Neuropsychopharmacology **34**, 2733–2745 (2009).
33.   Fegley, D. *et al.* Characterization of the fatty acid amide hydrolase inhibitor
cyclohexyl carbamic acid 3'-carbamoyl-biphenyl-3-yl ester (URB597): effects on
anandamide and oleoylethanolamide deactivation. *J. Pharmacol. Exp. Ther.* **313**, 352–
358 (2005).
34.   Piomelli, D. *et al.* Pharmacological profile of the selective FAAH inhibitor KDS-
4103 (URB597). *CNS Drug Rev* **12**, 21–38 (2006).
35.   Hill, M. N. *et al.* Endogenous cannabinoid signaling is essential for stress
adaptation. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 9406–9411 (2010).
36.   Imperatore, R. *et al.* Genetic deletion of monoacylglycerol lipase leads to
impaired cannabinoid receptor CB1R signaling and anxiety-like behavior. *Journal of*
Neurochemistry **135**, 799–813 (2015).
37.   Sihag, J. & Jones, P. J. H. Dietary fatty acid profile influences circulating and
tissue fatty acid ethanolamide concentrations in a tissue-specific manner in male Syrian
hamsters. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*
1864, 1563–1579 (2019).
38.   Artmann, A. *et al.* Influence of dietary fatty acids on endocannabinoid and N-
acylethanolamine levels in rat brain, liver and small intestine. *Biochimica et Biophysica*
Acta (BBA) - Molecular and Cell Biology of Lipids **1781**, 200–212 (2008).

39.   Xie, Z. *et al.* Analysis of transcriptome and metabolome profiles alterations in
fatty liver induced by high-fat diet in rat. *Metabolism* **59**, 554–560 (2010).
40.   Lee, K. N., Pariza, M. W. & Ntambi, J. M. Conjugated Linoleic Acid Decreases
Hepatic Stearoyl-CoA Desaturase mRNA Expression. *Biochemical and Biophysical*
Research Communications **248**, 817–821 (1998).
41.   Oosterveer, M. H. *et al.* High Fat Feeding Induces Hepatic Fatty Acid Elongation
in Mice. *PLOS ONE* **4**, e6066 (2009).
42.   Massillon, D., Arinze, I. J., Xu, C. & Bone, F. Regulation of Glucose-6-
phosphatase Gene Expression in Cultured Hepatocytes and H4IIE Cells by Short-chain
Fatty Acids ROLE OF HEPATIC NUCLEAR FACTOR-4α. *J. Biol. Chem.* **278**, 40694–
40701 (2003).
43.   Chen, G. Liver lipid molecules induce PEPCK-C gene transcription and attenuate
insulin action. *Biochemical and Biophysical Research Communications* **361**, 805–810
(2007).
44.   Bercik, P. *et al.* Chronic gastrointestinal inflammation induces anxiety-like
behavior and alters central nervous system biochemistry in mice. *Gastroenterology* **139**,
2102-2112.e1 (2010).
45.   Lyte, M., Li, W., Opitz, N., Gaykema, R. P. A. & Goehler, L. E. Induction of
anxiety-like behavior in mice during the initial stages of infection with the agent of
murine colonic hyperplasia Citrobacter rodentium. *Physiol. Behav.* **89**, 350–357 (2006).
46.   Gray, J. M. *et al.* Corticotropin-Releasing Hormone Drives Anandamide
Hydrolysis in the Amygdala to Promote Anxiety. *J. Neurosci.* **35**, 3879–3892 (2015).
47.   Lee, T. T.-Y., Hill, M. N., Hillard, C. J. & Gorzalka, B. B. Temporal changes in
N-acylethanolamine content and metabolism throughout the peri-adolescent period.
Synapse **67**, 4–10 (2013).
48.   Romeo, R. D. *et al.* Pubertal shifts in adrenal responsiveness to stress and
adrenocorticotropic hormone in male rats. *Psychoneuroendocrinology* **42**, 146–152
(2014).
49.   Netherton, C., Goodyer, I., Tamplin, A. & Herbert, J. Salivary cortisol and
dehydroepiandrosterone in relation to puberty and gender. *Psychoneuroendocrinology* **29**,
125–140 (2004).
50.   Gunnar, M. R., Wewerka, S., Frenn, K., Long, J. D. & Griggs, C. Developmental
changes in hypothalamus–pituitary–adrenal activity over the transition to adolescence:
Normative changes and associations with puberty. *Dev Psychopathol* **21**, 69–85 (2009).
51.   Siegel, R. S. & Dickstein, D. P. Anxiety in adolescents: Update on its diagnosis
and treatment for primary care providers. *Adolescent Health, Medicine and Therapeutics*
3, 1 (2012).
52.   McLean, C. P., Asnaani, A., Litz, B. T. & Hofmann, S. G. Gender differences in
anxiety disorders: Prevalence, course of illness, comorbidity and burden of illness.
Journal of Psychiatric Research **45**, 1027–1035 (2011).
53.   Mayo, L. M. *et al.* Elevated anandamide, enhanced recall of fear extinction, and
attenuated stress responses following inhibition of fatty acid amide hydrolase (FAAH): a
randomized, controlled experimental medicine trial. *Biological Psychiatry* (2019)
doi:10.1016/j.biopsych.2019.07.034.

54.   Sarkar, J., Wakefield, S., MacKenzie, G., Moss, S. J. & Maguire, J.
Neurosteroidogenesis Is Required for the Physiological Response to Stress: Role of
Neurosteroid-Sensitive GABAA Receptors. *J. Neurosci.* **31**, 18198–18210 (2011).
55.   Brown, D. H., Jr., Giusani, A. D., Chen, X. & Kumamoto, C. A. Filamentous
growth of Candida albicans in response to physical environmental cues and its regulation
by the unique CZF1 gene. *Molecular microbiology* **34**, 651–62 (1999).
56.   Can, A. *et al.* The Mouse Forced Swim Test. *J Vis Exp* (2012) doi:10.3791/3638.
57.   Caporaso, J. G. *et al.* Ultra-high-throughput microbial community analysis on the
Illumina HiSeq and MiSeq platforms. *ISME J* **6**, 1621–1624 (2012).
58.   Morena, M. *et al.* Training-Associated Emotional Arousal Shapes
Endocannabinoid Modulation of Spatial Memory Retrieval in Rats. *J. Neurosci.* **35**,
13962–13974 (2015).
59.   Qi, M., Morena, M., Vecchiarelli, H. A., Hill, M. N. & Schriemer, D. C. A robust
capillary liquid chromatography/tandem mass spectrometry method for quantitation of
neuromodulatory endocannabinoids. *Rapid Commun. Mass Spectrom.* **29**, 1889–1897
(2015).
653
654
655
656

Figure Legends

Figure 1: Gastrointestinal colonization with *Candida albicans* after a single inoculation without disruption of bacterial gut microbiota or invasive disease

A) Timeline for acute colonization model. 5-week-old female mice were cohoused for four days and handled daily. Mice were then moved into smaller groups in standard cages and orally inoculated with 5×10^7CFU of *C. albicans* or given buffer. On day one post-inoculation, fecal pellets were sterilely collected from mice and plated on YPD-SA to measure colonization levels. On day two post-inoculation, all mice underwent a behavioral test. On day three post-inoculation mice were anesthetized with isoflurane and sacrificed by decapitation. Some mice were subjected to restraint stress prior to sacrifice. B) *C. albicans* CFU/g of fecal pellets was measured on day 1 post-inoculation and CFU/g of cecum contents was measured after sacrifice on day three post-inoculation. No culturable fungi were measured on either day from mock-colonized mice. Day 1: N=56, Day 3: N=50. Figure includes data from 6 cohorts. Bars indicate geometric mean. C) A multiplex ELISA was used to measure inflammatory cytokines in the serum of mice. N.D, not detectable. Student's t-test, $p < 0.05$. Mock-colonized N=13, *C. albicans*-colonized N=14. Figure includes data from 3 cohorts. D-F) Microbiota analysis of the cecum contents of mice was performed using standard 16s rRNA DNA sequencing and QIIME analysis pipeline. Mock-colonized N=8 and *C. albicans*-colonized N=8. Figure includes data from 2 cohorts. D) The relative abundance of all bacterial families detected with a median relative abundance greater than 0 is shown. Bar shows geometric mean with SEM E) The average alpha diversity metric Chao1 was calculated to determine the diversity of the microbiota at different levels of sampling. The average Chao1 score of the experimental groups is shown. F) Beta diversity was calculated using weighted UniFrac scores and principal coordinates analysis was performed. For C-D, symbols indicate individual mice and bars indicate the average. Mock-colonized mice are represented as solid dots and bars; *C. albicans*-colonized mice are represented as open circles and bars.

Figure 2: *C. albicans* colonization increases anxiety-like behavior

On day two post-inoculation mice underwent either the Elevated Plus Maze (EPM) or the Forced Swim Test (FST). Trials in the behavioral tests were video-recorded and scored by a blinded observer after the fact. For the EPM data (A-F) that follows, mock-colonized N=22 and *C. albicans*-colonized N=19 and statistical analysis performed was Welch's t-test, $p<0.05$. Figures include data from two cohorts. A) Percentage of time spent in the open arms of the EPM. B) Total number of entries into the open arms of the EPM. C) Percentage of time spent in the closed arms of the EPM. D) Total entries into the closed arms of the EPM. E) Percentage of time spent in the neutral central square of the EPM (neither open nor closed arms). F) Total entries during the trial (sum of the open and closed arm entries), a metric for locomotor activity. G) Total time spent immobile during the six-minute FST trial. Immobility was defined as no movements beyond those required to stay afloat. For FST data: mock-colonized N=33, *C. albicans*-colonized N=24. Figure includes data from three cohorts. Throughout, symbols represent individual mice and bars indicate average. Mock-colonized mice are represented as solid dots and bars; *C. albicans*-colonized mice are represented as open circles and bars.

Figure 3: *C. albicans* colonization increases basal CORT and alters feedback inhibition after acute stress

A) Corticosterone (CORT) in trunk blood of mice sacrificed without stress was measured by ELISA. Mock-colonized N=15, *C. albicans*-colonized N=19. Figure includes data from four cohorts. Data was analyzed using Welch's t-test, p<0.05 B) CORT in trunk blood of mice sacrificed at different times of day. In addition to the data summarized in A (mice sacrificed from 13:00-16:00), B includes mice sacrificed without stress from 17:00-18:00. Data was binned in one-hour increments. Mock-colonized: 13:00 N=5, 14:00 N=9, 15:00 N=11, 16:00 N=8, 17:00 N=9, 18:00 N=5. *C. albicans*-colonized: 13:00 N=6, 14:00 N=13, 15:00 N=10, 16:00 N=11, 17:00 N=8, 18:00 N=9. C) Mice were subjected to 30 minutes of restraint stress and then sacrificed immediately (0m recovered) or after 30 minutes or 60 minutes of recovery as shown on the x-axis. Mock-colonized: 0m N=14, 30m N=15, 60m N=13. *C. albicans*-colonized: 0m N=15, 30m N=13, 60m N=12. D) Immunohistochemistry was used to visualize cFos protein with Alexa488 in the paraventricular nucleus (PVN) of mice using the 20X objective. DAPI was used to visualize nuclei and Alexa488 to visualize cFOS. The PVN of mice sacrificed unstressed (left) do not exhibit cFOS staining, while PVN of mice sacrificed after 30 minutes of stress and 30 minutes of recovery (right) are positive for cFOS. Representative images are shown. For all figures, symbols indicate individual mice and bars indicate the average. Mock-colonized mice are solid dots and bars and *C. albicans*-colonized mice are open circles and bars.

Figure 4: Neuroendocrine phenotypes of the *C. albicans*-colonized mice are
mediated through disruption of the endocannabinoid system
A-C) Mice were treated with the fatty acid amide hydrolase (FAAH) inhibitor
URB597, a well-characterized drug which increases AEA levels by blocking degradation.
Mice were given either URB597 or vehicle control by intraperitoneal injection 4-6 hours
prior to sacrifice (A) or testing in the EPM (B-C). A) Mice were treated with 1mg/kg of
URB597 or vehicle. ELISA was used to measure CORT in trunk blood of mice
sacrificed without stress. Welch's t-test, $p < 0.05$. Mock-colonized: vehicle N=15,
URB597 N=19. *C. albicans*-colonized: vehicle N=16, URB597 N=18. Figure includes
data from four cohorts. B-C) Mice were treated with 0.1mg/kg (triangles) or 0.15mg/kg
(circles) URB597 or vehicle, then tested in the EPM. Mock-colonized: vehicle N=17,
0.1mg/kg URB597 N=4, 0.15mg/kg URB597 N=8. *C. albicans*-colonized: vehicle
N=13, 0.1mg/kg URB597 N=4 0.15mg/kg URB597 N=8. B) Percentage time spent in
the open arms, normalized to the average percent open time of the mock-colonized
vehicle control, is shown. Welch's t-test, $p < 0.05$. C) Total arms entries, a metric for
overall locomotor activity is shown. D) AEA (pmol/g tissue) was measured in forebrain
samples by mass spectrometry. E-F) Linear regression of basal serum CORT and
forebrain AEA (D) in individual mice are shown. D-F Mock-colonized N=7, *C.*
albicans-colonized N=9. Untargeted mass spectrometry was used to measure relative
abundance of 735 compounds in the cecum contents of mice. N-acylethanolamides (E-
G), free fatty acids (H-I) and lysophospholipids (J) containing an 18-carbon chain with
varying degrees of unsaturation are shown here. Raw data was log-transformed. Mock-
colonized N=8, *C. albicans*-colonized N=8. Student's t-test, $p < 0.05$. Symbols indicate
individual mice and bars indicate average. Solid circles and bars are mock-colonized
mice and open circles and bars are *C. albicans*-colonized mice.

Figure 5: *C. albicans* colonization affects hepatic lipid-responsive gene expression

RNA was extracted from the liver of mice sacrificed unstressed on day three. cDNA was synthesized and qRT-PCR was used to quantify liver gene transcription. RNA gene expression is normalized to GAPDH and expressed relative to the average expression of the mock-colonized mice. A) Stearoyl-CoA desaturase 1 (*Scd1*) RNA expression in the liver is shown. Mock-colonized N=24, *C. albicans*-colonized N=27. Figure includes data from four cohorts. B) Expression of multiple lipogenic genes is shown: acyl-CoA 8-3 desaturase (*Fads1*), acyl-CoA 6 desaturase (*Fads2*), sterol regulatory element-binding protein 1 (*Srepbf1*) and fatty acid synthase (*Fasn*). Mock-colonized N=11, *C. albicans*-colonized N= 10. Figure includes data from two cohorts. C) Expression of the gluconeogenesis enzymes phosphoenolpyruvate carboxykinase (*Pck1*) (mock-colonized N=24, *C. albicans*-colonized N=23) and D) glucose-6-phosphatase (*G6Pc*) is shown (mock-colonized N=20, *C. albicans*-colonized N=18). Figures include data from four cohorts. Symbols represent individual mice and bars indicate average. Solid symbols and bars indicate mock-colonized mice and open symbols and bars indicate *C. albicans*-colonized mice.

    Figures

773   **Figure 1: Gastrointestinal colonization with *Candida albicans* after a single**

774   **inoculation without disruption of bacterial gut microbiota or invasive disease**

777 **Figure 2: *C. albicans* colonization increases anxiety-like behavior**

Figure 3: *C. albicans* **colonization increases basal CORT and alters feedback**
inhibition after acute stress

784 **Figure 4: Neuroendocrine phenotypes of the *C. albicans*-colonized mice are**
785 **mediated through disruption of the endocannabinoid system**

786
787
788

789 **Figure 5: *C. albicans* colonization affects hepatic lipid-responsive gene expression**