JACKIE: Fast enumeration of single- and multi-copy CRISPR binding sites

Jacqueline Jufen Zhu[1], Albert Wu Cheng[1,2,3,4,5]

1.   The Jackson Laboratory for Genomic Medicine, Farmington, CT 06032, USA

2.   The Jackson Laboratory Cancer Center, Bar Harbor, ME 04609, USA

3.   Department of Genetics and Genome Sciences, University of Connecticut Health

Center, Farmington, CT 06030, USA

4.   Institute for Systems Genomics, University of Connecticut Health Center, Farmington,

CT 06030, USA

5.   Correspondence: albert.cheng@jax.org

ABSTRACT

Summary: CRISPR-based methods for genome, epigenome editing and imaging have provided

powerful tools to interrogate the functions of the genome. The design of guide RNA (gRNA) is a

vital step of CRISPR experiments. We report here the implementation of JACKIE (Jackie and

Albert's CRISPR K-mer Instances Enumerator), a pipeline for enumerating all potential single-

and multi-copy CRISPR sites in the genome. We demonstrate the application of JACKIE to

identify locus-specific repetitive sequences for CRISPR/Casilio-based genomic labeling.

Availability: Source codes and CRISPR site databases (JACKIEdb) for hg38 and mm10 are

available for download at http://crispr.software/JACKIE

1. INTRODUCTION

CRISPR/Cas technologies have revolutionized genome research by providing versatile tools for

"writing" the genome, epigenome and transcriptome as well as "reading" the dynamics of the

genome through live cell genomic imaging techniques (Sander and Joung, 2014). Genomic

imaging by CRISPR/Cas-based technologies require clustered repetitive sequences (Cheng, et

al., 2016; Ma, et al., 2018; Maass, et al., 2018; Qin, et al., 2017). dCas9-based gene activation

and epigenome editing also benefited from synergistic binding events (Cheng, et al., 2016;

Cheng, et al., 2013; Maeder, et al., 2013; Mali, et al., 2013; Taghbalout, et al., 2019). CRISPR-

mediated gene deletion may be induced using a pair of gRNAs flanking the region of interest

(Canver, et al., 2017) while other CRISPR applications may benefit from a database of unique

CRISPR binding sites.

For fast enumeration of all potential CRISPR binding sites and their locations of

occurrences in the genome, we have implemented JACKIE (Jackie and Albert's CRISPR K-mer

Instances Enumerator) pipeline that is compatible with high performance computing (HPC)

clusters. We further provide helper scripts for extracting binding sites within regions of interest,

as well as a wrapper script to run Cas-OFFinder (Bae, et al., 2014) to identify off-target profiles

of selected CRISPR binding sites.

2. IMPLEMENTATION

JACKIE first scans the genome and enumerates any potential SpCas9 binding sites with the

PAM constraint (nGG). To allow memory and computation efficiency, sequences are encoded

into an unsigned 64-bit integer (NucKey) by encoding each nucleotide in 3 bits (Fig 1). The

chromosome and location for each instance are also encoded as integer values. The NucKey,

chromosome and location (KeyedPosition) are written into files prefixed by a predefined prefix

length (default to 6). The second step: Each bin file is loaded into a list in memory, and sorted

by NucKey, thus placing records of CRISPR sites with identical sequences adjacent to each

other in the list. The sequence-sorted list is then traversed to output CRISPR sites in bed

interval files. The extensive divide-and-conquer design allows JACKIE to be highly parallelized

on an HPC cluster (Fig 1).

Helper scripts were written to take the bed file outputs from JACKIE to further collapse

same-chromosome CRISPR binding sites into an extended bed interval file format with each

record containing the CRISPR binding sites encoded as blocks. Such extended bed files can

then be loaded onto genome browsers for visualization. Other helper scripts were written for

downstream processing, such as for the fast extraction of CRISPR sites overlapping defined set

of intervals and automated queries into an offline version of Cas-OFFinder (Bae, et al., 2014) to

compute off-target profiles for selected sgRNA sequences. sgRNA sequences, number of

occurrences, and span sizes (the distance between the 5' most site to 3' most site for CRISPR

sites on the same chromosome) are encoded in the output files so that CRISPR sites can be

filtered by simple text processing software (e.g., awk) to identify sites for specific purposes. We

used JACKIE to generate CRISPR binding site databases (JACKIEdb) for human genome

version hg38 and mouse genome mm10, which took ~35min and ~22min, respectively, on an

HPC cluster with 2,568 Intel Xeon compute cores and >38TB of system memory.

3. USE CASES

Some potential applications of JACKIE pipeline are illustrated in Fig 2. For example, highly

repetitive sequences (>30 copies) and low repetitive arrays (~6 copies) can be used for imaging

applications (Fig 2a) (Cheng, et al., 2016; Ma, et al., 2018; Maass, et al., 2018; Qin, et al., 2017).

These sites can be identified from JACKIE output by filtering for the desired range of repeat

number and span size. To demonstrate such use case, we have selected three CRISPR binding

sequences with >30 copies clustered specifically within 10kb region. We then applied Casilio

imaging to visualize the sequence clusters via fluorescence microscopy (Fig 2b), producing 3

fluorescence foci in each case, consistent with the ploidy of the HEK293T cells used in the

experiments. Similarly, promoter-specific repeats might be identified (e.g., >3 copies, within 1kb)

overlapping gene promoters for synergistic gene activation or repression by CRISPRa or

CRISPRi technologies (Fig 2c) (Cheng, et al., 2016; Cheng, et al., 2013; Maeder, et al., 2013;

Mali, et al., 2013; Taghbalout, et al., 2019). Two or more copy sites flanking or spanning a

region or interest might be identified for inducing deletion or complex chromosome

rearrangements (Fig 2d). (Canver, et al., 2017; Choi and Meyerson, 2014; Maddalo, et al., 2014;

Maresch, et al., 2016).

Although currently designed for SpCas9 binding site enumeration, JACKIE can be

modified to identify binding sites for other CRISPR/Cas systems by changing the code segment

written for the PAM requirement filtering. Alternatively, PAM requirement might be removed from

the codes to enable identification of all k-mer sites and their occurrences in the genome. The

use of one uint64 for encoding NucKey limits the enumeration of sequences up to 20mer. By

using multiple NucKey, or a variable-size vector of NucKey, longer k-mer sequences can be

enumerated.

Experimental Material & Methods

sgRNA spacer sequences (as listed in Fig 2b) were cloned into sgRNA-15xPBSc expression

vector via an oligo-annealing protocol as previously described (Cheng, et al., 2016). HEK293T

cells were cultivated in Dulbecco's modified Eagle's medium (DMEM) (Sigma) with 10% fetal

bovine serum (FBS)(Lonza), 4% Glutamax (Gibco), 1% Sodium Pyruvate (Gibco) and penicillin-

streptomycin (Gibco). Incubator conditions were 37 °C and 5% CO_2. Cells were seeded into 24-

well plates the day before being transfected with 50ng pAC1445-dCas9 plasmid (Addgene

#73169), 25ng pAC1447-Clover-PUFa plasmid (Addgene #73689) and 250ng sgRNA-15xPBSc

plasmid. Cells were fixed, mounted on slides, and imaged 48 hours after transfection on a Leica

SP8 confocal microscope.

Acknowledgement

This work has been supported by the Jackson Laboratory internal grants (to A.C.) and National

Human Genome Research Institute 1R01HG009900 (to A.C).

Figure Legends

Fig 1. Implementation of JACKIE. (a) JACKIE first scans the genome for potential SpCas9

binding sites, and encode each occurrence in a data structure called KeyedPosition, consisting

of NucKey (uint64), chrID (uint32), and position (int32). NucKey records a binary representation

of the 20-mer spacer sequence using 3 bits per nucleotide. chrID records an integer

representing a chromosome. A chrID-chromosome mapping file is generated automatically.

Position records the location of the binding site, with negative and positive integers representing

the minus strand and positive strand, respectively. JACKIE outputs KeyedPosition as it scans

the genome. For efficient sorting in the next step, KeyedPosition records are appended to a file

named by the 6-mer prefix (<6merPrefix>.bin). To parallelize this step, four processes are

started as separate cluster jobs, each focusing on the A, C, T, or G as the first nucleotide of

binding sites. **(b)** In the second step, JACKIE loads each <6merPrefix>.bin file and performs a

sort on the KeyedPosition records on the NucKey variable using std_sort routine and then

traverse the sorted list of KeyedPosition to output a bed interval file containing the coordinate

position   of   each   CRISPR   sites   with   item   name   encoded   by

<NucKey>.<copyNumber>/<sequence>,   and   score   field   recording   the   copy   number.

Parallelization is achieved by either starting separate job on individual <6merPrefix>.bin files, or

by starting batch jobs focusing on 2mer prefices (e.g., AA AT AC AG operating on AA*.bin,

AT*.bin, AC*.bin, AG*.bin, etc). The outputs (<6merPrefix>.bed) are then merged into a

combined bed file using cat unix command. Helper scripts are provided for downstream

processing of the bed file into collapsed bed file with each record encoding all binding sites of

the same sequence in the same chromosome, for extracting binding sites within defined

regions, or for running the Cas-OFFinder program to identify off-target profiles of selected

sequences.

Fig 2. Potential applications of JACKIE pipeline. (a) JACKIE can be used to identify

clustered CRISPR binding sites with high copy number or low copy number for CRISPR-based

genomic imaging experiments. **(b)** To demonstate genomic imaging use case, we filtered

sgRNA sequences with >30 copies clustered within specific <10kb regions and performed

Casilio imaging experiment using designed sgRNAs. UCSC genome browser screenshots are

shown on the left with the line-bar annotating the CRISPR sites, and fluorescence microscopy

images shown on the right. Microscopy images were derived by merging green (Casilio-

labeling) and blue (DAPI: nucleus stain). Arrows point to fluorescent loci of interest. Scale bars

indicate 5μm. **(c)** JACKIE can be used to identify clustered CRISPR binding sites for synergistic

activation or repression of target cis-regulatory elements (e.g., promoters) using CRISPR-based

epigenetic editing approaches (e.g., CRISPRa or CRISPRi). **(d)** JACKIE can identify unique

CRISPR binding sites in the genome for precise indel induction, dual CRISPR sites for targeted

deletion, or clustered sites spanning a genomic region to induce complex rearrangement

events.

Figure 2

a. High- and low-repeat imaging

Copy number>30 (high)
or ~6 (low)
Span size<10kb

b. Examples sgRNA design for imaging (hg38, >30 copies, <10kb)

i. GGTGTAGACAGTGGAGCAGC, chr10:1046518-1047610, 33 copies

ii. TGAGGACTGAATCTAGAACC, chr18:79033234-79037027, 52 copies

No gene features

iii. ACTAACACTGCAGCGACCCC, chr4 744703-747683, 53 copies

c. Synergistic CRISPRa/i

Copy number>=3
Span size<1kb
Overlap promoter

Gene X

d. Genome editing applications

Copy number=1

Copy number=2

Copy number>2

indels

deletion

Rearrangements

References

Bae, S., Park, J. and Kim, J.S. Cas-OFFinder: a fast and versatile algorithm that searches for

potential off-target sites of Cas9 RNA-guided endonucleases. *Bioinformatics (Oxford, England)*

2014;30(10):1473-1475.

Canver, M.C.*, et al.* Characterization of genomic deletion efficiency mediated by clustered

regularly interspaced short palindromic repeats (CRISPR)/Cas9 nuclease system in mammalian

cells. *The Journal of biological chemistry* 2017;292(6):2556.

Cheng, A.W.*, et al.* Casilio: a versatile CRISPR-Cas9-Pumilio hybrid for gene regulation and

genomic labeling. *Cell research* 2016;26(2):254-257.

Cheng, A.W.*, et al.* Multiplexed activation of endogenous genes by CRISPR-on, an RNA-guided

transcriptional activator system. *Cell research* 2013;23(10):1163-1171.

Choi, P.S. and Meyerson, M. Targeted genomic rearrangements using CRISPR/Cas

technology. *Nature communications* 2014;5:3728.

Ma, H.*, et al.* CRISPR-Sirius: RNA scaffolds for signal amplification in genome imaging. *Nature*

methods 2018;15(11):928-931.

Maass, P.G.*, et al.* Spatiotemporal allele organization by allele-specific CRISPR live-cell

imaging (SNP-CLING). *Nature structural & molecular biology* 2018;25(2):176-184.

Maddalo, D.*, et al.* In vivo engineering of oncogenic chromosomal rearrangements with the

CRISPR/Cas9 system. *Nature* 2014;516(7531):423-427.

Maeder, M.L.*, et al.* CRISPR RNA-guided activation of endogenous human genes. *Nature*

methods 2013;10(10):977-979.

Mali, P.*, et al.* CAS9 transcriptional activators for target specificity screening and paired

nickases for cooperative genome engineering. *Nature biotechnology* 2013;31(9):833-838.

Maresch, R.*, et al.* Multiplexed pancreatic genome engineering and cancer induction by

transfection-based CRISPR/Cas9 delivery in mice. *Nature communications* 2016;7:10770.

Qin, P.*, et al.* Live cell imaging of low- and non-repetitive chromosome loci using CRISPR-Cas9.

Nature communications 2017;8:14725.

Sander, J.D. and Joung, J.K. CRISPR-Cas systems for editing, regulating and targeting

genomes. *Nature biotechnology* 2014;32(4):347-355.

Taghbalout, A.*, et al.* Enhanced CRISPR-based DNA demethylation by Casilio-ME-mediated

RNA-guided coupling of methylcytosine oxidation and DNA repair pathways. *Nature*

communications 2019;10(1):4296.

174