Synthetic Agonist of Growth Hormone-Releasing Hormone as Novel Treatment for Heart Failure with Preserved Ejection Fraction

Raul A. Dulce[1], Rosemeire M. Kanashiro-Takeuchi[1,2], Lauro M. Takeuchi[1], Alessandro G. Salerno[1], Shathiyah Kulandavelu[1,6], Wayne Balkan[1,3], Marilia S.S.R. Zuttion[1], Renzhi Cai[4], Andrew V. Schally[3,4,5] and Joshua M. Hare[1,3]*

Affiliations:

[1]Interdisciplinary Stem Cell Institute, University of Miami Miller School of Medicine, Miami, Florida 33136, USA.

[2]Department of Molecular and Cellular Pharmacology, University of Miami Miller School of Medicine, Miami, Florida 33136, USA.

[3]Department of Medicine, University of Miami Miller School of Medicine, Miami, Florida 33136, USA.

[4]Endocrine, Polypeptide and Cancer Institute, Veteran Affairs Medical Center, Florida 33136, USA.

[5]Division of Hematology/Oncology, University of Miami Miller School of Medicine, Miami, Florida 33136, USA.

[6]Department of Pediatrics, University of Miami Miller School of Medicine, Miami, Florida 33136, USA.

*Corresponding Author: Joshua M. Hare, M.D. Director, Interdisciplinary Stem Cell Institute. Biomedical Research Building, 1501 N.W. 10th Avenue, Room 924, Miami, FL 33136, USA. Phone: +1-305-243-1999; Fax: +1-305-243-3906. Email: jhare@med.miami.edu.

Abstract

Heart failure with preserved ejection fraction (HFpEF) is characterized by a myocardium with impaired relaxation, fibrosis and ventricular stiffness. Despite the rapidly increasing prevalence of HFpEF, no effective therapies have emerged. Here we show that a potent synthetic agonist of growth hormone-releasing hormone (GHRH-A) both prevents and reverses the HFpEF phenotype generated in mice through continuous infusion of angiotensin-II (Ang-II). Animals treated with Ang-II had diastolic dysfunction, and isolated cardiomyocytes (ex vivo) exhibited incomplete relaxation, depressed contractile responses and altered myofibrillar protein phosphorylation. Calcium handling mechanisms were disturbed in cardiomyocytes from mice with HFpEF. The GHRH-A MR-356 prevented the development of the pathological phenotype and reversed the phenotype in vivo and ex vivo in mice with established HFpEF. Taken together these findings indicate that the GHRH receptor signaling pathway represents a new molecular target to counteract HFpEF-associated cardiomyocytye dysfunction by targeting myofilament phosphorylation. Accordingly, activation of the GHRH receptor with potent synthetic GHRH agonists may provide a novel therapeutic approach to management of HFpEF syndrome.

Heart failure (HF) remains a leading cause of death worldwide and is associated with two general clinical variants - heart failure with reduced ejection fraction (HFrEF), characterized by systolic dysfunction, and heart failure with preserved ejection fraction (HFpEF), predominantly due to diastolic dysfunction[1-3]. HFpEF accounts for ~50% of HF cases, a proportion that is predicted to increase at about 1% per year relative to HFrEF[4, 5]. Whereas there are numerous approved and guideline supported drugs for HFrEF, there are no approved therapies targeting HFpEF, and numerous clinical trials with several classes of drugs have failed to produce clinical benefits[6-10]. The HFpEF syndrome is characterized by endothelial dysfunction, hypertension, impaired relaxation and contractile reserve, ventricular stiffening and fibrosis[2, 11, 12]. The underlying molecular mechanisms of HFpEF are poorly understood due to its diverse etiology, which in most cases involves multi-organ pathological conditions[13-15]. Importantly, current therapeutic strategies fail to stop the progression of HFpEF[6, 8, 16-18], creating a healthcare crisis due to the unmet therapeutic need for HFpEF patients[19, 20].

The development of new potent synthetic growth hormone-releasing hormone agonists (GHRH-As) and their application in different models of myocardial injury (either ischemic[21-25] or non-ischemic[26]) demonstrates the effectiveness of myocardial GHRH receptors (GHRHRs) activation to reduce fibrosis or hypertrophy, respectively, and improve cardiac performance[22-26]. The signaling of GHRHR extends beyond the hypothalamic-pituitary axis[27] as the GHRHRs are present on cardiomyocytes[22, 23]. Based upon the effects of GHRH-As on myocardial fibrosis, hypertrophy and performance, we hypothesized they would improve cardiomyocyte function in HFpEF.

Among the limited number of animal models that reproduce the features of HFpEF observed in humans[14, 28-31], chronic infusion of a low dose of angiotensin II (Ang-II) in mice, reproduces the HFpEF cardiac phenotype in the absence of elevated systemic arterial blood pressure[32]. We used this model to study the effects of a GHRH-A, MR-356, on the performance of cardiomyocytes *ex vivo* and *in vivo*. Synthetic MR-356 is about 150 times more potent[21] than human GHRH(1-29)NH$_2$ and binds to the GHRHR with 7.5 fold greater affinity. Here we tested the hypotheses that activation of GHRHR signaling within the myocardium both prevents and reverses myocardial HFpEF features in vivo through mechanisms involving both myocardial fibrosis and impaired cardiomyocyte relaxation.

Results.

GHRH-A prevents the appearance of the HFpEF syndrome induced by in vivo Ang-II infusion.

Four-weeks of continuous infusion of Ang-II induced a cardiovascular phenotype that mimicked the phenotypic features of HFpEF[32]. Specifically, Ang-II-treated mice exhibited myocardial hypertrophy with increased heart weight normalized by body weight or tibia length (Figure 1A and Online Supplemental Table 1), increased relative wall thickness in diastole (RWTd) and increased cardiomyocyte width (Figure 1B-C) without changes in length (Online Supplemental Figure 1). Myocardial fibrosis was increased in the HFpEF hearts as evidenced by collagen III content, the transforming growth factor-β (TGF-β, a fibrosis mediator) and the myofibroblast biomarker α-smooth muscle actin (α-SMA) (Figure 1D-F). We also observed a trend to increase both the prolyl-4-hydroxylase-α-1 (P4HA1, a key enzyme for the hydroxylation of proline residues on collagen), and the hydroxyproline residues (a modified amino acid residue characteristic of the collagen primary structure) (Online Supplemental Figure 2A-B). Whereas collagen I and vimentin (a biomarker for undifferentiated fibroblasts) did not change (Online Supplemental Figure 2C-D). Co-administration of the GHRH analog, MR-356 prevented the increases in RWTd and cardiomyocyte width seen in the Ang-II treated group (Figure 1B-C). Moreover, MR-356 treatment protected against fibrosis, since collagen III content, TGF-β and α-SMA levels were reduced (Figure 1D-F).

Echocardiographic evaluation demonstrated that Ang-II infusion did not reduce ejection fraction (EF) (Online Supplemental Tables 2), either alone or under co-treatment with MR-356. Likewise, chamber dimensions were unaffected in systole between groups (Supplemental Table 2). Using invasive hemodynamic assessment (Table 1), we found increased arterial blood pressure, a characteristic of HFpEF, and clear evidence of diastolic dysfunction with increased end-diastolic pressure-volume relationship (EDPVR) (Figure 2A-D), relaxation time constant (tau) and left ventricular end-diastolic pressure (LVEDP) in Ang-II-treated mice compared to control (Figure 2E-F). Isovolumic relaxation time (IVRT) increased along the time course of the treatment (Figure 2H) (within group comparison - 4-week versus baseline), and this increase was greater in the Ang-II-treated group (as compared versus controls at 4-week time point) consistent with the initial description of this model[32]. Global longitudinal strain (GLS) increased (a

reduction in absolute values) with Ang-II treatment, similar to that observed in diastolic

dysfunction[33] (Figure 2I and Supplemental Table 2). Co-infusion of MR-356 completely

prevented the development of diastolic dysfunction (Figure 2C-I) and induced a robust

acceleration in dP/dt$_{min}$ (Figure 2G). Taken together, these results demonstrate that stimulation of

GHRH pathway with the potent agonist MR-356 effectively counteracted the development of

HFpEF induced by Ang-II.

GHRH-A prevents sarcomere and contractile dysfunction in HFpEF cardiomyocytes

In order to test whether GHRH agonists directly affect cardiomyocyte function, we

studied cardiomyocytes isolated from Ang-II-treated mice, which were electric field-stimulated

at 1, 2 and 4 Hz. They exhibited shorter resting sarcomere length compared to those from control

animals (Figure 3A and B), indicating an inability to relax, consistent with HFpEF[11, 32].

Quiescent HFpEF cardiomyocytes exposed to 1.8 mmol/L extracellular $CaCl_2$ were consistently

shorter compared to control cardiomyocytes, while there was no difference between groups in

Ca^{2+}-free buffer in the presence of 2,3-butanedione monoxime (BDM) (Online Supplemental

Figure 3A). HFpEF cardiomyocytes also exhibited depressed contractile response to pacing

(Figure 3A and C). However, concomitant treatment with MR-356 during Ang-II infusion, not

only prevented the impairment in sarcomere tension at resting intervals between twitches at all

frequencies (Figure 3A and B), but also preserved quiescent cardiomyocyte length in the

presence of extracellular Ca^{2+} (Online Supplemental Figure 3A, right panel). Treatment with

MR-356 also preserved sarcomere shortening and response to pacing at levels equivalent to those

in control cardiomyocytes (Figure 3A and C). Paradoxically, shorter sarcomere length in HFpEF

cardiomyocytes were associated with lower $[Ca^{2+}]$ at resting intervals (Figure 3D and E, *p*=0.028

compared with controls), which was prevented by treatment with MR-356. These findings

suggest a possible alteration in sensitivity of myofilaments for Ca^{2+}. There were no significant

differences in Ca^{2+} transient amplitude ($\Delta[Ca^{2+}]$) between groups (Figure 3D and F), but there

was a positive response to pacing in peak $[Ca^{2+}]$ in control cardiomyocytes while HFpEF

cardiomyocytes exhibited a depressed response (Online Supplemental Figure 3B). MR-356

reestablished the increasing $[Ca^{2+}]$-frequency relationship (Online Supplemental Figure 3B-C),

which is clearly shown by the normalized $\Delta[Ca^{2+}]$ in response to pacing, with respect to their

amplitude at 1 Hz (Figure 3G). The HFpEF cardiomyocytes exhibited a flat response compared

with the other groups (*p*<0.01), which correlates with their contractile response (Figure 3C).

In order to further characterizing this abnormal response of intracellular Ca^{2+} to pacing,

we assessed sarcoplasmic reticulum (SR) Ca^{2+} leak by using tetracaine (see Supplemental

methods and Figure 4A) and studied the SR Ca^{2+} leak-load relationship in cardiomyocytes.

Increased SR Ca^{2+} leak is a hallmark of heart failure. We found that Ang-II-treated myocytes

exhibited a steeper rise compared with controls ($[Ca^{2+}]$ constants k_{AngII}=0.0117±0.0011 µmol^{-1}.L

vs. $k_{control}$=0.0047±0.0029 µmol^{-1}.L, $p<0.05$) and, although it remained elevated at low SR Ca^{2+}

loads, the load-leak curve was flattened by treatment with MR-356 ($k_{AngII+MR-356}$=0.0016±0.0009

µmol^{-1}.L, $p<0.0001$ vs. Ang-II-HFpEF; Figure 4B). Thus, at matched SR Ca^{2+} load (SR Ca^{2+}

load =99.1±11.5 µmol/L), the Ca^{2+} leak was elevated in Ang-II-induced HFpEF cardiomyocytes

and partially prevented by treatment with MR-356 (Figure 4C).

Effect of GHRH-A on the impaired Ca^{2+} decline in HFpEF cardiomyocytes

This murine model of HFpEF is associated with a subtle but consistent reduction in Ca^{2+}

decay rate. We compared the time to different phases of the Ca^{2+} decay (time to 10%, tt10; time

to 50%, tt50; tau and time to 90%, tt90). At 1 Hz, we found slower $[Ca^{2+}]$ decay in HFpEF

cardiomyocytes compared with controls (Figure 5A and B). Treatment with MR-356 blocked the

delayed Ca^{2+} decline at tt90% in the Ang-II+MR-356 (Figure 5A and B). The main mechanism

involved in Ca^{2+} decay in mice is the re-uptake into the SR. Despite the slower Ca^{2+} decay in

Ang-II-treated cardiomyocytes, the Ca^{2+} content in the SR, estimated by a rapid caffeine pulse,

was not reduced compared to control. Ang-II+MR-356 co-treatment did increase the SR $[Ca^{2+}]$

in comparison to control cardiomyocytes (Figure 5C). To elucidate the molecular mechanisms

involved in such effects, we investigated the proteins participating in Ca^{2+} handling. Re-uptake

of cytosolic Ca^{2+} into the SR occurs through the SR Ca^{2+} ATPase, SERCA2. There was reduced

expression of SERCA2a protein in cardiomyocytes from Ang-II-treated mice (Figure 5D).

Phospholamban (PLB), a small Ca^{2+}-cycling regulatory protein that exerts an inhibitory effect on

SERCA2 by protein-protein interaction, can be phosphorylated at serine 16 (p-Ser16) or

threonine 17 (p-Thr17), and so releasing the repression on SERCA2a activity. We observed

increased p-Ser16 and p-Thr17 PLB following Ang-II treatment, which was further increased by

co-treatment with MR-356 (Figure 5E and F). We also assessed the protein expression of a

subunit of the L-type Ca^{2+} channel (L-TCC 1αC), an important voltage-induced entrance of Ca^{2+}

to the cardiomyocyte, which plays a main role in the excitation-contraction coupling. L-TCC

1αC was increased by Ang-II, suggesting a possible Ca^{2+} overload in the HFpEF model. MR-356

treatment partially prevented this molecular change (Online Supplemental Figure 4A). The

electrogenic Na^{+}/Ca^{2+} exchanger (NCX) plays a discrete role in the Ca^{2+} decay rate in murine

cardiomyocytes, but is primarily responsible for Ca^{2+} removal. NCX gains relevance under some

pathologic heart conditions, where its expression may appear to be up-regulated. We tested

protein expression of the cardiac isoform, NCX1, which was unchanged by Ang-II treatment but

was reduced by MR-356 co-treatment compared with control cardiomyocytes (Online

Supplemental Figure 4B).

Impaired cardiomyocyte relaxation in HFpEF is improved by GHRH-A

Myocardial stiffness and deficient relaxation are hallmarks of the HFpEF syndrome,

which may be due to factors external to the cardiomyocyte, such as fibrosis, or intrinsically

associated with sarcomeric proteins, such as defects in titin structure/function or changes in

sensitivity of myofilaments to Ca^{2+}. At the cardiomyocyte level, this Ang-II-induced HFpEF

model exhibited reduced ability to relax mainly during the latter part of the relaxation phase (tau

and tt90%, Figure 6A and B) in a greater extent than our observations on Ca^{2+} decay. Treatment

with MR-356 not only prevented the prolongation of these parameters in the Ang-II-treated

model, but also significantly improved all phases of sarcomere relaxation at all frequencies

(Figure 6A and B). The exaggerated effect of MR-356 on sarcomere relaxation in comparison

with that on Ca^{2+} cycling might imply direct targeting of myofilament proteins. Faster relaxation

is associated with desensitization of myofilaments to Ca^{2+}. Accordingly, we analyzed SL-$[Ca^{2+}]$

hysteresis loops of cardiomyocyte contractions induced by electric-field stimulation, in order to

detect differences in responsiveness to Ca^{2+}. In general terms, the morphology of the loops was

greatly affected in the HFpEF model (Figure 6C). In the relaxation phases (Figure 6C and D

left), there was chronic Ang-II-induced sensitization of myofilaments ($p<0.0001$, Figure 6D

right) consistent with the delayed Ca^{2+} decline in this HFpEF model. As expected, co-treatment

with MR-356 induced a robust desensitizing effect that correlated with the exacerbated lusitropic

effect of MR-356 on isolated cardiomyocytes (Figure 6D). Cardiac troponin I (cTnI) plays an

important role in regulating responsiveness of myofilaments to Ca^{2+}, and its phosphorylation is a

determining factor in the affinity of troponin C (TnC) for Ca^{2+}. We explored phosphorylation on

sensitive sites of cTnI with regulatory activity, serine 23/24. TnI phosphorylation was slightly

impaired in Ang-II-treated cardiomyocytes (p=0.071) while co-treatment with MR-356 promoted

increased phosphorylation (Figure 6E) showing a consistent correlation with the data of

responsiveness to Ca^{2+} in panel 6D. Myosin binding protein C (MyBPC) contributes

substantially in cross-bridge cycling and its phosphorylation is also a main regulator of

myofilament sensitivity[34, 35]. HFpEF cardiomyocytes exhibited depressed phosphorylation at

serine 282, but no alteration in phosphorylation was detected in cardiomyocytes of animals

treated with Ang-II+MR-356 (Figure 6F).

GHRH-A prevents impairment of the activation (contraction) phase of SL–[Ca^{2+}] loops

in HFpEF cardiomyocytes

An evident negative inotropism determined by lower dSL/dt_{max}, was seen in HFpEF

cardiomyocytes indicating a disturbed excitation-contraction coupling and MR-356 prevented

this effect (Supplemental Figure 5A). an alternative approach to further assess the interaction

Ca^{2+}-sarcomere shortening was through the analysis of the "force-activation" phase of the SL-

[Ca^{2+}] loops in order to demonstrate the effects of MR-356 at the level of cross-bridge cycling.

In accordance with the slower contraction rate induced by chronic Ang-II treatment, the slopes of

the activation phase were consistently reduced in HFpEF cardiomyocytes (Supplemental Figure

5B and C). This reduction indicates a deteriorated cross-bridge cooperativity (and contractile

performance) which was prevented by co-treatment with MR-356.

GHRH-A improves relaxation and reverts the HFpEF phenotype induced by eight-week

Ang-II infusion in CD1 mice.

Given the ability of GHRH-As to prevent the onset of HFpEF, we next performed

experiments to test whether GHRH-As reverses established HFpEF. Accordingly, we infused

mice with a low dose of Ang-II for 8 weeks and initiated either treatment with MR-356 or the

vehicle at week 4, once the HFpEF phenotype was already apparent. Ang-II infusions were

continued in both groups during the second 4-week period. In vivo hemodynamic assessments

revealed that MR-356 restored the EDPVR (Figure 7 A-C), as well as the relaxation time

constant (τ) (Figure 7D) and the LVEDP (Figure 7E) back towards normal in HFpEF animals.

On the other hand, the increase in IVRT during the first 4-week treatment with Ang-II in both

groups, showed a trend to continue increasing in the HFpEF group. This tendency appears to be

stopped or even reverted in the group treated with MR-356 (Figure 7F), although this effect was

not significant within the period of the study. The beneficial effects on diastolic function were

also present on isolated cardiomyocytes, since MR-356 restored the resting sarcomere length (in

diastole, Figure 7G-H). Moreover, sarcomere relaxation rate was accelerated by MR-356

treatment in comparison with the 8-week Ang-II group (Figure 7I), as represented by the traces

on Figure 7G. Treatment with MR-356 also reverted the delay in decay of Ca^{2+} as shown by the

tt50% in response to pacing, in consonance with the increased transcription (mRNA) of the gene

for SERCA2a (Supplemental Figure 6). Moreover, time to peak SL contraction (ttp) and tt50%

SL relaxation in response to pacing (Supplemental Figure 7A and B) were greatly improved by

MR-356 on HFpEF cardiomyocytes, which was associated with increased gene expression of

troponin I (TNNi1 and 3) and myosin heavy chain α (MHY6) (Supplemental Figure 7C-E).

Myocardial remodeling was also reverted by MR-356 in this model, at least in terms of the

reversal of the left ventricular hypertrophy (Supplemental Figure 8A) and the re-expression of

the fetal gene myosin heavy chain β (MHY7), since the ratio MHY6/MHY7 was substantially

increased (Supplemental Figure 8).

Discussion

Heart failure with preserved ejection fraction is a syndrome with rising incidence and prevalence that is unresponsive to conventional treatments for heart failure with reduced ejection fraction. Given the rising clinical burden, it is imperative that effective treatments are found. Here we show that a novel class of peptide compounds, GHRH-As, previously shown to ameliorate pathological myocardial remodeling, also possess a powerful effect to restore diastolic dysfunction, a hallmark of HFpEF, to normal in an animal model of HFpEF, acting broadly through mechanisms involving fibrosis as well as myocyte calcium cycling and myofilament calcium responses.

Chronic administration of Ang-II reproduces structural and functional myocardial changes associated with HFpEF in mice. In contrast to previous results published by Regan el al.[32], we found signs of hypertension, as the blood pressure in the aorta was increased in these Ang-II treated mice (Table 1). This present in the majority of patients with HFpEF and might represent a risk factor[36]. These animals showed preserved EF and other contractility indices but depressed overall function, as observed with GLS[33], and associated with elevated LVEDP, myocardial fibrosis and impaired diastolic function as measured by IVRT, EDPVR and the relaxation constant tau. Within this pathological context, HFpEF animals showed myocardial hypertrophy that was virtually redirected into a physiological hypertrophy by treatment with MR-356, since although the increase in heart mass was not prevented in mice co-treated with Ang II+MR-356, cardiomyocyte width and diastolic RWT were maintained as in control mice as well as the fibrosis markers and the overall myocardial performance. In consonance with this, MR-356 reverted the left ventricular hypertrophy in the 8-week HFpEF model (Supplemental figure 8A). Accordingly, our results show that the above-mentioned signs consistent with HFpEF syndrome, were both prevented or reversed by the treatment with the GHRH-A.

Isolated cardiomyocytes from HFpEF mice exhibited lusitropic impairment as well as alterations in Ca^{2+} handling consistent with deficient myocardial relaxation and diastolic dysfunction. In this sense, our compelling experimental data suggest that the deficient myofilament relaxation in HFpEF cardiomyocytes might be associated with higher affinity of myofilaments for Ca^{2+}, favoring binding of cytosolic Ca^{2+} to TnC and thus promoting a delayed release of the interaction between thin and thick filaments. This idea is supported by the finding

of myofilament sensitization in HFpEF cardiomyocytes (leftward shift in the relaxation phase of

SL-Ca^{2+} loops) associated with reduced phosphorylation of cTnI at serine 23/24 and cMyBPC at

serine 282. In this sense, Ca^{2+} bound to myofilaments represents a very high percentage of the

Ca^{2+} released from the SR in a typical heart beat (about 50%)[37]. Therefore, the increased affinity

of myofilaments for Ca^{2+} has a substantial impact on cytosolic [Ca^{2+}], being able to buffer large

amounts of Ca^{2+}, including the Ca^{2+} leaked from the SR in HFpEF cardiomyocytes, and therefore

reducing the detectable levels. Moreover, the higher affinity for Ca^{2+} would make slower its

detachment from the troponin complex, which would impact the cytosolic [Ca^{2+}] decay rate[38, 39].

The increased PLB phosphorylation in the HEpEF cardiomyocytes appears not being sufficient

to compensate against both, the mentioned effect of the higher affinity for Ca^{2+} of myofilaments

and the diminished SERCA2 expression. The enhanced phosphorylation induced by MR-356 on

PLB, together with the desensitization of myofilaments, counteracted more efficiently and

prevented Ca^{2+} decay rate to slow (as shown in Figure 5 for the 4-week HFpEF model and

Supplemental Figure 6 in the 8-week HFpEF model). Importantly, NCX1 working in a reverse

mode (NCX1$_{rev}$) can contribute to the slowing of the Ca^{2+} decay rate in our murine model of

HFpEF, since low concentrations of Ang-II indirectly induce such a mechanism[40] without

affecting its protein expression. A study using a rat model of chronic kidney disease

(nephrectomy), which exhibits characteristics of HFpEF[29], demonstrated a main role for this

NCX1$_{rev}$ in the setting of HFpEF. An inhibitor of the NCX1 (SEA400), both in vivo and in vitro,

abolished the typical alteration in myocardium relaxation and remodeling. In this regard, it may

be appropriate to note that SEA400 inhibits both NCX working modes, reverse and forward.

Consistently, in our study, co-treatment with MR-356 down-regulated NCX1 expression, which

would suggest a reduced NCX1$_{rev}$ activity in HFpEF+MR-356 animals. However, this effect of

MR-356 on NCX1 expression would also affect the forward mode of the exchanger, with the

consequent removal of less Ca^{2+} from the cardiomyocyte, which in our model, is evidenced by

the increased levels of intracellular Ca^{2+} (SR[Ca^{2+}]) in cells from mice treated with this GHRH-

A.

297   We observed that myofilament responsiveness to Ca^{2+} was greatly decreased in

cardiomyocytes from Ang-II+MR-356-treated mice. This lower affinity of myofilaments for

Ca^{2+} may not only explain the molecular basis for faster relaxation, but also be the reason why

treatment with MR-356 preserved diastolic [Ca^{2+}] in Ang-II-treated myocytes; i.e., by preventing

the increase in Ca^{2+} buffering capacity of myofilaments. It has been demonstrated that this

molecular mechanism facilitates Ca^{2+} dissociation from cardiac myofilaments in response to

increasing pacing by hastening cross-bridge detachment [41]. Phosphorylation of cTnI and

cMyBPC accelerates relaxation either under β-adrenergic stimulation or increasing pacing

frequency which has been suggested to prevent diastolic dysfunction[42-44, 45, 46].

In addition to the underlying effect of the elevated $SRCa^{2+}$ leak on the depressed

inotropic response of our HFpEF cardiomyocytes, an inability to properly cycle cross-bridges

might be also involved. cMyBPC directly promotes force generation by enhancing cross-bridge

binding and cooperative recruitment[47]. cMyBPC also acts as a brake on cross-bridge kinetics by

slowing down actin-myosin detachment[48, 49], an effect that is removed by phosphorylation of

cMyBPC [47, 50, 51]. MR-356 treatment prevented depressed phosphorylation of cMyBPC at serine

282 in HFpEF cardiomyocytes and might account, in part, for the preserved contractile reserve in

HFpEF cardiomyocytes treated with GHRH-A.

Among various synthetic GHRH agonists tested, MR-356 combines its high GHRH

activity and ventricular remodeling prevention in a rat model of acute myocardial infarction,

inducing the greatest reducing effect on infarct size[21]. At variance with those GHRH agonists

with high activation potency of GHRHR, but little or no myocardial repair capacity, downstream

activation of GHRHR by MR-356 might stimulate reparative pathways including adenylate

cyclase/PKA signaling. Likewise, MR-356 has been shown to promote survival of cardiac stem

cell and induce proliferation in vitro, possibly mediated by ERK1/2 and PI3K/AKT pathways[52].

JI-38, a GHRH-A comparable to MR-356 in terms of prevention of ventricular remodeling, has

been demonstrated to promote the up-regulation of the transcription factor GATA-4 and

proliferation within the myocardium[23]. A recent study using a model of pressure overload[26],

which exhibits characteristics of HFrEF, shows that the GHRH-A MR-409 attenuates remodeling

and improves myocardial function. These beneficial effects are consistent with previous

publications from our and other groups. However, while Gesmundo's work addresses a model

that has been studied for decades, transaortic constriction, in which several therapeutic strategies

have been confirmed as effective as they are in humans with HFrEF, our work focusses on a

highly relevant clinical problem with major unmet needs, the HFpEF syndrome. Our new results

show for first time the action of GHRH-As on the prevention and the reversal of the HFpEF

phenotype, especially diastolic dysfunction. Thus, based in the proven properties for cardiac

repair of GHRH-As and the robust and convincing data obtained in the present study, we

consider that GHRH-As are potentially useful for the management of HFpEF.

In summary, here we have shown that activation of GHRHR with a synthetic GHRH

agonist prevents the appearance of the HFpEF phenotype and reverses the signs of HFpEF once

established in mice, including dysfunctional diastolic relaxation in myocytes and intact animals

and myocardial remodeling. Increased SR Ca^{2+} leak associated with higher affinity of

myofilaments for Ca^{2+} and impaired cross-bridge cycling, underlie the dysfunctional contractile

response and lusitropy in these HFpEF cardiomyocytes, and may in part account for the

alterations in Ca^{2+} kinetics. Myofibrillar proteins appear to be a target for treatment with GHRH

agonists. Taken together these findings indicate that synthetic analogs of GHRH represent novel

therapeutic compounds for treating patients with the HFpEF syndrome.

Methods.

Experimental design and animal model

We used age matched (2-month old) CD1 mice (n=63, from Envigo). Mice were

implanted with a mini-osmotic pump (Alzet) to deliver Ang-II (0.2 mg/kg/day; Sigma-Aldrich

Co., Saint Louis, MO) for 4 weeks. Simultaneously, mice received daily injections of GHRH-

Agonist (GHRH-A [MR-356]: 200 μg/kg, n=23) or vehicle (DMSO+propylene-glycol, n=23)

during the same period of time. Unmanipulated (no mini-osmotic pump) age-matched CD1 mice

(n=17) were used as normal controls.

A separate group of CD1 mice (n=24, 2-month old) was implanted with the Alzet mini-

pumps containing Ang-II. After all mini-pumps content were delivered (4 weeks), the animals

were subjected to replacement of a mini-pumps for an additional 4-week Ang-II infusion. Upon

the second set of mini-pump implantation (day 29), the mice were randomized. Half of them

were daily injected with MR-356 and the remaining 12 mice received vehicle injections until the

day 56.

Materials

The GHRH-A, MR-356 ([N-Me-Tyr[1], Gln[8], Orn[12], Abu[15], Orn[21], Nle[27], Asp[28], Agm[29]]

hGH-RH(1-29)), was synthetized by solid phase methods and purified by HPLC in our

laboratories as described previously[21].

Echocardiography

363   Transthoracic echocardiography was performed as previously described with minor

364   modifications[53]. Briefly, mice were screened at baseline and 4 weeks following mini-osmotic

365   pump placement. All echocardiographic measurements were performed offline over 3 to 5 heart

366   beats.

Speckle Tracking Echocardiography (STE)

368   Strain analysis was conducted using speckle-tracking software, Vevostrain TM analysis

369   (Visual Sonics). Global longitudinal strain was calculated using B-mode cine images in the LV

370   parasternal long-axis view acquired at a high frame rate.

Hemodynamic Measurements

Hemodynamic studies were performed using a micro-tipped pressure-volume catheter

(SPR-839; Millar Instruments) as previously described with minor modifications. Briefly, the

catheter was inserted into the right carotid artery and advanced retrograde into the LV. Pressure-

volume loops were recorded at steady state, and during temporary inferior vena cava occlusion.

All analyses were performed using LabChart Pro version 8.1.5 software (ADInstruments). End-

diastolic pressure-volume pairs were plotted and then fitted by a linear regression. The reported

data representing the end-diastolic pressure-volume relationship (EDPVR) correspond to the

slope of the linear fitting: "stiffness coefficient" (mmHg/L).

Cardiomyocyte Isolation

Cardiac myocytes were isolated and prepared from hearts of mice by retrograde perfusion

through the aorta in a modified Langerdorf system with collagenase type 2 and protease type XIV

in an isolation solution (see the detailed procedure in the Supplemental Material online).

Cardiomyocytes were resuspended in Tyrode's buffer with 1.8 mmol/L $CaCl_2$, while a small

aliquot was kept in the presence of 10 mmol/L 2,3-butanedione monoxime (BDM) for cell length

and width measurement.

Intracellular Calcium and Sarcomere Length Measurement

Intracellular Ca^{2+} was measured using the Ca^{2+}-sensitive dye Fura-2 and a dual-excitation

(340/380 nm) spectrofluorometer (IonOptix LLC, Milton, MA, USA) in cardiomyocytes electric-

field stimulated at 1, 2 and 4 Hz. $[Ca^{2+}]_i$ was calculated as described previously[54]. $\Delta[Ca^{+2}]_i$

amplitude was considered as: peak $[Ca^{+2}]_i$ – resting $[Ca^{+2}]_i$(see detailed methodology in the online

Supplemental Material). Simultaneously, sarcomere length (SL) was recorded with an IonOptix

iCCD camera. Change in average SL was determined by Fast Fourier Transform analysis of the

Z-line density trace to the frequency domain, and SL shortening (SLS) was calculated as:

SLS(%)=100×(resting SL−peak SL)/resting SL.

Hysteresis Sarcomere Shortening-Calcium Loop Analysis

The hysteresis loops were obtained by plotting sarcomere length versus calcium

concentration of cardiomyocytes paced at 4 Hz. Two of the three main segments of such loops

were analyzed for this work: the "force-activation" and the "relaxation" phases[34, 55] (see detailed

explanation in the online Supplemental Material). The analyses of both, relaxation and force-

activation phases were performed by GraphPad Prism version 6.07 (San Diego, CA USA).

SR Ca^{2+} Leak and SR Ca^{2+} Load Measurement

SR Ca^{2+} leakage was assessed with 1 mmol/L tetracaine (Sigma-Aldrich) using a modified

protocol from the one described by Shannon *et al*[56]. Briefly, the observed decrease in the Fura-2

ratio in presence of tetracaine compared to the not-tetracaine treated condition was considered the

Ca^{2+} leak for a particular myocyte (detailed methodology is provided in the online Supplemental

Material). After tetracaine wash out, SR Ca^{2+} content was assessed on the same cell by a 20

mmol/L caffeine challenge. SR Ca^{2+} contents were calculated considering that SR represents 3.5%

and cytosol 65% of the myocyte volume. Ca^{2+} leak-SR $[Ca^{2+}]$ pairs were grouped by similar SR

$[Ca^{2+}]$ load, plotted and fitted by an exponential growth function using the Graph Pad Prism

software (version 6.07). Measurements were carried out at 37°C.

Western Blot Immunoanalysis

Isolated cardiomyocytes: Frozen cardiomyocytes were lysed in ice-cold RIPA buffer with

proteinase inhibitors (cOmplete protease inhibitor cocktail; Roche), sonicated and then

centrifuged at 12500 RPM for 15 minutes at 4°C.

Hearts: Frozen cardiac tissues were homogenized using a Pyrex glass-glass homogenizer

on liquid nitrogen using a urea-based lysing buffer. The samples were then centrifuged.

Supernatants from either procedure were collected and sampled for protein concentration.

Samples were electrophoresed and transferred to PVDF or nitrocellulose membranes (Bio-Rad

Laboratories). Detection of proteins was performed by immunoblotting using the adequate

antibodies as indicated (please, see detailed information in the Supplemental Material online).

Quantitative RT-PCR

Total RNA was isolated from cardiomyocytes using RNAeasy mini kit (Qiagen, Hilden,

Germany). RNA (1 μg) was reverse-transcribed according to instructions utilizing a cDNA

Synthesis kit (Applied Biosystems, Foster City, CA). Using Taqman® Universal Master Mix

(Applied Biosystems, Foster City, CA) in a Bio-Rad CFX96™ Real-Time PCR system (Bio-Rad

Laboratories, Hercules, CA, USA), qPCR was performed. All samples were run in duplicates and

normalized to 18S Taqman gene-expression as follows: ATP2a2: Mm01201431_m1; RYR2

Mm00465877_m1, TNNi1: Mm00502426_m1; TNNi3: Mm00437164_m1; MHY6:

Mm00440359_m1; MHY7: Mm00600555_m1and 18S Mm03928990_g1. The change in mRNA

expression was calculated compared to the change in normalized 18S, and values were expressed

as 2ΔCt.

Statistical Analysis

Data are reported as mean ± standard error of the mean (SEM). Statistical significance

between three groups was determined by one-way ANOVA, or two-way ANOVA followed by

Sidak's or Bonferonni's post hoc tests, as appropriate. For comparisons of two groups, Student's

t-test two-tailed was used. Analyses were performed using GraphPad Prism version 8.3.0. The

null hypothesis was rejected at $p<0.05$.

Study approval

All protocols and experimental procedures were approved by the University of Miami

Animal Care and Use Committee following the Guide for the Care and Use of Laboratory

Animals (NIH Publication No. 85-234, revised 2011).

References

444
445
1.   Berlot, B., Bucciarelli-Ducci, C., Palazzuoli, A. & Marino, P. Myocardial phenotypes and
dysfunction in HFpEF and HFrEF assessed by echocardiography and cardiac magnetic resonance.
Heart Fail Rev (2019).
2.   Lewis, G.A. et al. Biological Phenotypes of Heart Failure With Preserved Ejection Fraction. *J Am*
Coll Cardiol **70**, 2186-2200 (2017).
3.   Pieske, B. et al. How to diagnose heart failure with preserved ejection fraction: the HFA-PEFF
diagnostic algorithm: a consensus recommendation from the Heart Failure Association (HFA) of
the European Society of Cardiology (ESC). *Eur Heart J* **40**, 3297-3317 (2019).
4.   Owan, T.E. et al. Trends in prevalence and outcome of heart failure with preserved ejection
fraction. *N Engl J Med* **355**, 251-259 (2006).
5.   Oktay, A.A., Rich, J.D. & Shah, S.J. The emerging epidemic of heart failure with preserved
ejection fraction. *Curr Heart Fail Rep* **10**, 401-410 (2013).
6.   Silverman, D.N. et al. Association of beta-Blocker Use With Heart Failure Hospitalizations and
Cardiovascular Disease Mortality Among Patients With Heart Failure With a Preserved Ejection
Fraction: A Secondary Analysis of the TOPCAT Trial. *JAMA Netw Open* **2**, e1916598 (2019).
7.   Pitt, B. et al. Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med* **370**,
1383-1392 (2014).
8.   Solomon, S.D. et al. Angiotensin-Neprilysin Inhibition in Heart Failure with Preserved Ejection
Fraction. *N Engl J Med* **381**, 1609-1620 (2019).
9.   O'Connor, C.M. & deFilippi, C. PARAGON-HF - Why We Do Randomized, Controlled Clinical
Trials. *N Engl J Med* **381**, 1675-1676 (2019).
10. Takada, T. et al. Impact of elevated heart rate on clinical outcomes in patients with heart failure
with reduced and preserved ejection fraction: a report from the CHART-2 Study. *Eur J Heart*
Fail **16**, 309-316 (2014).
11. Melenovsky, V. et al. Cardiovascular features of heart failure with preserved ejection fraction
versus nonfailing hypertensive left ventricular hypertrophy in the urban Baltimore community:
the role of atrial remodeling/dysfunction. *J Am Coll Cardiol* **49**, 198-207 (2007).
12. Yang, J.H. et al. Endothelium-dependent and independent coronary microvascular dysfunction in
patients with heart failure with preserved ejection fraction. *Eur J Heart Fail* (2019).
13. Shah, S.J. et al. Phenotype-Specific Treatment of Heart Failure With Preserved Ejection Fraction:
A Multiorgan Roadmap. *Circulation* **134**, 73-90 (2016).
14. Schiattarella, G.G. et al. Nitrosative stress drives heart failure with preserved ejection fraction.
Nature **568**, 351-356 (2019).
15. Obokata, M. et al. The neurohormonal basis of pulmonary hypertension in heart failure with
preserved ejection fraction. *Eur Heart J* **40**, 3707-3717 (2019).
16. Kjeldsen, S.E. et al. Medical Therapies for Heart Failure With Preserved Ejection Fraction.
Hypertension **75**, 23-32 (2020).
17. Tarone, G. et al. Targeting myocardial remodelling to develop novel therapies for heart failure: a
position paper from the Working Group on Myocardial Function of the European Society of
Cardiology. *Eur J Heart Fail* **16**, 494-508 (2014).
18. Redfield, M.M. et al. Isosorbide Mononitrate in Heart Failure with Preserved Ejection Fraction. *N*
Engl J Med **373**, 2314-2324 (2015).
19. Banerjee, P. Heart failure with preserved ejection fraction: A clinical crisis. *Int J Cardiol* **204**,
198-199 (2016).
20. Fernandez-Ruiz, I. The search for an effective HFpEF treatment continues. *Nat Rev Cardiol* **16**,
647 (2019).
21. Cai, R. et al. Synthesis of new potent agonistic analogs of growth hormone-releasing hormone
(GHRH) and evaluation of their endocrine and cardiac activities. *Peptides* **52**, 104-112 (2014).

22. Granata, R. et al. Growth hormone-releasing hormone promotes survival of cardiac myocytes in
vitro and protects against ischaemia-reperfusion injury in rat heart. *Cardiovasc Res* **83**, 303-312
(2009).
23. Kanashiro-Takeuchi, R.M. et al. Activation of growth hormone releasing hormone (GHRH)
receptor stimulates cardiac reverse remodeling after myocardial infarction (MI). *Proc Natl Acad*
Sci U S A **109**, 559-563 (2012).
24. Kanashiro-Takeuchi, R.M. et al. Cardioprotective effects of growth hormone-releasing hormone
agonist after myocardial infarction. *Proc Natl Acad Sci U S A* **107**, 2604-2609 (2010).
25. Bagno, L.L. et al. Growth hormone-releasing hormone agonists reduce myocardial infarct scar in
swine with subacute ischemic cardiomyopathy. *J Am Heart Assoc* **4** (2015).
26. Gesmundo, I. et al. Growth hormone-releasing hormone attenuates cardiac hypertrophy and
improves heart function in pressure overload-induced heart failure. *Proc Natl Acad Sci U S A*
114, 12033-12038 (2017).
27. Matsubara, S., Sato, M., Mizobuchi, M., Niimi, M. & Takahara, J. Differential gene expression of
growth hormone (GH)-releasing hormone (GRH) and GRH receptor in various rat tissues.
Endocrinology **136**, 4147-4150 (1995).
28. Conceicao, G., Heinonen, I., Lourenco, A.P., Duncker, D.J. & Falcao-Pires, I. Animal models of
heart failure with preserved ejection fraction. *Neth Heart J* **24**, 275-286 (2016).
29. Primessnig, U. et al. Novel pathomechanisms of cardiomyocyte dysfunction in a model of heart
failure with preserved ejection fraction. *Eur J Heart Fail* **18**, 987-997 (2016).
30. Olver, T.D. et al. Western Diet-Fed, Aortic-Banded Ossabaw Swine: A Preclinical Model of
Cardio-Metabolic Heart Failure. *JACC Basic Transl Sci* **4**, 404-421 (2019).
31. Matyas, C. et al. Prevention of the development of heart failure with preserved ejection fraction
by the phosphodiesterase-5A inhibitor vardenafil in rats with type 2 diabetes. *Eur J Heart Fail*
19, 326-336 (2017).
32. Regan, J.A. et al. A mouse model of heart failure with preserved ejection fraction due to chronic
infusion of a low subpressor dose of angiotensin II. *Am J Physiol Heart Circ Physiol* **309**, H771-
778 (2015).
33. Wang, J., Khoury, D.S., Yue, Y., Torre-Amione, G. & Nagueh, S.F. Preserved left ventricular
twist and circumferential deformation, but depressed longitudinal and radial deformation in
patients with diastolic heart failure. *Eur Heart J* **29**, 1283-1289 (2008).
34. Tong, C.W., Gaffin, R.D., Zawieja, D.C. & Muthuchamy, M. Roles of phosphorylation of myosin
binding protein-C and troponin I in mouse cardiac muscle twitch dynamics. *J Physiol* **558**, 927-
941 (2004).
35. Tong, C.W., Nair, N.A., Doersch, K.M., Liu, Y. & Rosas, P.C. Cardiac myosin-binding protein-C
is a critical mediator of diastolic function. *Pflugers Arch* **466**, 451-457 (2014).
36. Tadic, M., Cuspidi, C., Frydas, A. & Grassi, G. The role of arterial hypertension in development
heart failure with preserved ejection fraction: just a risk factor or something more? *Heart Fail*
Rev **23**, 631-639 (2018).
37. Shannon, T.R., Ginsburg, K.S. & Bers, D.M. Reverse mode of the sarcoplasmic reticulum
calcium pump and load-dependent cytosolic calcium decline in voltage-clamped cardiac
ventricular myocytes. *Biophys J* **78**, 322-333 (2000).
38. Diaz, M.E., Trafford, A.W. & Eisner, D.A. The effects of exogenous calcium buffers on the
systolic calcium transient in rat ventricular myocytes. *Biophys J* **80**, 1915-1925 (2001).
39. Kataoka, A., Hemmer, C. & Chase, P.B. Computational simulation of hypertrophic
cardiomyopathy mutations in troponin I: influence of increased myofilament calcium sensitivity
on isometric force, ATPase and [Ca2+]i. *J Biomech* **40**, 2044-2052 (2007).
40. Perez, N.G. et al. A low dose of angiotensin II increases inotropism through activation of reverse
Na(+)/Ca(2+) exchange by endothelin release. *Cardiovasc Res* **60**, 589-597 (2003).
41. Baker, A.J. & Weiner, M.W. Force decline during muscle relaxation promotes calcium release to
the cytosol. *Am J Physiol* **273**, C85-91 (1997).

42.   Colson, B.A. et al. Myosin binding protein-C phosphorylation is the principal mediator of protein
kinase A effects on thick filament structure in myocardium. *J Mol Cell Cardiol* **53**, 609-616
(2012).
43.   Varian, K.D. & Janssen, P.M. Frequency-dependent acceleration of relaxation involves decreased
myofilament calcium sensitivity. *Am J Physiol Heart Circ Physiol* **292**, H2212-2219 (2007).
44.   Varian, K.D. et al. Impairment of diastolic function by lack of frequency-dependent myofilament
desensitization rabbit right ventricular hypertrophy. *Circ Heart Fail* **2**, 472-481 (2009).
45.   Dweck, D. et al. Long term ablation of protein kinase A (PKA)-mediated cardiac troponin I
phosphorylation leads to excitation-contraction uncoupling and diastolic dysfunction in a knock-
in mouse model of hypertrophic cardiomyopathy. *J Biol Chem* **289**, 23097-23111 (2014).
46.   Takimoto, E. et al. Frequency- and afterload-dependent cardiac modulation in vivo by troponin I
with constitutively active protein kinase A phosphorylation sites. *Circ Res* **94**, 496-504 (2004).
47.   Coulton, A.T. & Stelzer, J.E. Cardiac myosin binding protein C and its phosphorylation regulate
multiple steps in the cross-bridge cycle of muscle contraction. *Biochemistry* **51**, 3292-3301
(2012).
48.   Razumova, M.V., Bezold, K.L., Tu, A.Y., Regnier, M. & Harris, S.P. Contribution of the myosin
binding protein C motif to functional effects in permeabilized rat trabeculae. *J Gen Physiol* **132**,
575-585 (2008).
49.   Saber, W., Begin, K.J., Warshaw, D.M. & VanBuren, P. Cardiac myosin binding protein-C
modulates actomyosin binding and kinetics in the in vitro motility assay. *J Mol Cell Cardiol* **44**,
1053-1061 (2008).
50.   Shaffer, J.F., Kensler, R.W. & Harris, S.P. The myosin-binding protein C motif binds to F-actin
in a phosphorylation-sensitive manner. *J Biol Chem* **284**, 12318-12327 (2009).
51.   Sadayappan, S. et al. A critical function for Ser-282 in cardiac Myosin binding protein-C
phosphorylation and cardiac function. *Circ Res* **109**, 141-150 (2011).
52.   Florea, V. et al. Agonists of growth hormone-releasing hormone stimulate self-renewal of cardiac
stem cells and promote their survival. *Proc Natl Acad Sci U S A* **111**, 17260-17265 (2014).
53.   Yuan, C.C. et al. Constitutive phosphorylation of cardiac myosin regulatory light chain prevents
development of hypertrophic cardiomyopathy in mice. *Proc Natl Acad Sci U S A* **112**, E4138-
4146 (2015).
54.   Dulce, R.A. et al. Hydralazine and organic nitrates restore impaired excitation-contraction
coupling by reducing calcium leak associated with nitroso-redox imbalance. *J Biol Chem* **288**,
6522-6533 (2013).
55.   Ramirez-Correa, G.A., Cortassa, S., Stanley, B., Gao, W.D. & Murphy, A.M. Calcium sensitivity,
force frequency relationship and cardiac troponin I: critical role of PKA and PKC
phosphorylation sites. *J Mol Cell Cardiol* **48**, 943-953 (2010).
56.   Shannon, T.R., Ginsburg, K.S. & Bers, D.M. Quantitative assessment of the SR Ca2+ leak-load
relationship. *Circ Res* **91**, 594-600 (2002).

Acknowledgments

Sources of funding: Study was supported by NIH, grants 1R01 HL13735 and 1R01 HL107110
to JMH. JMH is also supported by NIH grants 5UM1 HL113460, 1R01 HL134558, 5R01
CA136387 and The Starr Foundation. São Paulo Research Foundation (FAPESP) grant#
2016/01044-0 (to AGS).

Author contributions: RAD was involved in designing research studies, and conducted
experiments, acquiring data, analyzing data, discussion and interpretation of results, and writing
the manuscript. RMKT, main responsible for designing research studies, she also conducted
experiments, acquiring and analyzing data, and discussing results. LMT conducted experiments,
acquired and analyzed data. AGS conducted experiments. SK analyzed and interpreted
echocardiographic measurements. WB participated in discussing results, writing and correction
of the manuscript. MSSRZ analyzed echocardiographic measurements. RC synthesized reagents.
AVS provided reagents and contributed to writing of the manuscript. JMH conceived the project,
participated in data and results interpretation, provided scientific and financial support for the
project, and contributed to writing of the manuscript.

Competing interests: Dr. Joshua Hare owned equity in Biscayne Pharmaceuticals, licensee of
intellectual property used in this study. Biscayne Pharmaceuticals did not provide funding for
this study. Dr. Joshua Hare is the Chief Scientific Officer, a compensated consultant and
advisory board member for Longeveron and holds equity in Longeveron. Dr. Hare is also the co-
inventor of intellectual property licensed to Longeveron. All other authors have declared that no
conflict of interest exists.

Materials & Correspondence: Additional information from the study is available and should be
requested to the corresponding author Dr. Joshua M. Hare: jhare@med.miami.edu

Tables:

- Table 1: Hemodynamic assessments on 4-week Ang-II infusion model

	Control (N=8)	Ang-II (N=9)	Ang-II+MR-356 (N=9)
HR (1/min)	535.9±11.7	543.9±7.2	523.5±14.2
EF (%)	51.0±4.0	55.7±5.8	59.6±2.9
Tau (1/sec)	5.21±0.33	7.12±0.57*	5.91±0.27†
dP/dt$_{max}$ (mmHg/sec)	7328±346	6866±519	7433±476
dP/dt$_{min}$ (mmHg/sec)	-5494±353	-4897±451	-7013±580*†
Ea (mmHg/µL)	1.65±0.25	1.95±0.29	1.83±0.13
LVEDP (mmHg)	4.13±0.89	6.46±0.38*	5.87±0.91
ESPVR (mmHg/µL)	2.71±0.28	1.57±0.19*	1.96±0.37*
EDPVR (mmHg/µL) [linear]	0.068±0.004	0.091±0.009*	0.059±0.004†
PRSW (mmHg/µL)	74.5±3.1	63.2±3.9	62.4±5.4
dP/dt$_{max}$-EDV (mmHg/µL.sec)	110.7±13.9	99.0±6.3	91.1±9.6
Systolic ABP (mmHg)	79.7±2.2	103.0±4.7*	88.5±4.6†
Diastolic ABP (mmHg)	54.7±2.7	75.5±3.7*	64.4±4.2

Table 1: All values represent mean±SEM. HR, heart rate; EF, ejection fraction; Tau, relaxation constant; dP/dt$_{max}$, maximal contraction rate; dP/dt$_{min}$, maximal relaxation rate; Ea, arterial elastance; LVEDP, left ventricular end-diastolic pressure; ESPVR, end-systolic pressure-volume relationship; EDPVR, slope of the end-diastolic pressure-volume relationship; PRSW, Preload recruitable stroke work; dP/dt$_{max}$-EDV, maximal contraction rate-end diastolic volume relationship. ABP, arterial blood pressure. * $p<0.05$ vs. control; † $p<0.05$ Ang-II+MR-356 vs. Ang-II, one-way ANOVA. N represents number of studied animals.

Figures:

Figure 1: Biometric, echocardiographic, cellular and molecular characterization of the
HFpEF model. GHRH-A MR-356 prevents cardiomyocyte growth and fibrosis. (A) Heart size
normalized to body weight. (B) Relative wall thickness in diastole (RWTd) in different groups (‡
p<0.05 4-week vs. baseline within groups; two-way ANOVA). (C) Width of isolated
cardiomyocytes from control (N=4), Ang-II-treated (N=5) and Ang-II+MR-356-treated (N=5)
mice. (D) Collagen III protein expression. Top, representative images of western blots. Bottom,
bar graph showing average results. (E) Transforming growth factor-β (TGF-β) and (F) α-smooth
muscle actin (α-SMA) protein expression in heart homogenates. *p<0.05 in Ang II vs. control and
† *p*<0.05 in Ang II+MR-356-treated vs. Ang II; One-way ANOVA. Numbers on bottom of bars
correspond to studied animals in panels A-B, while they indicate number of cells in panel C, or
tissue samples in panels D-F.

Figure 2: Effects of the GHRH-A MR-356 on diastolic dysfunction in the heart of Ang-II-induced HFpEF mice. (A) Representative pressure-volume (P-V) loops showing the linear regression for end-systolic pressure-volume relationship (ESPVR) and end-diastolic pressure-volume relationship (EDPVR) in a control mouse, or (B) Ang-II-treated mouse, or (C) Ang-II+MR-356 mouse. (D) Averaged slopes from the EDPVR linear fitting. (E) Hemodynamic assessment of the relaxation time constant (tau) at 4-week time point. (F) Left ventricular end-diastolic pressure (LVEDP). (G) Maximal rate of pressure decline (dP/dt_{min}). (H) Isovolumic relaxation time (IVRT) as measured by transthoracic echocardiography at baseline or after 4-week treatment. (I) Global longitudinal strain (GLS) analysis of the left ventricle assessed in same conditions. GLS represents the function of sub-endocardial longitudinal myocardial fibers that are more sensitive to reduced coronary perfusion and increased wall stress. Moreover, GLS reflects changes in the myocardial interstitium including the extent of myocardial fibrosis. ‡ $p<0.05$ at 4-week vs. baseline; ‡‡ $p<0.001$ at 4-week vs. baseline, two-way ANOVA. *$p<0.05$ vs. control; † $p<0.05$ in Ang-II+MR-356-treated vs. Ang-II; one or two-way ANOVA. Numbers on bottom of bars indicate number of analyzed animals.

Figure 3: Sarcomere and contractile dysfunction in the HFpEF model is prevented by co-treatment with GHRH-A MR-356. (A) Representative traces of sarcomere twitches of cardiomyocytes paced at 1 and 4 Hz obtained from CD1 control mice (N=4, 30 cells) or mice which underwent either Ang-II infusion (N=5, 41 cells) or Ang-II+MR-356 treatment (N=5, 38 cells). (B) Resting sarcomere length in cardiomyocytes from each group at different pacing rates. (C) Sarcomere shortening in response to pacing. (D) Representative traces of intracellular $[Ca^{2+}]$ in cardiomyocytes electric field-stimulated in same conditions as above. (E) $\Delta[Ca^{2+}]$ transient amplitude in response to pacing. (F) $\Delta[Ca^{2+}]$ transient-frequency relationship normalized respect to 1 Hz. (G) Diastolic $[Ca^{2+}]$ in response to increasing frequency of stimulation. From each mouse, a total of 5 to 9 cells were studied. *$p<0.05$ in Ang-II vs. control and † $p<0.05$ in Ang-II+MR-356-treated vs. Ang-II; two-way ANOVA.

Figure 4: Increased SR Ca^{2+} leak in HFpEF cardiomyocytes is prevented by treatment with the GHRH-A MR-356. (A) Representative traces of intracellular [Ca^{2+}] in cardiomyocytes illustrating the method for measuring SR Ca^{2+} leak using tetracaine. Blue trace corresponds to Ca^{2+} from a cardiomyocyte not exposed to tetracaine, while red trace is the Ca^{2+} from the same cardiomyocyte that was treated 70 seconds with tetracaine. (B) SR Ca^{2+} leak-load relationship in cardiomyocytes from control untreated (n=31 cells), Ang-II-treated (n=38 cells) or Ang-II+MR-356-treated (n=35 cells) CD1 mice. (C) Average SR Ca^{2+} leak in cardiomyocytes from all groups at matched SR Ca^{2+} load=99.1 µmol/L. *$p<0.05$ in Ang-II vs. control and † $p<0.05$ in Ang-II-treated vs. Ang-II+MR-356-treated; one or two-way ANOVA as appropriate. Numbers on bottom of bars indicate number of analyzed cells.

Figure 5: **Ca²⁺ decay is delayed in cardiomyocytes from the Ang-II-induced HFpEF model but preserved by treatment with the GHRH-A MR-356.** (A) Normalized Ca²⁺ transient reconstructed by using average dynamic parameters from control untreated, Ang-II-treated or Ang-II+MR-356-treated cardiomyocytes paced at 1 Hz. (B) Average times to different degrees of the Ca²⁺ peak decline: time to 10% (tt10%), time to 50% (tt50%), decay time constant (τ) and time to 90% (tt90%). (C) SR Ca²⁺ content in cardiomyocytes paced at 1 Hz from each group. Numbers on bottom of bars on panels B and C indicate number of analyzed cells. (D) SERCA2a protein expression; top, representative immunoblotting image with Hsp90 as loading control and bottom, average values of SERCA2a expression. (E) PLB phosphorylation at serine 16. (F) Phosphorylation of PLB at threonine 17. Top, representative images and bottom, average values. Numbers on bottom of bars on panels D, E and F indicate number of analyzed samples. $*p<0.05$ or $**p<0.01$ vs. control, $\dagger$ $p<0.05$ vs. Ang-II-treated; one-way ANOVA.

Figure 6: **Impairment in sarcomere relaxation due to myofilaments sensitization in HFpEF cardiomyocytes is prevented by treatment with the GHRH-A MR-356.** (A) Normalized sarcomere contraction re-constructed by using average SL at 1Hz from control untreated, Ang- II-treated or Ang-II+MR-356-treated cardiomyocytes. (B) Average times to different degrees of the relaxation: tt10%, tt50%, tau (τ) and tt90%. (C) SL-[Ca^{2+}] hysteresis loops in control, Ang-II- or Ang-II+MR-356-treated cardiomyocytes with overlapped sigmoidal fitting of the relaxation phases. (D) Normalized relaxation phase of SL-[Ca^{2+}] loops (left) and averaged effective [Ca^{2+}] of the relaxation phase ($_{relax}$EC$_{50}$s, right). Numbers on bottom of bars on panel D indicate number of analyzed cells from 4-5 hearts. (E) Phosphorylation of cTnI at serine 23/24 and total protein expression by western blot. (F) Phosphorylation of cMyBPC at serine 282 and total protein expression. Top, representative images; bottom, average values. Numbers on bottom of bars on panels E and F indicate number of analyzed samples. ******p*<0.05 vs. control, † *p*<0.05 vs. Ang-II + MR-356-treated; one-way ANOVA.

Figure 7: Effect of the GHRH-A MR-356 on the chronically installed HFpEF phenotype.

(A) Representative pressure-volume (P-V) loops showing the linear regression for end-systolic pressure-volume relationship (ESPVR) and end-diastolic pressure-volume relationship (EDPVR) in a mouse treated for 8 weeks with Ang-II, or (B) an Ang-II (8w)+MR-356-treated mouse. (C) Averaged slopes from the EDPVR linear fitting. (D) Hemodynamic assessment of the relaxation time constant (tau) at 8-week time point. (E) Averaged left ventricular end-diastolic pressure (LVEDP). (F) Isovolumetric relaxation time (IVRT) as measured by transthoracic echocardiography at baseline, 4-week and 8-week. (G) Representative SL traces comparing Ang-II (8w)-treated and Ang-II (8w) + MR-356 cardiomyocytes at 1 and 4 Hz pacing rates. (H) Resting sarcomere length of Ang-II (8w)-treated compared with Ang-II (8w) + MR-356-treated mouse cardiomyocytes paced at 1, 2 and 4 Hz. (I) Averaged times to different degrees of SL relaxation: time to 10% (tt10%), time to 50% (tt50%), relaxation time constant (τ) and time to 90% (tt90%) in 8-week-treated cardiomyocytes paced at 1 Hz. † $p<0.05$ vs. Ang-II (8w)+ MR-356-treated; Student's t-test. ‡ $p<0.05$ within group comparison, two-way ANOVA.