Time-resolved comparative molecular evolution of oxygenic photosynthesis

Thomas Oliver[1], Patricia Sánchez-Baracaldo[2], Anthony W. Larkum[3], A. William

Rutherford[1], Tanai Cardona[1]*

[1]Department of Life Sciences, Imperial College London, London, UK

[2]School of Geographical Sciences, University of Bristol, Bristol, UK

[3]University of Technology Sydney, Ultimo NSW, Australia

*Correspondence to: t.cardona@imperial.ac.uk

Abstract

Oxygenic photosynthesis starts with the oxidation of water to O_2, a light-driven reaction

catalysed by photosystem II. Cyanobacteria are the only prokaryotes capable of water

oxidation and therefore, it is assumed that relative to the origin of life and bioenergetics, the

origin of oxygenic photosynthesis is a late innovation. However, when exactly water

oxidation originated remains an unanswered question. Here we use relaxed molecular clocks

to compare one of the two ancestral core duplications that are unique to water-oxidizing

photosystem II, that leading to CP43 and CP47, with some of the oldest well-described events

in the history of life. Namely, the duplication leading to the Alpha and Beta subunits of the

catalytic head of ATP synthase, and the divergence of archaeal and bacterial RNA

polymerases and ribosomes. We also compare it with more recent events such as the

duplication of cyanobacteria-specific FtsH metalloprotease subunits, of CP43 variants used in

a variety of photoacclimation responses, and the speciation events leading to

Margulisbacteria, Sericytochromatia, Vampirovibrionia, and other clades containing

anoxygenic phototrophs. We demonstrate that the ancestral core duplication of photosystem

II exhibits patterns in the rates of protein evolution through geological time that are nearly

identical to those of the ATP synthase, RNA polymerase, or the ribosome. Furthermore, we

use ancestral sequence reconstruction in combination with comparative structural biology of

photosystem subunits, to provide additional evidence supporting the premise that water

oxidation had originated before the ancestral core duplications. Our work suggests that

photosynthetic water oxidation originated closer to the origin of life and bioenergetics than

can be documented based on species trees alone.

1. Introduction

1.1. Evolution of Cyanobacteria

The origin of oxygenic photosynthesis is considered a turning point in the history of life,

marking the transition from the ancient world of anaerobes into a productive aerobic world

that permitted the emergence of complex life [1]. Oxygenic photosynthesis starts with

photosystem II (PSII), the water-oxidising and O_2-evolving enzyme of Cyanobacteria and

photosynthetic eukaryotes. Today PSII is a highly conserved, multicomponent, membrane

protein complex, which was inherited by the *most recent common ancestor* (MRCA) of

Cyanobacteria in a form that is structurally and functionally quite similar to that found in all

extant species [2]. Thus, the origin of oxygenic photosynthesis antedates the MRCA of

Cyanobacteria by an undetermined amount of time.

Cyanobacteria's closest living relatives are the clades known as Vampirovibrionia

(formerly Melainabacteria) [3, 4], followed by Sericytochromatia [5] and Margulisbacteria

[6]. Currently, no photosynthetic strains have been described in these groups of uncultured

bacteria and this has led to the hypothesis that oxygenic photosynthesis arose during the time

spanning the divergence of Vampirovibrionia and the MRCA of Cyanobacteria, starting from

an entirely non-photosynthetic ancestral state [5, 7]. Molecular clock studies have suggested

that the span of time between the divergence of Cyanobacteria and Vampirovibrionia could

be of several hundred million years [8, 9]. However, it is still unclear from molecular clock

analyses and the microfossil record when exactly the MRCA of Cyanobacteria occurred [10].

For example, two recent independent analysis placed the same cyanobacterial ancestor

around a mean age younger than 1.5 Ga [11] and older than 3.5 Ga [12].

1.2. Evolution of photosystem II

The heart of PSII is made up of a heterodimeric reaction centre (RC) *core* coupled to a core

antenna. The two subunits of the RC core of PSII are known as D1 and D2, and these are

associated respectively with the antenna subunits known as CP43 and CP47. D1 and CP43

make up one monomeric half of the RC, and D2 and CP47, the other half. Water oxidation is

catalysed by a Mn_4CaO_5 cluster coordinated by ligands from both D1 and CP43 [13, 14]. The

cluster is functionally coupled to a redox active tyrosine-histidine pair (Y_Z-H190) also

located in D1, which relays electrons from Mn to the oxidised chlorophyll pigments of the

RC during charge separation [15]. In a cycle of four consecutive light-driven charge

separation events, O_2 is released in the decomposition of two water molecules.

Photosystems evolved first as homodimers [16, 17]: therefore, the core and the

antenna of PSII originated from ancestral gene duplication events that antedated the MRCA

of Cyanobacteria. In this way, CP43/D1 retain sequence and structural identity with

CP47/D2. The conserved structural and functional traits between CP43/D1 and CP47/D2

suggest that the ancestral PSII homodimer—prior to the duplication events—was not only

structurally similar to heterodimeric PSII, but also that it split water and had evolved

protective mechanisms against the formation of reactive oxygen species [17-19].

In a previous study, we attempted to measure the span of time between the duplication

that led to D1 and D2 (dD0) and a point that approximated the MRCA of Cyanobacteria: a

period of time that we called **ΔT** [18]. We observed that the magnitude of ΔT can be very

large, well over a billion years. Such large ΔT suggested that the origin of a water-oxidising

photosystem antedated Cyanobacteria themselves and potentially other groups of Bacteria

depending on how rapidly the domain diversified. This implies that an ancestral non-

photosynthetic state at the divergence of Margulisbacteria, Sericytochromatia, and

Vampirovibrionia (MSV) cannot be taken for granted despite their specialized heterotrophic

lifestyles. However, our study neither provided an absolute age for the MRCA of

Cyanobacteria nor the duplication event itself, as we simulated a comprehensive range of

scenarios. Instead, we showed that even when ΔT is over a billion years, the rate of protein

evolution at the duplication point (dD0) needed to be over 40 times greater than any rate ever

observed for D1 and D2, decreasing exponentially during the Archean and stabilising at

current rates during the Proterozoic. Thus, the shorter the ΔT, the faster the rate at dD0, with

the rate increasing following a power law function and reaching unrealistic values even when

ΔT is still in the order of several hundred million years [18]. It was still unclear if such

patterns of evolution were unique to the core subunits of PSII or whether other systems have

experienced similar evolutionary trajectories.

Here, to help in understanding the evolution of oxygenic photosynthesis,

Cyanobacteria and MSV as a function of time, we compared the duplication leading to the

RC antenna subunits, CP43 and CP47, to several well-defined ancient and more recent

events: including, but not limited to, the duplication of the core catalytic subunits of ATP

synthase, a very ancient event generally accepted to have occurred before the last universal

common ancestor (LUCA) [20-24]; the evolution of RNA polymerase catalytic subunit β

(RpoB) and ribosomal proteins, which are universally conserved and widely accepted to have

originated before the LUCA [25-28]. We further constrain our analysis using *in silico*

ancestral sequence reconstruction of PSII and through strict structural and functional

rationales. We show that the core subunits of PSII show molecular signatures that are usually

associated with some of the oldest transitions in the molecular evolution of life. We also

show that all events leading from the primordial homodimeric water-splitting photosystem to

Cyanobacteria's heterodimeric PSII can be reconstructed from a comparison of available

structural and genetic data.

2. Results

2.1. Phylogenetic overview

The phylogenies of CP43 and CP47 show that there is a much greater diversity of CP43 and

CP43-derived subunits than CP47 (Figure 1). This difference is the result of a greater number

of gene duplication in CP43 than CP47 and mirrors the evolution of D1 and D2 [2], in which

D1 has undergone more duplication events than D2. CP43 can be divided into two major

groups: those that are assembled into PSII and can bind the Mn_4CaO_5 cluster, and those

which have evolved to be used only as light harvesting complexes [29, 30], usually known as

chlorophyll binding proteins (CBP). The CBP are characterized by the loss of the extrinsic

loop between the 5^{th} and 6^{th} transmembrane helices, where the ligands to the cluster are

located (Figure 1b). This large extrinsic loop is found in both CP43 and CP47 and interacts

directly with the electron donor side of PSII, within D1 and D2 respectively. The unrooted

tree of CP43 is consistent with CBP having a single origin likely occurring before the MRCA

of Cyanobacteria (Supplementary Figure S1) but have undergone an extensive duplication-

driven diversification process. It also mirrors the evolution of D1 in that duplications appear

to have occurred before the MRCA of Cyanobacteria [2]. The earliest of these D1

duplications also led to variants that have lost the capacity to bind the Mn_4CaO_5 cluster [2,

31], but are likely used in other supporting functions. Most notably, during chlorophyll f

synthesis [32] in the far-red light acclimation response (farlip) [33].

Figure 1. Maximum Likelihood trees of PSII core antenna subunits and derived light harvesting proteins. **a** A

tree of CP43 and chlorophyll binding proteins (CBP). The unrooted tree splits into CP43 and CBP, with CP43

displaying a phylogeny roughly consistent with the evolution of Cyanobacteria, although several potential

duplications of CP43 are noticeable within heterocystous Cyanobacteria and closer relatives. Farlip variant

denotes the isoform used in the far-red light acclimation response. CBP shows a complex phylogeny strongly

driven by gene duplication events and fast evolution. The tree of CP47 was rooted at the divergence of

Gloeobacter spp. Scale bar represents number of substitutions per site. **b** Structural comparison of CP43, CP47,

and IsiA. The latter lacks the characteristic extrinsic loop (orange) of CP43 and CP47 that links the antenna with

the electron donor side of PSII.

CP43 and CP47 are also distantly related to the antenna domain of cyanobacterial PSI and

that of the type I RCs of phototrophic Chlorobi, Acidobacteria and Heliobacteria

(Supplementary Figure S2). We found that the level of sequence identity between any two

type I RC proteins is always greater than between a type I RC protein and CP43/CP47

(Supplementary Table S1). Therefore, the distance between CP43/CP47 and other type I

antenna domains is the largest distance in the molecular evolution of RC proteins after that

between type I and type II RC core domains. The phylogenetic relationships between type I

and type II RCs have been reviewed in detail before [19, 34, 35]. These are briefly

summarized and schematized in Supplementary Figure S3.

The phylogeny of Alpha and Beta subunits of the F-type ATP synthase showed that

all Cyanobacteria have a F-type ATP synthase, and a fewer number of strains have an

additional Na^+-translocating ATPase (N-ATPase) of the bacterial F-type, as had been

reported before [36]. We found that MSV have a standard F-type ATP synthase

(Supplementary Figure S4), but some N-ATPase Alpha and Beta sequences were also found

in Vampirovibrionia and Sericytochromatia datasets, but not in Margulisbacteria. In this

study we focused on the standard F-type ATP synthase of Cyanobacteria for further analysis.

The phylogeny of bacterial RNA polymerase subunit β (RpoB), with the intention of

molecular clock analysis, focused on Cyanobacteria and MSV, as well as phyla with known

phototrophic representatives and included Thermotogae and Aquificae as potential roots

(Supplementary Figure S5). The tree was largely consistent with previous observed

relationships between the selected groups [37], within Cyanobacteria and MSV, and within

other phototrophs and their non-phototrophic relatives [38, 39]. The only exception was

Aquificae, which branched as a sister clade to Acidobacteria, a feature that had been reported

before for RpoB [27], and likely represents an ancient horizontal gene transfer event.

2.2. Distances and rates

To gather temporal information, we compared the phylogenetic distances between CP43 and

CP47, Alpha and Beta subunits of the F-type ATP synthase, and archaeal and bacterial RpoB

(visualized in Figure 2, but see also Figures S1b, S2 and S3b). We found that the distances

between Alpha and Beta, and the divergence of archaeal and bacterial RpoB, are very large

relative to the distance between the divergence of Vampirovibrionia and Cyanobacteria. In

the case of RpoB, the distance between Vampirovibrionia and Cyanobacteria is about a fifth

of the distance between Archaea and Bacteria. However, the distance between CP43 and

CP47 (but also between D1 and D2 [18]) is of similar magnitude to that between Alpha and

Beta, and to that between archaeal and bacterial RpoB, but substantially surpasses the

distance between MVS and Cyanobacteria (Figure 2). These observations suggest that

ancestral proteins to CP43/CP47 and D1/D2 existed before the divergences of MVS.

Figure 2. Distance comparison of the core subunits of ATP synthase, RNA polymerase, and PSII. **a** Alpha and

Beta subunits of the F-type ATP synthase from Cyanobacteria, Vampirovibrionia, Sericytochromatia and

Margulisbacteria. *N* denotes Na$^+$ translocating N-type ATPase. *d*AB denotes the duplication event leading to

Alpha and Beta. The green dot marks the MRCA of Cyanobacteria, and the light-blue dot the MRCA of the

clade including Cyanobacteria and Vampirovibrionia. **b** Archaeal and Bacterial RpoB of RNA polymerase. **c**

CP43 and CP47 subunits of PSII. *f* denotes farlip variants. *d*CP marks the duplication event leading to CP43 and

CP47. Pink branches highlight the CPB subunits. **d** D1 and D2 of PSII showing a pattern that mimics that of

CP43 and CP47. Pink branches denote the atypical D1 forms and other variants that are thought to predate the

MRCA of Cyanobacteria. The sequences marked with *dominant* represent the standard D1 form of PSII,

We compared the within-group mean distances for Alpha, Beta, RpoB, and a concatenated

dataset of ribosomal proteins compiled in a previous independent study [38] (see

Supplementary Table S2). We found consistently, that Vampirovibrionia and

Margulisbacteria have larger within-group mean distances compared to Cyanobacteria, which

suggests greater rates of evolution in the non-photosynthetic clades. These were consistently

larger for Margulisbacteria relative to the other two groups. For example, RpoB in

Vampirovibrionia and Margulisbacteria showed 1.6x and 4.0x larger corrected mean

distances than Cyanobacteria, respectively (Supplementary Table S2). At the level of the

concatenated ribosomal proteins dataset, Margulisbacteria showed an almost 2-fold larger

within-group mean distance than Cyanobacteria.

We then compared the rates of evolution of CP43 and CP47 with those of Alpha and

Beta, using a Bayesian relaxed molecular clock approach with identical calibrations,

molecular clock parameters, and a simplified, but highly constrained sequence dataset (see

Materials and Methods for an expanded rationale). The goal of these experiment is not to use

the clock to estimate divergence times, but to measure and compare the rates of protein

evolution that are required to simulate any chosen span of time between the ancestral

duplications and the MRCA of Cyanobacteria. We used an autocorrelated log normal model

of rate variation with a non-parametric CAT+Γ model of amino acid substitutions to extract

rates of evolution. We will refer to the span of time between the duplication points leading to

Alpha and Beta (*d***AB**), or to CP43 and CP47 (*d***CP**), and the MRCA of Cyanobacteria as ΔT

(schematized in Figure 3).

In Figure 4a to d we examine the changes in the rate of evolution under specific

evolutionary scenarios. In the case of ATP synthase, we first assumed that the MRCA of

Cyanobacteria occurred after the GOE to simulate scenarios similar to those presented in [8]

or [11], at about 1.7 Ga; and that *d*AB occurred at 3.5 Ga (**ΔT = 1.8 Ga**). Under these

conditions the average rate of evolution of Alpha and Beta is calculated to be 0.28 ± 0.06

substitutions per site per Ga (δ Ga^{-1}). We will refer to the average rate through the

Proterozoic as ν_{min}. In this scenario, the rate of evolution at the point of duplication, which we

denote ν_{max}, is 7.32 ± 1.00 δ Ga^{-1}, making ν_{max}/ν_{min} 26. In other words, when the span of time

between the ancestral pre-LUCA duplication and the MRCA of Cyanobacteria is 1.8 Ga, the

rate of evolution at the point of duplication is about 26 times greater than any rate observed

through the diversification of Cyanobacteria or photosynthetic eukaryotes. To place these

rates in the larger context of protein evolution, we encourage the reader to refer to

Supplementary Text S1.

Figure 3. Schematic representation of ancestral duplication events. The MRCA of Cyanobacteria inherited a

standard F-type ATP synthase, with a heterohexameric catalytic head (F_1) made of alternating subunits Alpha

and Beta; and a PSII with a heterodimeric core and antenna. ΔT denotes the span of time between the ancestral

duplication events and the MRCA of Cyanobacteria, which in the case of ATP synthase, is suspected to antedate

the divergence of Bacteria and Archaea and their further diversification. FtsH contains an N-terminal

membrane-spanning domain attached to a soluble domain consisting of an AAA (ATPase associated with

diverse activities) module attached to a C-terminal protease domain. FtsH is universally conserved in Bacteria,

has a hexameric structure like that of ATP synthase's catalytic head, and can be found usually as

homohexamers, but also as heterohexamers. The MRCA of Cyanobacteria likely inherited three variant FtsH

subunit forms, one of which appears to have duplicated after the divergence of the genus *Gloeobacter*, and

possibly other early-branching Cyanobacteria [40]. This late duplication led to FtsH1 and FtsH2, which form

heterohexamers with FtsH3, following the nomenclature of Shao, et al. [40] FtsH1/FtsH3 is found in the

cytoplasmic membrane of Cyanobacteria, while FtsH2/FtsH3 is involved in the degradation of PSII and other

thylakoid membrane proteins.

Now, if we consider a scenario in which dAB is 4.0 Ga and leaving all other constraints

unchanged, v_{max} is 6.02 ± 0.9 δ Ga^{-1} resulting in a v_{max}/v_{min} of 21. If instead we keep the

duplication at 3.5 Ga but assume that the MRCA of Cyanobacteria occurred before the GOE

to simulate a more conservative scenario, at 2.6 Ga (**ΔT = 1.1 Ga**), we obtain that v_{min} is

consequently slower, 0.25 ± 0.06 δ Ga^{-1}, when compared to a post-GOE ancestor. This older

MRCA (smaller ΔT) thus leads to a rise in v_{max}, calculated to be 10.22 ± 1.37 δ Ga^{-1} and

leading to a v_{max}/v_{min} of 40. Given that the phylogenetic distance is a constant, the rate of

evolution increases with a decrease in ΔT following a power law function. We had observed

nearly identical evolutionary patterns for the core RC proteins D1 and D2 of PSII [18]. The

change in v_{max}/v_{min} as a function of ΔT is shown in Figure 4d.

Figure 4. Comparison of the changes in the rates of evolution as a function of time. **a** Rates of CP43 and CP47 modelled using two specific evolutionary scenarios. Orange trace was calculated under the assumption that the MRCA of Cyanobacteria occurred after the GOE, at ~1.7 Ga, while the duplication of CP43 and CP47 occurred at ~3.5 Ga. The fastest rate seen at the point of duplication is denoted as v_{max}, and stabilizes during the Proterozoic, v_{min}. Grey dots represent a scenario in which the MRCA of Cyanobacteria is thought to have occurred before the GOE, at ~2.6 Ga instead. **b** Same calculations as **a** but performed on Alpha and Beta sequences of cyanobacterial and plastid ATP synthase. **c** Rates of evolution of cyanobacterial and plastid FtsH subunits assuming a MRCA of Cyanobacteria at 1.7 and 2.6 Ga. **d** Changes in the rates of evolution with varying ΔT for ATP synthase (orange) and PSII subunits (grey). **e** Changes in the rates of evolution as a function of time for a relaxed molecular clock computed with the full dataset of CP43 (full circles) and CBP sequences (open circles). **f** Changes in the rate of evolution of bacterial RpoB showing an exponential decrease in the rate of evolution. The bar denotes the span of time between the MRCA of the clade containing Vampirovibrionia and Cyanobacteria and the MRCA of Cyanobacteria (0.44 Ga). **g** Changes in the rate of evolution of bacterial and archaeal RpoB. The longer bar represents the span of time between the divergence of Archaea and Bacteria and the MRCA of Cyanobacteria (1.21 Ga). **h** Changes in the rate of evolution of a dataset of concatenated ribosomal proteins showing similar patterns of evolution as RpoB. Error bars on each data point are standard errors on the rate and mean divergence time.

The core antenna of PSII, CP43 and CP47, showed patterns of divergence very similar to those of Alpha and Beta, both in terms of phylogenetic distances between paralogues and rates of evolution between orthologues (Figure 4a and b). The average rate of evolution of CP43 and CP47, assuming that the MRCA of Cyanobacteria occurred at 1.7 Ga, and the duplication (*d*CP) at 3.5 Ga (**ΔT = 1.8 Ga**), is 0.19 ± 0.05 δ Ga^{-1}. Slower than for Alpha and Beta under the same condition. This slower rate is consistent with the fact that CP43 and CP47 show less sequence divergence between orthologues at all taxonomic ranks of oxygenic phototrophs when compared to Alpha and Beta (see Supplementary Table S3). Furthermore, the rate at *d*CP, v_{max}, was 5.17 ± 0.84 δ Ga^{-1}, generating a v_{max}/v_{min} of 27, similar to Alpha and Beta (Figure 4). Thus, even when ΔT is 1.8 Ga, the rate at duplication point needs to be 27 times greater than the average rates observed during the Proterozoic. If we consider instead that the MRCA of Cyanobacteria occurred at 2.6 Ga and *d*CP at 3.5 Ga (**ΔT = 1.1 Ga**), this would slowdown v_{min} to 0.16 ± 0.04 δ Ga^{-1}, while v_{max} would increase to 7.81 ± 1.01 δ Ga^{-1} resulting in a v_{max}/v_{min} of 49. Therefore, the molecular evolution of the core subunits of PSII parallels that of ATP synthase both in terms of rates and distances through geological time.

We then studied a relatively recent gene duplication event (Figure 4c), which occurred long after the LUCA, but also after the MRCA of Cyanobacteria: that leading to

Cyanobacteria-specific FtsH1 and FtsH2 (*d*H0) [40]. This more recent duplication served as a

point of comparison and control (see Figure 3 for a scheme). In marked contrast to *d*AB, the

rate at the point of duplication was 0.66 ± 0.21 δ Ga^{-1}. We found that FtsH1 is evolving at an

average rate of 1.42 ± 0.29, while FtsH2 at a rate of 0.24 ± 0.06 δ Ga^{-1} under the assumption

that MRCA of Cyanobacteria occurred at 1.7 Ga. Thus, under the assessed conditions, FtsH1

is evolving about 5.3 times faster than FtsH2, while the latter is evolving at a rate similar to

that of Alpha and Beta. If the MRCA of Cyanobacteria is assumed to have occurred at 2.6

Ga, all rates slowdown respectively, but the rate of FtsH1 remains over five times faster than

FtsH2. Unlike *d*AB, *d*H0 is consistent with classical neofunctionalization, in which the copy

that gains new function experiences an acceleration of the rate of evolution [41, 42]. Like

PSII and ATP synthase, the calculated rates of evolution match observed distances as

estimated by the change in the level of sequence identity as a function of time, in which the

fastest evolving FtsH1 accumulated greater sequence change than FtsH2 in the same period

(Supplementary Table S3).

Given that the complex evolution of CP43 and CBP involved several major

duplication events and potentially large variations in the rate of evolution (Figure 1 and

Supplementary Figure S1), we carried out a molecular clock analysis of a large dataset of 392

CP43 and 465 CBP proteins using cross-calibrations across paralogues. Clocks were executed

with no constraint on the MRCA of Cyanobacteria. The estimated mean divergence time for

the oldest node, the duplication at the origin of CBP, is 2.23 Ga (95% confidence interval, CI:

1.90 - 2.69 Ga) using an autocorrelated log normal model (see Figure 4e, Supplementary

Figure S6 for a chronogram, and Supplementary Table S4 for a comparison of estimated ages

under different models). The mean divergence time for the node representing the CP43

inherited by the MRCA of Cyanobacteria was calculated to be 2.22 Ga (95% CI: 1.88 - 2.68

Ga). Thus, a span of time of only 15 Ma is measured between these two mean ages. The

average rate of evolution of CP43, not including CBP sequences, was found to be 0.14 ± 0.05

δ Ga^{-1}, which is in the same range as determined in the simplified, but highly constrained

experiment above. We noted a 6-fold increase in the rate of evolution associated with the

duplication leading to the farlip-CP43 variant (Figure 4e). This duplication led to an

acceleration of the rate similar in magnitude to that of FtsH1/FtsH2 and is consistent with a

neofunctionalization as the photosystems evolved to use far-red light and bind chlorophyll *f*.

CBP sequences, on average, display rates of evolution about three times faster than

CP43 (Figure 4e). However, the serial duplications that led to the evolution of CP43-derived

light harvesting complexes resulted in accelerations in the rate of evolution of a similar

magnitude as observed for *d*AB and *d*CP. The largest of these is associated with the origin of

PcbC [30], a variant commonly found in heterocystous Cyanobacteria and Cyanobacteria that

use alternative pigments, such as chlorophyll *b*, *d* and *f*. The ancestral node of PcbC was

timed at 2.07 Ga (95% CI: 1.76 - 2.50 Ga) with a rate of $11.7 \pm 2.42 \, \delta \, Ga^{-1}$, decelerating

quickly, but stabilizing at about four times faster rates than the average rate of CP43. We find

it noteworthy that the fast rates of evolution associated with the origin of CBP are not

associated with very large spans of time between these and CP43, nor did it result in very old

root node ages despite the use of very broad constraints.

2.3. Species divergence

To understand the evolution of MSV relative to Cyanobacteria we wished to apply a

molecular clock to a system where the calculated rates could be compared to observed rates

as determined by distances between species of known divergence times or at similar

taxonomic ranks. We found RpoB to be suitable for this because it has been inherited

vertically with few instances of horizontal gene transfer and had enough signal to resolve

known phylogenetic relationships between and within clades. In collecting the RpoB

sequences, we noted for the first time that Margulisbacteria and Vampirovibrionia share a

comparatively greater level of divergence at similar taxonomic ranks than Cyanobacteria. For

example, the level of sequence divergence of RpoB from two species of *Termititenax*

(Margulisbacteria) [43] is about 40% greater than the distance between *Gloeobacter* spp. and

any other cyanobacterium (not including gaps and insertions), the latter being the largest

distance between oxygenic phototrophs. In the case of Gastranaerophilales

(Vampirovibrionia) [5], which are specialised gut bacteria and should therefore not be much

older than animals, the level of sequence identity of RpoB was found to be 70% for the two

most distant strains in this group, contrasted to 84% for *Gloeobacter* spp. when compared to

any other cyanobacterium. As listed in Supplementary Table S2, within-group mean

distances suggest that faster rates are widespread and not just unique to RpoB.

We implemented a set of 12 calibrations across bacteria, including two calibrations on

Margulisbacteria and two in Vampirovibrionia with the aim of covering both slower and

faster evolving lineages. The following results are based on an autocorrelated log normal

molecular clock using CAT+Γ, a root constrained with a broad interval ranging from between

4.52 and 3.41 Ga, and as described in Materials and Methods (Figure 5). We found this to

perform well and provided results comparable to other independent studies that did not

combine a full set of MVS sequences and other clades with phototrophs in a single tree

(Table 1). Nonetheless, a pipeline of sensitivity experiments tested the dependency of these

results on models and prior assumptions: these are shown and described in Supplementary

Figure S7 and S8.

The root of the tree (divergence of Thermotogae) was timed at 3.64 Ga (95% CI: 3.42

- 4.11 Ga) and the divergence of Cyanobacteria at 2.74 Ga (95% CI: 2.46 - 3.12 Ga). Thus,

the span of time between the mean age of the root and the mean age of the MRCA of

Cyanobacteria was calculated to be 0.89 Ga. The span of time between Margulisbacteria and

Cyanobacteria was found to be 0.67 Ga; and between Vampirovibrionia and Cyanobacteria

0.44 Ga (Figure 4f and 5). The latter is a value that is consistent with previous studies using

entirely different rationales, datasets and calibrations [8, 9].

We also noted an exponential decrease in the rates of evolution of RpoB through the

Archean, which stabilised at current levels in the Proterozoic (Figure 4f). The rate at the root

node was calculated to be $2.37 \pm 0.45\ \delta\ Ga^{-1}$ and the average rate of evolution of RpoB

during the Proterozoic was found to be $0.19 \pm 0.06\ \delta\ Ga^{-1}$. The average rate of cyanobacterial

RpoB was $0.14 \pm 0.04\ \delta\ Ga^{-1}$; for Margulisbacteria was $0.44 \pm 0.17\ \delta\ Ga^{-1}$, and for

Vampirovibrionia $0.19 \pm 0.05\ \delta\ Ga^{-1}$: about 3.1x and 1.3x the mean cyanobacterial rate

respectively. These rates agree reasonably well with the observed distances (Supplementary

Table S2), further indicating that the calibrations used in these clades performed well and

recapitulated patterns of evolution consistent with lifestyle and trophic modes. Nevertheless,

we suspect that the values for MSV represent underestimations of the true rates of evolution

(slower than they should be), as some of the clades that include symbionts still appear much

older than anticipated from their hosts (Supplementary Text S2).

Furthermore, a more complex, but commonly used model like CAT+GTR+Γ

implementing a birth-death prior with 'soft bounds' on the calibrations, resulted in rates that

were smoothed out, which translated into spread-out divergence times with Margulisbacteria

and Vampirovibrionia evolving at 1.9x and 0.7x times the cyanobacterial rates, respectively

(Supplementary Figure S8, model **n** to **p**). These weird rate effects are thus translated into a

Mesoproterozoic, very late, age for Cyanobacteria, and a relatively older divergence time for

Vampirovibrionia: results that replicate those presented recently in ref. [11].

Figure 5. Bayesian relaxed molecular clock of bacterial RpoB. The tree highlights the spans of time between the MRCA of Cyanobacteria (green dot) and their closest relatives. Calibrated nodes are marked with red dots. Anoxygenic phototrophs are highlighted in red branches and non-phototrophic bacteria in grey, with the exception of Margulisbacteria, Sericytochromatia and Vampirovibronia, which are coloured as indicated in the figure. Superimposed at the top are the implied distribution and divergence time for ATP synthase and PSII. Horizontal bars within the tree mark 95% confidence intervals. These are shown in selected nodes of interest for clarity but see Table 1.

Table 1. Divergence time estimates

Event	RpoB (Ga) (95% CI)	Concatenated ribosomal proteins	TimeTree compilation[a]	Shih et al.[b]	Magnabosco et al.[c]
Divergence of Thermotogae	3.64 (3.43 - 4.11)	3.01 (2.64 - 3. 51)	3.81 (3.51 - 4.16)		
Divergence of Margulisbacteria	3.42 (3.17 - 3.86)	2.93 (2.58 - 3.44)			
Divergence of Sericytochromatia	3.26 (3.01 - 3.68)				
Divergence of Vampirovibrionia/Stem Cyanobacteria	3.19 (2.92 - 3.61)	2.76 (2.40 - 3.23)	3.19 (2.26 - 3.55)	2.54 (2.09 - 3.06)	2.77 (2.39 -2.93)
MRCA of Cyanobacteria	2.75 (2.46 - 3.13)	2.22 (1.86 - 2.64)	2.24 (1.81 - 2.82)	2.02 (1.72 - 2.37)	2.24 (1.91 -2.41)
Divergence of Heliobacteria	2.28 (1.74 - 2.78)		2.23 (1.99 - 2.48)		
MRCA of Heliobacteria	1.78 (1.13 - 2.36)				
Divergence of phototrophic Chloroflexi	1.83 (1.41 - 2.25)		1076	1.098 (0.80 - 1.41)	2.86 (2.49 - 3.00)
MRCA of phototrophic Chloroflexi	1.71 (1.24 - 2.16)			0.86 (0.61 - 1.14)	1.98 (1.68 - 2.43)
Divergence of phototrophic Chlorobi	2.80 (2.45 - 3.20)				2.56 (2.26 - 2.85)
MRCA of phototrophic Chlorobi	1.91 (1.53 - 2.31)		525		1.71 (1.64 -1.95)
Earliest phototrophic Proteobacteria	2.82 (2.55 - 3.21)		2.48 (2.04 - 2.62)		
Divergence of *Chloracidobacterium*	2.04 (1.53-2.51)				

[a]Data from TimeTree compiling estimated divergence times from independent studies as described in ref. [44] For the divergence of Thermotogae the estimated age was taken from eight different studies (n=8). For the divergence of Melaibacteria (or divergence from Cyanobacteria's closest relatives), n=10. For stem Heliobacteria, n=2. For stem phototrophic Chloroflexi and Chlorobi, the values were reported from a single study, Marin, et al. [45] For earliest potential phototrophic Protepbacteria, n=3. This latter was taken as the node made by the clades including *Niveispirillum lacus* and *Rhodobacter sphaeroides* to match our RpoB tree. The independent studies used by TimeTree to generate the estimated ages are listed in Supplementary Table S8.

[b]Data taken from ref. [8] Model T68 for the cyanobacterial dates. That is, no GOE calibration with a calibration on *Bangiomorpha*. The dates on Chloroflexi were taken from ref. [46]

[c]Data taken from ref. [9] Model A. This used a 1.2 Ga calibration on heterocystous Cyanobacteria.

To investigate the effect of the age of the root on the divergence time of MVS and Cyanobacteria we also varied the root prior from 3.2 to 4.4 Ga (Supplementary Figure S7). We noted that regardless of the time of origin of Bacteria (approximated by the divergence of Thermotogae in our analysis), a substantially faster rate is required during the earliest diversification events, decreasing through the Archaean and stabilizing in the Proterozoic.

This matches well the patterns of evolution of ATP synthase and PSII core subunits as shown

in the previous section.

We then compared the above RpoB molecular clock with a different clock that

included a set of 112 diverse sequences from Archaea, in addition to the sequences from

Thermotogae, MSV, and Cyanobacteria, but removing all other bacterial phyla (Figure 6a).

We found that the calculated average rate of evolution of bacterial RpoB during the

Proterozoic was slower (0.09 ± 0.03 δ Ga^{-1}) than in the absence of archaeal sequences ($0.19 \pm$

0.06 δ Ga^{-1}), resulting in overall older mean ages (see Figure 4g). However, the rate at the

Bacteria/Archaea divergence point was 6.87 ± 1.17 δ Ga^{-1}, similar to the rate for dAB and

dCP, requiring therefore an exponential decrease in the rates similar to that observed for ATP

synthase and the PSII core subunits. Notably, this rate and exponential decrease is associated

with a span of time between the mean estimated ages of the LUCA and the MRCA of

Cyanobacteria of 1.21 Ga. The span between Vampirovibrionia and Cyanobacteria was found

to be 0.46 Ga (Figure 6a).

Figure 5 also highlights that the MRCA of none of the groups containing anoxygenic

phototrophs nor their divergence from their closest non-phototrophic relatives appears to be

older than one of the oldest and best accepted geochemical evidence for photosynthesis at

3.41 Ga [47] (Table 1). For example, the MRCA of Heliobacteria or of phototrophic

Chlorobi, containing homodimeric type I RCs, which have traditionally been described as to

harbour primitive forms of photosynthesis, are likely to have existed after the GOE, even

allowing for large uncertainties in the calculations.

Finally, we performed a similar molecular clock analysis on a subset of concatenated

ribosomal proteins published by Hug et al. [38] that included Archaea, Thermotagae,

Margulisbacteria, Vampirovibrionia and Cyanobacteria. Even though this dataset generated

younger ages compared to RpoB (see Table 1 and Figure 6), a similar exponential decrease in

the rates was observed with a rate at the Bacteria/Archaea divergence point measured at 4.62

± 0.71 δ Ga^{-1} and an average rate of bacterial ribosomal protein evolution of 0.30 ± 0.07 δ Ga^{-}

^{1} (Figure 4h). The span of time between the mean estimated ages of the LUCA and the

MRCA of Cyanobacteria was 1.45 Ga; and between Vampirovibrionia and Cyanobacteria

0.54 Ga (Figure 6b). If one assumes therefore that the span of time between the LUCA and

the diversification of Bacteria is narrower, that would imply a faster rate at the root node and

a steeper deceleration of the rates.

Figure 6. Comparison of relaxed molecular clocks of RpoB (**a**) and a concatenated dataset of ribosomal proteins (**b**) that include sequences from Archaea and as described in the main text. RNAP stands for RNA polymerase. Clocks were calculated using a calibration on the root with a maximum of 4.52 Ga and a minimum of 3.41 Ga and applying a log normal autocorrelated clock with CAT+Γ model. Bars on selected nodes denote 95% confidence intervals.

2.4. Structural analysis

A fundamental premise of our investigation is that water oxidation started before the duplication of D1 and D2, and CP43 and CP47. The rationale behind this premise has been laid out before [17, 48], and more extensively recently [18]. This rationale raises the question of how the D2/CP47 side of the RC lost its capacity to carry out water-splitting catalysis. To gain further insight on the nature of the structural site around the water-oxidising complex in the ancestral photosystem, we used ancestral sequence reconstruction to predict the most

probable ancestral states. We will refer to the ancestral protein to D1 and D2 as D0

(Supplementary Figure S9). We generated 14 predicted D0 sequences using a combination of

three ASR methods and amino acid substitution models. On average the 14 D0 sequences had

$87.12 \pm 0.55\%$ sequence identity indicating that the different algorithms provided largely

consistent results. While the regions that include all transmembrane helices are aligned

unambiguously, the N-terminal and C-terminal ends were aligned less confidently due to

greater sequence variability at both ends. Nonetheless, we found that the predicted D0

sequences retain more identity with D1 than D2 along the entire sequence. The level of

sequence identity of D0 compared to the D1 (PsbA1) of *Thermosynechococcus vulcanus* was

found to be $69.58 \pm 0.55\%$, and $36.32 \pm 0.15\%$ compared to D2.

The ligands to the Mn_4CaO_5 in PSII are provided from three different structural

domains (Supplementary Figure S10): 1) the D1 ligands D170 and E189, located near the

redox Y_Z. These are in the lumenal loop between the 3^{rd} and 4^{th} transmembrane helices. 2)

The D1 ligands H332, E333, D342, and A344 located at the C-terminus. 3) The CP43

ligands, E354 and R357, located in the extrinsic loop between the 5^{th} and 6^{th} helices, with the

latter residue less than 4 Å from the Ca. Remarkably, there is structural and sequence

evidence supporting the loss of ligands in these three different regions of CP47/D2.

In all the D0 sequences, at position D1-170 and 189, located in the unambiguously

aligned region, the calculated most likely ancestral states were E170 and E189, respectively.

The mutation D170E results in a PSII phenotype with activity similar to that of the wild-type

[49]. At position D1-170, a glutamate was predicted with average posterior probabilities

(PPs) of 44.2% (FastML), 67.2% (MEGA) and 77.0% (PAML). At position D1-E189, the

average PPs for glutamate were 31.1% (FastML), 35.2% (MEGA) and 40.2% (PAML). The

distribution of PPs across a selection of D0 sequences and key sites is presented in

Supplementary Figure S11 and Table S5. In contrast, D2 has strictly conserved phenylalanine

residues at these positions, but the PP of phenylalanine being found at either of these

positions was less than about 5% for all predicted D0 sequences. As a comparison, the redox

active tyrosine residues Y_Z (D1-Y161) and Y_D (D2-Y160), which are strictly conserved

between D1 and D2, have a predicted average PPs of 68.8% (FastML), 98.8% (MEGA) and

98.6% (PAML). Therefore, the ligands to the catalytic site in the ancestral protein leading to

D2 were likely lost by direct substitutions to phenylalanine residues, while retaining the

redox active D2-Y160 (Y_D) and H189 pair (Supplementary Figure S10).

Figure 7. Structural rearrangements of the large extrinsic loop of CP43 and CP47. **a** The antenna domain of heliobacterial PshA is shown in three different rotated perspectives. Only the first six transmembrane helices of the antenna are shown for clarity. A Ca at the electron donor side is bound from an extrinsic loop between the 5th (E) and 6th (F) helices. This extrinsic loop, EF_1 (blue), is made of two small alpha helices. The fourth molecular view furthest to the right shows the link between the electron donor site and EF_1 in closer detail. **b** The CP43 subunit of PSII with the extrinsic loop shown in colours. **c** The CP47 subunit of PSII. Immediately after the 5th helix (E), a long alpha helix protrudes outside the membrane in both CP43 and CP47 and showing structural and sequence identity (orange). We denote this helix EF_2. After EF_2 structural differences are noticed

between CP43 and CP47 as schematised in panel **d**. In CP43, after helix EF_2 a loop is found (shown in red

ribbons), which we denote EF_3. This contains the residues that bind the Mn_4CaO_5 cluster and it is followed by a

domain that resembles EF_1 in the HbRC at a structural level. In CP47, EF_3 and EF_1 retain sequence identity with

the respective regions in CP43. CP47 has additional sequence that is not found in CP43 (EF_4, purple). The green

arrows mark the position at which the domain swap occurred in CP43 relative to CP47. We found that the

CP43-E354 and R357 are found in the equivalent domain in CP47 as E436 and N438 coordinating a Ca atom.

N438 (EF_3) links to EF_1 via K332. It is unclear if the EF_1 region in the HbRC is strictly homologous to that in

CP43 and CP47 as very little sequence identity is found between the two: however, a couple of conserved

residues between all EF_1 may suggest it emerged from structural domains present in the ancestral RC protein

(see Supplementary Figure S13).

Prompted by the finding of a Ca-binding site at the electron donor site of the homodimeric

type I RC of Heliobacteria (Firmicutes) with several similarities to the Mn_4CaO_5 cluster of

PSII, including a link to the antenna domain and the C-terminus [19], we revisited the

sequences and structural overlaps of CP43 and CP47. We found that a previously unnoticed

structural rearrangement within the extrinsic loop occurred in one subunit relative to the other

(marked EF_3 and EF_4 in Figure 7, Supplementary Figure S12 and S13). CP43 retains the

simplest domain, being about 60 residues shorter than CP47. If CP43 retains the ancestral

fold, the additional sequence in CP47's swapped domain (EF_4 in Figure 7d) would have

contributed to the loss a catalytic cluster as it inserted one phenylalanine residue (CP47-

F360) into the electron donor site, less than 4Å from Y_D. An equivalent residue does not exist

in CP43.

We then noted that in the swapped region (EF_3 in Figure 7d), sequence identity is

retained between CP43 and CP47 (Supplementary Figure S12). We found that CP43-E354

and R357 are equivalent to CP47-E435 and N438. An inspection of the crystal structure of

cyanobacterial PSII showed that these two residues specifically bind a Ca of unknown

function in (Figure 7c and d). The presence of an equivalent glutamate to CP43-E354 in

CP47 is consistent with this being already present before duplication.

Finally, a peculiar but well-known trait conserved across Cyanobacteria and

photosynthetic eukaryotes is that the 5' end of the *psbC* gene (CP43) overlaps with the 3' end

of the *psbD* gene (D2) usually over 16 bp (Supplementary Table S6). The *psbD* gene contains

a well-defined Shine-Dalgarno ribosomal binding site downstream of the *psbC* gene and over

the coded D2 C-terminal sequence [50-52]. The evolution of this unique gene overlap has no

current explanation in the literature, but its origin could have disrupted the C-terminal ligands

of the ancestral protein leading to the modern D2, an event that would have contributed to

and favoured heterodimerization.

3. Discussion

3.1. Origin of oxygenic photosynthesis and rates of evolution

The duplication of ATP synthase's ancestral catalytic subunit, and the archaeal/bacterial

divergence of RNA polymerase and the ribosome, are some of the oldest evolutionary events

known in biology [20-28]. When taken on their own, their phylogenetic features are often

interpreted as evidence of the earliest origin. We have shown in here and in our previous

work [18] that PSII show patterns of molecular evolution that closely parallel those of these

very ancient systems and independently of the exact time of origin of Cyanobacteria or their

closest relatives. Therefore, it cannot be taken for granted that there was ever a long period of

time between the origin of life and the origin of anoxygenic photosynthesis, followed by

another long period of time between the origin of anoxygenic photosynthesis and the origin

of oxygenic photosynthesis.

The exponential decrease in the rates of evolution observed in the studied systems,

even when the span of time between the ancestral duplications (or the LUCA) and the MRCA

of Cyanobacteria was well over a billion years, are consistent with our assessment that a large

ΔT for the evolution of the core subunits of ATP synthase and PSII cannot be explained

entirely by duplication-driven effects. Instead, we speculate that these effects are the result of

the faster rates of evolution that are expected to have occurred during the earliest history of

life due to unrestricted positive selection at the origin of bioenergetics, accelerated rates of

mutations caused by environmental factors [53-55], lack of sophisticated DNA repair

mechanisms [56], or other genetic properties attributed to early life [57, 58], yet in

combination with very large spans of time.

It is worth highlighting that ATP synthase, RNA polymerase, the ribosome, and the

photosystems, are all complex molecular machines, with crucial functions and under strict

regulation. These features largely explain their slow rates of evolution and the high degree of

sequence (structural and functional) conservation through geological time. A similar level of

complexity, however, can be traced back to the last common ancestor of each one of these

molecular systems regardless of how ancient they truly are. Now, given that the rates of

evolution of the core of PSII have remained slower than those of ATP synthase for billions of

years, even through evolutionary processes that could be associated with acceleration in the

rates of evolution such as: 1) primary endosymbiosis at the origin of photosynthetic

eukaryotes [59], 2) radiations within clades of flowering plants [60], or 3) the radiation of

marine *Prochlorococcus* and *Synechococcus* [61, 62], just to name a few; if those rates have

remained proportional since the origin of the enzymes, it could imply that the duplications of

the core of PSII occurred before the duplication leading to Alpha and Beta.

In consequence, to argue that oxygenic photosynthesis is a late evolutionary

innovation relative to the origin of life, one must first demonstrate that the rates of protein

evolution of PSII and ATP synthase catalytic core subunits, RNA polymerase, and the

ribosome, have not remained proportional to each other through geological time. In such a

way that PSII—exclusively—experienced unprecedentedly faster rates of evolution. One

could potentially argue that the origin of water oxidation itself could account for this

hypothetical unprecedentedly faster rates. However, because of the relationships between

speed, *distance*, and *time*, a decrease in ΔT of about 60% (from 1.1 to 0.46 Ga for example),

would result in a 30-fold additional increase in the rates at the earliest stages of

diversification, resulting in photosystems that would be evolving at rates that surpass those of

the fastest evolving peptide toxins (Figure 4d and Supplementary Text S1). It should be

noted, that a ΔT of over a billion years already accounts for a period of fast evolution at the

origin of all these ancient systems.

The reader should also be reminded that these patterns of evolution, though they

might appear somewhat unusual, do not emerge from the application of any particular

molecular clock approach or computational analysis, but from the inherently long

phylogenetic distance that separates Alpha and Beta, Archaea and Bacteria, or the core

subunits of PSII, together with relatively slow rates of evolution throughout their entire

multibillion-year diversification process.

3.2. Diversification of Bacteria

It has been postulated before that the diversification of the major phyla of Bacteria occurred

very rapidly, starting at about 3.4 Ga and peaking at about 3.2 Ga, in what was dubbed as the

Archean Genetic Expansion [63]. Another recent independent molecular clock analysis put

the MRCA of Bacteria at about 3.4 Ga [11]; and Marin et al. [45] placed the major bacterial

radiation, except for the divergence of Thermotogae and Aquifecae, starting at a mean

divergence time of about 3.2 Ga too. Our clocks of RpoB or ribosomal proteins are also in

agreement with these patterns. A recent study suggested that the major groups of Bacteria and

Archaea diversified rapidly after the LUCA and hypothesized that the long distance between

archaeal and bacterial ribosomal proteins could be attributed to fast rates of ribosome

evolution during their divergence [64], but see also [65]. Our analyses suggest that the more

explosive the diversification of prokaryotes, the greater the chance that photosynthetic water-

splitting is an ancestral trait of all life. And yet, if phototrophic communities—whether

oxygenic or not—already existed 3.2 to 3.4 Ga ago, as it is supported by the geochemical

record [47, 66], then the earliest bacteria were likely photosynthetic. That the earliest bacteria

were photosynthetic is entirely consistent with the evolution of RC proteins, which indicates

that the structural and functional specialization that led to the two photosystem types,

antedated the diversification processes leading to the known phyla containing photosynthetic

bacteria [34]. This is also consistent with the fact that none of the groups of extant

photosynthetic bacteria appear to be older than the earliest geochemical evidence for

photosynthesis.

The presented data is also consistent with recent studies of the oxygenation of the

planet, which suggests that even if photosynthetic O_2 evolution started as early as the oldest

rocks, the properties of early Earth biogeochemistry would have maintained very low

concentrations of O_2 over the Archean without the need to invoke any particular biological

innovation or trigger to coincide with the GOE [67-69].

3.2. Structural constraints

We showed that the phylogenetic distance between CP43/CP47 and other type I RC proteins

is the second largest distance after that between type I and type II (Supplementary Figure S2).

It is conventionally considered that the first six transmembrane helices of the photosystems

make up the antenna domain, while the photochemical core encompasses the last five helices.

Structurally and functionally the antenna domain extends to the 8^{th} helix, both in type I RCs

and in PSII; with the latter retaining one antenna chlorophyll in the equivalent 8^{th} helix

(marked Z in Figure 8), as well as substantial sequence identity around this chlorophyll's

binding site as shown before [34, 70]. These conserved features indicate that there is not a

moment in time during the evolution of PSII in which it was devoid of core antenna proteins.

In other words, CP43/CP47 are a descendant of the ancestral core antenna of type II RCs,

now lost and replaced in anoxygenic type II RCs of phototrophic Proteobacteria, Chloroflexi

and others. Both structural and phylogenetic evidence are in agreement and indicate that the

ancestral type II RC, at the dawn of photosynthesis, was architecturally like water-splitting

PSII. The above disproves and supersedes a previous hypothesis by Cardona [35] where the

antenna of PSII was claimed to have originated from a refolding of an entire type I RC

protein.

Figure 8. Structural comparisons of the antenna and core domains of photosynthetic RCs. **a** A monomeric unit

(PshA) of the homodimeric RC of Heliobacteria (HbRC, Firmicutes). The RC protein is made up of 11

transmembrane helices. The first six N-terminal helices are traditionally considered as the antenna domain

(orange ribbons), while the last five helices are the RC core domain (grey and light-blue ribbons). Below the

structure, the organization of the 11 helices is laid down linearly for guidance: 1^N denotes the first N-terminal

helix and 11_C the last C-terminal helix. P denotes the "special pair" pigment; M, the "monomeric"

bacteriochlorophyll electron donor; and A, the primary electron acceptor. F_X is the Fe_4S_4 cluster characteristic of

type I RCs. Antenna pigments bound by the antenna domain are shown in green lines. Antenna pigments bound

by the 7^{th} and 8^{th} helices are shown as purple sticks, except for the bacteriochlorophyll molecule denoted as Z, in

olive green and bound by the 8^{th} helix. The antenna domain connects the electron donor side of the RC via N263

and a Ca-binding site. **b** A monomeric unit of PSI (PsaB, Cyanobacteria) following a similar structural

organization and nomenclature as in panel **a**. Unlike the HbRC, PSI binds quinones (Q) as intermediary electron

acceptors between A and F_X. **c** A monomeric unit of PSII (CP43/D1 and CP47/D2 in the inset). It has a

structural organization similar to that of type I RCs. However, the monomeric unit is split into two proteins after

the 6[th] transmembrane helix. The Mn_4CaO_5 cluster is coordinated by D1, but is directly connected to the antenna

domain via E354 and R357 in manner that resembles the HbRC. **d** A monomeric unit of an anoxygenic type II

reaction (PbRC, Proteobacteria). Unlike type I RCs, type II RCs lack F_X. Instead, a non-heme Fe^{2+} is found

linking the RC proteins. The PbRC lacks antenna domain and the antenna pigment Z bound at the equivalent 8[th]

helix.

Sequence reconstruction of the ancestral subunit to D1 and D2 is consistent with the

existence of a highly oxidizing homodimeric photosystem, though transient and short-lived

[18], that could split water in either side of the RC [48]. The structural comparisons between

CP43/D1 and CP47/D2 suggest a mechanism for heterodimerization and loss of catalysis on

one side that accounts for all ligands. These include direct amino acid substitutions, the

domain swap within the extrinsic region of the ancestral protein to CP47, and the gene

overlap between *psbC* (CP43) and *psbD* (D2). It would be difficult to reconcile these unique

structural and genetic features with a scenario in which PSII evolved water oxidation at a late

stage, or starting as a purple anoxygenic type II RC, once the heterodimerization process was

well underway or completed, or in the absence of core antenna domains, given that these

interact directly with the electron donor side of PSII, in a manner strikingly similar to that of

the homodimeric type I RC of the Firmicutes [19]. What is more, these structural and

functional features also suggest that the atypical forms of D1, lacking ligands to the Mn_4CaO_5

cluster, such as the so-called chlorophyll *f* synthase [32, 71], or the so-called "rogue" or

"sentinel" D1 [31, 72], diverged from forms of D1 that were able to support water oxidation.

Even if they originated from early duplications that could have antedated the MRCA of

Cyanobacteria [2] and as additionally supported by ASR analysis (Supplementary Figure S9

and S10).

We have calculated that (oxygenic) PSII has experienced the slowest rates of

evolution between type II RCs, with the core of the anoxygenic type II RC of Proteobacteria

and Chloroflexi evolving approximately five times faster than the core of PSII [18]. That

considerably faster rate has led to conspicuous structural changes of the anoxygenic RC

relative to PSII and type I photosystems. The consequence of these differences in the rates are

visually apparent (Figure 8 and Supplementary Figure S3) as the anoxygenic RC has

experienced greater sequence and structural change than PSII. That is, PSII retains a greater

number of ancestral traits that can be traced to before the ancestral core duplications. For

example, like type I, PSII retains: 1) substantially greater structural symmetry at the core, 2)

the antenna functionally linked to the core and the electron donor side, 3) lack of histidine

ligands to the monomeric photochemical chlorophylls (marked M in Figure 8), or 4) the use

of a chlorophyll *a* derived pigment as the primary electron acceptor. This list is not meant to

be exhaustive, but see refs. [19, 73] for additional detail. It follows then, that because these

conserved traits can be traced to the common ancestor of type I and type II, then the rates of

evolution of PSII should have remained slow and constrained relative to that of other

photosystems, since before the core duplications. These structural constraints demonstrate

that the core subunits of PSII did not experience unprecedentedly faster rates of evolution. In

fact, and not surprisingly, the anoxygenic type II RC which experienced the highest rates of

evolution, has accumulated greater change and greater loss of ancestral traits.

699   4. Conclusion

Both phylogenetic and structural evidence converge towards a scenario in which

photosynthetic water-splitting started at an early stage during the evolution of life and long

before the rise of what we understand today as Cyanobacteria. Nonetheless, greater resolution

of divergence time estimation in deep-time evolutionary studies could be achieved if

molecular clocks are complemented with experimentally validated rates of protein and

genome evolution across taxa of interest.

We think it is plausible that there was never a discrete origin of photosynthesis, but

that the process may trace back to abiogenic photochemical reactions, some of which may

have resulted in the oxidation of water, and at the interface of nascent membranes, membrane

proteins, photoactive tetrapyrroles and other inorganic cofactors: much in the same way that

ribosomes may have originated at the interface of nascent genetics and protein synthesis [74].

A photosynthetic origin of life is not a new idea [75-77] and abiotic photosynthesis-like

chemistry has been recently proposed to have occurred at Gale Crater on Mars [78], and to

occur even on Earth [79].

715   5. Materials and methods

5.1. Sequence alignments and phylogenetic analysis

The first dataset was retrieved on the 31[st] of October 2017 to initiate this project. It included a

total of 1389 type I RC, CP43 and CP47 protein sequences from the NCBI refseq database

using BLAST. From Cyanobacteria, 675 PsaA and PsaB and 685 CP43 and CP47 subunits

were retrieved. From anoxygenic phototrophs, 24 PscA and 4 PshA were obtained. This

dataset did not contain CBP proteins.

A second dataset was retrieved on the 10th of October 2018 from the same database.

This dataset focused on CP43 and CP47 subunits and consisted of a total of 1232 sequences.

It included 392 CP43, 465 CBP proteins, and 375 CP47 subunits. 40 CP43 and 40 CP47

sequences from a diverse set of photosynthetic eukaryotes were selected manually and

included. In addition, a selection of cyanobacterial and plastid CP43, CP47, AtpA (Alpha),

AtpB (Beta), and FtsH were manually selected for an analysis of the rates of evolution as

illustrated in Supplementary Figure S14 and described below.

A dataset of Alpha and Beta subunits belonging to cyanobacterial F-type ATP

synthase were retrieved from the NCBI refseq on the 31st of August 2019. 507 and 529

cyanobacterial Alpha and Beta sequences were obtained respectively. We retrieved Alpha

and Beta homologous for Margulisbacteria (19 and 18 sequences respectively),

Sericytochromatia (4 and 2), and Vampirovibrionia (66 and 49) using AnnoTree [80] and

searching with the KEGG codes K02111 (F-type Alpha) and K02112 (F-type Beta). A total

of 111 AtpA (subunit A) and 176 AtpB (subunit B) from archaeal V-type ATP synthase were

retrieved using the sequences from *Methanocaldococcus jannaschii* as BLAST queries.

A dataset of 366 bacterial RNA polymerase subunit β (RpoB) were collected from the

NCBI refseq database on the same date as above. We focused on Cyanobacteria (65

sequences) and other phyla known to contain phototrophic representatives, as well as for their

potential for allocating calibration points as described below. These included sequences from

Firmicutes (32), Chloroflexi (32), Proteobacteria (102), Acidobacteria (34), Chlorobi (26),

Gemmatimonadetes (3), Aquifecae (12) and Thermotogae (24). Sequences for *Candidatus*

Eremiobacterota (2) recently reported to include phototrophic representatives [81],

Margulisbacteria (13), Sericytochromatia (2), and Vampirovibrionia (11) were obtained from

AnnoTree using KEGG code K03043 as query. In addition, three extra margulisbacterial

RpoB sequences were collected: from *Termititenax persephonae* and *Termititenax aidoneus*

retrieved from the refseq database, and from margulisbacterium *Candidatus* Ruthmannia

eludens obtained from https://www.ebi.ac.uk/ena/data/view/PRJEB30343 [82]. A second

dataset containing only sequences from Cyanobacteria and MSV, in addition to sequences

from Archaea was also used. To retrieve the archaeal sequences the subunit B' (RpoB1) of

Methanocaldococcus jannaschii was used as a BLAST query. A total 213 sequences across

the entire diversity of Archaea were obtained. Two version of these were observed, those

with split B (RpoB1 and RpoB2) and those with full-length B subunits: from the latter, 112

full-length B subunits were selected for molecular clock analysis.

All sequences were aligned with Clustal Omega using 5 combined guided trees and

Hidden Markov Model Iterations [83]. Given that RpoB sequences are known to contain

many clade specific indels [27], we further processed this particular dataset using Gblocks

[84] to remove these indels and poorly aligned positions allowing smaller final blocks, gap

positions within the final blocks, and less strict flanking positions. This procedure left a total

of 903 well-aligned sites for the dataset with only bacterial sequences and 788 for the dataset

with archaeal sequences.

Maximum Likelihood phylogenetic analysis was performed with the PhyML online

service using the Smart Model Selection mode implementing the Bayesian Information

Criterion for parameter selection [85, 86]. Tree searching operations were calculated with the

Nearest Neighbour Interchange model. Support was computed using the average likelihood

ratio test [87]. Trees were visualised with Dendroscope V3.5.9 [88].

5.2. Distance estimation

Distance estimation was performed as a straightforward and intuitive approach to detect

variations in the rate of evolution across taxa with known divergence time of at similar

taxonomic ranks. This was done in several ways. Distance trees were plotted using BioNJ

[89] as implemented in Seaview V4.7 [90] using observed distances and 100 bootstrap

replicates. Within-group mean distances were calculated using three different substitution

models as implemented in the package MEGA-X [91]: no. of differences, a Poisson model,

and a JTT model. A gamma distribution to model rates among sites was used with a

parameter of 1.00 and 500 bootstrap replicates.

To compare changes in the rates of evolution as a function of time of key proteins

used in this study, we compared the percentage of sequence identity between orthologues in a

total of 20 pairs of sequences from representative photosynthetic eukaryotes and

Cyanobacteria with known or approximate divergence times. These are listed in

Supplementary Table S3. We compared CP43, CP47, Alpha, Beta and FtsH. Of these 17

pairs, the first 15 were comparisons between land plants. The approximated divergence times

were based on those recommended by Clarke, et al. [92] after their extensive review of the

plant fossil record. These were taken as the average of the hard minimum and soft maximum

fossil ages suggested by the authors. The soft maximum age for the earliest land plants was

taken as 515 Ma as discussed extensively in Morris, et al. [93]

Another sequence comparison was made between the unicellular red alga

Cyanidioschyzon merolae and *Arabidopsis thaliana*. The earliest well-accepted fossil

evidence for red algae is that of the multicellular *Bangiomorpha* [94, 95] dated to about 1.0

Ga [96]. Recently fossils of multicellular eukaryotic algae have been reported at 1.6 Ga [97-

99]. The distance between heterocystous Cyanobacteria and their closest non-heterocystous

relatives is represented by a pairwise comparison between *Chroococcidiopsis thermalis* and

Nostoc sp. PCC 7120. Excellently preserved heterocystous Cyanobacteria have been reported

from Tonian deposits older than 0.72 Ga [100]. The largest distance is that between

Gloeobacter spp. and any other cyanobacterium or photosynthetic eukaryote, representing the

distance since the MRCA of Cyanobacteria of uncertain divergence time.

To compare the level of sequence identity between RC proteins, two datasets of 10

random amino acid sequences were generated using the Sequence Manipulation Suit [101].

The datasets contained sequences of 350 and 750 residues. These were independently aligned

as described above, resulting in 45 pairwise sequence identity comparisons for each dataset.

These random sequence datasets were used as a rough minimum threshold of identity.

Alignments of RC proteins were generated using three representative sequences spanning

known diversity. Cyanobacterial CP43, CP47, standard D1 and D2 sequences were from

Gloeobacter violaceus, *Stanieria cyanosphaera*, and *Nostoc* sp. PCC 7120; Heliobacterial

PshA from *Heliobacterium modesticaldum*, *Heliobacillus mobilis*, and *Heliorestis convoluta*;

PscA from *Chlorobium tepidum*, *Prosthecochloris aestuarii*, *Chlorobium* sp. GBchlB,

Chloracidobacterium thermophilum and *Chloracidobacterium* sp. CP2-5A; Protoebacterial L

and M from *Methylobacterium* sp. 88A, *Roseivivax halotolerans*, and *Blastochloris*

sulfoviridis; and L and M from *Chloroflexus* sp. Y-400-fl, *Roseiflexus castenholzii*, and

Oscillochloris trichoides.

5.3. Quantification of rates of evolution (rationale)

Molecular clocks are conventionally used to estimate divergence times. In general terms,

given: 1) a tree topology, which sets the relationship between taxa; 2) a sequence alignment,

which sets the phylogenetic distance between taxa; and 3) some known events (calibrations),

which set the rates of evolution, the molecular clock can then estimate divergence times. This

means that if the tree topology and divergence times for two sets of protein sequences are the

same, any differences in phylogenetic distances between these two should only reflect

differences in the rate of evolution. Thus, assuming that CP43/CP47 and Alpha/Beta have

mainly been inherited vertically in Cyanobacteria and photosynthetic eukaryotes, any

difference in phylogenetic distance between the two is the result of differences in the rates of

evolution. For example, the level of sequence identity between CP43 in *Cyanidioschizon* and

Arabidopsis is 78%, and the level of sequence identity between Alpha in the same species is

69%. Given that these plastid-encoded subunits have mostly been inherited vertically since

the MRCA of Archaeplastida, then one can argue that Alpha is evolving somewhat faster

than CP43. This is because faster rates of protein evolution should lead to faster rates of

change resulting in a faster decrease in the level of sequence identity (increase in

phylogenetic distance). Now, because CP43 and CP47 are paralogues, we can then use this

approach to estimate the rates of evolution at the moment of duplication under a given

number of specific scenarios. For example, assuming that the MRCA of Cyanobacteria

occurred at 2.0 Ga, we can then model how the rates of evolution would change at the

moment of duplication if this occurred at 2.5, 3.0, 3.5 Ga, or at any other time point of

interest. Because these set of proteins have likely achieved mutational saturation at the largest

distance, the rate of evolution at the point of duplication will be an underestimation (slower

than it should be), given that saturation would hide substitution events. Similarly, keeping the

time of duplication constant, we can then model how the rates of evolution would change

across the tree if the MRCA of Cyanobacteria is assumed to have occurred at 2.0, 2.5, 3.0 Ga,

or at any other time of interest. Therefore, even though it is difficult to determine the absolute

time of origin of Cyanobacteria, or of the early duplications of the core subunits of ATP

synthase, it is possible to determine what rates of evolution are required to fulfil any

particular scenario.

5.4. Quantification of rates of evolution

To measure rates of evolution of CP43/CP47 and Alpha/Beta a total of 19 sequences from

photosynthetic eukaryotes and 4 cyanobacterial sequences per subunit were selected. A

standardized tree topology was constructed from consensus evolutionary relationships as

illustrated in Supplementary Figure S14a: 1) Relationships between land plants was taken

from ref. [92] 2) It is well established that the divergence of red algae predates the MRCA of

land plants. 3) Ponce-Toledo, Deschamps [102] recently suggested that *Gloeomargarita* is

the closest living cyanobacterial relative to the plastid ancestor and predated the emergence

of heterocystous cyanobacteria, but see also [103]. 4) The clade containing

Chroococcidiopsis thermalis PCC 7203 is one of heterocystous Cyanobacteria's closest non-

heterocystous relatives [104, 105]. 5) *Gloeobacter* spp. is the earliest branching and well-

described genus of Cyanobacteria capable of oxygenic photosynthesis [105-109].

Calibration points were allocated as shown in Supplementary Figure S14a and listed

in Table 2. Nodes 1 to 15 were applied following the justifications and best practice listed in

ref. [92] and using the 515 Ma older calibration as suggested in ref. [93] Node 16 represents

here the MRCA of red and green lineages of photosynthetic eukaryotes. The minimum age

was set to 1.0 Ga based on the *Bangiomorpha* fossil as described above [94, 95]. The

maximum age was set to 1.8 Ga, which is similar the oldest ages reported in recent molecular

clock analyses for the MRCA of photosynthetic eukaryotes and it is also similar to the age of

the earliest plausible fossilised unicellular eukaryotes [110, 111]. That said, compelling fossil

evidence for eukaryotes could go as far back as 2.1 Ga [112]. Node 17 marks the divergence

of heterocystous Cyanobacteria and it was given a minimum age of 0.72 Ga based on the

recently described fossils of filaments bearing clear heterocysts from the Tonian period [100].

A maximum age of 1.65 Ga was given to this node, based on the recent report of

heterocystous cyanobacteria from the Gaoyuzhuang Formation [113]. Although, it is unclear

if reported akinetes that are older than 1.65 Ga are truly that [10], we considered that the

0.72-1.65 Ma range for the origin of heterocystous Cyanobacteria was a reasonably broad

constraint for the purpose of this experiment, but see below.

Node 18 denotes the MRCA of Cyanobacteria. The age for this node is highly debated

ranging from before to after the GOE. Node 19 represents the duplication events leading to

CP43 and CP47, and to Alpha and Beta. To calculate the rates of evolution under different

scenarios, node 18 and node 19 were varied. Firstly, a molecular clock was run using a

scenario that assumed that the MRCA of Cyanobacteria postdated the GOE. To do this, node

18 was set to be between 1.6 and 1.8 Ga, which emulates results reported in recent studies [8,

11]. This was compared to a scenario that assumed that the MRCA of Cyanobacteria

antedated the GOE, and thus node 18 was set to be between 2.6 and 2.8 Ga, which simulates

other evolutionary scenarios [103, 114]. In both cases, the duplication event (node 19) was

set to be 3.5 Ga old, or changed as stated in the main text, by assigning a gamma prior at the

desired time fixed with a narrow standard deviation of 0.05 Ga. In a separate experiment, the

age of the duplication was varied while maintaining node 18 restricted to between 1.6 and 1.8

Ga, while node 19, the root, was set with a gamma prior with an average varied from 0.8 to

4.2 and with a narrow standard deviation of 0.05 Ga.

The period of time between the duplication event (node 19), which led to the

divergence of CP43 and CP47, and the MRCA of Cyanobacteria (node 18), we define as ΔT.

ΔT is calculated as the subtraction of the mean age of node 19 and node 18. For PSII, we

used node 18 from the CP43 subunit and for ATP synthase we used node 18 from the Alpha

subunit. In consequence, varying the age of the duplication from 0.8 to 4.2 Ga allows changes

in the rate of evolution to be simulated with varying ΔT, ranging from 0.2 to 2.5 Ga. That is

to say, it allows the rate of evolution at the point of duplication to be calculated if it occurred

at any point in time before the MRCA of Cyanobacteria.

Table 2. Calibration points used in this study

Node label	Relationship	Max age (Ma)	Min age (Ma)
1	*Arabidopsis* v *Populus*	127	82
2	*Arabidopsis* v *Sorghum*	248	124
3	*Chloranthus* v *Liriodendron*	248	98
4	*Arabidopsis* v *Illicium*	248	124
5	*Arabidopsis* v *Nymphaea*	248	124
6	*Arabidopsis* v *Amborella*	248	124
7	*Welwitschia* v *Pinus*	309	121
8	*Welwitschia* v *Cryptomeria*	309	147
9	*Welwitschia* v *Ginkgo*	366	107
10	*Welwitschia* v *Cycas*	366	306
11	*Arabidopsis* v *Cycas*	366	306
12	*Arabidopsis* v *Psilotun*	454	388
13	*Arabidopsis* v *Anthoceros*	515	420
14	*Arabidopsis* v *Physcomitrella*	515	420
15	*Arabidopsis* v *Marchantia*	515	449
16	*Arabidopsis* v *Cyanidioschyzon*	—	1000
17	*Nostoc* v closest non-heterocystous strain	1560 (or none)	720
18	*Arabidopsis* v *Gloeobacter*	Variable (or none)	Variable (or none)
19	CP43 v CP47 or AtpA v AtpB duplications	Variable	Variable
20	*Richelia intracelullaris* v closest neighbouring sequence	250	100
21	Chroococcales cyanobacteria	—	2016
22	*Termititenax*	396	140
23	*Candidatus* Ruthmannia eludens v GCA 002716725.1	660	542
24a	GB GCA 001899365.1 Koala v GB GCA 001917115.1 Homo	201 (or 55)	—
24b	GB GCA 001917115.1 *Homo* v GB GCA 000980455.1 Homo	201	—
25	GB GCA 001917115.1 *Homo* v *Vampirovibrio chlorellavorus*	1800	—
26	*Polynucleobacter necessarius*	444	—
27	*Bradyrhizobium*	86	—
28	Rickettsias	—	1800
29	*Wolbachia* and *Anaplasma*	395	—
30	Phototrophic Chlorobi	—	1640
31	Chromatiaceae	—	1640

Molecular clocks were calculated with Phylobayes 3.3f using a log normal autocorrelated

molecular clock model under the CAT+Γ non-parametric model of amino acid substitution

and a uniform distribution of equilibrium frequencies. We preferred a CAT model instead of

a CAT+GTR as the latter only outperforms the former on sequence alignments of over 1000

sites and it is also much less computationally expensive. Four discrete categories for the

gamma distribution were used and four chains were executed in parallel through all

experiments carried out in this study. The instant rates of evolution, which are the rates at

each internal node in the tree, were retrieved from the output files of Phylobayes. These rates

are calculated by the software as described by the developers elsewhere [115, 116] and are

expressed as amino acid changes per site per unit of time.

As an additional comparison, we calculated rates of evolution for cyanobacterial FtsH

subunits, which AAA ATPase domain is structurally similar to the catalytic head of ATP

synthase [117]. Cyanobacterial FtsH are involved in membrane protein quality control, with

some isoforms targeting specifically PSII subunits. It features a late duplication event leading

to cyanobacterial FtsH1 and FtsH2 subunits (using the nomenclature of Shao, et al. [40]). It

was shown previously that the duplication leading to FtsH1 and FtsH2 occurred after the

divergence of *Gloeobacter* spp. Genes encoding plastid FtsH subunits are encoded in the

nuclear genome of photosynthetic eukaryotes. Because not all the selected species had fully

sequenced nuclear genomes, only those with available FtsH sequences at the time of this

study were used. The tree topology shown in Supplementary Figure S14b was used as

template for the calculation of the rate of evolution and was based on the topology presented

by Shao, et al. [40] It was concluded that the Cyanobacteria-inherited closest paralog to

FtsH1 and FtsH2 in photosynthetic eukaryotes was also acquired before their initial

duplication. Therefore, from all FtsH paralogs in photosynthetic eukaryote genomes, those

with greater sequence identity to cyanobacterial FtsH1/2 were used. Because this duplication

is specific to Cyanobacteria, a few additional strains were included in this tree following

well-established topologies [103, 105]. Calibrations were placed as indicated in

Supplementary Figure S14b. To test the change in the rate of evolution at the time of

duplication in comparison with CP43/CP47, node 19 was set to 1.6-1.8 Ga or 2.6-2.8 Ga and

molecular clocks were run as described in the preceding paragraph.

Finally, we conducted a large molecular clock using the combined 897 CP43 and CBP

sequences, including 40 eukaryotic CP43 sequences, to test whether using a more complex

phylogeny would result in rates of evolution substantially different to those calculated with

the method described above. Calibrations were assigned as illustrated in Supplementary

Figure S15. Cross-calibrations were used across paralogs constraining the origin of

heterocystous Cyanobacteria. In this case, only the minimum constraint of 0.72 Ga was used

with no maximum constraint to allow greater flexibility. Additional calibrations were

assigned also across paralogs (point 20 in Supplementary Figure S15), this was considered as

the node made by *Richelia intracellularis* and its closest sister sequence, as implemented in

ref. [114] This strain is a specific endosymbiont of a diatom and its divergence was set to be

no older than the earliest discussed age for diatoms [118]. The root equivalent to the MRCA

of Cyanobacteria (divergence of *Gloeobacter* in CP43) was not calibrated. The root of the

tree was varied: first it was given a maximum age of 4.52 Ga as recently implemented and

justified by Betts, et al. [11] as the earliest plausible time in which the planet was inhabitable

after the moon forming impact [119], and no minimum age was used. A second tree was

executed with no constraint on the root and no root prior. A third root was implemented

constrained to be between 2.3 Ga (the GOE) and 3.2 Ga. The latter date represents the age of

the cyanobacteria-like well-preserved microbial mats of the Berberton Greenstone Belt in

South Africa and neighbouring Eswatini [66]. Rates were obtained using the autocorrelated

CAT+Γ model as described above. Because these root constraints did not have a strong effect

in the overall estimated rates, we carried out an additional control applying an uncorrelated

gamma clock model [120] with a root constrained at 4.52 Ga and no minimum age.

5.5. Molecular clock of RpoB and concatenated ribosomal proteins

The primary objective of this experiment was not to determine the absolute time of origin of

Cyanobacteria, but to understand the spans of time between Cyanobacteria and their relatives.

We also wanted to understand what rates of evolution are associated with those spans of time

and how these change under different evolutionary scenarios. To do this, we applied a

molecular clock to the phylogeny of RpoB sequences described above. We implemented 12

calibrations. The calibrations were assigned on the phylogeny as shown in Supplementary

Figure S16 and listed in Table 2. A set of calibrations consisted of the earliest unambiguous

evidence for Chroococcales Cyanobacteria of the Belcher group (point 21), the age of which

has been recently revisited to 2.01 Ga [121]. This was assigned to the younger node from

where Chroococcales strains branch out in the tree, with no maximum restrictions. The

appearance of heterocystous Cyanobacteria were restricted from 0.72 Ga and 1.56 Ga as

described above. No constraints on the node representing the MRCA of Cyanobacteria were

used. However, for rigor, we also tested an alternative single calibration on the node

representing the MRCA of Cyanobacteria with a maximum of 2.01 Ga and no minimum, and

with no other calibrations in the clade. This considered a scenario in which crown group

Cyanobacteria are younger than the Belcher fossils.

In addition to cyanobacterial calibrations, we also applied the often-used biomarker

evidence for phototrophic Chlorobi and Chromatiaceae at 1.64 Ga [122], see for example

refs. [9, 63] These were used as a minimum with no maximum constraints (node 30 and 31

respectively in Supplementary Figure S16).

A set of calibrations were chosen from well-described symbiotic relationships. Two of

these are known in the phylum Margulisbacteria. The first one is *Termititenax*, these are

specific ectosymbionts of spirochetes that live within oxymonad protists in the gut of diverse

termites and cockroaches [43]. The two *Termititenax* sequences used in this analysis

clustered together and therefore we calibrated this node to be between 396 Ma, some of the

oldest fossil evidence of insects [123] and 140 Ma for *Mastotermes nepropadyom*, a Jurassic

fossil termite [124] (node 22). This was done under the assumption that this symbiotic

relationship may have started before the divergence of termites and cockroaches, as

suggested in ref. [43] The second one is *Candidatus* Ruthmannia eludens, a cell-type specific

endosymbiont of placozoans, early-branching metazoans. This symbiont has been detected in

all haplotypes examined, regardless of geographical location or sampling time [82, 125],

which suggest that this association may be as old as placozoans. Therefore, we used a

minimum calibration of 542 Ma as the Cambrian explosion of animal diversity and 660 Ma,

the earliest biomarker evidence for desmosponges [126] (node 23), which should antedate the

divergence of placozoans. This calibration was assigned to the node separating *Candidatus*

Ruthmannia eludens from its closest sister sequence.

Similar to Margulisbacteria, members of the clade Vampirovibrionia have been

reported to form close associations with eukaryotes. The clade Gastranaerophilales is thought

to be composed mostly of strains that inhabit the animal gut. Thus, we calibrated the node

separating two strains isolated from human and koala faeces that clustered together with a

maximum age of 201 Ma, representing the Jurassic split of marsupials and placental

mammals [127] (node 24a). However, the koala and human sequences were embedded in the

Gastranaerophilales clade within other sequences from the human gut. Because of this, we

trialled changing this calibration to 55 Ma instead, the oldest primate fossil [128] and

assuming that the retrieved sequences from the human gut had a common ancestor younger

than the MRCA of primates. Alternatively, we tested moving this calibration to the ancestral

nodes of the clade that included all the human gut sequences (node 24b). Gastranaerophilales

is closely related to the order Vampirovibrionales, which include *Vampirovibrio*

chlorellavorus. This strain is a predator of the eukaryotic green algae *Chlorella* [129], and

therefore we trialled a calibration assuming that Gastranaerophilales and Vampirovibrionales

radiated after the MRCA of eukaryotes (node 25). We thus assigned a maximum calibration

to this node of 1.8 Ga representing the earliest described plausible eukaryote fossils [110] and

no minimum age.

Another highly specific obligate symbiosis is that of the betaproteobacterium

Polynucleobacter necessarius and ciliates of the genus *Euplotes* (Spirotrichea) [130].

Polynucleobacter has close free-living phototrophic relatives within the same genus [130].

We set the node separating the phototrophic and non-phototrophic *Polynucleobacter* (node

26) a maximum age of 444 Ma for the oldest fossil evidence of spirotrichs, as implemented in

Parfrey et al. [131], and which predates the radiation of the genus *Euplotes* [132].

Another well-known association is that of the soil bacteria *Bradyrhizobium* and

legumes. Thus we gave the node separating *Bradyrhizobium* spp. from its closest relative in

the RpoB tree, *Xanthobacter autotrophicus*, a maximum age of 86 Ma for Rosids, which

contain legumes [93] (node 27).

The Rickettsiales are Alphaproteobacteria that exists in very close association with

eukaryotes [133]. An association that may reach to the lineage leading to the origin of

mitochondria [134]. Therefore, we assumed that the divergence of Rickettsiales occurred

before the MRCA of eukaryotes and gave this node a minimum age of 1.8 Ga [110] (node

28). Finally, the family Anaplasmataceae contains bacteria that exists in close association

with insects as endosymbionts (e.g. *Wolbachia*) or as parasite vectors (e.g. *Anaplasma*).

Therefore, we set a maximum constraint for the MRCA of *Wolbachia* and *Anaplasma* (node

29)*, excluding *Neorickettsia*, to be as old as the earliest evidence for insects about 395 Ma

ago [123].

To constrain the age of the root, we first set a broad gamma prior with an average of

3.8 Ga and a standard deviation of 0.5 Ga. We found this to perform well and used it as

benchmark to compare with a range of evolutionary models and the effects of key

calibrations (Supplementary Figure S8). Alternatively, we applied a broad calibration on the

root with a maximum of 4.52 Ga as described above and a minimum of 3.41 Ga, which is the

earliest well-accepted evidence for photosynthesis [47]. This evidence was hypothesized to

be anoxygenic in ref. [47] Therefore, Bacteria should be at least as old as the earliest

evidence for photosynthesis under conventional evolutionary scenarios.

To further understand the effect that a distant outgroup would have on the estimated

divergence time, and to measure the rate of evolution of RpoB at the divergence of Bacteria

and Archaea, we repeated the clock including 112 diverse archaeal sequences, in addition to

Thermotogae, MSV, and Cyanobacteria, but removing all other clades as a compromise

between robustness and computing time. Calibrations were assigned on MSV and

Cyanobacteria as described above. Relaxed molecular clocks were computed using an

autocorrelated log normal clock and applying a CAT+Γ model. This was compared against a

CAT+Γ and a CAT+GTR+Γ using birth-death priors and soft bounds on the calibrations

allowing for 2.5% tail probability falling outside the minimum and maximum boundary, or

5% in the case of a single boundary. In addition, we also compared to an uncorrelated gamma

model (Supplementary Figure S8).

We compared the RpoB molecular clock (CAT+Γ) with that of a clock executed using

a dataset of concatenated ribosomal proteins of 2596 aligned sites, obtained from an

independent study [38]. In total, 157 sequences were used including 56 cyanobacterial

sequences, 14 sequences from Vampirovibrionia, 9 from Margulisbacteria, 9 from

Thermotogae and 78 from Archaea. A set of calibrations were used including those two in

Cyanobacteria (point 17 and 21), a calibration on the MRCA of Gastranaerophilales with a

maximum 201 Ma (point 24b), and a root calibration between 4.52 and 3.41 Ga as described

above.

1048

5.6 Ancestral sequence reconstruction

Ancestral sequence reconstruction (ASR) of D1 and D2 amino acid sequences was carried

out with a dataset collected on the 17[th] of November 2017. Duplicates and partial sequences

were removed, leaving 755 D1 and 248 D2 sequences. CD-HIT [135] was used to remove

sequences with greater than 92% sequence identity to create a representative sample. The L

and M RC subunits from 5 strains of Proteobacteria were used as outgroup. The final

alignment did not include the atypical variant D1 sequence from *Gloeobacter kilaueensis*

(NCBI accession AGY58976.1) as this showed an unstable phylogenetic position in this

dataset. Maximum likelihood trees used as input for ASR were computed with PhyML using

Smart Model Selection [86]. The LG substitution model with observed amino acid

frequencies and four gamma rate categories exhibited the best log likelihood (LG+Γ+F)

(Supplementary Table S7). We used the top four models for tree reconstruction (LG+Γ+F,

LG+Γ+I+F, LG+Γ and LG+ Γ+I) in addition to another tree computed using the WAG

substitution model with observed amino acid frequencies and four gamma rate categories

(WAG+Γ+F). These trees were used as input trees to calculate maximum likelihood ancestral

states at each site for the node corresponding to the homodimeric reaction protein, D0. Three

ASR programs were used for the reconstructions: FastML [136], Lazarus [137] (a set of
Python scripts which wraps PAML [138]) and MEGA-7 [139]. The substitution model used
by all three programs corresponded to the substitution model used for the specific input tree
in PhyML and the branch lengths of the trees were fixed. In FastML, a maximum likelihood
method of indel reconstruction using a probability cut-off of 0.7 was used. In MEGA-7, all
sites were used for analysis with no branch swap filter. In Lazarus, the '—gapcorrect' option
was used to parsimoniously place indels.
1072

5.7. Structural analysis
The following crystal structures were used in this work: the crystal structure of PSII from
Thermosynechococcus vulcanus, PDB ID: 3wu2 [13]; the anoxygenic type II RC of
Thermochromatium tepidum, PDB ID: 5y5s [140]; the homodimeric type I RC from
Heliobacterium modesticaldum, 8v5k [141]; PSI from *Thermosynechococcus elongatus*, PDB
ID: 1jb0 [142]; and the cryo-EM IsiA structure from *Synechocystis* sp. PCC 6803 [143].
Structures were visualised using Pymol^{TM} V. 1.8.2.2 (Schrodinger, LLC) and structural
overlaps were carried out with the CEAlign plugin [144].
1081

6. Acknowledgements
This work was done with the support of a Leverhulme Trust grant (RPG-2017-223 to T.C.
and A.W.R.), a UKRI Future Leaders Fellowship (MR/T017546/1 to T.C.) a Royal Society
University Research Fellowship (P.S.-B.), and the Biotechnology and Biological Sciences
Research Council (BB/K002627/1 and BB/L011206/1 to A.W.R.). The authors are grateful to
Dr. Travis J. Lawrence and Prof. Christoph Heubeck for feedback on a version of this
manuscript. Computing resources were provided by the HPC facility at Imperial College
London. The authors declare no conflict of interests.
1090

7. References
1.   Sánchez-Baracaldo, P. and T. Cardona, *On the origin of oxygenic photosynthesis and*
Cyanobacteria. New Phytol, 2020. **225**: 1440-1446. DOI: 10.1111/nph.16249.
2.   Cardona, T., J.W. Murray, and A.W. Rutherford, *Origin and evolution of water*
oxidation before the last common ancestor of the Cyanobacteria. Mol. Biol. Evol.,
2015. **32**: 1310-1328. DOI: 10.1093/molbev/msv024.
3.   Di Rienzi, S.C., I. Sharon, K.C. Wrighton, O. Koren, L.A. Hug, B.C. Thomas, J.K.
Goodrich, J.T. Bell, T.D. Spector, J.F. Banfield, and R.E. Ley, *The human gut and*
groundwater harbor non-photosynthetic bacteria belonging to a new candidate
phylum sibling to Cyanobacteria. Elife, 2013. **2**: e01102. DOI: 10.7554/eLife.01102.

1101	4.	Soo, R.M., J. Hemp, and P. Hugenholtz, *Evolution of photosynthesis and aerobic*
1102		*respiration in the cyanobacteria.* Free Radical Biol. Med., 2019. **140**: 200-205. DOI:
1103		10.1016/j.freeradbiomed.2019.03.029.

4.   Soo, R.M., J. Hemp, and P. Hugenholtz, *Evolution of photosynthesis and aerobic*
respiration in the cyanobacteria. Free Radical Biol. Med., 2019. **140**: 200-205. DOI:
10.1016/j.freeradbiomed.2019.03.029.

5.   Soo, R.M., J. Hemp, D.H. Parks, W.W. Fischer, and P. Hugenholtz, *On the origins of*
oxygenic photosynthesis and aerobic respiration in Cyanobacteria. Science, 2017.
355: 1436-1440. DOI: 10.1126/science.aal3794.

6.   Anantharaman, K., C.T. Brown, L.A. Hug, I. Sharon, C.J. Castelle, A.J. Probst, B.C.
Thomas, A. Singh, M.J. Wilkins, U. Karaoz, E.L. Brodie, K.H. Williams, S.S.
Hubbard, and J.F. Banfield, *Thousands of microbial genomes shed light on*
interconnected biogeochemical processes in an aquifer system. Nat. Commun., 2016.
7: 13219. DOI: 10.1038/ncomms13219.

7.   Carnevali, P.B.M., F. Schulz, C.J. Castelle, R.S. Kantor, P.M. Shih, I. Sharon, J.M.
Santini, M.R. Olm, Y. Amano, B.C. Thomas, K. Anantharaman, D. Burstein, E.D.
Becraft, R. Stepanauskas, T. Woyke, and J.F. Banfield, *Hydrogen-based metabolism*
as an ancestral trait in lineages sibling to the Cyanobacteria. Nat. Commun., 2019.
10: 463. DOI: 10.1038/s41467-018-08246-y.

8.   Shih, P.M., J. Hemp, L.M. Ward, N.J. Matzke, and W.W. Fischer, *Crown group*
Oxyphotobacteria postdate the rise of oxygen. Geobiology, 2017. **15**: 19-29. DOI:
10.1111/gbi.12200.

9.   Magnabosco, C., K.R. Moore, J.M. Wolfe, and G.P. Fournier, *Dating phototrophic*
microbial lineages with reticulate gene histories. Geobiology, 2018. **16**: 179-189.
DOI: 10.1111/gbi.12273.

10.   Schirrmeister, B.E., P. Sánchez-Baracaldo, and D. Wacey, *Cyanobacterial evolution*
during the Precambrian. Int. J. Astrobiol., 2016. **15**: 187-204. DOI:
10.1017/S1473550415000579.

11.   Betts, H.C., M.N. Puttick, J.W. Clark, T.A. Williams, P.C.J. Donoghue, and D. Pisani,
Integrated genomic and fossil evidence illuminates life's early evolution and
eukaryote origin. Nat. Ecol. Evol., 2018. **2**: 1556-1562. DOI: 10.1038/s41559-018-
0644-x.

12.   Garcia-Pichel, F., J. Lombard, T. Soule, S. Dunaj, S.H. Wu, and M.F. Wojciechowski,
Timing the Evolutionary Advent of Cyanobacteria and the Later Great Oxidation
Event Using Gene Phylogenies of a Sunscreen. Mbio, 2019. **10**. DOI:
10.1128/mBio.00561-19.

13.   Umena, Y., K. Kawakami, J.R. Shen, and N. Kamiya, *Crystal structure of oxygen-*
evolving Photosystem II at a resolution of 1.9 Å. Nature, 2011. **473**: 55-60. DOI:
10.1038/nature09913.

14.   Ferreira, K.N., T.M. Iverson, K. Maghlaoui, J. Barber, and S. Iwata, *Architecture of*
the photosynthetic oxygen-evolving center. Science, 2004. **303**: 1831-1838. DOI:
10.1126/science.1093087.

15.   Cardona, T., A. Sedoud, N. Cox, and A.W. Rutherford, *Charge separation in*
Photosystem II: A comparative and evolutionary overview. Biochim. Biophys. Acta,
2012. **1817**: 26-43. DOI: 10.1016/j.bbabio.2011.07.012.

16.   Rutherford, A.W., T. Mattioli, and W. Nitschke, *The FeS-type photosystems and the*
evolution of photosynthetic reaction centers, in *Origin and evolution of biological*
energy conversion, H. Baltscheffsky, Editor. 1996, VCH: New York, N. Y. 177-203.

17.   Rutherford, A.W. and W. Nitschke, *Photosystem II and the quinone–iron-containing*
reaction centers, in *Origin and evolution of biological energy conversion*, H.
Baltscheffsky, Editor. 1996, VCH: New York, N. Y. 143–175.

18.   Cardona, T., P. Sánchez-Baracaldo, A.W. Rutherford, and A.W.D. Larkum, *Early*
1150   *Archean origin of Photosystem II.* Geobiology, 2019. **17**: 127-150. DOI:
1151   10.1111/gbi.12322.
1152   19.   Cardona, T. and A.W. Rutherford, *Evolution of photochemical reaction centres: more*
1153   *twists?* Trends Plant Sci., 2019. **24**: 1008-1021. DOI: 10.1016/j.tplants.2019.06.016.
1154   20.   Mulkidjanian, A.Y., K.S. Makarova, M.Y. Galperin, and E.V. Koonin, *Inventing the*
1155   *dynamo machine: the evolution of the F-type and V-type ATPases.* Nat. Rev.
1156   Microbiol., 2007. **5**: 892-9. DOI: 10.1038/nrmicro1767.
1157   21.   Gogarten, J.P., H. Kibak, P. Dittrich, L. Taiz, E.J. Bowman, B.J. Bowman, M.F.
1158   Manolson, R.J. Poole, T. Date, T. Oshima, J. Konishi, K. Denda, and M. Yoshida,
1159   *Evolution of the vacuolar H^+-ATPase: Implications for the origin of eukaryotes.* Proc.
1160   Natl. Acad. Sci. U.S.A., 1989. **86**: 6661-6665. DOI: 10.1073/pnas.86.17.6661.
1161   22.   Lane, N., J.F. Allen, and W. Martin, *How did LUCA make a living? Chemiosmosis in*
1162   *the origin of life.* Bioessays, 2010. **32**: 271-280. DOI: 10.1002/bies.200900131.
1163   23.   Ouzounis, C.A., V. Kunin, N. Darzentas, and L. Goldovsky, *A minimal estimate for*
1164   *the gene content of the last universal common ancestor: Exobiology from a terrestrial*
1165   *perspective.* Res. Microbiol., 2006. **157**: 57-68. DOI: 10.1016/j.resmic.2005.06.015.
1166   24.   Muller, V. and G. Gruber, *ATP synthases: structure, function and evolution of unique*
1167   *energy converters.* Cell. Mol. Life Sci., 2003. **60**: 474-494. DOI:
1168   10.1007/s000180300040.
1169   25.   Werner, F. and D. Grohmann, *Evolution of multisubunit RNA polymerases in the three*
1170   *domains of life.* Nat. Rev. Microbiol., 2011. **9**: 85-98. DOI: 10.1038/nrmicro2507.
1171   26.   Kyrpides, N., R. Overbeek, and C. Ouzounis, *Universal protein families and the*
1172   *functional content of the Last Universal Common Ancestor.* J. Mol. Evol., 1999. **49**:
1173   413-423. DOI: 10.1007/Pl00006564.
1174   27.   Lane, W.J. and S.A. Darst, *Molecular evolution of multisubunit RNA polymerases:*
1175   *sequence analysis.* J. Mol. Biol., 2010. **395**: 671-85. DOI: 10.1016/j.jmb.2009.10.062.
1176   28.   Harris, J.K., S.T. Kelley, G.B. Spiegelman, and N.R. Pace, *The genetic core of the*
1177   *universal ancestor.* Genome Res., 2003. **13**: 407-412. DOI: 10.1101/gr.652803.
1178   29.   Chen, M., Y. Zhang, and R.E. Blankenship, *Nomenclature for membrane-bound light-*
1179   *harvesting complexes of cyanobacteria.* Photosynth. Res., 2008. **95**: 147-154. DOI:
1180   10.1007/s11120-007-9255-0.
1181   30.   Chen, M., R.G. Hiller, C.J. Howe, and A.W.D. Larkum, *Unique origin and lateral*
1182   *transfer of prokaryotic chlorophyll-b and chlorophyll-d light-harvesting systems.* Mol.
1183   Biol. Evol., 2005. **22**: 21-28. DOI: 10.1093/molbev/msh250.
1184   31.   Murray, J.W., *Sequence variation at the oxygen-evolving centre of Photosystem II: a*
1185   *new class of 'rogue' cyanobacterial D1 proteins.* Photosynth. Res., 2012. **110**: 177-84.
1186   DOI: 10.1007/s11120-011-9714-5.
1187   32.   Ho, M.Y., G. Shen, D.P. Canniffe, C. Zhao, and D.A. Bryant, *Light-dependent*
1188   *chlorophyll f synthase is a highly divergent paralog of PsbA of Photosystem II.*
1189   Science, 2016. **353**: aaf9178. DOI: 10.1126/science.aaf9178.
1190   33.   Gan, F., G. Shen, and D.A. Bryant, *Occurrence of far-red light photoacclimation*
1191   *(FaRLiP) in diverse cyanobacteria.* Life (Basel), 2014. **5**: 4-24. DOI:
1192   10.3390/life5010004.
1193   34.   Cardona, T., *A fresh look at the evolution and diversification of photochemical*
1194   *reaction centers.* Photosynth. Res., 2015. **126**: 111-134. DOI: 10.1007/s11120-014-
1195   0065-x.
1196   35.   Cardona, T., *Reconstructing the origin of oxygenic photosynthesis: Do assembly and*
1197   *photoactivation recapitulate evolution?* Front. Plant Sci., 2016. **7**: 257. DOI:
1198   10.3389/fpls.2016.00257.

36. Dibrova, D.V., M.Y. Galperin, and A.Y. Mulkidjanian, *Characterization of the N-*
ATPase, a distinct, laterally transferred Na$^+$-translocating form of the bacterial F-
type membrane ATPase. Bioinformatics, 2010. **26**: 1473-1476. DOI:
10.1093/bioinformatics/btq234.
37. Battistuzzi, F.U., A. Feijao, and S.B. Hedges, *A genomic timescale of prokaryote*
evolution: insights into the origin of methanogenesis, phototrophy, and the
colonization of land. BMC Evol. Biol., 2004. **4**: 44. DOI: 10.1186/1471-2148-4-44.
38. Hug, L.A., B.J. Baker, K. Anantharaman, C.T. Brown, A.J. Probst, C.J. Castelle, C.N.
Butterfield, A.W. Hernsdorf, Y. Amano, K. Ise, Y. Suzuki, N. Dudek, D.A. Relman,
K.M. Finstad, R. Amundson, B.C. Thomas, and J.F. Banfield, *A new view of the tree*
of life. Nat. Microbiol., 2016. **1**: 16048. DOI: 10.1038/nmicrobiol.2016.48.
39. Parks, D.H., C. Rinke, M. Chuvochina, P.A. Chaumeil, B.J. Woodcroft, P.N. Evans,
P. Hugenholtz, and G.W. Tyson, *Recovery of nearly 8,000 metagenome-assembled*
genomes substantially expands the tree of life. Nat. Microbiol., 2017. **2**: 1533-1542.
DOI: 10.1038/s41564-017-0012-7.
40. Shao, S., T. Cardona, and P.J. Nixon, *Early emergence of the FtsH proteases involved*
in photosystem II repair. Photosynthetica, 2018. **56**: 163-177. DOI: 10.1007/s11099-
018-0769-9.
41. Innan, H. and F. Kondrashov, *The evolution of gene duplications: classifying and*
distinguishing between models. Nature Reviews Genetics, 2010. **11**: 97-108. DOI:
10.1038/nrg2689.
42. Lynch, M. and J.S. Conery, *The evolutionary fate and consequences of duplicate*
genes. Science, 2000. **290**: 1151-1155. DOI: 10.1126/science.290.5494.1151.
43. Utami, Y.D., H. Kuwahara, K. Igai, T. Murakami, K. Sugaya, T. Morikawa, Y.
Nagura, M. Yuki, P. Deevong, T. Inoue, K. Kihara, N. Lo, A. Yamada, M. Ohkuma,
and Y. Hongoh, *Genome analyses of uncultured TG2/ZB3 bacteria in*
'Margulisbacteria' specifically attached to ectosymbiotic spirochetes of protists in the
termite gut. ISME J., 2019. **13**: 455-467. DOI: 10.1038/s41396-018-0297-4.
44. Kumar, S., G. Stecher, M. Suleski, and S.B. Hedges, *TimeTree: A resource for*
timelines, timetrees, and divergence times. Mol. Biol. Evol., 2017. **34**: 1812-1819.
DOI: 10.1093/molbev/msx116.
45. Marin, J., F.U. Battistuzzi, A.C. Brown, and S.B. Hedges, *The timetree of*
prokaryotes: New insights into their evolution and speciation. Mol. Biol. Evol., 2017.
34: 437-446. DOI: 10.1093/molbev/msw245.
46. Shih, P.M., L.M. Ward, and W.W. Fischer, *Evolution of the 3-hydroxypropionate*
bicycle and recent transfer of anoxygenic photosynthesis into the Chloroflexi. Proc.
Natl. Acad. Sci. U.S.A., 2017. **114**: 10749-10754. DOI: 10.1073/pnas.1710798114.
47. Tice, M.M. and D.R. Lowe, *Photosynthetic microbial mats in the 3,416-Myr-old*
ocean. Nature, 2004. **431**: 549-52. DOI: 10.1038/nature02888.
48. Rutherford, A.W. and P. Faller, *Photosystem II: evolutionary perspectives.* Philos.
Trans. Royal Soc. B, 2003. **358**: 245-253. DOI: 10.1098/rstb.2002.1186.
49. Nixon, P.J. and B.A. Diner, *Aspartate 170 of the Photosystem II reaction center*
polypeptide D1 is involved in the assembly of the oxygen-evolving manganese cluster.
Biochemistry, 1992. **31**: 942-948. DOI: 10.1021/Bi00118a041.
50. Adachi, Y., H. Kuroda, Y. Yukawa, and M. Sugiura, *Translation of partially*
overlapping psbD-psbC mRNAs in chloroplasts: the role of 5'-processing and
translational coupling. Nucleic Acids Res., 2012. **40**: 3152-8. DOI:
10.1093/nar/gkr1185.

51.   Carpenter, S.D., J. Charite, B. Eggers, and W.F. Vermaas, *The psbC start codon in*
1248   *Synechocystis sp. PCC 6803.* FEBS Lett., 1990. **260**: 135-7. DOI: 10.1016/0014-
1249   5793(90)80085-w.
1250   52.   Chisholm, D. and J.G. Williams, *Nucleotide sequence of psbC, the gene encoding the*
1251   *CP-43 chlorophyll a-binding protein of Photosystem II, in the cyanobacterium*
1252   *Synechocystis 6803.* Plant Mol. Biol., 1988. **10**: 293-301. DOI: 10.1007/BF00029879.
1253   53.   Cockell, C.S., *Ultraviolet radiation and the photobiology of earth's early oceans.*
1254   Orig. Life Evol. Biospheres, 2000. **30**: 467-99. DOI: 10.1023/a:1006765405786.
1255   54.   Lewis, C.A., J. Crayle, S.T. Zhou, R. Swanstrom, and R. Wolfenden, *Cytosine*
1256   *deamination and the precipitous decline of spontaneous mutation during Earth's*
1257   *history.* Proc. Natl. Acad. Sci. U.S.A., 2016. **113**: 8194-8199. DOI:
1258   10.1073/pnas.1607580113.
1259   55.   Levy, M. and S.L. Miller, *The stability of the RNA bases: implications for the origin*
1260   *of life.* Proc. Natl. Acad. Sci. U.S.A., 1998. **95**: 7933-8. DOI:
1261   10.1073/pnas.95.14.7933.
1262   56.   Eisen, J.A. and P.C. Hanawalt, *A phylogenomic study of DNA repair genes, proteins,*
1263   *and processes.* Mutat. Res./DNA Repair, 1999. **435**: 171-213. DOI: 10.1016/s0921-
1264   8777(99)00050-6.
1265   57.   Koonin, E.V. and M.Y. Galperin, *The major transitions in evolution: A comparative*
1266   *genomic perspective* in *Sequence — Evolution — Function: Computational*
1267   *Approaches in Comparative Genomics.* 2003, Springer-Science+Business Media,
1268   B.V>. 252-292. DOI: 10.1007/978-1-4757-3783-7.
1269   58.   Woese, C., *The universal ancestor.* Proc. Natl. Acad. Sci. U.S.A., 1998. **95**: 6854-9.
1270   DOI: 10.1073/pnas.95.12.6854.
1271   59.   Moore, K.R., C. Magnabosco, L. Momper, D.A. Gold, T. Bosak, and G.P. Fournier,
1272   *An Expanded Ribosomal Phylogeny of Cyanobacteria Supports a Deep Placement of*
1273   *Plastids.* Front. Microbiol., 2019. **10**. DOI: 10.3389/fmicb.2019.01612.
1274   60.   Soltis, P.S., R.A. Folk, and D.E. Soltis, *Darwin review: angiosperm phylogeny and*
1275   *evolutionary radiations.* P Roy Soc B-Biol Sci, 2019. **286**. DOI:
1276   10.1098/rspb.2019.0099.
1277   61.   McCutcheon, J.P. and N.A. Moran, *Extreme genome reduction in symbiotic bacteria.*
1278   Nat. Rev. Microbiol., 2011. **10**: 13-26. DOI: 10.1038/nrmicro2670.
1279   62.   van der Staay, G.W., S.Y. Moon-van der Staay, L. Garczarek, and F. Partensky, *Rapid*
1280   *evolutionary divergence of Photosystem I core subunits PsaA and PsaB in the marine*
1281   *prokaryote Prochlorococcus.* Photosynth Res, 2000. **65**: 131-9. DOI:
1282   10.1023/A:1006445810996.
1283   63.   David, L.A. and E.J. Alm, *Rapid evolutionary innovation during an Archaean genetic*
1284   *expansion.* Nature, 2011. **469**: 93-96. DOI: 10.1038/Nature09649.
1285   64.   Zhu, Q., U. Mai, W. Pfeiffer, S. Janssen, F. Asnicar, J.G. Sanders, P. Belda-Ferre,
1286   G.A. Al-Ghalith, E. Kopylova, D. McDonald, T. Kosciolek, J.B. Yin, S. Huang, N.
1287   Salam, J.-Y. Jiao, Z. Wu, Z.Z. Xu, K. Cantrell, Y. Yang, E. Sayyari, M. Rabiee, J.T.
1288   Morton, S. Podell, D. Knights, W.-J. Li, C. Huttenhower, N. Segata, L. Smarr, S.
1289   Mirarab, and R. Knight, *Phylogenomics of 10,575 genomes reveals evolutionary*
1290   *proximity between domains Bacteria and Archaea.* Nat. Commun., 2019. **10**: 5477.
1291   DOI: 10.1038/s41467-019-13443-4.
1292   65.   Berkemer, S.J. and S.E. McGlynn, *A New Analysis of Archaea–Bacteria Domain*
1293   *Separation: Variable Phylogenetic Distance and the Tempo of Early Evolution.* Mol.
1294   Biol. Evol., 2020. DOI: 10.1093/molbev/msaa089.
1295   66.   Homann, M., C. Heubeck, A. Airo, and M.M. Tice, *Morphological adaptations of*
1296   *3.22 Ga-old tufted microbial mats to Archean coastal habitats (Moodies Group,*

1297   *Barberton Greenstone Belt, South Africa).* Precambrian Res., 2015. **266**: 47-64. DOI:
1298   10.1016/j.precamres.2015.04.018.

1299   67.   Alcott, L.J., B.J.W. Mills, and S.W. Poulton, *Stepwise Earth oxygenation is an*
1300   *inherent property of global biogeochemical cycling.* Science, 2019. **366**: 1333-1337.
1301   DOI: 10.1126/science.aax6459.

1302   68.   Kadoya, S., D.C. Catling, R.W. Nicklas, I.S. Puchtel, and A.D. Anbar, *Mantle data*
1303   *imply a decline of oxidizable volcanic gases could have triggered the Great*
1304   *Oxidation.* Nat. Commun., 2020. **11**: 2774. DOI: 10.1038/s41467-020-16493-1.

1305   69.   Reinhard, C.T. and N.J. Planavsky, *Biogeochemical Controls on the Redox Evolution*
1306   *of Earth's Oceans and Atmosphere.* Elements, 2020. **16**: 191-196. DOI:
1307   10.2138/gselements.16.3.191.

1308   70.   Cardona, T., *Photosystem II is a chimera of reaction centers.* J. Mol. Evol., 2017. **84**:
1309   149-151. DOI: 10.1007/s00239-017-9784-x.

1310   71.   Trinugroho, J.P., M. Beckova, S. Shao, J. Yu, Z. Zhao, J.W. Murray, R. Sobotka, J.
1311   Komenda, and P.J. Nixon, *Chlorophyll f synthesis by a super-rogue photosystem II*
1312   *complex.* Nat Plants, 2020. **6**: 238-244. DOI: 10.1038/s41477-020-0616-4.

1313   72.   Wegener, K.M., A. Nagarajan, and H.B. Pakrasi, *An atypical psbA gene encodes a*
1314   *sentinel D1 protein to form a physiologically relevant inactive Photosystem II*
1315   *complex in Cyanobacteria.* J. Biol. Chem., 2015. **290**: 3764-74. DOI:
1316   10.1074/jbc.M114.604124.

1317   73.   Cardona, T., *Thinking twice about the evolution of photosynthesis.* Open Biology,
1318   2019. **9**: 180246. DOI: 10.1098/rsob.180246.

1319   74.   Petrov, A.S., C.R. Bernier, C.L. Hsiao, A.M. Norris, N.A. Kovacs, C.C. Waterbury,
1320   V.G. Stepanov, S.C. Harvey, G.E. Fox, R.M. Wartell, N.V. Hud, and L.D. Williams,
1321   *Evolution of the ribosome at atomic resolution.* Proc. Natl. Acad. Sci. U.S.A., 2014.
1322   **111**: 10251-10256. DOI: 10.1073/pnas.1407205111.

1323   75.   Granick, S., *Speculations on the origins and evolution of photosynthesis.* Ann. N. Y.
1324   Acad. Sci., 1957. **69**: 292-308. DOI: 10.1111/j.1749-6632.1957.tb49665.x.

1325   76.   Mauzerall, D., *Light, iron, Sam Granick and the origin of life.* Photosynth. Res., 1992.
1326   **33**: 163-170. DOI: 10.1007/BF00039178.

1327   77.   Mulkidjanian, A.Y., A.Y. Bychkov, D.V. Dibrova, M.Y. Galperin, and E.V. Koonin,
1328   *Origin of first cells at terrestrial, anoxic geothermal fields.* Proc. Natl. Acad. Sci.
1329   U.S.A., 2012. **109**: E821-E830. DOI: 10.1073/pnas.1117774109.

1330   78.   Franz, H.B., P.R. Mahaffy, C.R. Webster, G.J. Flesch, E. Raaen, C. Freissinet, S.K.
1331   Atreya, C.H. House, A.C. McAdam, C.A. Knudson, P.D. Archer, J.C. Stern, A.
1332   Steele, B. Sutter, J.L. Eigenbrode, D.P. Glavin, J.M.T. Lewis, C.A. Malespin, M.
1333   Millan, D.W. Ming, R. Navarro-Gonzalez, and R.E. Summons, *Indigenous and*
1334   *exogenous organics and surface-atmosphere cycling inferred from carbon and oxygen*
1335   *isotopes at Gale crater.* Nat Astron, 2020. DOI: 10.1038/s41550-019-0990-x.

1336   79.   Lu, A.H., Y. Li, H.R. Ding, X.M. Xu, Y.Z. Li, G.P. Ren, J. Liang, Y.W. Liu, H.
1337   Hong, N. Chen, S.Q. Chu, F.F. Liu, H.R. Wang, C. Ding, C.Q. Wang, Y. Lai, J. Liu,
1338   J. Dick, K.H. Liu, and M.F. Hochella, *Photoelectric conversion on Earth's surface via*
1339   *widespread Fe- and Mn-mineral coatings.* Proc. Natl. Acad. Sci. U.S.A., 2019. **116**:
1340   9741-9746. DOI: 10.1073/pnas.1902473116.

1341   80.   Mendler, K., H. Chen, D.H. Parks, B. Lobb, L.A. Hug, and A.C. Doxey, *AnnoTree:*
1342   *visualization and exploration of a functionally annotated microbial tree of life.*
1343   Nucleic Acids Res., 2019. **47**: 4442-4448. DOI: 10.1093/nar/gkz246.

1344   81.   Ward, L.M., T. Cardona, and H. Holland-Moritz, *Evolutionary implications of*
1345   *anoxygenic phototrophy in the bacterial phylum Candidatus Eremiobacterota (WPS-*
1346   *2).* Front. Microbiol., 2019. **10**: 1658. DOI: 10.3389/fmicb.2019.01658.

1347   82.   Gruber-Vodicka, H.R., N. Leisch, M. Kleiner, T. Hinzke, M. Liebeke, M. McFall-
1348   Ngai, M.G. Hadfield, and N. Dubilier, *Two intracellular and cell type-specific*
1349   *bacterial symbionts in the placozoan Trichoplax H2.* Nat. Microbiol., 2019. **4**: 1465-
1350   1474. DOI: 10.1038/s41564-019-0475-9.
1351   83.   Sievers, F., A. Wilm, D. Dineen, T.J. Gibson, K. Karplus, W.Z. Li, R. Lopez, H.
1352   McWilliam, M. Remmert, J. Soding, J.D. Thompson, and D.G. Higgins, *Fast,*
1353   *scalable generation of high-quality protein multiple sequence alignments using*
1354   *Clustal Omega.* Mol. Syst. Biol., 2011. **7**: 539. DOI: 10.1038/msb.2011.75.
1355   84.   Castresana, J., *Selection of conserved blocks from multiple alignments for their use in*
1356   *phylogenetic analysis.* Mol. Biol. Evol., 2000. **17**: 540-552. DOI:
1357   10.1093/oxfordjournals.molbev.a026334.
1358   85.   Guindon, S., J.F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk, and O. Gascuel,
1359   *New algorithms and methods to estimate maximum-likelihood phylogenies: assessing*
1360   *the performance of PhyML 3.0.* Syst. Biol., 2010. **59**: 307-21. DOI:
1361   10.1093/sysbio/syq010.
1362   86.   Lefort, V., J.E. Longueville, and O. Gascuel, *SMS: Smart model selection in PhyML.*
1363   Mol. Biol. Evol., 2017. **34**: 2422-2424. DOI: 10.1093/molbev/msx149.
1364   87.   Anisimova, M. and O. Gascuel, *Approximate likelihood-ratio test for branches: A*
1365   *fast, accurate, and powerful alternative.* Syst. Biol., 2006. **55**: 539-52. DOI:
1366   10.1080/10635150600755453.
1367   88.   Huson, D.H. and C. Scornavacca, *Dendroscope 3: an interactive tool for rooted*
1368   *phylogenetic trees and networks.* Syst. Biol., 2012. **61**: 1061-7. DOI:
1369   10.1093/sysbio/sys062.
1370   89.   Gascuel, O., *BIONJ: An improved version of the NJ algorithm based on a simple*
1371   *model of sequence data.* Mol. Biol. Evol., 1997. **14**: 685-695. DOI:
1372   10.1093/oxfordjournals.molbev.a025808.
1373   90.   Gouy, M., S. Guindon, and O. Gascuel, *SeaView version 4: A multiplatform graphical*
1374   *user interface for sequence alignment and phylogenetic tree building.* Mol. Biol.
1375   Evol., 2010. **27**: 221-224. DOI: 10.1093/molbev/msp259.
1376   91.   Kumar, S., G. Stecher, M. Li, C. Knyaz, and K. Tamura, *MEGA X: Molecular*
1377   *evolutionary genetics analysis across computing platforms.* Mol. Biol. Evol., 2018.
1378   **35**: 1547-1549. DOI: 10.1093/molbev/msy096.
1379   92.   Clarke, J.T., R.C.M. Warnock, and P.C.J. Donoghue, *Establishing a time-scale for*
1380   *plant evolution.* New Phytol, 2011. **192**: 266-301. DOI: 10.1111/j.1469-
1381   8137.2011.03794.x.
1382   93.   Morris, J.L., M.N. Puttick, J.W. Clark, D. Edwards, P. Kenrick, S. Pressel, C.H.
1383   Wellman, Z.H. Yang, H. Schneider, and P.C.J. Donoghue, *The timescale of early land*
1384   *plant evolution.* Proc. Natl. Acad. Sci. U.S.A., 2018. **115**: E2274-E2283. DOI:
1385   10.1073/pnas.1719588115.
1386   94.   Butterfield, N.J., *Bangiomorpha pubescens n. gen., n. sp.: implications for the*
1387   *evolution of sex, multicellularity, and the Mesoproterozoic/Neoproterozoic radiation*
1388   *of eukaryotes.* Paleobiology, 2000. **26**: 386-404. DOI: 10.1666/0094-
1389   8373(2000)026<0386:Bpngns>2.0.Co;2.
1390   95.   Knoll, A.H., S. Worndle, and L.C. Kah, *Covariance of microfossil assemblages and*
1391   *microbialite textures across an upper mesoproterozoic carbonate platform.* Palaios,
1392   2013. **28**: 453-470. DOI: 10.2110/palo.2013.p13-005r.
1393   96.   Gibson, T.M., P.M. Shih, V.M. Cumming, W.W. Fischer, P.W. Crockford, M.S.W.
1394   Hodgskiss, S. Worndle, R.A. Creaser, R.H. Rainbird, T.M. Skulski, and G.P.
1395   Halverson, *Precise age of Bangiomorpha pubescens dates the origin of eukaryotic*
1396   *photosynthesis.* Geology, 2017. **46**: 135-138. DOI: 10.1130/G39829.1.

97.   Bengtson, S., T. Sallstedt, V. Belivanova, and M. Whitehouse, *Three-dimensional*
preservation of cellular and subcellular structures suggests 1.6 billion-year-old
crown-group red algae. Plos Biol, 2017. **15**: e2000735. DOI:
10.1371/journal.pbio.2000735.
98.   Sallstedt, T., S. Bengtson, C. Broman, P.M. Crill, and D.E. Canfield, *Evidence of*
oxygenic phototrophy in ancient phosphatic stromatolites from the Paleoproterozoic
Vindhyan and Aravalli Supergroups, India. Geobiology, 2018. **16**: 139-159. DOI:
10.1111/gbi.12274.
99.   Qu, Y.G., S.X. Zhu, M. Whitehouse, A. Engdahl, and N. McLoughlin, *Carbonaceous*
biosignatures of the earliest putative macroscopic multicellular eukaryotes from 1630
Ma Tuanshanzi Formation, north China. Precambrian Res., 2018. **304**: 99-109. DOI:
10.1016/j.precamres.2017.11.004.
100.   Pang, K., Q. Tang, L. Chen, B. Wan, C. Niu, X. Yuan, and S. Xiao, *Nitrogen-fixing*
heterocystous cyanobacteria in the tonian period. Curr Biol, 2018. **28**: 616-622 e1.
DOI: 10.1016/j.cub.2018.01.008.
101.   Stothard, P., *The sequence manipulation suite: JavaScript programs for analyzing*
and formatting protein and DNA sequences. BioTechniques, 2000. **28**: 1102-1104.
DOI: 10.2144/00286ir01.
102.   Ponce-Toledo, R.I., P. Deschamps, P. Lopez-Garcia, Y. Zivanovic, K. Benzerara, and
D. Moreira, *An early-branching freshwater cyanobacterium at the origin of plastids.*
Curr Biol, 2017. **27**: 386-391. DOI: 10.1016/j.cub.2016.11.056.
103.   Sánchez-Baracaldo, P., J.A. Raven, D. Pisani, and A.H. Knoll, *Early photosynthetic*
eukaryotes inhabited low-salinity habitats. Proc. Natl. Acad. Sci. U.S.A., 2017. **114**:
E7737-E7745. DOI: 10.1073/pnas.1620089114.
104.   Fewer, D., T. Friedl, and B. Budel, *Chroococcidiopsis and heterocyst-differentiating*
cyanobacteria are each other's closest living relatives. Mol. Phylogen. Evol., 2002.
23: 82-90. DOI: 10.1006/mpev.2001.1075.
105.   Shih, P.M., D. Wu, A. Latifi, S.D. Axen, D.P. Fewer, E. Talla, A. Calteau, F. Cai, N.
Tandeau de Marsac, R. Rippka, M. Herdman, K. Sivonen, T. Coursin, T. Laurent, L.
Goodwin, M. Nolan, K.W. Davenport, C.S. Han, E.M. Rubin, J.A. Eisen, T. Woyke,
M. Gugger, and C.A. Kerfeld, *Improving the coverage of the cyanobacterial phylum*
using diversity-driven genome sequencing. Proc. Natl. Acad. Sci. U.S.A., 2013. **110**:
1053-8. DOI: 10.1073/pnas.1217107110.
106.   Schirrmeister, B.E., A. Antonelli, and H.C. Bagheri, *The origin of multicellularity in*
cyanobacteria. BMC Evol. Biol., 2011. **11**: 45. DOI: 10.1186/1471-2148-11-45.
107.   Ciccarelli, F.D., T. Doerks, C. von Mering, C.J. Creevey, B. Snel, and P. Bork,
Toward automatic reconstruction of a highly resolved tree of life. Science, 2006. **311**:
1283-7. DOI: 10.1126/science.1123061.
108.   Nakamura, Y., T. Kaneko, S. Sato, M. Mimuro, H. Miyashita, T. Tsuchiya, S.
Sasamoto, A. Watanabe, K. Kawashima, Y. Kishida, C. Kiyokawa, M. Kohara, M.
Matsumoto, A. Matsuno, N. Nakazaki, S. Shimpo, C. Takeuchi, M. Yamada, and S.
Tabata, *Complete genome structure of Gloeobacter violaceus PCC 7421, a*
cyanobacterium that lacks thylakoids. DNA Res, 2003. **10**: 137-145. DOI:
10.1093/dnares/10.4.137.
109.   Honda, D., A. Yokota, and J. Sugiyama, *Detection of seven major evolutionary*
lineages in cyanobacteria based on the 16S rRNA gene sequence analysis with new
sequences of five marine Synechococcus strains. J. Mol. Biol., 1999. **48**: 723-739.
DOI: 10.1007/Pl00006517.

110.   Knoll, A.H., E.J. Javaux, D. Hewitt, and P. Cohen, *Eukaryotic organisms in*
1446   *Proterozoic oceans.* Philos. Trans. Royal Soc. B, 2006. **361**: 1023-1038. DOI:
1447   10.1098/rstb.2006.1843.
1448   111.   Han, T.M. and B. Runnegar, *Megascopic eukaryotic algae from the 2.1-billion-year-*
1449   *old Negaunee iron-formation, Michigan.* Science, 1992. **257**: 232-235. DOI:
1450   10.1126/science.1631544.
1451   112.   El Albani, A., M.G. Mangano, L.A. Buatois, S. Bengtson, A. Riboulleau, A. Bekker,
1452   K. Konhauser, T. Lyons, C. Rollion-Bard, O. Bankole, S.G.L. Baghekema, A.
1453   Meunier, A. Trentesaux, A. Mazurier, J. Aubineau, C. Laforest, C. Fontaine, P.
1454   Recourt, E.C. Fru, R. Macchiarelli, J.Y. Reynaud, F. Gauthier-Lafaye, and D.E.
1455   Canfield, *Organism motility in an oxygenated shallow-marine environment 2.1 billion*
1456   *years ago.* Proc. Natl. Acad. Sci. U.S.A., 2019. **116**: 3431-3436. DOI:
1457   10.1073/pnas.1815721116.
1458   113.   Shi, M., Q.L. Feng, M.Z. Khan, and S.X. Zhu, *An eukaryote-bearing microbiota from*
1459   *the early mesoproterozoic Gaoyuzhuang Formation, Tianjin, China and its*
1460   *significance.* Precambrian Res., 2017. **303**: 709-726. DOI:
1461   10.1016/j.precamres.2017.09.013.
1462   114.   Sánchez-Baracaldo, P., *Origin of marine planktonic cyanobacteria.* Sci. Rep., 2015.
1463   **5**: 17418. DOI: 10.1038/srep17418.
1464   115.   Lepage, T., D. Bryant, H. Philippe, and N. Lartillot, *A general comparison of relaxed*
1465   *molecular clock models.* Mol. Biol. Evol., 2007. **24**: 2669-80. DOI:
1466   10.1093/molbev/msm193.
1467   116.   Kishino, H., J.L. Thorne, and W.J. Bruno, *Performance of a divergence time*
1468   *estimation method under a probabilistic model of rate evolution.* Mol. Biol. Evol.,
1469   2001. **18**: 352-361. DOI: 10.1093/oxfordjournals.molbev.a003811.
1470   117.   Langklotz, S., U. Baumann, and F. Narberhaus, *Structure and function of the*
1471   *bacterial AAA protease FtsH.* Biochim. Biophys. Acta, 2012. **1823**: 40-8. DOI:
1472   10.1016/j.bbamcr.2011.08.015.
1473   118.   Medlin, L.K., W.H.C.F. Kooistra, R. Gersonde, P.A. Sims, and U. Wellbrock, *Is the*
1474   *origin of the diatoms related to the end-Permian mass extinction?* Nova Hedwigia,
1475   1997. **65**: 1-11.
1476   119.   Barboni, M., P. Boehnke, B. Keller, I.E. Kohl, B. Schoene, E.D. Young, and K.D.
1477   McKeegan, *Early formation of the Moon 4.51 billion years ago.* Science Advances,
1478   2017. **3**: e1602365. DOI: 10.1126/sciadv.1602365.
1479   120.   Drummond, A.J., S.Y.W. Ho, M.J. Phillips, and A. Rambaut, *Relaxed phylogenetics*
1480   *and dating with confidence.* Plos Biol, 2006. **4**: 699-710. DOI:
1481   10.1371/journal.pbio.0040088.
1482   121.   Hodgskiss, M.S.W., O.M.J. Dagnaud, J.L. Frost, G.P. Halverson, M.D. Schmitz, N.L.
1483   Swanson-Hysell, and E.A. Sperling, *New insights on the Orosirian carbon cycle,*
1484   *early Cyanobacteria, and the assembly of Laurentia from the Paleoproterozoic*
1485   *Belcher Group.* Earth Planet. Sci. Lett., 2019. **520**: 141-152. DOI:
1486   10.1016/j.epsl.2019.05.023.
1487   122.   Brocks, J.J., G.D. Love, R.E. Summons, A.H. Knoll, G.A. Logan, and S.A. Bowden,
1488   *Biomarker evidence for green and purple sulphur bacteria in a stratified*
1489   *Palaeoproterozoic sea.* Nature, 2005. **437**: 866-870. DOI: 10.1038/Nature04068.
1490   123.   Engel, M.S. and D.A. Grimaldi, *New light shed on the oldest insect.* Nature, 2004.
1491   **427**: 627-630. DOI: 10.1038/nature02291.
1492   124.   Legendre, F., A. Nel, G.J. Svenson, T. Robillard, R. Pellens, and P. Grandcolas,
1493   *Phylogeny of Dictyoptera: Dating the origin of cockroaches, praying mantises and*

1494		*termites with molecular data and controlled fossil evidence.* PloS one, 2015. **10**:
1495		e0130127. DOI: 10.1371/journal.pone.0130127.
1496	125.	Eitel, M., H.J. Osigus, R. DeSalle, and B. Schierwater, *Global diversity of the*
1497		*Placozoa.* PloS one, 2013. **8**: e57131. DOI: 10.1371/journal.pone.0057131.
1498	126.	Zumberge, J.A., G.D. Love, P. Cardenas, E.A. Sperling, S. Gunasekera, M. Rohrssen,
1499		E. Grosjean, J.P. Grotzinger, and R.E. Summons, *Demosponge steroid biomarker 26-*
1500		*methylstigmastane provides evidence for Neoproterozoic animals.* Nat. Ecol. Evol.,
1501		2018. **2**: 1709-1714. DOI: 10.1038/s41559-018-0676-2.
1502	127.	Grossnickle, D.M., S.M. Smith, and G.P. Wilson, *Untangling the multiple ecological*
1503		*radiations of early mammals.* Trends Ecol Evol, 2019. **34**: 936-949. DOI:
1504		10.1016/j.tree.2019.05.008.
1505	128.	Williams, B.A., R.F. Kay, and E.C. Kirk, *New perspectives on anthropoid origins.*
1506		Proc. Natl. Acad. Sci. U.S.A., 2010. **107**: 4797-804. DOI: 10.1073/pnas.0908320107.
1507	129.	Soo, R.M., B.J. Woodcroft, D.H. Parks, G.W. Tyson, and P. Hugenholtz, *Back from*
1508		*the dead; the curious tale of the predatory cyanobacterium Vampirovibrio*
1509		*chlorellavorus.* PeerJ, 2015. **3**: e968. DOI: 10.7717/peerj.968.
1510	130.	Hahn, M.W., J. Schmidt, A. Pitt, S.J. Taipale, and E. Lang, *Reclassification of four*
1511		*Polynucleobacter necessarius strains as representatives of Polynucleobacter*
1512		*asymbioticus comb. nov., Polynucleobacter duraquae sp. nov., Polynucleobacter*
1513		*yangtzensis sp. nov and Polynucleobacter sinensis sp.s nov., and emended description*
1514		*of Polynucleobacter necessarius.* Int. J. Syst. Evol. Microbiol., 2016. **66**: 2883-2892.
1515		DOI: 10.1099/ijsem.0.001073.
1516	131.	Parfrey, L.W., D.J. Lahr, A.H. Knoll, and L.A. Katz, *Estimating the timing of early*
1517		*eukaryotic diversification with multigene molecular clocks.* Proc. Natl. Acad. Sci.
1518		U.S.A., 2011. **108**: 13624-9. DOI: 10.1073/pnas.1110633108.
1519	132.	Fernandes, N.M. and C.G. Schrago, *A multigene timescale and diversification*
1520		*dynamics of Ciliophora evolution.* Mol. Phylogen. Evol., 2019. **139**: 106521. DOI:
1521		10.1016/j.ympev.2019.106521.
1522	133.	Yu, X.-J. and D.H. Walker, *The Order Rickettsiales*, in *The prokaryotes: A handbook*
1523		*on the biology of bacteria*, M. Dworkin, Editor. 2006, Springer: Singapure. 493-528.
1524		DOI: 10.1007/0-387-30745-1_20.
1525	134.	Roger, A.J., S.A. Munoz-Gomez, and R. Kamikawa, *The Origin and Diversification*
1526		*of Mitochondria.* Curr Biol, 2017. **27**: R1177-R1192. DOI:
1527		10.1016/j.cub.2017.09.015.
1528	135.	Li, W. and A. Godzik, *Cd-hit: a fast program for clustering and comparing large sets*
1529		*of protein or nucleotide sequences.* Bioinformatics, 2006. **22**: 1658-9. DOI:
1530		10.1093/bioinformatics/btl158.
1531	136.	Ashkenazy, H., O. Penn, A. Doron-Faigenboim, O. Cohen, G. Cannarozzi, O. Zomer,
1532		and T. Pupko, *FastML: a web server for probabilistic reconstruction of ancestral*
1533		*sequences.* Nucleic Acids Res., 2012. **40**: W580-W584. DOI: 10.1093/nar/gks498.
1534	137.	Hanson-Smith, V., B. Kolaczkowski, and J.W. Thornton, *Robustness of ancestral*
1535		*sequence reconstruction to phylogenetic uncertainty.* Mol. Biol. Evol., 2010. **27**:
1536		1988-1999. DOI: 10.1093/molbev/msq081.
1537	138.	Yang, Z.H., *PAML 4: Phylogenetic analysis by maximum likelihood.* Mol. Biol. Evol.,
1538		2007. **24**: 1586-1591. DOI: 10.1093/molbev/msm088.
1539	139.	Kumar, S., G. Stecher, and K. Tamura, *MEGA7: Molecular evolutionary genetics*
1540		*analysis version 7.0 for bigger datasets.* Mol. Biol. Evol., 2016. **33**: 1870-1874. DOI:
1541		10.1093/molbev/msw054.

140.   Yu, L.J., M. Suga, Z.Y. Wang-Otomo, and J.R. Shen, *Structure of photosynthetic*
LH1-RC supercomplex at 1.9 Å resolution. Nature, 2018. **556**: 209-213. DOI:
10.1038/s41586-018-0002-9.
141.   Gisriel, C., I. Sarrou, B. Ferlez, J.H. Golbeck, K.E. Redding, and R. Fromme,
Structure of a symmetric photosynthetic reaction center-photosystem. Science, 2017.
357: 1021-1025. DOI: 10.1126/science.aan5611.
142.   Jordan, P., P. Fromme, H.T. Witt, O. Klukas, W. Saenger, and N. Krauss, *Three-*
dimensional structure of cyanobacterial Photosystem I at 2.5 Å resolution. Nature,
2001. **411**: 909-17. DOI: 10.1038/35082000.
143.   Toporik, H., J. Li, D. Williams, P.L. Chiu, and Y. Mazor, *The structure of the stress-*
induced photosystem I-IsiA antenna supercomplex. Nat Struct Mol Biol, 2019. **26**:
443-449. DOI: 10.1038/s41594-019-0228-8.
144.   Jia, Y.T., G. Dewey, I.N. Shindyalov, and P.E. Bourne, *A new scoring function and*
associated statistical significance for structure alignment by CE. J Comput Biol,
2004. **11**: 787-799. DOI: Doi 10.1089/1066527042432260.

1557