Non-redundant roles of caspase-4 and caspase-5 in heme driven- IL-1β release and cell death

Beatriz E. Bolívar[1, 2], Brittany A. Rohrman[1], Chloé I. Charendoff[1], Vanda Yazdani [1, 2], John D. Belcher[3], Gregory M. Vercellotti[3], Jonathan M. Flanagan[1, 2] and Lisa Bouchier-Hayes[1, 2,4*]

[1]Department of Pediatrics, Division of Hematology-Oncology, Baylor College of Medicine, Houston, TX 77030, USA.

[2]William T. Shearer Center for Human Immunobiology, Texas Children's Hospital, Houston, TX 77030, USA.

[3]Department of Medicine, Division of Hematology, Oncology and Transplantation, University of Minnesota, Minneapolis, MN 55455, USA.

[4]Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, TX 77030, USA.

*Corresponding author:

Lisa Bouchier-Hayes   E-mail: bouchier@bcm.edu

Running Title

Heme activates inflammatory caspases

Abstract

Excessive release of heme from red blood cells is a key pathophysiological feature of several disease states including bacterial sepsis, malaria, and sickle cell disease. This hemolysis results in an increased level of free heme that has been implicated in the inflammatory activation of monocytes, macrophages, and endothelium. Here we show that extracellular heme engages human inflammatory caspases, caspase-1, caspase-4, and caspase-5, resulting in the release of IL-1β. Heme-induced IL-1β release was further increased in macrophages from patients with sickle cell disease. Heme activates caspase-1 in an inflammasome-dependent manner, but heme-induced activation of caspase-4 and caspase-5 is inflammasome-independent. Furthermore, we show that caspase-5, but not caspase-4, is essential for heme-induced IL-1β release, while caspase-4 contributes to heme-induced cell death. Together, we have identified that extracellular heme acts as a DAMP that can engage canonical and non-canonical inflammasome activation as a key mediator of inflammation in macrophages.

Key words

Heme, inflammasome, caspase-1, caspase-4, caspase-5

Introduction

The interactions between inflammatory caspases and inflammasomes are critical for preventing uncontrolled inflammation and mediating appropriate inflammation under infectious and sterile conditions. Inflammasomes are multi-protein complexes that provide the platform for recruitment and activation of inflammatory caspases and are essential for cellular inflammatory responses (1). The inflammatory caspases include the human caspases-1, -4, and -5 and murine caspase-11 (1). This subset of the broader caspase protease family does not mediate apoptosis but specifically regulates inflammation by facilitating the activation and release of the pro-inflammatory cytokines interleukin (IL)-1β and IL-18 (2). While the essential nature of inflammatory caspases in pathogen clearance is well established, their role in sterile inflammation (inflammation in the absence of infection) is less clear. Sterile inflammation occurs when non-pathogenic inflammatory stimuli activate inflammasomes. These stimuli are known as damage-associated molecular patterns (DAMPs) and are generally endogenous signals released by cells. This type of inflammation is important for wound healing and tissue regeneration but, if unchecked, can contribute to tissue damage associated with conditions like ischemic stroke, myocardial infarction, and neurodegeneration (3). Despite the importance of DAMPs for triggering inflammation, the endogenous signaling molecules that trigger sterile inflammation are not fully resolved (4).

Heme has the features of a DAMP because it is released following red blood cell destruction and triggers an inflammatory response. Extracellular hemoglobin and heme are highly pro-oxidant molecules that are assiduously scavenged by haptoglobin (Hp) and hemopexin (Hpx), respectively. Excessive hemolysis can saturate and deplete the Hp and Hpx systems, resulting in free heme with strong pro-inflammatory capabilities (5). Heme has been shown to activate caspase-1 in mouse macrophages via assembly of the NLRP3 inflammasome (6). Heme has also

been shown to activate toll-like receptor (TLR) 4 in endothelial cells to activate NFκB (7). For IL-

1β release to proceed, two signals are needed. Signal 1 activates NFκB to induce pro-IL-1β

expression and expression of additional inflammasome proteins including NLRP3, in a process

known as priming. Signal 2 provides an intracellular signal that induces inflammasome assembly

and caspase-1 activation that cleaves pro-IL-1β to its mature form, which is released from the cell

(8). Heme is naturally taken up and recycled by macrophages, providing a physiological

intracellular Signal 2 (44). Due to its ability to activate both caspase-1 and TLR4 and to be

internalized by macrophages, extracellular heme has the properties of a DAMP that could

potentially provide both Signal 1 and Signal 2 to initiate an effective inflammatory response.

Excessive release of heme from red blood cells is a key feature of several pathological states,

including sepsis, malaria, and sickle cell disease (SCD). SCD is the most prevalent inherited blood

disorder, affecting approximately 100,000 Americans and millions worldwide (9-11). The clinical

manifestations of SCD arise from a complex pathophysiology including chronic hemolytic anemia,

increased susceptibility to infection, and vaso-occlusive events (12). Chronically elevated heme

levels induce the inflammatory activation of monocytes, macrophages, and the endothelium (7,

13, 14). This inflammation can result in vaso-occlusion, acute chest syndrome, and organ damage

(5, 7, 15). Heme-induced activation of monocytes and macrophages contributes to these severe

complications through release of inflammatory cytokines, like IL-1β, that trigger endothelial

activation, upregulation of adhesion factors, and vaso-occlusion (16, 17). Indeed, in a study of

children with SCD, patients having a vaso-occlusive pain crisis demonstrated elevated levels of

proinflammatory cytokines IL-1β, IL-6, IL-10, tumor necrosis factor (TNF)-α, and free heme (18).

The role of the inflammatory caspases in heme-induced inflammation in the context of hemolytic

disorders such as SCD has not been well studied.

Mice deficient in caspase-1 or the inflammasome proteins NLRP3 or ASC survive following

hemolysis induced by a lethal dose of phenylhydrazine (6). Thus, caspase-1 is essential for an

effective inflammatory response to hemolysis. However, the roles of caspase-4 and caspase-5 in

this process are unknown. Caspase-4 and caspase-5 are the human orthologues of murine

caspase-11. Caspase-4, -5, and -11 have been shown to be activated by intracellular LPS

independent of inflammasomes and they each have been shown to cleave the pore-forming

protein gasdermin D (GSDMD) (19-21). Cleavage of GSDMD allows its N-terminal fragment to

insert in the plasma membrane forming a pore predicted to be 180Å in diameter (22). This pore

is of sufficient size to allow release of mature IL-1β, but also permits influx of ions leading to cell

swelling and a necrotic form of cell death called pyroptosis (23-25). Although caspase-1 can also

cleave GSDMD, blocking caspase-1 dependent cleavage delays, but does not inhibit, pyroptosis

(19). Therefore, current thinking is that caspase-1 cleaves proIL-1β and proIL-18, while caspase-

4, caspase-5, and caspase-11 cleave GSDMD to induce pyroptosis, allowing active cytokine

release. It has been proposed that cytokine release and pyroptosis cannot be uncoupled (26).

However, contradicting this theory, some reports show living cells releasing IL-1β (27, 28). Here,

we show that heme induces inflammasome dependent caspase-1 activation and inflammasome-

independent caspase-4 and caspase-5 activation. Furthermore, we show that caspase-5 is

essential for IL-1β release and caspase-4 contributes to heme-induced cell death.

  Results

  Heme-induced IL-1β release from human macrophages requires priming

  It has been previously shown that heme can induce IL-1β release in LPS-primed murine bone

  marrow derived macrophages (mBMDM) and that heme can activate TLR4 to induce NFκB

  activation in endothelial cells (6, 7). This suggests that heme can provide both the priming signal

  (Signal 1) required to induce pro-IL-1β expression and the inflammasome activating signal (Signal

  2) required to induce inflammasome-dependent caspase-1 activation. To test this in human

  macrophages, we isolated CD14+ monocytes from peripheral blood from healthy donors and

  differentiated them into macrophages with granulocyte-macrophage colony-stimulating factor

  (GM-CSF). We exposed the macrophages to heme in the presence or absence of prior priming

  with LPS. Treatment with heme in non-primed macrophages induced a modest increase in IL-1β.

  However, in order for heme to induce maximal IL-1β release, prior priming with LPS was required.

  Treatment with LPS alone resulted in similar IL-1β release to heme alone (Figure 1A). To show

  that the IL-1β release was specific to heme, we increased the amount of serum in the culture

  media. Hemopexin and albumin are present in FBS and both bind to heme and to prevent its

  cellular uptake (29-32). As little as 1% FBS concentration in the cellular media was sufficient to

  block heme-induced IL-1β release (Figure 1B).

  Sickle cell macrophages are constantly exposed to high levels of extracellular heme due to

  persistent hemolysis of sickled red blood cells (13). To determine the effect of this on IL-1β

  release, we compared human monocyte-derived macrophages (hMDM) from healthy donors to

  hMDM isolated from SCD patients. We noted that heme-treated hMDM from SCD patients were

  more sensitive to heme-induced IL-1β release relative to the control hMDM (Figure 1C). However,

  priming with LPS was required for strong IL-1β release in both groups. Together, these results

  suggest that while heme is able to activate caspase-1 to induce IL-1β release, it is not sufficient

to prime the cells for IL-1β release.

It has been shown that macrophages from sickle mice express higher levels of M1 markers (33).

To investigate IL-1β release by heme in the two extreme macrophage subtypes, we polarized the

CD14+ monocytes towards M1 (pro-inflammatory) by treating with GMCSF for seven days or

towards M2 (anti-inflammatory) by treating with IL-4 for a further day after the incubation with

GMCSF. LPS-primed M1 or M2 hMDM were stimulated with heme or with ATP. We showed lower

levels of IL-1β release in M2 compared to M1 hMDM in response to both heme and ATP (Figure

1D). In response to heme, M1 macrophages underwent cell death with a necrotic morphology. In

contrast, M2 macrophages remained viable with an intact plasma membrane but they appeared

more flattened and granular than the unstimulated cells (Figure 1E). Together, these results

suggest that SCD macrophages have increased inflammatory caspase activity resulting in

increased IL-1β activation and that this may be due, in part, to the increased proportion of M1

macrophage population in patients with SCD.

Caspase-1, caspase-4, and caspase-5 are activated by heme in the absence of priming

Given that heme induces IL-1β release in macrophages, we next investigated the ability of heme

to activate caspase-1 and the other human inflammatory caspases, caspase-4 and caspase-5.

Caspase-1 is activated upon proximity-induced dimerization following recruitment to

inflammasomes (1, 34). The recruitment of caspase-1 to an inflammasome is mediated by specific

interactions between specific protein interaction domains in the inflammasome proteins.

Specifically, the caspase recruitment domain (CARD) in ASC or NLRC4 binds to the CARD in the

prodomain of caspase-1 to recruit it to the inflammasome and facilitate induced proximity of the

caspase (35, 36). To measure the induced proximity of each caspase, we used caspase

bimolecular fluorescence complementation (BiFC) (37). BiFC uses non-fluorescent fragments of

the yellow fluorescent protein, Venus ("split Venus") that can associate to reform the fluorescent

Venus complex when fused to interacting proteins (38). When the CARD-containing prodomain

of caspase-1 is fused to each half of split Venus, recruitment of caspase-1 to inflammasomes and

the subsequent induced proximity results in enforced association of the two Venus halves (39).

Thus Venus fluorescence (BiFC) acts as a read-out for caspase induced proximity, the proximal

step for activation. We transiently expressed the prodomain of caspase-1 (C1-Pro, aa 1-102)

fused to each of the split Venus fragments, Venus C (aa 155-239) and Venus N (aa 1-173) in

hMDM (Figure 2A). Following exposure to heme, we noted a significant increase in the proportion

of cells that became Venus positive. Surprisingly, and unlike the requirement for IL-1β release,

this induction of caspase-1 BiFC did not require prior priming with LPS. When we transfected

macrophages with the caspase-4 prodomain (C4-Pro) or caspase-5 prodomain (C5-Pro) BiFC

pairs, we observed a similar pattern, where caspase-4 and caspase-5 BiFC were induced by

heme independent of LPS priming (Figure 2B,C). This suggests that caspase-1, caspase-4, and

caspase-5 are each activated by heme in macrophages. When we analyzed the appearance and

localization of the fluorescent complexes produced by heme in each case, we noticed some

differences (Figure 2D). The caspase-1 complex appeared as a single green punctum typical of

ASC specks and of ASC-induced caspase-1 BiFC that we previously reported (39, 40). Caspase-

4 and caspase-5 BiFC appeared as a series of punctate spots located throughout the cytoplasm

of the cell that did not co-localize with mitochondria. This indicates that there may be different

mechanisms of action for caspase-1 compared to caspase-4 or caspase-5.

Caspase-5 activation is impaired in M2 macrophages

Next, we used the BiFC system to measure inflammatory caspase activation in M1 and M2

macrophages. We polarized monocytes into M1 and M2 as before and transfected donor-matched

M1 and M2 cells with each of the inflammatory caspase BiFC pairs. Following exposure to heme,

caspase-1, caspase-4, and caspase-5 BiFC was increased as before in M1 hMDM (Figure 3A).

Interestingly, while caspase-1 and caspase-4 BiFC was induced to the same extent in M1 and

M2 macrophages, the M2 macrophages had impaired heme-induced caspase-5 BiFC. To explore

this further, we measured caspase-1, caspase-4, and caspase-5 expression in M1 and M2

macrophages by immunoblot (Figure 3B). Endogenous caspase-1 and caspase-4 were similarly

expressed in both subtypes and their expression was unchanged with the addition of heme.

Caspase-5 expression was readily detected in M1 macrophages, but was lowly expressed in

unstimulated M2 macrophages. Heme did not induce caspase-5 expression in the M2 subgroup,

but pre-treatment with LPS restored the level of caspase-5 to the level detected in M1 cells.

Heme activates caspase-4 and caspase-5 independently of inflammasomes

Intracellular LPS has been shown to bind and induce oligomerization of caspase-4, caspase-5

and caspase-11 independently of inflammasomes (21). The oligomerization is mediated by CARD

clustering, which leads to induced proximity and caspase dimerization. Since heme can promote

caspase-4 and caspase-5 induced proximity and is naturally taken up and recycled by

macrophages, we reasoned that heme may represent an intracellular trigger for non-canonical

inflammatory caspase activation (activation independent of inflammasomes). To test this, we

investigated if heme-induced caspase activation requires interactions with inflammasomes.

Recruitment of caspases to inflammasomes is dependent on an intact CARD. The aspartate

residue (D59) in caspase-1 is essential for the ASC-caspase-1 interaction and mutation of this

D59 residue blocks ASC-induced caspase-1 BiFC (39, 41). We modified the conserved CARD-

binding residue found in caspase-4 (D59) and caspase-5 (D117) and showed that disruption of

this residue also blocked caspase-4 and caspase-5 BiFC induced by NLRC4 (Supplemental

Figure S1A, B). Using the CARD-disrupting mutant in caspase-1 (D59R), caspase-5 (D117R),

and caspase-4 (D59R), we tested if heme-induced inflammatory caspase BiFC is inflammasome-

dependent. We transfected hMDM with the caspase-1, caspase-4 or caspase-5 Pro BiFC pairs

or with the corresponding CARD-disrupting mutant BiFC pairs. Following exposure to heme,

caspase-1 (D59R) BiFC was significantly decreased compared to the wild type caspase-1

reporter (Figure 4A). This indicates that heme triggers recruitment of caspase-1 to the

inflammasome for its subsequent activation. In contrast, heme-induced caspase-4 and capase-5

BiFC was not changed when the CARD mutant BiFC reporters were expressed (Figure 4A). This

suggests that heme activates caspase-4 and caspase-5 independently of inflammasomes.

To confirm these results, we used siRNA to silence the inflammasome adaptor protein ASC. ASC

is essential to the assembly of most characterized inflammasomes including the NLRP1, NLRP3

and AIM2 inflammasomes (1). It has been shown to be dispensable for the NLRC4

inflammasome, but ASC enhances NLRC4 inflammasome activity (42, 43). Therefore, if heme

induces inflammatory caspase activation via the NLRP3 or alternative inflammasome complex,

loss of ASC is predicted to block this activity. Si-RNA-mediated silencing of ASC in hMDM

reduced heme-induced caspase-1 BiFC to background levels (Figure 4B, Supplemental Figure

S1C). In contrast, ASC depletion had no effect on the levels of caspase-4 or caspase-5 BiFC

induced by heme (Figure 4B). Together, these results show that caspase-4 and caspase-5 are

activated by heme in an inflammasome independent manner.

Cytokine release and cell death are differentially regulated by caspase-4 and caspase-5

Having identified caspase-4 and caspase-5 induced proximity following heme exposure, we set

out to explore the functional outcomes of this mechanism. We used CRISPR/Cas9 to delete

caspase-4 and caspase-5 from the monocytic THP-1 cell line. We compared caspase-4 and

caspase-5-deficient cells to parental THP-1 cells and to caspase-1 deficient THP-1 cells

(purchased from Invivogen). The cells were first treated with PMA to differentiate them to

macrophage-like and to prime them for pro-IL-1β expression, and then exposed to heme or

additional pro-inflammatory stimuli (26). As expected, caspase-1 deficiency completely blocked

IL-1β release induced by heme and by nigericin, a bacterial potassium ionophore and potent

inducer of the NLRP3 inflammasome (Figure 5A) (44, 45). Surprisingly, LPS also induced

substantial IL-1β release in PMA-primed THP-1 cells that was caspase-1 dependent. The LPS

we used was ultrapure, so it lacks contaminants that could trigger inflammasome activation (46).

Therefore, we reasoned that the prior treatment of PMA facilitated delivery of LPS to the cell. A

similar phenomenon has been shown in murine macrophages, where cholera toxin B, through

binding to its receptor GM1 gangliosaccharide, promotes endocytosis of LPS (47). Consistent

with this, loss of caspase-4 reduced LPS-induced IL-1β release, but had no effect on nigericin-

induced IL-1β release. Thus, LPS treatment of PMA-primed THP-1 cells appears to represent a

non-canonical inflammasome stimulus. Loss of caspase-4 had no effect on heme-induced IL-1β

release. In contrast, deficiency of capase-5 resulted in a significant decrease in heme-induced IL-

1β release. Loss of capase-5 also potently blocked IL-1β release induced by nigericin and LPS.

These results show that caspase-5 is essential for IL-1β release by heme as well as canonical

and non-canonical inflammasome inducers.

Next, we investigated the requirement of each caspase for heme-induced cell death. Heme

treatment of THP-1 cells induced substantial cell death as measured by 7-AAD uptake. In contrast

to IL-1β release, priming with PMA or LPS was not required to induce cell death. This death was

not reduced in cells deficient in caspase-1 or caspase-5. However, loss of caspase-4 significantly

decreased heme-induced cell death (Figure 5B). This suggests that heme-induced caspase-4

activation functions to trigger cell death. Despite the reduction of heme-induced cell death,

caspase-4 loss did not completely block the death, suggesting that caspase-independent cell

death mechanisms also contribute to this process.

Caspase-1, caspase-4, and caspase-5 have been shown to cleave GSDMD, the proposed

mechanism by which inflammatory caspases induce both pyroptosis and release of mature

cytokines. Given that we noted a difference between the requirements of caspase-4 and caspase-

5 in heme-induced cell death and IL-1β release respectively, we next investigated GSDMD

cleavage in PMA-primed THP-1 cells. Surprisingly, we were not able to detect the N-terminal

GSDMD cleavage fragment in cells treated with heme alone (Figure 5C). This suggests that either

heme-induced IL-1β release is GSDMD-independent or that the amount of cleavage is below the

detection limits of this assay. We did detect the cleavage product when cells were treated with

LPS followed by heme. Loss of caspase-1 or caspase-4 had no effect on GSDMD cleavage. If

anything, we noted a minor increase in cleavage and were able to detect the cleaved fragment in

the LPS-treated lanes. In complete contrast, loss of caspase-5 blocked GSDMD cleavage. This

suggests that the caspase-4 dependent cell death induced by heme is mediated by an alternative

mechanism that is independent of GSDMD cleavage.

To investigate the kinetics of heme-induced caspase-1 activation relative to cell death, we used

time-lapse confocal microscopy. For this, we generated THP-1 cells stably expressing the

caspase-1 pro BiFC components. We designed a bicistronic construct, where the caspase-1 Pro-

VC and caspase-1 Pro-VN are expressed in a single vector separated by the viral 2A self-cleaving

peptide, similar to a caspase-2 reporter we previously described (48). This design ensures that

the caspase-1 BiFC components are expressed at equal levels, because they are translated from

a single mRNA transcript. These cells also expressed a linked mCherry gene as a reporter for

expression of the BiFC components. In addition, loss of mCherry fluorescence can be used to

detect cell lysis. The caspase-1 BiFC complex was detected approximately 18 h following the

addition of heme and its appearance was followed by lysis of the cell that was detected by loss of

the mCherry protein upon cell rupture (Figure 6A, Supplemental Movie S1). Cell death occurred

within 5 minutes of detection of the C1-Pro BiFC complex (Figure 6B). Thus, recruitment of

caspase-1 to inflammasomes in response to heme immediately precedes cell death. It is

important to note that this death occurred in the presence of the pan-caspase inhibitor qVD-OPH.

This, together with the results from Figure 5B suggests that there is a substantial caspase-

independent component to heme-induced cell death. However, the close timing of caspase-1

recruitment to inflammasomes and cell lysis indicates that these events are mechanistically

linked, which could imply that caspases contribute to the death in a manner that does not require

its catalytic activity.

Discussion

Our data show that heme activates caspase-1, caspase-4, and caspase-5. However, there are significant differences in the upstream requirements for activation, the localization of the activation complexes; and the outcomes of activation of each caspase. We found that caspase-5, but not caspase-4, is essential for heme-induced IL-1β release, while only caspase-4 contributes to heme-induced cell death. Our results indicate that caspase-4 and caspase-5 have non-overlapping functions in heme-induced inflammation and caspase-1 activation. Together, these data underscore the important functions of inflammatory caspases in heme-induced sterile inflammation.

While SCD is primarily a disease characterized by anemia, many of its clinical complications are exacerbated by chronic inflammation. Sickle shaped red blood cells are more prone to hemolysis releasing excess heme into the blood stream that overwhelms the body's heme scavenging systems. Consistent with this, we noted that heme-treated macrophages derived from SCD patients released more IL-1β when compared to healthy controls. This suggests that SCD macrophages are more sensitive to heme-induced inflammation. Because heme can activate TLR4, we originally suspected that this would be due to a lifetime exposure to heme in patients with SCD resulting from higher rates of hemolysis, which would prime the cells for inflammasome activation. However, heme alone was insufficient to trigger IL-1β release in macrophages from patients with SCD or from healthy donors indicating that an extra priming step is required. Monocytes from SCD patients have been shown to have increased levels of the macrophage markers CD14 and CD11b (16). In addition, liver macrophages from HbS sickle mice had increased surface expression of the M1 macrophage markers CD86, MHCII, iNOS, and IL-6 (13). This suggests that sickle macrophages are likely to skew to the M1-like pro-inflammatory phenotype. Treatment of HbS sickle mice with the heme scavenger hemopexin reverted the M1

polarization, indicating that heme is the inducer of M1 polarization (13). Thus, we propose that

the increased proportion of activated and pro-inflammatory M1 macrophages in patients with SCD

is the reason why the cells, once primed by LPS, are more sensitive to heme-induced IL-1β

release rather than the effect of heme priming. Interestingly, heme alone was sufficient to induce

caspase-1, caspase-4 and caspase-5 activation. Caspase-1 activation by heme has been

reported to be dependent on the NLRP3 inflammasome (6). Unlike the other inflammasome

receptors, NLRP3 requires a priming step for activation (49, 50). Our results may suggest that

heme can prime for NLRP3 expression, but does not provide enough of a priming signal to induce

sufficient pro-IL-1β expression. Induction of NLRP3 expression has been shown to be dependent

on reactive oxygen species (ROS) (8). Heme induces ROS and blocking ROS has been shown

to inhibit both caspase-1 cleavage and IL-1β processing. Therefore, heme-induced ROS

generation may be a mechanism by which heme can prime for NLRP3 expression but not for pro-

IL-1β expression.

Our observation that heme-induced caspase-4 and caspase-5 induced proximity does not require

LPS priming could be explained in one of two ways. Similar to caspase-1, heme may be sufficient

to prime the cells for caspase activation. However, we also show that, in contrast to caspase-1,

caspase-4 and caspase-5 induced proximity is activated by heme independently of CARD-

inflammasome interactions and of the common inflammasome adaptor protein ASC. Caspase-5

is lowly expressed in many cell types and its expression is induced by LPS and interferon (51).

Despite this, primary M1-polarized macrophages express abundant caspase-5. M2 macrophages

have reduced caspase-5 expression and display reduced caspase-5 induced proximity in

response to heme. This observation may suggest that full length, endogenous caspase-5 is

required to promote assembly of the caspase-5 signaling platform. This is consistent with studies

that have shown that caspase-4 and caspase-5 are direct intracellular receptors for LPS (21).

LPS binds to the CARD in these caspases to trigger oligomerization of caspase-4 or caspase-5

in the absence of any known inflammasome protein. Our data indicate that heme acts in an

analogous fashion to induce caspase-4 or caspase-5 oligomerization either directly or through a

cytosolic mediator. Further supporting a mechanistic difference between heme-induced caspase-

1 activation and heme-induced caspase-4 or caspase-5 activation, we observed distinct

differences in the size and localization of the caspase-4 and caspase-5 signaling complexes

induced by heme compared to the single ASC-like speck observed for caspase-1. Heme is

naturally taken up and recycled by macrophages (44), representing a physiological stimulus and,

importantly, a trigger of sterile inflammation that directly engages caspase-4 and caspase-5.

Thus, heme acts as a canonical stimulus by engaging inflammasome-dependent caspase-1

activation and also as a non-canonical stimulus by activating caspase-4 and caspase-5

independent of inflammasomes.

Caspase-4 and caspase-5 are often assumed to be redundant proteins that directly phenocopy

caspase-11 in mice. Studies using a transgenic mouse generated to express human caspase-4

indicated that caspase-4 does not completely phenocopy caspase-11 and a recent study showed

a broader reactivity of caspase-4 to LPS compared to caspase-11 (52, 53). In addition, humans

are more sensitive to endotoxemia than rodents (40). These studies highlight the important

differences between the human and murine innate immune response and suggest that the

presence of an additional inflammatory caspase in humans may contribute to these differences.

We show that caspase-5 is not only essential for heme-induced IL-1β release but for IL-1β release

induced by canonical and non-canonical stimuli. In contrast, caspase-4 is not required for heme

induced IL-1β release. Although loss of caspase-5 reduced GSDMD cleavage, it did not impact

cell death. Caspase-4 was the only inflammatory caspase to contribute to heme-induced cell

death. These results suggest that additional death mechanisms are engaged by heme. Heme has

been shown to induce RIPK3-dependent necrosis (54) and, therefore, this may be a contributing

mechanism to the death we observed. RIPK3-dependent necrosis has been shown to limit

pathogenic inflammation and associated tissue damage without impairing IL-1β release (55).

Given that cell death occurs immediately following detection of the caspase-1 activation platform,

it is possible that a similar mechanism occurs here. Whether caspase-4 actively contributes to the

alternative cell death pathway or if there are indeed multiple independent cell death pathways

engaged by heme is yet to be determined. Nevertheless, the association between caspase-11,

excess pyroptosis, and inflammation-associated tissue damage is well established (56, 57). In a

mouse model of SCD, the consequences of heme-induced inflammation is tissue damage that

manifests as vaso-occlusion and death (7). Our results suggest that caspase-4 and not caspase-

1 or caspase-5 contribute to heme-induced pyroptosis. Therefore, in humans caspase-4 may be

an effector of heme-induced tissue damage in SCD while caspase-5 is an essential regulator of

heme-induced inflammation. Our model predicts that blocking caspase-4 would protect from

tissue damage, while blocking caspase-5 would protect from uncontrolled inflammation (fever,

pain, etc.). Further exploration of the distinct roles of these two caspases is required to fully

understand how chronic inflammation can be controlled in hemolytic disorders.

Materials and Methods

Chemicals and antibodies

The following antibodies were used: anti-Caspase-1 (D7F10 from Cell Signaling Technology, Danvers, MA, USA), anti-Caspase-4 (4450 from Cell Signaling Technology, Danvers, MA, USA), anti-Caspase-5 (D3G4W from Cell Signaling Technology, Danvers, MA, USA); anti-actin (C4 from MP Biomedicals); anti-GSDMD (G7422 from Sigma-Aldrich, St. Louis, MO, USA), anti-GSDMD N-term (E7H9G from Cell Signaling Technology, Danvers, MA, USA), anti-IL-1β (MAB601, R&D Systems, Inc., Minneapolis, MN, USA). All cell culture media reagents were purchased from Thermo Fisher (Waltham, MA, USA). Ultrapure LPS (from E. coli O111:B4) was purchased from Invivogen (San Diego, CA, USA). Unless otherwise indicated, all other reagents were purchased from Sigma-Aldrich (St. Louis, MO, USA).

Plasmids

The pBiFC.VC155 and pBiFC.VN173 plasmids encoding C1-Pro, C4-Pro, or C5-Pro were described previously (39). Single mutations were introduced using QuikChange Site-Directed Mutagenesis Kit (Agilent Technologies, Santa Clara, CA, USA). The bicistronic vector consists of C1-Pro VC and C1-Pro VN linked with a 2A peptide. Silent mutations were introduced into the second C1-Pro nucleotide sequence to prevent the sequence recombining out during cloning due to the presence of two identical C1 Pro sequences. The sequence was generated by IDT (Coralville, IA, USA) and cloned into pRRL-MND-MCS-2A-mCherry-2A-Puro. pDsRed-mito was purchased from Clontech (Takara). Each construct was verified by sequencing.

Preparation and culture of primary human monocyte-derived macrophages

Whole blood samples were obtained from patients with SCD who attend our hematology clinic as

part of their routine care. The study was approved by the Baylor College of Medicine Institutional

Review Board, and informed consent was obtained from all participants, or legal guardian (if

participants were minor). To isolate peripheral blood mononuclear cells, whole blood obtained

from healthy blood donors or SCD patients was separated using the Ficoll Paque (GE Healthcare)

gradient protocol (58). CD14+ monocytes were isolated from PBMCs using magnetic bead

selection (Miltenyi Biotec). To differentiate cells into macrophages, monocytes were seeded at

5×10^6 -1×10^7/10 cm dish in RPMI-1640 medium supplemented with FCS (10% (v/v)), glutamax (2

mM), and Penicillin/ Streptomycin (50 I.U./50 µg/ml) and GM-CSF (50 ng/mL). Cells were allowed

to adhere overnight and culture media was exchanged the following day for fresh GM-CSF-

supplemented media. Media was exchanged every 2-3 days and were considered fully

differentiated at 7 days, as determined by morphology. To differentiate into M2 macrophages,

media was exchanged and cells were incubated for additional 24 hours in media supplemented

with IL-4 (50 ng/mL). For the matched M1 macrophages, cells were incubated in fresh GM-CSF

supplemented media for an additional day.

Heme preparation and administration

Heme solution was prepared immediately before use by solubilizing 3.3 mg of porcine hemin,

(oxidized version of heme) in 100 µL of NaOH solution (0.1-1.0 M). The mixture was vortexed for

5 min in the dark to allow reduction of hemin to heme. 900 µL of serum-free RPMI-1640 media

was added to the resulting solution and vortexed for an additional 5 min in the dark followed by

filtration through a sterile 0.22 µm spin filter. The term "heme" is used generically to refer to both

heme and "hemin." Unless otherwise indicated, cells were treated with heme in the presence of

0.1% FBS in the culture media to prevent FBS sequestering and inhibiting heme. Heme is rapidly

recycled by circulating macrophages, therefore to mimic the transient exposure of cells to heme

and to limit its toxicity, where indicated, cells were exposed to heme for 1 h followed by addition

of an equal volume of culture media containing 10% FBS to inactivate the heme.

Cell culture and generation of cell lines

THP-1 cells were grown in RPMI medium containing FCS (10% (v/v)), glutamax (2 mM), and

Penicillin/ Streptomycin (50 I.U./50 µg/ml). MCF-7 cells were grown in Dulbecco's Modified

Essential Medium (DMEM) containing FCS (10% (v/v)), L-glutamine (2 mM), and Penicillin/

Streptomycin (50 I.U./50 µg/ml). *CASP1*-deficient THP-1 cells were purchased from Invivogen

(San Diego, CA, USA). Caspase-4 and caspase-5 were deleted from THP-1 cells using an

adaptation of the CRISPR/Cas9 protocol described in (59). Protospacer sequences for each

target gene were identified using the CRISPRscan scoring algorithm (www.crisprscan.org (60)).

DNA templates for sgRNAs were made by PCR using pX459 plasmid containing the sgRNA

scaffold sequence and using the following primers:

∆CASP4(40)   sequence:

TTAATACGACTCACTATAGGGAAACAACCGCACACGCCgttttagagctagaaatagc;

∆CASP5(46)   sequence:

TTAATACGACTCACTATAGGTCCTGGAGAGACCGCACAgttttagagctagaaatagc;

∆CASP5(53)   sequence:

TTAATACGACTCACTATAGGTCAAGGTTGCTCGTTCTAgttttagagctagaaatagc ; universal

reverse primer: AGCACCGACTCGGTGCCACT. sgRNAs were generated by *in vitro* transcription

using the Hiscribe T7 high yield RNA synthesis kit (New England Biolabs). Purified sgRNA (0.5

µg) was incubated with Cas9 protein (1 µg, PNA Bio) for 10 min at room temperature. THP-1 cells

were electroporated with the sgRNA/Cas9 complex using the Neon transfection system (Thermo

Fisher Scientific) at 1600V, 10ms, and 3 pulses. For the caspase-5 deficient cell line two sgRNA

were selected at either end of the gene to delete the intervening region. Deletion of *CASP4* or

CASP5 was confirmed by western blot or by PCR. Single cell clones were generated by single

cell plating of the parental cell line. Gene deletion in the single cell clones was confirmed by

sequencing.

Transient Transfection and siRNA

For transfection of hMDMs, 1x 10^5 cells were transfected with the appropriate plasmid

combinations or plasmid/siRNA combinations using the Neon Transfection System (Thermo

Fisher) and a 10 µL Neon tip at 1000 V, 40 ms, and 2 pulses. Cells were transfected with amounts

of the relevant expression plasmids as described in the figure legends. A total of 4 wells were

transfected for every plasmid and incubated in 200 µL of antibiotic-free media. After 1 h, 200 µL

of complete growth media containing Penicillin/ Streptomycin (50 I.U./50 µg/ml) was added.

Expression was allowed for 24 h and media was exchanged for fresh media prior to treatment.

Control siRNAs were siCyclophilin B (ON-TARGETplus SMARTpool from GE Dharmacon). ASC

siRNAs were ON-TARGETplus ASC SMARTpool (GE Dharmacon). 1 x 10^5 MCF-7 cells were

transfected with appropriate plasmid combinations using Lipofectamine 2000 transfection reagent

(Invitrogen) according to manufacturer's instructions.

Microscopy

Cells were imaged using a spinning disk confocal microscope (Zeiss, Thornwood, NY, USA),

equipped with a CSU-X1A 5000 spinning disk unit (Yokogowa Electric Corporation, Japan), multi

laser module with wavelengths of 458 nm, 488 nm, 514 nm, and 561 nm, and an Axio Observer

Z1 motorized inverted microscope equipped with a precision motorized XY stage (Carl Zeiss

MicroImaging, Thornwood, NY, USA). Images were acquired with a Zeiss Plan-Neofluar 40x 1.3

NA or 64x 1.4 NA objective on an Orca R2 CCD camera using Zen 2012 software (Zeiss,

Thornwood, NY, USA). Cells were plated on dishes containing coverslips (Mattek Corp. Ashland,

MA, USA) coated with poly-D-lysine hydrobromide 24 h prior to treatment. For time-lapse

experiments media on the cells was supplemented with Hepes (20 mM) and 2-mercaptoethanol

(55 μM). Cells were allowed to equilibrate to 37°C in 5% CO_2 prior to focusing on the cells.

Image analysis

Images were analyzed by drawing regions around individual cells and then computing average

intensity the pixels for each fluor using Zen 2012 software (Zeiss, Thornwood, NY, USA). Data

was scaled by the following formula: scaled point = (Max − x)/MaxDifference, where Max equals

the maximum value in the series, x equals the point of interest, and MaxDifference equals the

maximum minus the minimum value in the series.

ELISA IL-1β Measurements

THP-1 cells were plated at 1x10^6 cells/mL in the presence of PMA (phorbol 12-myristate 13-

acetate, 10 ng/mL) for three days to prime the cells and to induce them to acquire a macrophage-

like phenotype. On the third day, supernatants were removed and replaced for fresh media

without PMA and cells were allowed to rest for an additional 24 h. Cells were washed prior to

treatment as indicated in the Figure Legends. IL-1β concentration in harvested clarified

supernatants was measured with the IL-1β Duoset ELISA Kit (R&D Systems, Minneapolis, MN,

USA) according to the manufacturer's instructions.

Flow cytometry and measurement of cell death

Cells were treated as indicated and collected by centrifugation. Cells were washed with PBS and

resuspended in 100 µl of PBS supplemented with 5% FCS, 0.5% BSA, 2 mM EDTA and 1 µl of

7-AAD (Thermo Fisher Rockford, IL, USA). 7-AAD cells were quantitated by flow cytometry and

analyzed with FloJo software (FloJo LLC., Ashland, OR, USA).

Immunoblotting

Cells were treated as indicated. Cells were lysed in IP-lysis buffer (50 mM Tris pH 7.4, 150 mM

NaCl, 0.1% SDS, 1% NP-40 containing protease inhibitors (Complete Mini Protease Inhibitor

Cocktail, Roche). Protein concentration was determined by BCA assay (Pierce). 25-35 µg total

protein was resolved by SDS-PAGE and transferred onto 0.45 µm nitrocellulose membrane

(Thermo Scientific, Rockford, IL, USA), and immunodetected using appropriate primary and

peroxidase-coupled secondary antibodies (GE Healthcare Life Sciences, Pittsburg, PA, USA).

Proteins were visualized by West Pico and West Dura chemiluminescence Substrate (Thermo

Fisher, Rockford, IL, USA).

Statistical Analysis

Statistical comparisons were performed using two-tailed Student's t test or 1-way ANOVA

calculated using Prism 6.0 (Graph Pad) software.

Acknowledgements

Funding for the project includes NIH/NIDDK T32DK060445 (BB), NIH/NIDDK F32DK121479 (BB), NIH/NIGMS R01GM121389 (LBH), NIH/NHLBI R01HL114567 (JDB, GMV), NIH/NHLBI R01-HL136415 (JMF). This project was supported by the Cytometry and Cell Sorting Core at Baylor College of Medicine with funding from the CPRIT Core Facility Support Award (CPRIT-RP180672), the NIH (CA125123 and RR024574) and the assistance of Joel M. Sederstrom

Figures

Figure 1: Heme induces IL-1β release that is increased in SCD patients. **(A)** CD14+ monocytes were isolated from 5 healthy donors and differentiated into macrophages using GM-CSF for 7 days. When fully mature, cells were primed with or without LPS (100 ng/mL) for 3 h, washed and treated with or without heme (50 μM) in 0.1% FBS. Mature IL-1β levels were measured in cellular supernatants by ELISA at the indicated times. Error bars represent standard deviation of 5 independent experiments. ***p<0.001, ****p<0.0001 calculated by 1-way ANOVA. **(B)** GM-CSF-differentiated human macrophages from healthy donors were primed with or without LPS for 3 h (100 ng/mL), followed by treatment with or without heme (50 μM) in 0.1, 1, 5, or 10%

FBS. After 20 h, IL-1β concentration was measured in cultured supernatants by ELISA. Error bars represent standard deviation of 4 independent biological replicates. **(C)** GMCSF-differentiated human macrophages were isolated from healthy donors (control) or sickle cell disease (SCD) patients and primed with or without LPS (100 ng/mL) for 3 h followed by treatment with or without heme (50 μM) in 0.1% FBS. IL-1β concentration was measured in cultured supernatants at 20 h by ELISA. Error bars represent standard deviation of 4 control and 4 SCD samples across 4 independent experiments. **p<0.01 calculated by 1-way ANOVA. **(D)** CD14+ monocytes were isolated from 3-5 healthy donors and differentiated in GM-CSF for 7 days. On day 7, macrophages were polarized either into the M2-like phenotype with IL-4 (50 ng/mL) or into the M1-like phenotype with GM-CSF for one additional day. On day 8, both groups were washed and primed with LPS (100 ng/mL) for 3 h followed by treatment with or without heme (50 μM) in 0.1% FBS or ATP (5 mM) in 10% FBS. IL-1β concentration was measured in cultured supernatants at 20 h by ELISA. Error bars represent standard deviation of 3-5 independent experiments. *p<0.05 calculated by Student's t test. **(E)** Representative phase-contrast images of M1 and M2 macrophages from (D) are shown. Scale bar represents 100 μm.

Figure 2: Heme activates the inflammatory caspases. GMCSF-differentiated human macrophages isolated from healthy donors were transfected with the C1-Pro VC (300 ng) and C1-Pro VN (300 ng) **(A)**; C4-Pro VC (500 ng) and of C4-Pro VN (500 ng) **(B)**; or C5-Pro VC (1000 ng) and C5-Pro VN (1000 ng) **(C)**; along with dsRedmito (50 ng) as a reporter for transfection. 24 h after transfection, cells were treated with or without LPS (100 ng/mL) for 3 h followed by treatment with or without heme (50 μM) in 0.1% FBS. After 1 h, FBS was reconstituted to 5% to inhibit extracellular heme. Cells were assessed for the percentage of dsRed-positive transfected cells that were Venus-positive at 20 h, determined from a minimum of 300 cells per well. Results are represented as percent Venus-positive cells over background (untreated cells). Error bars represent standard deviation of four independent experiments. *p< 0.05; **p< 0.01; calculated by

Student's t test. **(D)** Representative images show caspase BiFC in green and mitochondria in red.

Scale bar represents 10 µm.

Figure 3: Heme-induced caspase-5 activation and expression is reduced in M2 macrophages. (A) GM-CSF-differentiated human macrophages isolated from healthy donors were polarized into M1 or M2 macrophages and transfected with the C1-Pro BiFC pair (300 ng of each), the C4-Pro BiFC pair (500 ng of each), or the C5-Pro BiFC pair (1000 ng of each) along with dsRedmito (50 ng) as a reporter for transfection. 24 h after transfection, cells were treated with or without heme (50 µM) in 0.1% FBS. After 1 h, FBS was reconstituted to 5% to inhibit extracellular heme. Cells were assessed for the percentage of dsRed-positive transfected cells that were Venus-positive at 20 h, determined from a minimum of 300 cells per well. Error bars represent standard deviation of three independent experiments. ***p=0.001; calculated by Student's t test. **(B)** Cells were polarized to M1 or M2 macrophages as in (A) and treated with or

without LPS (100 ng/mL) for 3 h followed by heme (50 µM) in 0.1% FBS. After 1 h, FBS was reconstituted to 5% to inhibit extracellular heme and 20 h later cell lysates were immunoblotted for caspase-1, caspase-4, caspase-5, or actin as a loading control.

A

B

Figure 4: Heme activates caspase-4 and caspase-5 independently of inflammasomes. (A) GM-CSF-differentiated human macrophages isolated from healthy donors were transfected with the C1-Pro BiFC pair (300 ng of each), the D59R mutant C1-Pro BiFC pair (300 ng of each), the C4-Pro BiFC pair (500 ng of each), the D59R mutant C4-Pro BiFC pair (500 ng of each), the C5-Pro BiFC pair (1000 ng of each), or the D117R mutant C5-Pro BiFC pair (1000 ng of each) along with dsRedmito (50 ng) as a reporter for transfection. 24 h after transfection, cells were treated with or without heme (50 µM) in 0.1% FBS. After 1 h, FBS was reconstituted to 5% to inhibit extracellular heme. Cells were assessed for the percentage of dsRed-positive transfected cells

that were Venus-positive at 20 h, determined from a minimum of 300 cells per well. Error bars represent standard deviation of four independent experiments. *p=0.05; calculated by Student's t test. **(B)** GM-CSF-differentiated human macrophages isolated from healthy donors were transfected with the C1-Pro BiFC pair (300 ng of each), the C4-Pro BiFC pair (500 ng of each), or the C5-Pro BiFC pair (1000 ng of each), along with dsRedmito (50 ng) as a reporter for transfection with either siRNA against ASC or a control siRNA (25 nM). 24 h after transfection, cells were treated with or without heme (50 μM) in 0.1% FBS. After 1 h, FBS was reconstituted to 5% to inhibit extracellular heme. Cells were assessed for the percentage of dsRed-positive transfected cells that were Venus-positive at 20 h, determined from a minimum of 300 cells per well. Error bars represent standard deviation of three independent experiments. **p<0.01.

Figure 5: Cytokine release and cell death are differentially regulated by caspase-4 and caspase-5. (A) THP-1 cells (WT), THP-1 cells deficient in caspase-1 (ΔC1, generated by shRNA), caspase-4 (ΔC4, generated by CRISPR/Cas9), or caspase-5 (ΔC5, generated by CRISPR/Cas9) were primed with PMA (10 ng/mL) for three days and allowed to recover for one day. Cells were treated with a 1 h heme pulse (50 μM), nigericin (5 μM) or LPS (100 ng/mL). IL-1β concentration

was measured in cultured supernatants by ELISA at 20 h. Error bars represent standard deviation of 5-14 independent experiments. *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001; calculated by Student's t test. **(B)** THP-1 cells or THP-1 cells deficient in the indicated caspases were treated with or without heme (50 μM) for a 1 h pulse. Cell death was assessed by flow cytometry for 7-AAD uptake 20 h later. Error bars represent standard deviation of 3 independent experiments. **p=0.01; calculated by Student's t test. **(C)** PMA-primed THP-1 cells or THP-1 cells deficient in the indicated caspases were treated with or without LPS (100 ng/mL) for 3 h followed by treatment with or without heme (50 μM) for a 1 h pulse. 20 h later, cell lysates were immunoblotted for the indicated proteins or actin as a loading control. Results are representative of two independent experiments.

Figure 6. Heme-induced inflammasome assembly immediately precedes cell death. (A) PMA-primed THP-1 cells stably expressing the C1-Pro BiFC pair were treated with heme in the presence of qVD-OPH (5 μM) and images were taken by confocal microscopy every 5 min for 24 h. Frames from the time-lapse show representative cells undergoing BiFC (green) prior to cell lysis as measured by the loss mCherry (red). Scale bars represent 5 μm. **(B)** Graph of the cells from (A) that became Venus-positive is shown. Each point on the mCherry graph (*red)* is scaled and aligned to each point on the caspase-1 BiFC graph (*green*) that represents the average intensity of mCherry or Venus in the cell at 5 min intervals where time=0 is the point of onset of

mCherry loss representing cell lysis. Arrow shows the point of onset of caspase-1 BiFC immediately prior to cell lysis. Error bars represent SEM of 9 individual cells.

References

1. Bolivar, B. E., T. P. Vogel, and L. Bouchier-Hayes. 2019. Inflammatory caspase regulation: maintaining balance between inflammation and cell death in health and disease. *FEBS J* 286: 2628-2644.
2. van de Veerdonk, F. L., M. G. Netea, C. A. Dinarello, and L. A. Joosten. 2011. Inflammasome activation and IL-1beta and IL-18 processing during infection. *Trends Immunol* 32: 110-116.
3. Rock, K. L., E. Latz, F. Ontiveros, and H. Kono. 2010. The sterile inflammatory response. *Annual review of immunology* 28: 321-342.
4. Martin, S. J. 2016. Cell death and inflammation: the case for IL-1 family cytokines as the canonical DAMPs of the immune system. *FEBS J* 283: 2599-2615.
5. Wagener, F. A., A. Eggert, O. C. Boerman, W. J. Oyen, A. Verhofstad, N. G. Abraham, G. Adema, Y. van Kooyk, T. de Witte, and C. G. Figdor. 2001. Heme is a potent inducer of inflammation in mice and is counteracted by heme oxygenase. *Blood* 98: 1802-1811.
6. Dutra, F. F., L. S. Alves, D. Rodrigues, P. L. Fernandez, R. B. de Oliveira, D. T. Golenbock, D. S. Zamboni, and M. T. Bozza. 2014. Hemolysis-induced lethality involves inflammasome activation by heme. *Proc Natl Acad Sci U S A* 111: E4110-4118.
7. Belcher, J. D., C. Chen, J. Nguyen, L. Milbauer, F. Abdulla, A. I. Alayash, A. Smith, K. A. Nath, R. P. Hebbel, and G. M. Vercellotti. 2014. Heme triggers TLR4 signaling leading to endothelial cell activation and vaso-occlusion in murine sickle cell disease. *Blood* 123: 377-390.
8. Bauernfeind, F., E. Bartok, A. Rieger, L. Franchi, G. Nunez, and V. Hornung. 2011. Cutting edge: reactive oxygen species inhibitors block priming, but not activation, of the NLRP3 inflammasome. *J Immunol* 187: 613-617.
9. Hematology, A. S. o. Sickle Cell Disease.
10. Prevention, C. f. D. C. a. 2017. Sickle Cell Disease.
11. Wankhade, V., R. B. Andhale, and S. Lodhan. 2013. Prediction of Physiological Status of Sickle Cell Anemic Patients by Quantitative Observations of Microscopic Components of Urine. *Trends in Medical Research* 8: 27-31.
12. Al-Jafar, H. 2017. Sickle Cell Disease Clinical Classification. *Blood* 130: 4771.
13. Vinchi, F., M. Costa da Silva, G. Ingoglia, S. Petrillo, N. Brinkman, A. Zuercher, A. Cerwenka, E. Tolosano, and M. U. Muckenthaler. 2016. Hemopexin therapy reverts heme-induced proinflammatory phenotypic switching of macrophages in a mouse model of sickle cell disease. *Blood* 127: 473-486.
14. Vinchi, F., L. De Franceschi, A. Ghigo, T. Townes, J. Cimino, L. Silengo, E. Hirsch, F. Altruda, and E. Tolosano. 2013. Hemopexin therapy improves cardiovascular function by preventing heme-induced endothelial toxicity in mouse models of hemolytic diseases. *Circulation* 127: 1317-1329.
15. Ghosh, S., O. A. Adisa, P. Chappa, F. Tan, K. A. Jackson, D. R. Archer, and S. F. Ofori-Acquah. 2013. Extracellular hemin crisis triggers acute chest syndrome in sickle mice. *The Journal of clinical investigation* 123: 4809-4820.
16. Belcher, J. D., P. H. Marker, J. P. Weber, R. P. Hebbel, and G. M. Vercellotti. 2000. Activated monocytes in sickle cell disease: potential role in the activation of vascular endothelium and vaso-occlusion. *Blood* 96: 2451-2459.
17. Safaya, S., M. H. Steinberg, and E. S. Klings. 2012. Monocytes from sickle cell disease patients induce differential pulmonary endothelial gene expression via activation of NF-kappaB signaling pathway. *Molecular immunology* 50: 117-123.
18. Carvalho, M. O. S., T. Araujo-Santos, J. H. O. Reis, L. C. Rocha, B. A. V. Cerqueira, N. F. Luz, I. M. Lyra, V. M. Lopes, C. G. Barbosa, L. M. Fiuza, R. P. Santiago, C. V. B.

Figueiredo, C. C. da Guarda, M. Barral Neto, V. M. Borges, and M. S. Goncalves. 2018. Inflammatory mediators in sickle cell anaemia highlight the difference between steady state and crisis in paediatric patients. *British journal of haematology* 182: 933-936.

19. Kayagaki, N., I. B. Stowe, B. L. Lee, K. O'Rourke, K. Anderson, S. Warming, T. Cuellar, B. Haley, M. Roose-Girma, Q. T. Phung, P. S. Liu, J. R. Lill, H. Li, J. Wu, S. Kummerfeld, J. Zhang, W. P. Lee, S. J. Snipas, G. S. Salvesen, L. X. Morris, L. Fitzgerald, Y. Zhang, E. M. Bertram, C. C. Goodnow, and V. M. Dixit. 2015. Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. *Nature* 526: 666-671.

20. Shi, J., Y. Zhao, K. Wang, X. Shi, Y. Wang, H. Huang, Y. Zhuang, T. Cai, F. Wang, and F. Shao. 2015. Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. *Nature* 526: 660-665.

21. Shi, J., Y. Zhao, Y. Wang, W. Gao, J. Ding, P. Li, L. Hu, and F. Shao. 2014. Inflammatory caspases are innate immune receptors for intracellular LPS. *Nature* 514: 187-192.

22. Ruan, J., S. Xia, X. Liu, J. Lieberman, and H. Wu. 2018. Cryo-EM structure of the gasdermin A3 membrane pore. *Nature* 557: 62-67.

23. Sborgi, L., S. Ruhl, E. Mulvihill, J. Pipercevic, R. Heilig, H. Stahlberg, C. J. Farady, D. J. Muller, P. Broz, and S. Hiller. 2016. GSDMD membrane pore formation constitutes the mechanism of pyroptotic cell death. *EMBO J* 35: 1766-1778.

24. Liu, X., Z. Zhang, J. Ruan, Y. Pan, V. G. Magupalli, H. Wu, and J. Lieberman. 2016. Inflammasome-activated gasdermin D causes pyroptosis by forming membrane pores. *Nature* 535: 153-158.

25. Aglietti, R. A., A. Estevez, A. Gupta, M. G. Ramirez, P. S. Liu, N. Kayagaki, C. Ciferri, V. M. Dixit, and E. C. Dueber. 2016. GsdmD p30 elicited by caspase-11 during pyroptosis forms pores in membranes. *Proc Natl Acad Sci U S A* 113: 7858-7863.

26. Cullen, S. P., C. J. Kearney, D. M. Clancy, and S. J. Martin. 2015. Diverse Activators of the NLRP3 Inflammasome Promote IL-1beta Secretion by Triggering Necrosis. *Cell Rep* 11: 1535-1548.

27. Conos, S. A., K. E. Lawlor, D. L. Vaux, J. E. Vince, and L. M. Lindqvist. 2016. Cell death is not essential for caspase-1-mediated interleukin-1beta activation and secretion. *Cell Death Differ* 23: 1827-1838.

28. Evavold, C. L., J. Ruan, Y. Tan, S. Xia, H. Wu, and J. C. Kagan. 2018. The Pore-Forming Protein Gasdermin D Regulates Interleukin-1 Secretion from Living Macrophages. *Immunity* 48: 35-44 e36.

29. Bunn, H. F., and J. H. Jandl. 1968. Exchange of heme among hemoglobins and between hemoglobin and albumin. *The Journal of biological chemistry* 243: 465-475.

30. Balla, J., H. S. Jacob, G. Balla, K. Nath, J. W. Eaton, and G. M. Vercellotti. 1993. Endothelial-cell heme uptake from heme proteins: induction of sensitization and desensitization to oxidant damage. *Proceedings of the National Academy of Sciences* 90: 9285.

31. Adams, P. A., and M. C. Berman. 1980. Kinetics and mechanism of the interaction between human serum albumin and monomeric haemin. *Biochem J* 191: 95-102.

32. Kanno, T., K. Yasutake, K. Tanaka, S. Hadano, and J. E. Ikeda. 2017. A novel function of N-linked glycoproteins, alpha-2-HS-glycoprotein and hemopexin: Implications for small molecule compound-mediated neuroprotection. *PLoS One* 12: e0186227.

33. Vinchi, F., M. Costa da Silva, G. Ingoglia, S. Petrillo, N. Brinkman, A. Zuercher, A. Cerwenka, E. Tolosano, and M. U. Muckenthaler. 2016. Hemopexin therapy reverts heme-induced proinflammatory phenotypic switching of macrophages in a mouse model of sickle cell disease. *Blood* 127: 473-486.

34. Aaronson, D. S., and C. M. Horvath. 2002. A road map for those who don't know JAK-STAT. *Science* 296: 1653-1655.

35. Poyet, J. L., S. M. Srinivasula, M. Tnani, M. Razmara, T. Fernandes-Alnemri, and E. S. Alnemri. 2001. Identification of Ipaf, a human caspase-1-activating protein related to Apaf-1. *J Biol Chem* 276: 28309-28313.

36. Srinivasula, S. M., J. L. Poyet, M. Razmara, P. Datta, Z. Zhang, and E. S. Alnemri. 2002. The PYRIN-CARD protein ASC is an activating adaptor for caspase-1. *J Biol Chem* 277: 21119-21122.

37. Bouchier-Hayes, L., A. Oberst, G. P. McStay, S. Connell, S. W. Tait, C. P. Dillon, J. M. Flanagan, H. M. Beere, and D. R. Green. 2009. Characterization of cytoplasmic caspase-2 activation by induced proximity. *Mol Cell* 35: 830-840.

38. Shyu, Y. J., H. Liu, X. Deng, and C. D. Hu. 2006. Identification of new fluorescent protein fragments for bimolecular fluorescence complementation analysis under physiological conditions. *Biotechniques* 40: 61-66.

39. Sanders, M. G., M. J. Parsons, A. G. Howard, J. Liu, S. R. Fassio, J. A. Martinez, and L. Bouchier-Hayes. 2015. Single-cell imaging of inflammatory caspase dimerization reveals differential recruitment to inflammasomes. *Cell Death Dis* 6: e1813.

40. Masumoto, J., S. Taniguchi, K. Ayukawa, H. Sarvotham, T. Kishino, N. Niikawa, E. Hidaka, T. Katsuyama, T. Higuchi, and J. Sagara. 1999. ASC, a novel 22-kDa protein, aggregates during apoptosis of human promyelocytic leukemia HL-60 cells. *J Biol Chem* 274: 33835-33838.

41. Proell, M., M. Gerlic, P. D. Mace, J. C. Reed, and S. J. Riedl. 2013. The CARD plays a critical role in ASC foci formation and inflammasome signalling. *Biochem J* 449: 613-621.

42. Broz, P., J. von Moltke, J. W. Jones, R. E. Vance, and D. M. Monack. 2010. Differential requirement for Caspase-1 autoproteolysis in pathogen-induced cell death and cytokine processing. *Cell host & microbe* 8: 471-483.

43. Mariathasan, S., K. Newton, D. M. Monack, D. Vucic, D. M. French, W. P. Lee, M. Roose-Girma, S. Erickson, and V. M. Dixit. 2004. Differential activation of the inflammasome by caspase-1 adaptors ASC and Ipaf. *Nature* 430: 213-218.

44. Pelegrin, P., and A. Surprenant. 2006. Pannexin-1 mediates large pore formation and interleukin-1beta release by the ATP-gated P2X7 receptor. *EMBO J* 25: 5071-5082.

45. Perregaux, D., J. Barberia, A. J. Lanzetti, K. F. Geoghegan, T. J. Carty, and C. A. Gabel. 1992. IL-1 beta maturation: evidence that mature cytokine formation can be induced specifically by nigericin. *J Immunol* 149: 1294-1303.

46. Faustin, B., L. Lartigue, J. M. Bruey, F. Luciano, E. Sergienko, B. Bailly-Maitre, N. Volkmann, D. Hanein, I. Rouiller, and J. C. Reed. 2007. Reconstituted NALP1 inflammasome reveals two-step mechanism of caspase-1 activation. *Mol Cell* 25: 713-724.

47. Kayagaki, N., M. T. Wong, I. B. Stowe, S. R. Ramani, L. C. Gonzalez, S. Akashi-Takamura, K. Miyake, J. Zhang, W. P. Lee, A. Muszynski, L. S. Forsberg, R. W. Carlson, and V. M. Dixit. 2013. Noncanonical inflammasome activation by intracellular LPS independent of TLR4. *Science* 341: 1246-1249.

48. Ando, K., M. J. Parsons, R. B. Shah, C. I. Charendoff, S. L. Paris, P. H. Liu, S. R. Fassio, B. A. Rohrman, R. Thompson, A. Oberst, S. Sidi, and L. Bouchier-Hayes. 2017. NPM1 directs PIDDosome-dependent caspase-2 activation in the nucleolus. *J Cell Biol* 216: 1795-1810.

49. Bauernfeind, F. G., G. Horvath, A. Stutz, E. S. Alnemri, K. MacDonald, D. Speert, T. Fernandes-Alnemri, J. Wu, B. G. Monks, K. A. Fitzgerald, V. Hornung, and E. Latz. 2009. Cutting edge: NF-kappaB activating pattern recognition and cytokine receptors license NLRP3 inflammasome activation by regulating NLRP3 expression. *J Immunol* 183: 787-791.

50. Franchi, L., T. Eigenbrod, and G. Nunez. 2009. Cutting edge: TNF-alpha mediates sensitization to ATP and silica via the NLRP3 inflammasome in the absence of microbial stimulation. *J Immunol* 183: 792-796.
51. Lin, X. Y., M. S. Choi, and A. G. Porter. 2000. Expression analysis of the human caspase-1 subfamily reveals specific regulation of the CASP5 gene by lipopolysaccharide and interferon-gamma. *J Biol Chem* 275: 39920-39926.
52. Lagrange, B., S. Benaoudia, P. Wallet, F. Magnotti, A. Provost, F. Michal, A. Martin, F. Di Lorenzo, B. F. Py, A. Molinaro, and T. Henry. 2018. Human caspase-4 detects tetra-acylated LPS and cytosolic Francisella and functions differently from murine caspase-11. *Nature communications* 9: 242.
53. Kajiwara, Y., T. Schiff, G. Voloudakis, M. A. Gama Sosa, G. Elder, O. Bozdagi, and J. D. Buxbaum. 2014. A critical role for human caspase-4 in endotoxin sensitivity. *J Immunol* 193: 335-343.
54. Fortes, G. B., L. S. Alves, R. de Oliveira, F. F. Dutra, D. Rodrigues, P. L. Fernandez, T. Souto-Padron, M. J. De Rosa, M. Kelliher, D. Golenbock, F. K. Chan, and M. T. Bozza. 2012. Heme induces programmed necrosis on macrophages through autocrine TNF and ROS production. *Blood* 119: 2368-2375.
55. Kitur, K., S. Wachtel, A. Brown, M. Wickersham, F. Paulino, H. F. Penaloza, G. Soong, S. Bueno, D. Parker, and A. Prince. 2016. Necroptosis Promotes Staphylococcus aureus Clearance by Inhibiting Excessive Inflammatory Signaling. *Cell Rep* 16: 2219-2230.
56. Kayagaki, N., S. Warming, M. Lamkanfi, L. Vande Walle, S. Louie, J. Dong, K. Newton, Y. Qu, J. Liu, S. Heldens, J. Zhang, W. P. Lee, M. Roose-Girma, and V. M. Dixit. 2011. Non-canonical inflammasome activation targets caspase-11. *Nature* 479: 117-121.
57. Broz, P., T. Ruby, K. Belhocine, D. M. Bouley, N. Kayagaki, V. M. Dixit, and D. M. Monack. 2012. Caspase-11 increases susceptibility to Salmonella infection in the absence of caspase-1. *Nature* 490: 288-291.
58. Riedhammer, C., D. Halbritter, and R. Weissert. 2016. Peripheral Blood Mononuclear Cells: Isolation, Freezing, Thawing, and Culture. *Methods in molecular biology (Clifton, N.J.)* 1304: 53-61.
59. Gundry, M. C., L. Brunetti, A. Lin, A. E. Mayle, A. Kitano, D. Wagner, J. I. Hsu, K. A. Hoegenauer, C. M. Rooney, M. A. Goodell, and D. Nakada. 2016. Highly Efficient Genome Editing of Murine and Human Hematopoietic Progenitor Cells by CRISPR/Cas9. *Cell reports* 17: 1453-1461.
60. Moreno-Mateos, M. A., C. E. Vejnar, J. D. Beaudoin, J. P. Fernandez, E. K. Mis, M. K. Khokha, and A. J. Giraldez. 2015. CRISPRscan: designing highly efficient sgRNAs for CRISPR-Cas9 targeting in vivo. *Nature methods* 12: 982-988.