Temporal learning among prefrontal and striatal ensembles

Eric Emmons[1], Gabriela Tunes-Chuiffa[2], Jeeyu Choi[3], R. Austin Bruce[4], Matthew A. Weber[4], Youngcho Kim[4], Nandakumar S. Narayanan[4*]

Affiliations:
[1]Department of Psychiatry, Yale University, New Haven, CT, 06515
[2]Universidade Federal do ABC, Sao Paulo, Brazil
[3]Busan National University, Busan, South Korea
[4]Department of Neurology, University of Iowa, Iowa City, IA 52242.

Abstract: 145
Main text: 4545
Figures: 6
Tables: 4

*Corresponding Author
Nandakumar Narayanan
nandakumar-narayanan@uiowa.edu
169 Newton Road
Pappajohn Biomedical Discovery Building—1336
University of Iowa, Iowa City, 52242
319-353-5698

Abstract

Behavioral flexibility requires the prefrontal cortex and striatum. Here, we investigate neuronal

ensembles in the medial frontal cortex (MFC) and the dorsomedial striatum (DMS) during one

form of behavioral flexibility: learning a new temporal interval. We studied corticostriatal

neuronal activity as rodents trained to respond after a 12-second fixed interval (FI12) learned to

respond at a shorter 3-second fixed interval (FI3). On FI12 trials, we discovered time-related

ramping was reduced in the MFC but not in the DMS in two-interval vs. one-interval sessions.

We also found that more DMS neurons than MFC neurons exhibited differential interval-related

activity on the first day of two-interval performance. Finally, MFC and DMS ramping was

similar with successive days of two-interval performance but DMS temporal decoding increased

on FI3 trials. These data suggest that the MFC and DMS play distinct roles during temporal

learning and provide insight into corticostriatal circuits.

Introduction

Behavioral flexibility requires learning to adapt to uncertainty. Two forebrain structures critical for flexibility are the prefrontal cortex and striatum (Fuster, 2008; Kehagia et al., 2010). Prefrontal cortical neurons densely innervate the striatum (Gabbott et al., 2005; Wall et al., 2013) and disruptions of either structure profoundly impact the learning of new goals, rules, and strategies (Hart et al., 2018; Ragozzino, 2007). Dysfunctional corticostriatal circuits and connectivity are implicated in a range of psychiatric and neurological disorders (Deutch, 1993; Shepherd, 2013). However, the relative roles of prefrontal and striatal networks during behavioral flexibility are unclear.

One task that provides an ideal window into behavioral flexibility is interval timing, which requires participants to estimate an interval of several seconds via a motor response. Across species, interval timing requires the prefrontal cortex and striatum (Coull et al., 2011; Dallérac et al., 2017; Emmons et al., 2017, 2016; Matell and Meck, 2004; Merchant and de Lafuente, 2014). Work from our group and others has shown that both prefrontal and striatal neurons encode temporal information via 'time-related ramping' activity—or monotonic changes in firing rate over a temporal interval (Bakhurin et al., 2017; Donnelly et al., 2015; Emmons et al., 2017; Kim et al., 2018; Narayanan, 2016; Wang et al., 2018). Our past work suggested that ramping activity in neurons of the medial frontal cortex (MFC) and the dorsomedial striatum (DMS) is very similar, with ~40% of neurons in each area exhibiting such activity (Emmons et al., 2017). We have also found that MFC inactivation attenuates DMS ramping (Emmons et al., 2019, 2017) and that MFC stimulation is sufficient to increase DMS ramping (Emmons et al., 2019). These data suggest that DMS ramping is closely linked to MFC ramping and suggest the hypothesis that MFC and DMS ensembles respond similarly as animals learn new temporal

intervals. By contrast, recordings from primate lateral prefrontal cortex and caudate indicate that

striatal ensembles encode stimulus-response associations earlier than prefrontal ensembles,

leading to the hypothesis that prefrontal and striatal ensembles play differential roles during

learning (Antzoulatos and Miller, 2011; Histed et al., 2009; Pasupathy and Miller, 2005).

We tested these hypotheses by recording MFC and DMS activity in rodents as they

learned to respond to a new interval. Specifically, rodents previously trained to perform a 12-

second fixed-interval task learned a new version of the task that included two fixed-intervals—3

seconds and 12 seconds. We report three main results. First, time-related ramping activity in the

MFC decreased on 12-second interval trials during two-interval sessions compared to one-

interval sessions, whereas activity in the DMS on 12-second interval trials did not change.

Second, DMS neurons were more likely to have distinct firing patterns during the 12-second vs.

3-seconds interval than those of the MFC, but this interval-related activity normalized between

MFC and DMS over subsequent days of two-interval performance. Finally, MFC and DMS

ramping did not change consistently over subsequent two-interval sessions, but temporal

decoding by DMS ensembles improved for FI3 trials. These data suggest that the MFC and DMS

play distinct roles during temporal learning.

Methods

Rodents

All procedures were approved by the University of Iowa IACUC, and all methods were performed in accordance with the relevant guidelines and regulations (protocol #7072039). Seven male Long-Evans rats were trained on the 12-second fixed-interval timing task (FI12) according to procedures described in detail previously (Emmons et al., 2017, 2016). In brief, the rats were autoshaped to press a lever for water reward using a fixed-ratio task before being trained on 12-second fixed-interval timing (FI12). Trials began with the presentation of a house light, and the first response made after 12 seconds resulted in the delivery of a water reward, a concurrent click, and termination of the house light (Fig. 1A; video S1). Responses made before the interval ended were unreinforced. Trials were separated by a randomly chosen 6-, 8-, 10-, or 12-second intertrial interval. After animals behaved consistently, the MFC and DMS were each implanted with recording electrodes (Fig. 1B; see below). Animals were then acclimatized to the recording procedures and recordings were made during behavior in the FI12 task (Day 0). The following day, an additional 3-s interval (FI3) was added to the task and cued by a light distinct from the one used to indicate FI12. Behavior and simultaneous neuronal activity in the MFC and DMS were recorded over the following three days (Day 1, Day 2, and Day 3). Some data from subsequent recording sessions in these rodents were included in prior manuscripts (Emmons et al., 2017, 2016).

Surgical and histological procedures

Animals were anesthetized using ketamine (100 mg/kg) and xylazine (10 mg/kg), and a

surgical level of anesthesia was maintained using ketamine supplements (10 mg/kg).

Craniotomies were drilled above the left MFC and left DMS and four holes were drilled for skull

screws, which were connected to electrode recording arrays via a separate ground wire.

Microelectrode arrays were composed of 4x4 50-µm stainless steel wires (250 µm between wires

and rows; impedance measured *in vitro* at ~400 kΩ; Plexon, Dallas, TX). These arrays were

positioned in the MFC (coordinates from bregma: AP +3.2, ML ±1.2, DV -3.6 @ 12° in the

lateral plane) and the DMS (coordinates from bregma: AP +0.0, ML ±4.2, DV -3.6 @ 12° in the

posterior or lateral plane) while recording neuronal activity to verify that implantation was in

correct brain area. The craniotomy was sealed with cyanoacrylate ('SloZap', Pacer Technologies,

Rancho Cucamonga, CA), and the reaction was accelerated by 'ZipKicker' (Pacer Technologies)

and methyl methacrylate (AM Systems, Port Angeles, WA). Rats recovered for one week before

being acclimatized to behavioral and recording procedures.

Following these experiments, the rats were anesthetized and sacrificed by injection with

100 mg/kg sodium pentobarbital and transcardially perfused with 4% formalin. Brains were post-

fixed in a solution of 4% formalin and 20% sucrose before being sectioned on a freezing

microtome. Brain slices were mounted on Superfrost Plus microscope slides and stained for cell

bodies using either DAPI or Cresyl violet. Histological reconstruction was completed using

postmortem analysis of electrode placement by slide-scanning fluorescent microscopy

(Olympus).

Neurophysiological recordings and neuronal analyses

Neuronal ensemble recordings were made using a multi-electrode recording system (Plexon). In each animal, one electrode without single units was reserved for local referencing, yielding 15 electrodes per animal. Offline Sorter (Plexon) was used to analyze the signals after the experiments and to remove artifacts. Spike activity was analyzed for all cells that fired at rates above 0.1 Hz. Principal component analysis (PCA) and waveform shape were used for spike sorting. DMS neurons were classified as either medium spiny neurons (MSNs) or interneurons based on peak-to-trough ratio and the spike half-peak width of spike waveforms (Fig. 1C; Berke, 2011). Single units were defined as those 1) having a consistent waveform shape, 2) being a separable cluster in PCA space, and 3) having a consistent refractory period of at least 2 ms in interspike interval histograms.

Statistics

Basic analyses were performed via ANOVA, Wilcoxon rank-sum tests, and Cohen's D. As in our prior work, we quantified temporal control of action during fixed-interval performance in two ways. We calculated the curvature of time-response histograms (Emmons et al., 2019; Fry et al., 1960; Narayanan et al., 2012). Curvature values range between -1 and 1 and are calculated from the cumulative response record by deviation from a straight line; 0 indicates a constant response rate throughout the interval. Curvature indices are resistant to differences in response rate, smoothing, or binning. Second, we modeled each response using generalized linear mixed-effects models (GLMM; *fitglme.m* in MATLAB) where the outcome is response time, the predictor variable was Day, and the random effect was animal. For two-interval trials, single-trial analyses were used to find start times and coefficients of variation for FI3 and FI12 trials (Church et al., 1994). All data and statistical approaches were reviewed by the Biostatistics,

Epidemiology, and Research Design Core (BERD) at the Institute for Clinical and Translational

Sciences (ICTS) at the University of Iowa.

Analyses of neuronal activity and basic firing properties were carried out using

NeuroExplorer (Nex Technologies, Littleton, MA) and custom routines for MATLAB, as

described in detail previously (Emmons et al., 2019, 2017; Parker et al., 2014). Peri-event rasters

and time-histograms were constructed around houselight and lever press. As in our past work,

neuronal modulations were quantified in two ways. First, we used principal component analysis

(PCA), a data-driven set of orthogonal basis vectors that captures patterns of activity in

multivariate neuronal ensembles. PCA was calculated from average peri-event time histograms

computed from kernel-density estimates (*ksdensity.m*; bandwidth 0.5) and normalized using

zscore. As in our past work, we used absolute values of PC1 scores (indicated by |PC1|) compare

ramping strength across areas and days (Emmons et al., 2017; Kim et al., 2017; Parker et al.,

2017, 2014).

We also used *fitglme.m* to construct GLMMs to analyze neuronal modulations. For

neuron-by-neuron analysis, we used GLMMs where the outcome variable was the firing rate

(binned at 0.1 s) and the predictor was time in the interval (0 to 3 seconds on FI3, or 0 to 12

seconds on FI12), interval (FI3 or FI12), or responses. Neurons with a main effect of time were

considered 'ramping' neurons, neurons with a main effect of interval-type were considered

interval-modulated, and neurons with a main effect of response were considered response-

modulated. Neurons with time-related ramping were defined as those with a main-effect of time

in the interval via GLMMs where the outcome was firing rate binned at 0.1 s, and the predictor

was time in the interval. We also performed trial-by-trial GLMMs for all trials and neurons

where the outcome was firing rate, and the predictors were area (MFC or DMS) or Day 0 or Day

1, and random effects were lever presses and neurons (Table 1). To examine interval-related

modulation for all trials and neurons, we used an equation where the outcome was firing rate, the

predictors were interval type (FI3 or FI12), area, or Day, and random effects were lever presses

and neurons (Table 2). For two-interval performance, we used GLMMs for FI3 trials (Table 3)

and FI12 trials (Table 4) on Days 1-3. Poisson distributions were used for all firing rate models.

We used a naïve Bayesian classifier to examine neuronal ensemble decoding, as we have

in our past work (Emmons et al., 2017; Kim et al., 2017). We calculated kernel density estimates

(bandwidth: 1.2) of trial-by-trial firing rates from MFC and DMS neurons. To prevent edge

effects that might bias classifier performance, we included data from 6 seconds prior to trial start

and 6 seconds after interval end. We used leave-one-out cross-validation to predict objective

time from firing rate within a trial. We evaluated classifier performance by computing the R^2 of

objective time and predicted time only for bins during the interval. With perfect classification,

the R^2 would approach 1. Classifier performance was compared to ensembles with time-shuffled

firing rates. For each area and interval, performance was compared via GLMMs of R^2 vs. each

day.

Results

We studied temporal learning in the MFC and DMS by introducing a new 3-s fixed

interval (FI3) to rats after they had been trained on a task with 12-second fixed-intervals (FI12;

Fig. 1A). We compared behavior on FI12 trials only in sessions with one interval ("Day 0") to

two-interval sessions in which FI12 trials were randomly intermixed with FI3 trials ("Day 1").

We quantified the temporal control of action via a 'curvature' index calculated from the

cumulative distribution of time-response histograms. We have used this index extensively in the

past (Emmons et al., 2019; Narayanan et al., 2012). On FI12 trials, the curvature index trended

towards being lower on Day 1 vs. Day 0 (Fig. 2A-B; Day 0 curvature: 0.29 ± 0.04, mean $\pm$ SEM;

Day 1 curvature: 0.17 ± 0.06; signrank $p = 0.08$; Cohen's d $= 0.86$). Response times were

significantly shorter on Day 1 vs. Day 0 (Fig. 2C-D; Day 0: 10.14 ± 0.09 vs. Day 1: 9.61 ± 0.10

seconds; main effect of Day: $F_{(1,3784)} = 12.28$, $p = 0.0005$; $R^2 = 0.003$). These results suggest that

the timing of FI12 responses was shifted earlier on FI12 trials on Day 1 when they were

intermixed with FI3 trials compared to Day 0, when FI12 trials were presented alone.

We recorded neuronal ensembles simultaneously in the MFC and DMS as animals well-

trained on FI12 trials on Day 0 learned to perform a two-interval task with FI3 and FI12 trials

randomly intermixed on Day 1. As in our past work, we found that neurons in both brain regions

exhibited time-dependent ramping, i.e. monotonic increases in firing across the interval

(Emmons et al., 2017; Fig. 3A-B). On FI12 trials 35 of 59 (59%) MFC neurons exhibited

ramping activity on Day 0, surprisingly, there were only 19 of 47 (40%) ramping neurons in

MFC on Day 1 (Fig. 3C; $X^2 = 3.74$, $p = 0.05$). In the DMS, 32 of 67 (48%) neurons ramped Day

0 and 29 of 58 (50%) ramped on Day 1, ($X^2 = 0.06$, $p = 0.80$).

To further compare ramping activity, we turned to principal component analysis (PCA)

as a data-driven approach to compare neuronal activity patterns (Chapin and Nicolelis, 1999;

Emmons et al., 2017; Narayanan and Laubach, 2009). Consistent with our prior work, we found

that principal component 1 (PC1) exhibited time-related ramping (Fig 3H; Emmons et al., 2017;

Kim et al., 2017; Parker et al., 2015, 2014; Zhang et al., 2019). For the MFC, the strength of

|PC1| was lower on Day 1 than Day 0 (Fig. 3I; signrank $p = 0.03$; Cohen's d = 0.50), whereas

there was no consistent difference for the DMS (Fig 3I; signrank $p = 0.60$). Trial by-trial analysis

of firing rate on FI12 trials revealed that time-related ramping interacted with both the brain area

(MFC and DMS) as well one- vs. two-interval sessions (i.e., Day 0 vs. Day 1). There was a

three-way interaction between time-related ramping, brain area, and one vs. two intervals (Table

1). For FI12 trials, these data suggest that time-related ramping in the MFC was stronger on Day

0 compared to Day 1 when FI12 trials were intermixed with FI3 trials. Our results suggest that

neuronal ensembles in the MFC, but not in the DMS, are sensitive to the temporal context

(Jazayeri and Shadlen, 2010; Shi et al., 2013).

Next, we compared MFC and DMS activity on FI3 and FI12 trials. First, we analyzed

fixed-interval behavior using single-trial analysis, which was developed for peak-interval timing

but can be useful to analyze start times during fixed-interval tasks (Church et al., 1994; Emmons

et al., 2019). On the first day of the shorter 3-second interval, single-trial analysis revealed that

animals had shorter start times on FI3 trials compared to FI12 trials (FI3: 3.18 ± 0.69 vs. FI12

7.06 ± 0.34, signrank $p = 0.03$; Cohen's d = 2.9; single-trial analysis could not compute FI3 start

times from one animal, which was subsequently removed from this analysis). One indication that

timing processes are scalar is that the coefficient of variation (CV—the ratio of standard

deviation of temporal estimates to the mean) is relatively constant at different intervals (Gibbon

et al., 1984; Rakitin et al., 1998). Accordingly, we found that during fixed-interval performance,

start time CVs were similar on FI3 and FI12 trials (FI3: 0.59+/-0.12 vs. FI12: 0.41+/-0.03

signrank p: 0.56). These data suggest that start times during fixed-interval timing exhibit scalar

properties (Gibbon et al., 1984).

Ramping neurons can have distinct slopes of firing rates vs. time on FI3 and FI12 trials

(Fig 4A). We ran GLMMs where firing-rate slope vs. time was the outcome variable, and FI3 vs.

FI12 interval and Day were predictor variables; note that we were interested in the magnitude of

the slope and we focus on its absolute value, indicated by |slope|. Consistent with past work by

our group and others, ramping neuron |slopes| were consistently steeper on FI3 vs. FI12 trials for

both the MFC (Fig. 4B; main effect of interval: $F_{(143)} = 4.52$, $p = 0.04$, $R^2 = 0.22$) and for the

DMS (main effect of interval: $F_{(214)} = 6.91$, $p = 0.01$; $R^2 = 0.40$(Emmons et al., 2017; Mello et

al., 2015; Wang et al., 2018). There was no effect of Day or higher interactions for either MFC

or DMS. These data are consistent with influential drift-diffusion models of interval timing,

suggesting that drift rates increase with shorter intervals (Simen et al., 2011).

Next, we searched for neurons in which firing rates were a function of interval duration

(Fig 4C). Specifically, we used GLMMs to identify neurons with a main effect of interval on

firing rate (Fig. 4E; FI3 vs. FI12 trials). Interval-modulated neurons were more common in the

DMS than the MFC on Day 1 of two-interval performance (Fig 4D; MFC 10 of 47 vs. DMS: 28

of 58; $X^2 = 8.20$; $p = 0.004$). Interestingly, ~50% of interval-modulated neurons also had ramping

activity in MFC (5 of 10) and DMS (15 of 28). Of note, the number of interval-modulated

neurons was not different between MFC and DMS on Days 2 and 3 (Fig 4D). Consistent with

these analyses, GLMMs revealed a significant interaction between interval-modulation, brain

area, and Days 1-3 (Table 2).

A comparison of behavior across the three days of two-interval performance revealed that

response times shortened for both FI3 and FI12 trials (Fig. 5A-B: GLMM FI3: $F_{(1741)} = 8.35$, $p =$

0.0002; $R^2 = 0.05$; FI12: $F_{(4796)} = 6.98$, $p = 0.0009$; $R^2 = 0.002$), but the curvature of time-

response histograms did not reliably change (FI3: $F_{(40)} = 0.03$, $p = 0.87$, $R^2 = 0.17$; FI12: $F_{(40)} =$

1.05, $p = 0.31$, $R^2 = 0.61$). As in prior work demonstrating temporal scaling in the MFC and

DMS, PCA revealed that the principal components for FI3 and FI12 trials were very similar (Fig.

5C-D; Pearson's *rho* correlation: PC1: = 0.99 $p < 0.001$; PC2: -0.95, $p < 0.001$; PC3: 0.95, $p <$

0.001; (Emmons et al., 2017; Mello et al., 2015; Wang et al., 2018). However, |PC1| did not

change consistently over the three days of two-interval performance for MFC or DMS (Fig. 5E-

H; Table 3-4). Taken together, these data indicate that ramping-related patterns of activity of

corticostriatal ensembles did not consistently change as animals performed two-interval tasks on

Days 1-3.

We turned to decoding analyses based on machine learning to capture more complex

features of MFC and DMS ensembles (Fig 6; Emmons et al., 2017; Gouvea et al., 2015; Kim et

al., 2017). Specifically, we constructed neuron-dimensional arrays of smoothed trial-by-trial

firing rates over the interval binned at 0.1 seconds. We decoded time in the interval from

ensemble firing rates using naïve Bayesian classifiers. Classifier performance was assessed by

computing the variance explained (R^2) of predicted vs. observed time. For all sessions, R^2 was

much less for time-shuffled ensembles—i.e., ensembles constructed from neuronal activity

shuffled in time (Fig. 6B-G; signrank $p = 4*10^{-47}$; Cohen's d = 1.57). We found that temporal

decoding had a main effect of Day only for the DMS Ensembles on FI3 trials (Fig. 6F&G; $F_{(108)}$

= 6.07, $p = 0.02$, $R^2 = 0.06$). These results suggest that as the response times shortened with two-

interval performance, temporal decoding improved only for DMS ensembles on FI3 trials.

Discussion

We found three key distinctions between MFC and DMS during temporal learning. First,

time-related ramping in the MFC decreased as animals that had been trained on a one-interval

task learned to respond to a second interval that was novel and shorter. Second, interval-

modulated neurons were more common in the DMS early in two-interval performance. Third,

time-related ramping in the MFC and DMS did not change with two-interval performance, but

temporal decoding improved only for the DMS on FI3 trials. Our data suggest that MFC

ensembles are sensitive to the context or 'rules' in the task – i.e., FI12 vs. FI3/FI12, while the

DMS optimizes behavior particularly on FI3 trials. These data provide insight into the relative

roles of prefrontal and striatal networks during temporal learning.

These results contradicted our hypothesis that time-related ramping in the MFC and DMS

would be similarly affected by the introduction of a new temporal interval. Our hypothesis was

based on five lines of evidence: 1) the existence of strong projections from the MFC to the DMS

(Gabbott et al., 2005; Han et al., 2017; Wall et al., 2013), 2) clear roles for both structures in

interval timing (Coull et al., 2011; De Corte et al., 2019; Emmons et al., 2017; Meck, 2006), 3)

similarities in time-related ramping in the MFC and DMS (Emmons et al., 2017), 4) the necessity

of MFC activity for DMS ramping (Emmons et al., 2019, 2017), and 5) our recent demonstration

that the stimulation of axons that project from the MFC to the DMS is sufficient to increase time-

related ramping in the DMS (Emmons et al., 2019). Given these data, it is notable that the MFC

and DMS play distinct roles during temporal learning, although this observation is concordant

with the vastly different connectivity and synaptic organization of these two structures

(Shepherd, 2003). Nevertheless, decreases in MFC ramping after the introduction of a shorter

interval suggest that MFC ramping is sensitive to the temporal context of the one-interval vs.

two-interval task, and they may reflect Bayesian priors of temporal probabilities (Jazayeri and

Shadlen, 2010; Shi et al., 2013).

Differences between the MFC and DMS were anticipated based on a recent comparison

of neuronal ensembles during a temporal categorization task that involved maze running (Kim et

al., 2018). This study indicated that ramping was more prevalent in the MFC than the dorsal

striatum. In this task as the intervals became longer, temporal decoding by the MFC was less

effective than that by the striatum. Although this task was more complex than ours and a number

of others (Bakhurin et al., 2017, 2016; Donnelly et al., 2015; Narayanan, 2016; Wang et al.,

2018), the strong temporal encoding across corticostriatal ensembles is consistent with our

findings here.

We found that the DMS contained more neurons in which there was a main effect of

interval compared to the MFC on the early days of two-interval performance. Notably, half of

interval-modulated neurons were not ramping. These data suggest that patterns beyond time-

related ramping encode information about temporal intervals. On progressive days of two-

interval performance, interval-related activity between the MFC and DMS equalized. Because

our task design involved a second cue for FI3 intervals, we cannot distinguish whether this

activity was related to working memory for temporal intervals, cue-related processing, or other

aspects of interval timing. Future work using more advanced learning paradigms may clarify

these patterns of activity.

Our findings are in line with drift-diffusion models of two-interval tasks, as we find that

time-related ramping scales with the interval duration (Simen et al., 2011). We find that MFC

ramping is sensitive to temporal context whereas DMS ramping is not, and that non-ramping

interval-related modulations and temporal predictions in the DMS change with two-interval

performance. These results suggest that time-related ramping reflects distinct processes in MFC

and DMS. Given that MFC activity influences ramping in DMS (Emmons et al., 2019, 2017),

DMS ramping activity might integrate aspects of MFC ramping as well as non-ramping activity.

Because time-related ramping activity in MFC and DMS ensembles did not change

during two-interval performance, ramping activity may be remarkably stable in both brain

regions when the temporal context does not change. It is unclear how ramping might change with

extended periods of behavior over several days or weeks (Barnes et al., 2005; Graybiel, 2008;

Yin et al., 2005). However, we did find that on FI3 trials, temporal decoding in the DMS

improved even though DMS ramping was stable. In the DMS patterns beyond ramping activity

might change during two-interval performance and contribute to improved temporal decoding

(Paton and Buonomano, 2018). The improvement in temporal prediction despite unchanged

ramping activity supports improved 'population clock'-based temporal predictions during FI3

trials (Karmarkar and Buonomano, 2007; Laje and Buonomano, 2013).

To our knowledge, our study is one of the first to record from corticostriatal ensembles

during temporal learning. The striatum has a well-established role in learning other contexts

including habit formation, reversal learning, and instrumental learning (Graybiel and Grafton,

2015; Kimchi and Laubach, 2009; Yin and Knowlton, 2006). However, direct comparisons of

cortical and striatal learning in rodents, in any context, are rare. One exception is a recent study

of prenatal alcohol exposure, which showed that the orbitofrontal cortex disengages and the

dorsal striatum updates reward contingencies (Marquardt et al., 2020). These findings parallel

the changes in the MFC and DMS that we report here. The observation that prenatal exposure to

alcohol leads to changes in cortical activity underscores the clinical significance of this brain

circuit.

During associative learning in primates, corticostriatal ensembles are highly sensitive to

learning, with striatal neurons rapidly encoding new associations and the prefrontal cortex

learning more slowly (Pasupathy and Miller, 2005). Primate striatal neurons rapidly encoded

stimulus-response associations, whereas primate prefrontal neurons encoded category abstraction

(Antzoulatos and Miller, 2011). In line with these results, we found that time-related ramping

decreased in two-interval vs. one-interval sessions, suggesting that prefrontal ensembles may be

sensitive to temporal categories or context. It is important to note that these primate studies

recorded from lateral prefrontal areas, which lack a clear rodent analogue (Laubach et al., 2018),

and that they employed vastly different task conditions. Nevertheless, our work provides insight

into the dynamics of rodent corticostriatal ensembles during an elementary temporal learning

paradigm.

Our study has several limitations. First, we used fixed-interval timing; peak-interval

timing tasks might enable more precise dissection of start and stop times (Rakitin et al., 1998).

Second, our techniques cannot identify the genetic or molecular identity of recorded neurons.

This detail would be of particular interest in the case of the DMS, which contains D1 and D2

MSNs (Kreitzer, 2009). Third, we are unsure if the MFC and DMS neurons we captured were

connected, because of the sparsity of cortical projections and constraints of our recording

techniques (Wall et al., 2013). This limitation might be overcome in future work by exploiting

optogenetic tagging and retrograde viral tracing to isolate corticostriatal projections (Otis et al.,

2017). Studying how MFC-DMS connectivity changes with learning might provide further

insight in corticostriatal circuits. Fourth, we cannot reliably follow neurons over separate

sessions. Finally, we were unable to clearly identify clear correlates of temporal learning during

the first two-interval session on Day 1. Corticostriatal ensembles may rapidly learn the new

interval. Such rapid learning has been observed in the striatum, but capturing it might require a

different task design to capture trial-by-trial neuronal dynamics during learning (Kimchi and

Laubach, 2009).

In summary, we investigated corticostriatal ensembles in rodents that had been well-

trained to perform a single fixed-interval timing task while they learned to incorporate a new

interval. We discovered that time-related ramping activity in the MFC decreased following

introduction of the shorter interval, whereas ramping activity in the DMS was unchanged. We

also found that more DMS neurons fired differentially on each interval compared to the MFC

early in two-interval performance. Finally, corticostriatal ramping activity did not change on the

days following the initiation of two-interval performance, yet DMS temporal decoding

improved. Taken together, our data suggest that the MFC and DMS play distinct roles in

temporal learning.

Acknowledgements: We thank Morgan Kennedy and Tomas Lence for technical help.

Funding: NIMH R01 to NN and Titan Neurological Fund to NN, NIMH F31 to EE, CAPES to
GT

Contributions: EE, YK, and NN designed experiments; EE and GTC collected data, EE, YK,
KC, and NN analyzed data, AB independently checked the code and data, and EE, GTC, MW,
AB, YK, and NN wrote the manuscript.

Data and Code: Available at narayanan.lab.uiowa.edu

References

Antzoulatos EG, Miller EK. 2011. Differences between Neural Activity in Prefrontal Cortex and
Striatum during Learning of Novel Abstract Categories. *Neuron* **71**:243–249.
doi:10.1016/j.neuron.2011.05.040
Bakhurin KI, Goudar V, Shobe JL, Claar LD, Buonomano DV, Masmanidis SC. 2017.
Differential Encoding of Time by Prefrontal and Striatal Network Dynamics. *J Neurosci*
37:854–870. doi:10.1523/JNEUROSCI.1789-16.2017
Bakhurin KI, Goudar V, Shobe JL, Claar LD, Buonomano DV, Masmanidis SC. 2016.
Differential encoding of time by prefrontal and striatal network dynamics. *J Neurosci*.
doi:10.1523/JNEUROSCI.1789-16.2016
Barnes TD, Kubota Y, Hu D, Jin DZ, Graybiel AM. 2005. Activity of striatal neurons reflects
dynamic encoding and recoding of procedural memories. *Nature* **437**:1158–1161.
doi:10.1038/nature04053
Berke JD. 2011. Functional properties of striatal fast-spiking interneurons. *Front Syst Neurosci*
5:45. doi:10.3389/fnsys.2011.00045
Chapin JK, Nicolelis MA. 1999. Principal component analysis of neuronal ensemble activity
reveals multidimensional somatosensory representations. *J Neurosci Methods* **94**:121–
140.
Church RM, Meck WH, Gibbon J. 1994. Application of scalar timing theory to individual trials.
J Exp Psychol Anim Behav Process **20**:135–155.
Coull JT, Cheng R-K, Meck WH. 2011. Neuroanatomical and neurochemical substrates of
timing. *Neuropsychopharmacology* **36**:3–25. doi:10.1038/npp.2010.113
Dallérac G, Graupner M, Knippenberg J, Martinez RCR, Tavares TF, Tallot L, El Massioui N,
Verschueren A, Höhn S, Bertolus JB, Reyes A, LeDoux JE, Schafe GE, Diaz-Mataix L,
Doyère V. 2017. Updating temporal expectancy of an aversive event engages striatal
plasticity under amygdala control. *Nat Commun* **8**:13920–13920.
doi:10.1038/ncomms13920
De Corte BJ, Wagner LM, Matell MS, Narayanan NS. 2019. Striatal dopamine and the temporal
control of behavior. *Behav Brain Res* **356**:375–379. doi:10.1016/j.bbr.2018.08.030
Deutch AY. 1993. Prefrontal cortical dopamine systems and the elaboration of functional
corticostriatal circuits: implications for schizophrenia and Parkinson's disease. *J Neural*
Transm Gen Sect **91**:197–221.
Donnelly NA, Paulsen O, Robbins TW, Dalley JW. 2015. Ramping single unit activity in the
medial prefrontal cortex and ventral striatum reflects the onset of waiting but not
imminent impulsive actions. *Eur J Neurosci* n/a-n/a. doi:10.1111/ejn.12895
Emmons EB, Corte BJD, Kim Y, Parker KL, Matell MS, Narayanan NS. 2017. Rodent Medial
Frontal Control of Temporal Processing in the Dorsomedial Striatum. *J Neurosci*
37:8718–8733. doi:10.1523/JNEUROSCI.1376-17.2017
Emmons EB, Kennedy M, Kim Y, Narayanan NS. 2019. Corticostriatal stimulation compensates
for medial frontal inactivation during interval timing. *Scientific Reports* **9**:1–9.
doi:10.1038/s41598-019-50975-7
Emmons EB, Ruggiero RN, Kelley RM, Parker KL, Narayanan NS. 2016. Corticostriatal Field
Potentials Are Modulated at Delta and Theta Frequencies during Interval-Timing Task in
Rodents. *Front Psychol* **7**. doi:10.3389/fpsyg.2016.00459
Fry W, Kelleher RT, Cook L. 1960. A mathematical index of performance on fixed-interval
schedules of reinforcement. *J Exp Anal Behav* **3**:193–199. doi:10.1901/jeab.1960.3-193

Fuster J. 2008. The Prefrontal Cortex, 4th Edition. ed. New York, NY: Academic Press.

Gabbott PLA, Warner TA, Jays PRL, Salway P, Busby SJ. 2005. Prefrontal cortex in the rat:
projections to subcortical autonomic, motor, and limbic centers. *J Comp Neurol* **492**:145–
177. doi:10.1002/cne.20738

Gibbon J, Church RM, Meck WH. 1984. Scalar timing in memory. *Ann N Y Acad Sci* **423**:52–77.

Graybiel AM. 2008. Habits, Rituals, and the Evaluative Brain. *Annual Review of Neuroscience*
31:359–387. doi:10.1146/annurev.neuro.29.051605.112851

Graybiel AM, Grafton ST. 2015. The striatum: where skills and habits meet. *Cold Spring Harb*
Perspect Biol **7**:a021691–a021691. doi:10.1101/cshperspect.a021691

Han S-W, Kim Y-C, Narayanan NS. 2017. Projection targets of medial frontal D1DR-expressing
neurons. *Neuroscience Letters* **655**:166–171. doi:10.1016/j.neulet.2017.06.057

Hart G, Bradfield LA, Balleine BW. 2018. Prefrontal Corticostriatal Disconnection Blocks the
Acquisition of Goal-Directed Action. *J Neurosci* **38**:1311–1322.
doi:10.1523/JNEUROSCI.2850-17.2017

Histed MH, Pasupathy A, Miller EK. 2009. Learning substrates in the primate prefrontal cortex
and striatum: sustained activity related to successful actions. *Neuron* **63**:244–253.
doi:10.1016/j.neuron.2009.06.019

Jazayeri M, Shadlen MN. 2010. Temporal context calibrates interval timing. *Nat Neurosci*
13:1020–1026. doi:10.1038/nn.2590

Karmarkar UR, Buonomano DV. 2007. Timing in the Absence of Clocks: Encoding Time in
Neural Network States. *Neuron* **53**:427–438. doi:10.1016/j.neuron.2007.01.006

Kehagia AA, Murray GK, Robbins TW. 2010. Learning and cognitive flexibility: frontostriatal
function and monoaminergic modulation. *Curr Opin Neurobiol* **20**:199–204.
doi:10.1016/j.conb.2010.01.007

Kim J, Kim D, Jung MW. 2018. Distinct Dynamics of Striatal and Prefrontal Neural Activity
During Temporal Discrimination. *Front Integr Neurosci* **12**:34.
doi:10.3389/fnint.2018.00034

Kim Y-C, Han S-W, Alberico SL, Ruggiero RN, De Corte B, Chen K-H, Narayanan NS. 2017.
Optogenetic Stimulation of Frontal D1 Neurons Compensates for Impaired Temporal
Control of Action in Dopamine-Depleted Mice. *Curr Biol* **27**:39–47.
doi:10.1016/j.cub.2016.11.029

Kimchi EY, Laubach M. 2009. The dorsomedial striatum reflects response bias during learning.
J Neurosci **29**:14891–14902. doi:10.1523/JNEUROSCI.4060-09.2009

Kreitzer AC. 2009. Physiology and Pharmacology of Striatal Neurons. *Annual Review of*
Neuroscience **32**:127–147. doi:10.1146/annurev.neuro.051508.135422

Laje R, Buonomano DV. 2013. Robust timing and motor patterns by taming chaos in recurrent
neural networks. *Nat Neurosci* **16**:925–933. doi:10.1038/nn.3405

Laubach M, Amarante LM, Swanson K, White SR. 2018. What, If Anything, Is Rodent
Prefrontal Cortex? *eNeuro* **5**. doi:10.1523/ENEURO.0315-18.2018

Marquardt K, Cavanagh JF, Brigman JL. 2020. Alcohol exposure in utero disrupts cortico-
striatal coordination required for behavioral flexibility. *Neuropharmacology* **162**:107832–
107832. doi:10.1016/j.neuropharm.2019.107832

Matell MS, Meck WH. 2004. Cortico-striatal circuits and interval timing: coincidence detection
of oscillatory processes. *Brain Res Cogn Brain Res* **21**:139–170.
doi:10.1016/j.cogbrainres.2004.06.012

Meck WH. 2006. Neuroanatomical localization of an internal clock: A functional link between mesolimbic, nigrostriatal, and mesocortical dopaminergic systems. *Brain Research* **1109**:93–107. doi:10.1016/j.brainres.2006.06.031

Mello GBM, Soares S, Paton JJ. 2015. A Scalable Population Code for Time in the Striatum. *Current Biology* **25**:1113–1122. doi:10.1016/j.cub.2015.02.036

Merchant H, de Lafuente V. 2014. Introduction to the neurobiology of interval timing. *Adv Exp Med Biol* **829**:1–13. doi:10.1007/978-1-4939-1782-2_1

Narayanan NS. 2016. Ramping activity is a cortical mechanism of temporal control of action. *Curr Opin Behav Sci* **8**:226–230. doi:10.1016/j.cobeha.2016.02.017

Narayanan NS, Land BB, Solder JE, Deisseroth K, DiLeone RJ. 2012. Prefrontal D1 dopamine signaling is required for temporal control. *Proc Natl Acad Sci USA* **109**:20726–20731. doi:10.1073/pnas.1211258109

Narayanan NS, Laubach M. 2009. Delay activity in rodent frontal cortex during a simple reaction time task. *J Neurophysiol* **101**:2859–2871. doi:10.1152/jn.90615.2008

Otis JM, Namboodiri VMK, Matan AM, Voets ES, Mohorn EP, Kosyk O, McHenry JA, Robinson JE, Resendez SL, Rossi MA, Stuber GD. 2017. Prefrontal cortex output circuits guide reward seeking through divergent cue encoding. *Nature* **advance online publication**. doi:10.1038/nature21376

Parker KL, Chen K-H, Kingyon JR, Cavanagh JF, Narayanan NS. 2014. D1-Dependent 4 Hz Oscillations and Ramping Activity in Rodent Medial Frontal Cortex during Interval Timing. *J Neurosci* **34**:16774–16783. doi:10.1523/JNEUROSCI.2772-14.2014

Parker KL, Kim YC, Kelley RM, Nessler AJ, Chen K-H, Muller-Ewald VA, Andreasen NC, Narayanan NS. 2017. Delta-frequency stimulation of cerebellar projections can compensate for schizophrenia-related medial frontal dysfunction. *Mol Psychiatry* **22**:647–655. doi:10.1038/mp.2017.50

Parker KL, Ruggiero RN, Narayanan NS. 2015. Infusion of D1 Dopamine Receptor Agonist into Medial Frontal Cortex Disrupts Neural Correlates of Interval Timing. *Front Behav Neurosci* **9**:294. doi:10.3389/fnbeh.2015.00294

Pasupathy A, Miller EK. 2005. Different time courses of learning-related activity in the prefrontal cortex and striatum. *Nature* **433**:873–876.

Paton JJ, Buonomano DV. 2018. The Neural Basis of Timing: Distributed Mechanisms for Diverse Functions. *Neuron* **98**:687–705. doi:10.1016/j.neuron.2018.03.045

Ragozzino ME. 2007. The Contribution of the Medial Prefrontal Cortex, Orbitofrontal Cortex, and Dorsomedial Striatum to Behavioral Flexibility. *Annals of the New York Academy of Sciences* **1121**:355–375. doi:10.1196/annals.1401.013

Rakitin BC, Gibbon J, Penney TB, Malapani C, Hinton SC, Meck WH. 1998. Scalar expectancy theory and peak-interval timing in humans. *J Exp Psychol Anim Behav Process* **24**:15–33.

Shepherd G. 2003. Synaptic Organization of The Brain, 34d ed. Oxford: Oxford University Press.

Shepherd GMG. 2013. Corticostriatal connectivity and its role in disease. *Nat Rev Neurosci* **14**:278–291. doi:10.1038/nrn3469

Shi Z, Church RM, Meck WH. 2013. Bayesian optimization of time perception. *Trends Cogn Sci* **17**:556–564. doi:10.1016/j.tics.2013.09.009

Simen P, Balci F, de Souza L, Cohen JD, Holmes P. 2011. A model of interval timing by neural integration. *J Neurosci* **31**:9238–9253. doi:10.1523/JNEUROSCI.3121-10.2011

Wall NR, De La Parra M, Callaway EM, Kreitzer AC. 2013. Differential Innervation of Direct-
and Indirect-Pathway Striatal Projection Neurons. *Neuron* **79**:347–360.
doi:10.1016/j.neuron.2013.05.014
Wang J, Narain D, Hosseini EA, Jazayeri M. 2018. Flexible timing by temporal scaling of
cortical responses. *Nature Neuroscience* **21**:102. doi:10.1038/s41593-017-0028-6
Yin HH, Knowlton BJ. 2006. The role of the basal ganglia in habit formation. *Nat Rev Neurosci*
7:464–476. doi:10.1038/nrn1919
Yin HH, Ostlund SB, Knowlton BJ, Balleine BW. 2005. The role of the dorsomedial striatum in
instrumental conditioning. *European Journal of Neuroscience* **22**:513–523.
doi:10.1111/j.1460-9568.2005.04218.x
Zhang Q, Jung D, Larson T, Kim Y, Narayanan NS. 2019. Scopolamine and Medial Frontal
Stimulus-Processing during Interval Timing. *Neuroscience* **414**:219–227.
doi:10.1016/j.neuroscience.2019.07.004
511

Figures:

Figure 1: Fixed-interval timing tasks and recording locations: A) On Day 0, rodents

performed fixed-interval timing tasks in which a reward was given for the first lever press after a

12-second interval (FI12). Interval start was cued by a house light, motivation was a liquid

reward, and presses before interval end were unreinforced. On Day 1, a second, shorter 3-second

interval (FI3) was introduced and cued by a distinct light. FI3 trials were randomly intermixed

with FI12 trials. Recordings were performed for two days following the initial two-interval

performance (Day 2 and Day 3). B) Animals were implanted with neuronal ensemble arrays

targeting the medial frontal cortex (MFC) and dorsomedial striatum (DMS). C) MSNs within the

DMS were identified based on waveform shape.

Figure 2: Response times reflect temporal context. A) Kernel density estimates of time-

response histograms across animals on FI12 trials on Day 0 (black) when FI12 trials were

presented alone to Day 1, with FI12 trials were presented alongside randomly intermixed FI3

trials. B) Curvature indices of time-response histograms from 7 animals on Day 0 vs. Day 1. C)

Compilation of every FI12 response from every animal on Day 0 (darker colors) vs. Day 1

(lighter colors); each animal is represented by dots of a different color. D) Violin plot of all

responses; vertical lines denote the mean and thicker horizontal gray lines span the interquartile

range. Data from FI12 trials from seven animals; ~ indicates a trend via Wilcoxon rank-sum; *

indicates $p < 0.05$ via GLMMs.

Figure 3: MFC ramping reflects temporal context. Peri-event rasters for single neurons in the

A) MFC (blue) and B) DMS (green). Top panels: each row represents a trial; each tick is an

action potential; darker colors represent Day 0 (FI12) and lighter colors represent Day 1

(FI12/FI3); all data are from FI12 trials only. C) Quantification of neurons that underwent time-

related ramping via GLMMs in the MFC and DMS for Day 0 and Day 1; * indicates $p < 0.05$ via

a chi-squared test. Peri-event histograms from all neurons in the MFC on Day 0 (D) and Day 1

(E) and all MSNs in the DMS on Day 0 (F) and Day 1(G). H) Principal component analyses

revealed three main components; percentage of variance is indicated in parentheses. I) |PC1

scores| for MFC and DMS ensembles on Day 1 vs. Day 0. Each circle represents the |PC1 score|

from a single neuron; horizontal lines denote the mean and thicker vertical lines span the

545   interquartile range. * indicates $p < 0.05$ via Wilcoxon sign-rank. Data from MFC and DMS

546   recordings in 7 animals.

Figure 4: MFC and DMS activity is distinct on FI3 vs. FI12 trials. A) An exemplar ramping neuron from the DMS; the slope of firing rate vs. time was steeper on FI3 vs. FI12 trials. B) |Slopes| for MFC (blue) and DMS (green) neurons on Day1, Day 2, and Day 3 for FI3 and FI12 trials; for both MFC and DMS |slopes| were higher on FI3 trials. C) An exemplar neuron from the DMS that fired differentially on FI3 vs. FI12 trials. D) The number of neurons with a main effect of firing rate vs. interval for the MFC and DMS. * indicates $p < 0.05$ via a chi-squared test; Data from MFC and DMS recordings in 7 animals.

Figure 5: MFC and DMS ramping is stable with two-interval performance. A) Response

times by day for FI3 and B) FI12 trials over the three days of two-interval performance. Principal

components for C) FI3 and D) FI12 trials. Ramping activity as measured by |PC1| scores for E)

FI3 trials in the MFC (blue), F) FI12 trials in the MFC, G) FI3 trials in the DMS, and H) FI12 in

the DMS. Horizontal lines denote the mean and thicker vertical gray lines span the interquartile

range. * indicates $p < 0.05$ via GLMMs. Data were from MFC and DMS recordings in 7 animals.

563

Figure 6: Decoding reveals that DMS improves temporal predictions with two-interval performance. A) We trained decoders (naïve Bayesian classifiers) to predict time from firing rate. B) Decoder performance for DMS ensembles for FI3 trials and C) for FI12 trials. Predicted time is on the y axis and observed time is on the x axis; decoded time is in yellow. D) Decoded performance for the same DMS ensembles with time-shuffled data for FI3 trials and E) FI12 trials. F). We measured decoder performance by calculating the variance explained (R^2) of predicted vs. observed time. For FI3 trials, DMS ensembles increased R^2 with two-interval performance, while MFC and G) FI12 MFC and DMS decoding was unchanged. * indicates a main effect of Day 1 → Day 3 via GLMMs; data from MFC and DMS recordings in 7 animals.

 Tables (significant values in **bold)**

Table 1: Day 0-Day 1 effects on firing rate in MFC and DMS
Model: FiringRate~Times*Area*Learning+(1|Response)+(1|Neurons)
Obs 2294280 Model R^2 0.19

Predictor	*F*	*p*
Times	111.69	**4.00E-26**
Area	6.06	**0.014**
Learning	1.77	0.183
Times:Area	686.31	**3.00E-151**
Times:Learning	96.96	**7.00E-23**
Area:Learning	1.39	0.238
Times:Area:Learning	97.5	**5.00E-23**

Table 2: Firing rate on FI3 vs. FI12 trials
Model: FiringRate~Area*Learning*Interval+(1|Response)+(1|Neurons)
Obs 3018780 Model R^2 0.15

Predictor	*F*	*p*
Area	0.48	0.49
Learning	0.58	0.45
Interval	3.89	**0.05**
Area:Learning	0.81	0.37
Area:Interval	0.32	0.57
Learning:Interval	13.16	**0.0003**
Area:Learning:Interval	6.35	**0.01**

Table 3: FI3 Day 1-Day 3 effects on firing rate
Model: FiringRate~Times*Area*Learning+(1|Response)+(1|Neurons)
Obs 533580 R^2 0.15

Predictor	*F*	*p*
Times	0	0.959
Area	0.5	0.481
Learning	0.56	0.456
Times:Area	5.41	**0.02**
Times:Learning	0.04	0.846
Area:Learning	0.69	0.407
Times:Area:Learning	1.2	0.274

Table 4: FI12 Day 1-Day 3 effects on firing rate
Model: FiringRate~Times*Area*Learning+(1|Response)+(1|Neurons)
Obs 2485200 R^2 0.16

Predictor	*F*	*p*
Times	21.84	**3.00E-06**
Area	0.84	0.361
Learning	0.16	0.69
Times:Area	34.34	**5.00E-09**
Times:Learning	7.41	**0.006**
Area:Learning	0.74	0.388
Times:Area:Learning	7.08	**0.008**

574