Increased population density depresses activity but does not influence dispersal in the snail *Pomatias*

elegans

Maxime Dahirel[1,2], Loïc Menut[1], Armelle Ansart[1]

[1]Univ Rennes, CNRS, ECOBIO (Ecosystèmes, biodiversité, évolution) - UMR 6553, F-35000 Rennes,

France

[2]INRAE, Université Côte d'Azur, CNRS, ISA, France

Abstract

Dispersal is a key trait linking ecological and evolutionary dynamics, allowing organisms to optimize

fitness expectations in spatially and temporally heterogeneous environments. Some organisms can

both actively disperse or enter a reduced activity state in response to challenging conditions, and

both responses may be under a trade-off. To understand how such organisms respond to changes in

environmental conditions, we studied the dispersal and activity behaviour in the gonochoric land

snail *Pomatias elegans*, a litter decomposer that can reach very high local densities, across a wide

range of ecologically relevant densities. We found that crowding up to twice the maximal recorded

density had no detectable effect on dispersal tendency in this species, contrary to previous results in

many hermaphroditic snails. *Pomatias elegans* is nonetheless able to detect population density; we

show they reduce activity rather than increase dispersal in response to crowding. We discuss these

results based on the ability of many land snails to enter dormancy for extended periods of time in

case of unfavourable conditions. We propose that dormancy ("dispersal in time") may be more

advantageous in species with especially poor movement abilities, even by land mollusc standards,

like *P. elegans*. Interestingly, dispersal and activity were correlated independently of density; this

dispersal syndrome may reflect a dispersal-dormancy trade-off at the individual level. Additionally,

we found snails with heavier shells relative to their size tended to be less mobile, which may reflect

physical and metabolic constraints on movement and/or survival during inactivity. We finally discuss

how the absence of density-dependent dispersal may explain why *P. elegans* is often found at very

high local densities, and the possible consequences for ecosystem functioning and litter

decomposition.

Keywords

Introduction

Dispersal, i.e. movement potentially leading to gene flow in space, is a key organismal trait influencing and linking ecological and evolutionary dynamics in sometimes complex ways (Ronce, 2007; Clobert *et al.*, 2012; Bonte & Dahirel, 2017). As the cost-benefit balance of dispersal is expected to vary depending on individual phenotype and experienced conditions, dispersal is often highly context- and phenotype-dependent (Bowler & Benton, 2005; Clobert *et al.*, 2012; Baines, Ferzoco & McCauley, 2019; Endriss *et al.*, 2019). This multicausality of dispersal likely explains why even well-studied environmental factors, like increased population density and competition, may lead to very different dispersal responses depending on populations and species (Bowler & Benton, 2005; Matthysen, 2005; Kim, Torres & Drummond, 2009; Fronhofer, Kropf & Altermatt, 2015; Baines *et al.*, 2019; Harman *et al.*, 2020). To understand the context-dependency of dispersal, it is thus necessary to study a broader range of life histories and contexts, including organisms that have alternative strategies to escape negative environmental conditions, such as dormancy, sometimes seen as a form of "dispersal in time" (Buoro & Carlson, 2014).

The land winkle *Pomatias elegans* (Müller 1774)(**Fig. 1**) is a common gastropod species in forests and hedgerows across Europe (Kerney & Cameron, 1999; Falkner *et al.*, 2001; Welter-Schultes, 2012). As a caenogastropod, it has separate sexes, contrary to the majority of land molluscs which are hermaphrodite (Barker, 2001). Its active dispersal capacities are poorly studied but expected to be very low, even by land mollusk standards (Kramarenko, 2014); average dispersal distances after 42 days are below 20 cm, according to one of the only (indirect) records available in the peer-reviewed literature (Pfenninger, 2002). Like many other land snails, it is able to survive up to several months under unfavourable conditions by entering dormancy (Falkner *et al.*, 2001). *P. elegans* is often distributed in a very aggregated way across landscapes, and when a population is present, local densities can be highly variable, going from fewer than one to several hundred individuals.m^{-2} (Pfenninger, 2002). When it is present, *P. elegans* plays a key role in ecosystem functioning, by decomposing litter and altering its chemical and physical characteristics (Coulis *et al.*, 2009; De Oliveira, Hättenschwiler & Tanya Handa, 2010).

Figure 1. *Pomatias elegans* shell showing the measurements used in the present study (photograph:

Armelle Ansart).

We here tested the hypothesis that active dispersal and activity were density-dependent in *P.*

elegans. As in many other species, including other land molluscs (Baur, 1993; Bowler & Benton, 2005;

Matthysen, 2005; Dahirel *et al.*, 2016), we expected dispersal to overall increase with population

density and associated competition levels. We expected dispersers to be more active than residents,

in agreement with general predictions about dispersal-dormancy syndromes and the pace-of-life

hypothesis (Cote *et al.*, 2010; Réale *et al.*, 2010; Buoro & Carlson, 2014). If activity merely is a

correlate of dispersal, or vice-versa, we would expect them to both change with density in the same

way. If low activity is also/instead a correlate of dormancy, we may expect the opposite response to

density: indeed, some previous snail studies showed depression of activity in response to density

(Cameron & Carter, 1979). In accordance with the phenotypic compensation hypothesis, which

predicts that risk-taking behaviour should be associated with morphological traits limiting

vulnerability to predators (e.g. Ahlgren *et al.*, 2015), we also expected dispersers/active individuals to

be better defended, by having heavier shells. Finally, we accounted for sex differences, given sex-

biased dispersal is often observed in other taxa with separated sexes (Trochet *et al.*, 2016).

Material and methods

We collected *P. elegans* snails in spring 2017 in leaf litter and at the bottom of calcareous walls

(Falkner *et al.*, 2001) in Arcais, France (approximate location: 46°17'60"N, 0°41'21"W). They were

then kept under controlled conditions (20°C, L:D 16:8) in plastic boxes (25 × 25 × 9 cm, about 500

snails.m^{-2}) with a 1 cm layer of soil and a roughly 2-3 cm thick layer of leaf litter (predominantly oaks

Quercus robur and *Q. petraea* and beech *Fagus sylvatica*) collected in urban and suburban woodlots

and hedgerows on and close to the University of Rennes Beaulieu campus. We provided humidity by

watering soil and litter every 10 to 15 days.

Dispersal experiment

In June 2017, we created two-patch systems by adapting the system we developed for slugs in

Fronhofer et al. (2018), but with a shorter inter-patch gap and longer dispersal period to account for

the more limited dispersal capacities of *P. elegans*. We divided 32×12 cm^2 transparent plastic boxes

(height: 9 cm) in two 12×12 cm^2 patches ("release" and "arrival") and one central 10×12 cm

matrix, using plastified cardboard walls. A 2×12 cm^2 slot was left open in each wall to allow snails to

leave and enter patches. Both patches were filled with watered soil and litter as in maintenance

boxes (see above), with the matrix was kept dry and empty.

We then placed between 1 and 30 snails in each "departure" patch, and first closed the patches for

24 h to let the snails habituate. We then let them free to move for 4 days, and recorded which snails

stayed in their release patch (hereafter "residents") and which snails dispersed to the arrival patch.

We tested the following group sizes: 1, 2, 3, 4, 5, 10, 15, 20, and 30 snails per box, with respectively

30, 20, 10, 9, 6, 4, 3, 3, and 2 replicates (total: 371 snails in 87 replicates). We chose these replicate

numbers to have at least 30 individuals and 2 replicates per group size. The selected group sizes

correspond to initial densities ranging from 69.4 to 2083.3 snails.m^{-2} (taking only departure patch

surface area into account), and likely provide a good coverage of natural densities. Indeed, densities

of up to 1072 detected snails.m^{-2} have been found using 25×25 cm^2 quadrats in natural populations

(Pfenninger, 2002), and a separate study indicates this species has a field detectability in the 0.5 to

0.9 range when hand collected, especially because many individuals may have buried themselves in

the substrate at any one time (Albano *et al.*, 2015). Tests were spread haphazardly over the month

due to logistical constraints (start dates ranging from June 2 to June 26); no reproduction was

observed during this interval.

Sexing

We aimed to have balanced sex ratios across densities (in line with natural populations (Boycott,

1916)). However, reproductive organs are not externally visible in *Pomatias elegans*, and body size is

not an accurate enough correlate, meaning a classification in males and females can only be done by

dissection (Creek, 1951). We nonetheless used the fact that females are on average slightly bigger

and wider than males (Boycott, 1916; Jordaens, Platts & Backeljau, 2001) to putatively sex snails

before the experiment, and then assigned equal numbers of putative males and females to densities

and boxes. To predict sex, we first measured the shell height, shell width and aperture height (**Fig. 1**)

to the nearest 0.1 mm in a set of 26 individuals not used in dispersal experiments, and sexed them by

dissection after freezing at -20°C. We used these individuals to fit an initial Linear Discriminant

Analysis (LDA) predicting sex as a function of shell characteristics, and used this LDA to predict the

sex of our focal snails. Experimental individuals were then added to the LDA dataset once dissected

after each test session, in order to improve its predictive ability. Overall, 96 individuals out of 371

(25.88%) were misclassified prior to experiments (49 out of 185 males, based on reproductive

organs, were first classified as females, 47 females out of 186 first classified as males). There was no

evidence of sex bias in the final dataset, whether overall or relative to test density (**Supplementary**

Material 1).

Post-dispersal phenotyping

Within 24h after dispersal tests, we placed each snail on a small (8 cm diameter) clean patch of wet

substrate (synthetic upholstery foam) and recorded the latency to activity, snails being defined as

active if their operculum was open and both their foot and tentacles were visible out of the shell. All

snails that had not moved after 20 minutes remained inactive, operculum closed, for at least two

hours, indicating they were not simply slower individuals, but actually behaved differently from the

active ones. Indeed, a limited number was further monitored and remained fully inactive even after 7

hours. We therefore analysed activity as a binary variable (active within 20 minutes/ not active for at

least two hours). After sexing (see above), shells were dried at room temperature before being

weighed twice (to the nearest 0.1 mg; CP224S scale, Sartorius, Göttingen, Germany). Twenty-four out

of 371 shells were partly broken during dissection and could not be weighed.

Statistical analyses

All analyses were done in a Bayesian framework, using Stan (Carpenter *et al.*, 2017; Stan

Development Team, 2018) with R (R Core Team, 2019) and the *brms* package (Bürkner, 2017).

Scripting, analysis and plotting relied on the *tidybayes*, *bayesplot*, and *patchwork* packages, as well as

the *tidyverse* family of packages (Gabry *et al.*, 2019; Kay, 2019; Pedersen, 2019; Wickham *et al.*,

2019).

We used a multivariate generalized linear mixed model (GLMM) framework (Dingemanse &

Dochtermann, 2013), which allowed us not only to draw inferences on dispersal and activity, but also

on their correlations with each other and with (relative) shell mass, while handling missing shell mass

data by in-model imputation (McElreath, 2016). A full write up of the model is given in

Supplementary Material 2. Briefly, we analysed the probability of dispersing and the probability of

activity as individual-level binary variables (dispersed/did not disperse; active/not active) using
Bernoulli GLMMs with a logit link. We tested whether dispersal or activity state depended on
experienced population density, sex and body size (defined as shell height; **Fig. 1**). The models also
included a random effect of storage box, as well as a random effect of individual identity.
Importantly, the latter was set to be identical (same values) in both the dispersal and activity models.
In effect, this means it corresponds to a latent individual "behaviour" variable that would influence
both dispersal and activity. This allowed us to estimate the effect of dispersal on activity (and vice-
versa), while accounting for the fact both are estimated with uncertainty and the causal path from
one to the other is unknown. Note that using instead dispersal status as a covariate in the activity
model leads to similar inferences (**Supplementary Material 3);** it however prevents us from doing the
analyses described in the next few sentences. To be able to separate the effects of relative shell mass
from those of body size, we used a third submodel. We fitted a linear mixed model in which ln(Shell
mass) was a function of ln(Shell height), sex, and a random effect of individual identity. The use of
repeated measures (see "Post-dispersal phenotyping") of shell mass allowed us here to separate
individual variation from residual error, and facilitated the estimation of the former. Finally,
individual-level random effects were included in a variance-covariance matrix, allowing us to
evaluate the correlation between "relative" shell mass (after accounting for body size) and the latent
behavioural variable. All continuous variables were centred and scaled to unit standard deviation
(**Supplementary Material 4**), and the "sex" variable was converted to a centred dummy numerical
variable (-0.5 = "Male"; +0.5 = "Female")(Schielzeth, 2010).

We set weakly informative priors by following McElreath (2016)'s suggestions for continuous and
proportion variables. We used a Normal(μ = 0, σ = 1.5) prior for the dispersal and activity fixed
effects intercepts, and a Normal(0, 1) prior for all the other fixed effects. We set a half-Normal(0, 1)
prior for all standard deviations (random effects as well as residual variance of shell mass), and a
LKJ(η = 2) prior for the random effects correlation matrix.

We ran four chains for 25000 iterations, with the first 5000 of each chain used for warmup. We
checked mixing graphically and confirmed chain convergence using the improved $\hat{R}$ statistic (Vehtari
et al., 2020). The chains were run longer than the default number of iterations (2000), to ensure
effective sample size was satisfactory for all parameters (bulk- and tail-ESS sensu Vehtari *et al.*, 2020
> 1000). All posterior summaries are given as: mean [95% Higher Posterior Density Intervals].

Results

144 snails out of 371 (38.81%) dispersed across all test boxes. There was no detectable effect of box
population density (**Fig. 2A**), sex or shell height on dispersal probability (**Table 1**).

Figure 2. Effect of population density (in snails per box) on *Pomatias elegans* dispersal rate (A) and probability of activity (B). Grey points are observed proportions for each test box, black points averages for each density; black curves give posterior predicted means and grey bands 95% credible intervals.

Snails exposed to higher population densities were less likely to be active afterwards (**Table 1**, **Fig. 2B**). There were no clear effects of sex or size (**Table 1**). Analysis of the latent behavioural variable shows dispersers were also more likely to be active on average, and vice-versa (mean differences on the logit scale: 0.65 [0.12; 1.19] and 0.69 [0.12; 1.26], respectively; **Fig. 3 and Supplementary Material 3**). After accounting for shell length, sex and measurement error, individual shell mass was negatively correlated with the behavioural latent variable: dispersers/active snails tended to have lighter shells, relative to their size, than residents/inactive individuals (**Fig. 4, Table 1**).

Figure 3. Relationship between dispersal status and activity post-dispersal. White bars correspond to observed frequencies, grey "eyes" to posterior predictions based on model intercepts and individual-level random effects, i.e. averaging out the effects of sex, size and population density (black dots: posterior means; segments: 95% credible intervals).

Discussion

We studied the relationship between *Pomatias elegans* dispersal and population density over most of the known range of natural densities (Pfenninger, 2002), yet found no evidence of density-dependent dispersal. There was however a negative relationship between density and activity, and dispersal, activity and (relative) shell mass co-vary in this species.

Figure 4. Relationship between dispersal status, activity post-dispersal, and mean predicted shell mass, that is relative to the shell mass predicted for an otherwise average snail of same shell height and sex (black dots: posterior means; segments: 95% credible intervals).

We additionally found no evidence of sex bias in dispersal probability or activity. While sex-biased dispersal is abundantly documented in non-molluscan groups (Bowler & Benton, 2005; Trochet *et al.*, 2016), there are numerous species with no bias. Broadly, sex-dependent dispersal is predicted to arise when there are sex-differences in spatio-temporal fitness variation due to e.g. sex differences in resource use and/or in local mate competition (Hovestadt, Mitesser & Poethke, 2014; Trochet *et al.*, 2016; Li & Kokko, 2019). Our results, in light of dispersal theory, imply that male and female *P. elegans* use resources similarly and that they do not present polygyny or any mating system leading to predictable sex differences in spatio-temporal fitness variation. While sex ratio is, overall, balanced in this species (Boycott, 1916), the other predictions remain to be tested by the collection of further natural history data. We should additionally note that our experimental setup (one measure of dispersal, after 4 days, and groups that were sex-balanced on average) may not have allowed us to detect finer sex differences in dispersal, i.e. in timing (Li & Kokko, 2019), or conditional strategies dependent on locally experienced sex-ratio (Hovestadt *et al.*, 2014).

The absence of density-dependent dispersal was more surprising, as other land mollusc species have been shown to alter their dispersal in response to population density (Oosterhoff, 1977; Hamilton & Wellington, 1981; Livshits, 1985; Baur, 1993; Dahirel *et al.*, 2016), and density-dependent dispersal is common in many other groups (Bowler & Benton, 2005; Matthysen, 2005; Mathieu *et al.*, 2010;

Bitume *et al.*, 2013; Fronhofer *et al.*, 2015; De Bona *et al.*, 2019, among many others). Additionally,
the costs of competition, which can trigger dispersal, are well-documented in land molluscs
(Oosterhoff, 1977; Hamilton & Wellington, 1981; Dan & Bailey, 1982). However, density-independent
dispersal also exists in this group, and despite population density likely being one of the most studied
drivers of dispersal, we do not have a systematic understanding of how species traits or
environmental effects drive differences in the dispersal-density reaction norm (Baines *et al.*, 2019). *P.*
elegans, contrary to most land molluscs has separate sexes, and differences in reproductive systems
may drive the evolution of different dispersal strategies (Eppley & Jesson, 2008). Indeed, all things
equal, hermaphrodites are *a priori* more likely than species with separate sexes to find a mate in
small populations (Tomlinson, 1966). This means the density threshold at which the costs of
competition outweigh the benefits of grouping (Kim *et al.*, 2009; Fronhofer *et al.*, 2015) and the costs
of dispersal (Bonte *et al.*, 2012) is likely lower for hermaphrodites. We hypothesize that context
being equal (including e.g. movement costs), this may make hermaphrodites more likely than
gonochoric species to exhibit positive density-dependent dispersal over a range of natural densities.
In any case, our study shows that further comparisons between hermaphroditic and gonochoric
species sharing ecological and life-history characteristics are crucially needed to clarify the role of
reproductive system in shaping context-dependent dispersal.

Alternatively (but not exclusively), the absence of density response in dispersal may reflect the fact
Pomatias snails possess other mechanisms to respond to unfavorable densities. Indeed, by contrast
with dispersal, activity propensity did decrease with increasing population density (**Fig. 2B**). Like
many other shelled land gastropods, *P. elegans* is able of reduced activity and dormancy when
context is unfavourable (up to several months in response to dry conditions; Falkner *et al.*, 2001).
Dormancy has mostly been studied with respect to abiotic factors in molluscs (Cook, 2001), but
activity reduction in response to increased competition has been observed in several species
(Oosterhoff, 1977; Cameron & Carter, 1979; Dan & Bailey, 1982). This is generally thought to result
from interference (mediated through mucus and/or faeces) by the most competitive individuals,
which remain active. However, when the experimental/observational setup also allowed for
dispersal, species that reduced activity in response to density (*Cepaea nemoralis, Cornu aspersum,*
Xerocrassa seetzenii) did however also exhibit density-dependent dispersal (Oosterhoff, 1977;
Shachak, Leeper & Degen, 2002; Dahirel *et al.*, 2016). Given the very limited dispersal capacities of *P.*
elegans (Pfenninger, 2002) even compared to other snails (Kramarenko, 2014), the least competitive
Pomatias snails may well be more successful reducing activity to wait out more favourable
conditions, in a form of "temporal dispersal", than by attempting to find better by moving away
(Buoro & Carlson, 2014). The fact there was no effect of body size on activity suggests the loss of

activity is not simply an inhibition of activity by competitive individuals, leading to competitive

exclusion. Indeed, in that case we would have expected larger individuals to stay active at high

densities while smaller less competitive snails are inhibited (Dan & Bailey, 1982). Note that we only

studied emigration, the first step of dispersal, and density-dependent responses may also happen

during transience (Shachak et al., 2002; Poethke, Gros & Hovestadt, 2011; Dahirel et al., 2016). In any

case, our results show that studying the effect of population density across life histories may help

shed light on its influence on dispersal in space and/or time.

All three phenotypic traits we studied were linked. First, dispersers were more likely to be active

post-dispersal test, and vice versa (**Fig. 3**, **Supplementary Material 3**), independently of density

context. These results line up with predictions on dispersal syndromes and the pace-of-life

hypothesis (Cote et al., 2010; Réale et al., 2010), which predicts a metabolically-underpinned link

between activity and risk taking behaviour (such as dispersal). They can also be interpreted as a

trade-off between dispersal in space and in time (Buoro & Carlson, 2014), in which individuals that do

not disperse are more likely to be dormant at any point in time, even after accounting for density

effects. Second, we found that more mobile snails (higher activity and/or dispersal) had on average

lighter shells (**Fig. 4**, **Table 1**), compared to inactive/resident snails. This result goes against our

predictions based on the phenotypic compensation hypothesis, which states that risk-taking

individuals should have better morphological defences against predators (Hulthén et al., 2014;

Ahlgren et al., 2015; Kuo, Irschick & Lailvaux, 2015). However, phenotypic compensation is not a hard

rule (De Winter et al., 2016), and while thicker/heavier shells are harder to break (see e.g. Rosin et

al., 2013), variation in shell characteristics depends on many abiotic and biotic parameters

(Goodfriend, 1986). Indeed, thicker shells may also better conserve humidity (Goodfriend, 1986),

which may be adaptive for snails more prone to be dormant for extended periods of time. Heavier

shells are likely to increase the costs of movement (Herreid & Full, 1986); given the already high costs

of movement in land molluscs (Denny, 1980), this may explain why mobile individuals have lighter

shells, despite the potentially increased predation risk. Experimental manipulations of the costs and

benefits of heavier shells (by e.g. testing snails under varying humidity conditions, and/or by adding

weights to shells), may help disentangle the underpinnings of this dispersal/shell mass syndrome.

Although P. elegans is overall much less abundant than other macroinvertebrate decomposers at the

landscape scale (David, 1999), it can locally reach very high densities (Pfenninger, 2002). Our study

shows that increased densities do not trigger increased movement from crowded to empty habitats

in this species. This helps explain the maintenance of this strong spatial heterogeneity and high local

densities, as dispersal cannot play its "equalizing" role as well. Given decomposition by P. elegans can

have marked effects on litter characteristics (Coulis et al., 2009; De Oliveira et al., 2010), and

facilitate decomposition by other co-occuring macroinvertebrates (De Oliveira *et al.*, 2010), this

absence of a common context-dependency in dispersal may have broader implications for local

heterogeneity in soil ecosystem functioning and soil communities.

Acknowledgments

294 We thank Valérie Briand for her help in obtaining some of the cited scientific articles.

Data availability

296 Data and analysis script are archived on GitHub (https://github.com/mdahirel/pomatias-dispersal-
297 2017) and Zenodo (doi: 10.5281/zenodo.3691977) (latest release at time of submission: v1.0)

References

Ahlgren, J., Chapman, B.B., Nilsson, P.A. & Brönmark, C. (2015). Individual boldness is linked to
protective shell shape in aquatic snails. *Biology Letters* **11**, 20150029.

Albano, P.G., Strazzari, G., D'Occhio, P. & Succetti, F. (2015). Field estimates of detectability and site
occupancy show that northern Italy forest molluscs are spatially rare and poorly detectable.
Italian Journal of Zoology **82**, 592–608.

Baines, C.B., Ferzoco, I.M.C. & McCauley, S.J. (2019). Phenotype-by-environment interactions
influence dispersal. *J. Anim. Ecol.* **88**, 1263–1274.

Barker, G.M. (2001). Gastropods on land: phylogeny, diversity and adaptive morphology. In *The
biology of terrestrial molluscs*: 1–146. Barker, G.M. (Ed.). Wallingford, UK: CABI.

Baur, A. (1993). Effects of food availability and intra-and interspecific interaction on the dispersal
tendency in the land snail *Chondrina clienta*. *J. Zool.* **230**, 87–100.

Bitume, E.V., Bonte, D., Ronce, O., Bach, F., Flaven, E., Olivieri, I. & Nieberding, C.M. (2013). Density
and genetic relatedness increase dispersal distance in a subsocial organism. *Ecol. Lett.* **16**,
430–437.

Bonte, D. & Dahirel, M. (2017). Dispersal: a central and independent trait in life history. *Oikos* **126**,
472–479.

Bonte, D., Van Dyck, H., Bullock, J.M., Coulon, A., Delgado, M., Gibbs, M., Lehouck, V., Matthysen, E.,
Mustin, K., Saastamoinen, M., Schtickzelle, N., Stevens, V.M., Vandewoestijne, S., Baguette,
M., Barton, K., Benton, T.G., Chaput-Bardy, A., Clobert, J., Dytham, C., Hovestadt, T., Meier,
C.M., Palmer, S.C.F., Turlure, C. & Travis, J.M.J. (2012). Costs of dispersal. *Biol. Rev.* **87**, 290–
312.

Bowler, D.E. & Benton, T.G. (2005). Causes and consequences of animal dispersal strategies: relating
individual behaviour to spatial dynamics. *Biological Reviews* **80**, 205–225.

Boycott, A.E. (1916). On sexual characteristics in the shell and radula of *Pomatias elegans* (Müller).
Proceedings of the Malacological Society of London. **v.12-13 (1916-1919)**.

Buoro, M. & Carlson, S.M. (2014). Life-history syndromes: Integrating dispersal through space and
time. *Ecology Letters* **17**, 756–767.

Bürkner, P.-C. (2017). brms: an R package for bayesian multilevel models using Stan. *Journal of*
Statistical Software **80**, 1–28.

Cameron, R.A.D. & Carter, M.A. (1979). Intra- and interspecific effects of population density on
growth and activity in some helicid land snails (Gastropoda: Pulmonata). *The Journal of*
Animal Ecology **48**, 237.

Carpenter, B., Gelman, A., Hoffman, M.D., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M., Guo,
J., Li, P. & Riddell, A. (2017). Stan: a probabilistic programming language. *Journal of Statistical*
Software **76**, 1–32.

Clobert, J., Baguette, M., Benton, T.G. & Bullock, J.M. (Eds.). (2012). *Dispersal ecology and evolution*.
Oxford, UK: Oxford University Press.

Cook, A. (2001). Behavioural ecology: on doing the right thing, in the right place at the right time. In
The biology of terrestrial molluscs: 447–488. Barker, G.M. (Ed.). Wallingford, UK: CABI.

Cote, J., Clobert, J., Brodin, T., Fogarty, S. & Sih, A. (2010). Personality-dependent dispersal:
characterization, ontogeny and consequences for spatially structured populations.
Philosophical Transactions of the Royal Society of London B: Biological Sciences **365**, 4065–
4076.

Coulis, M., Hättenschwiler, S., Rapior, S. & Coq, S. (2009). The fate of condensed tannins during litter
consumption by soil animals. *Soil Biology and Biochemistry* **41**, 2573–2578.

Creek, G.A. (1951). The reproductive system and embryology of the snail *Pomatias elegans* (Müller).
Proceedings of the Zoological Society of London **121**, 599–640.

Dahirel, M., Vardakis, M., Ansart, A. & Madec, L. (2016). Density-dependence across dispersal stages
in a hermaphrodite land snail: insights from discrete choice models. *Oecologia* **181**, 1117–
1128.

Dan, N. & Bailey, S.E.R. (1982). Growth, mortality, and feeding rates of the snail *Helix aspersa* at
different population densities in the laboratory, and the depression of activity of helicid
snails by other individuals, or their mucus. *J. Mollus. Stud.* **48**, 257–265.

David, J.-F. (1999). Abundance, biomass and functional structure of the saprophagous macrofauna in
the litter and soil of Mediterranean oak forests. *Pedobiologia* **43**, 319–327.

De Bona, S., Bruneaux, M., Lee, A.E.G., Reznick, D.N., Bentzen, P. & López-Sepulcre, A. (2019). Spatio-
temporal dynamics of density-dependent dispersal during a population colonisation. *Ecology*
Letters.

De Oliveira, T., Hättenschwiler, S. & Tanya Handa, I. (2010). Snail and millipede complementarity in
decomposing Mediterranean forest leaf litter mixtures: Snail and millipede interactions.
Functional Ecology **24**, 937–946.

De Winter, G., Ramalho Martins, H., Trovo, R.A. & Chapman, B.B. (2016). Knights in shining armour
are not necessarily bold: defensive morphology correlates negatively with boldness, but
positively with activity, in wild threespine stickleback,. *Evolutionary Ecology Research* **13**.

Denny, M. (1980). Locomotion: the cost of gastropod crawling. *Science* **208**, 1288–1290.

Dingemanse, N.J. & Dochtermann, N.A. (2013). Quantifying individual variation in behaviour: mixed-
365   effect modelling approaches. *Journal of Animal Ecology* **82**, 39–54.

366   Endriss, S.B., Vahsen, M.L., Bitume, E.V., Monroe, J.G., Turner, K.G., Norton, A.P. & Hufbauer, R.A.
367   (2019). The importance of growing up: juvenile environment influences dispersal of
368   individuals and their neighbours. *Ecology Letters* **22**, 45–55.

369   Eppley, S.M. & Jesson, L.K. (2008). Moving to mate: the evolution of separate and combined sexes in
370   multicellular organisms. *Journal of Evolutionary Biology* **21**, 727–736.

371   Falkner, G., Obrdlik, P., Castella, E. & Speight, M.C.D. (2001). *Shelled Gastropoda of western Europe*.
372   München, Germany: Friedrich Held Gesellschaft.

373   Fronhofer, E.A., Kropf, T. & Altermatt, F. (2015). Density-dependent movement and the
374   consequences of the Allee effect in the model organism *Tetrahymena*. *Journal of Animal*
375   *Ecology* **84**, 712–722.

376   Fronhofer, E.A., Legrand, D., Altermatt, F., Ansart, A., Blanchet, S., Bonte, D., Chaine, A., Dahirel, M.,
377   Laender, F.D., Raedt, J.D., Gesu, L. di, Jacob, S., Kaltz, O., Laurent, E., Little, C.J., Madec, L.,
378   Manzi, F., Masier, S., Pellerin, F., Pennekamp, F., Schtickzelle, N., Therry, L., Vong, A.,
379   Winandy, L. & Cote, J. (2018). Bottom-up and top-down control of dispersal across major
380   organismal groups. *Nature Ecology & Evolution* **2**, 1859.

381   Gabry, J., Simpson, D., Vehtari, A., Betancourt, M. & Gelman, A. (2019). Visualization in Bayesian
382   workflow. *Journal of the Royal Statistical Society: Series A (Statistics in Society)* **182**, 389–402.

383   Goodfriend, G.A. (1986). Variation in land-snail shell form and size and its causes: a review.
384   *Systematic Zoology* **35**, 20.

385   Hamilton, P.A. & Wellington, W.G. (1981). The effects of food and density of the movement of *Arion*
386   *ater* and *Ariolimax columbianus* (Pulmonata: Stylommatophora) between habitats. *Res Popul*
387   *Ecol* **23**, 299–308.

388   Harman, R.R., Ii, J.G., Shivaji, R. & Cronin, J.T. (2020). Frequency of occurrence and population-
389   dynamic consequences of different forms of density-dependent emigration. *The American*
390   *Naturalist*.

391   Herreid, C.F. & Full, R.J. (1986). Energetics of hermit crabs during locomotion: the cost of carrying a
392   shell. *Journal of Experimental Biology* **120**, 297–308.

393   Hovestadt, T., Mitesser, Oliver & Poethke, H.-J. (2014). Gender-specific emigration decisions sensitive
394   to local male and female density. *The American Naturalist* **184**, 38–51.

395   Hulthén, K., Chapman, B.B., Nilsson, P.A., Hollander, J. & Brönmark, C. (2014). Express yourself: bold
396   individuals induce enhanced morphological defences. *Proceedings of the Royal Society B:*
397   *Biological Sciences* **281**, 20132703.

398   Jordaens, K., Platts, E. & Backeljau, T. (2001). Genetic and morphological variation in the land winkle
399   *Pomatias elegans* (Müller) (Caenogastropoda: Pomatiasidae). *J Molluscan Stud* **67**, 145–152.

Kay, M. (2019). *tidybayes: Tidy Data and Geoms for Bayesian Models*.

Kerney, M.-P. & Cameron, R.-A.-D. (1999). *Guide des escargots et limaces d'Europe.* Lonay (Suisse):
Delachaux et Niestlé.

Kim, S.-Y., Torres, R. & Drummond, H. (2009). Simultaneous positive and negative density-dependent
dispersal in a colonial bird species. *Ecology* **90**, 230–239.

Kramarenko, S. (2014). Active and passive dispersal of terrestrial mollusks: a review. *Ruthenica* **24**.

Kuo, C.-Y., Irschick, D.J. & Lailvaux, S.P. (2015). Trait compensation between boldness and the
propensity for tail autotomy under different food availabilities in similarly aged brown anole
lizards. *Functional Ecology* **29**, 385–392.

Li, X.-Y. & Kokko, H. (2019). Intersexual resource competition and the evolution of sex-biased
dispersal. *Front. Ecol. Evol.* **7**.

Livshits, G.M. (1985). Ecology of the terrestrial snail *Brephulopsis bidens* (Pulmonata: Enidae):
mortality, burrowing and migratory activity. *Malacologia* **26**, 213–223.

Mathieu, J., Barot, S., Blouin, M., Caro, G., Decaëns, T., Dubs, F., Dupont, L., Jouquet, P. & Nai, P.
(2010). Habitat quality, conspecific density, and habitat pre-use affect the dispersal
behaviour of two earthworm species, *Aporrectodea icterica* and *Dendrobaena veneta*, in a
mesocosm experiment. *Soil Biology and Biochemistry* **42**, 203–209.

Matthysen, E. (2005). Density-dependent dispersal in birds and mammals. *Ecography* **28**, 403–416.

McElreath, R. (2016). *Statistical Rethinking: A Bayesian Course with Examples in R and Stan.* 1st edn.
Boca Raton: Chapman and Hall/CRC.

Oosterhoff, L.M. (1977). Variation in growth rate as an ecological factor in the landsnail *Cepaea
nemoralis* (L.). *Netherlands Journal of Zoology* **27**, 1–132.

Pedersen, T.L. (2019). *patchwork: The Composer of Plots.*

Pfenninger, M. (2002). Relationship between microspatial population genetic structure and habitat
heterogeneity in *Pomatias elegans* (O.F. Müller 1774) (Caenogastropoda, Pomatiasidae).
Biological Journal of the Linnean Society **76**, 565–575.

Poethke, H.J., Gros, A. & Hovestadt, T. (2011). The ability of individuals to assess population density
influences the evolution of emigration propensity and dispersal distance. *Journal of
Theoretical Biology* **282**, 93–99.

R Core Team. (2019). *R: a language and environment for statistical computing.* Vienna, Austria: R
Foundation for Statistical Computing.

Réale, D., Garant, D., Humphries, M.M., Bergeron, P., Careau, V. & Montiglio, P.-O. (2010).
Personality and the emergence of the pace-of-life syndrome concept at the population level.
Philosophical Transactions of the Royal Society of London B: Biological Sciences **365**, 4051–
4063.

Ronce, O. (2007). How does it feel to be like a rolling stone? Ten questions about dispersal evolution.
Annual Review of Ecology, Evolution, and Systematics **38**, 231–253.

Rosin, Z.M., Kobak, J., Lesicki, A. & Tryjanowski, P. (2013). Differential shell strength of *Cepaea*
nemoralis colour morphs—implications for their anti-predator defence. *Naturwissenschaften*
100, 843–851.

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients.
Methods in Ecology and Evolution **1**, 103–113.

Shachak, M., Leeper, A. & Degen, A. (2002). Effect of population density on water influx and
distribution in the desert snail *Trochoidea seetzenii*. *Ecoscience* **9**, 287–292.

Stan Development Team. (2018). *RStan: the R interface to Stan*.

Tomlinson, J. (1966). The advantages of hermaphroditism and parthenogenesis. *Journal of*
Theoretical Biology **11**, 54–58.

Trochet, A., Courtois, E.A., Stevens, V.M., Baguette, M., Chaine, A., Schmeller, D.S., Clobert, J. &
Wiens, J.J. (2016). Evolution of sex-biased dispersal. *The Quarterly Review of Biology* **91**, 297–
320.

Vehtari, A., Gelman, A., Simpson, D., Carpenter, B. & Bürkner, P.-C. (2020). Rank-normalization,
folding, and localization: An improved \widehat{R} for assessing convergence of MCMC.
arXiv:1903.08008 [stat].

Welter-Schultes, F. (2012). *European non-marine molluscs, a guide for species identification*.
Göttingen: Planet Poster Editions.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A.,
Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J.,
Robinson, D., Seidel, D., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K. & Yutani, H.
(2019). Welcome to the Tidyverse. *Journal of Open Source Software* **4**, 1686.

Tables

Table 1. Posterior means [and 95% credible intervals] for the parameters of the multivariate model

of dispersal, activity and shell mass. See **Supplementary Material 2** for the full write-up of the model,

and **Supplementary Material 4,** for original means and SDs of transformed variables (*N* = 371 snails,

except for Shell mass, *N* = 347 snails, measured twice)

	Dispersal probability	Activity probability	Shell mass (ln-transformed then scaled)
Fixed effects			
Intercept	-0.57 [-0.94 ; -0.21]	0.66 [0.29 ; 1.06]	-0.01 [-0.09 ; 0.06]
Population density (scaled)	-0.01 [-0.37; 0.38]	-0.60 [-1.00 ; -0.21]	--
Sex (effect of being female)	-0.16 [-0.77 ; 0.47]	0.24 [-0.41 ; 0.86]	0.18 [-0.02 ; 0.37]
Body size (shell height, ln-transformed then scaled)	0.01 [-0.31 ; 0.33]	0.18 [-0.16 ; 0.51]	0.62 [0.52 ; 0.71]
Random effects			
Box-level SD	0.64 [0.01 ; 1.13]	0.68 [0.11; 1.22]	--
Individual-level SD	0.93 [0.43 ; 1.41]		0.74 [0.68 ; 0.79]
Correlation between box-level SDs	0.05 [-0.69; 0.74]		--
Correlation between individual-level SDs	-0.39 [-0.66; -0.14]		
Residual SD	--	--	0.03 [0.03 ; 0.03]

Figure legends

Figure 1. *Pomatias elegans* shell showing the measurements used in the present study (photograph:
Armelle Ansart).

Figure 2. Effect of population density (in snails per box) on *Pomatias elegans* dispersal rate (A) and
probability of activity (B). Grey points are observed proportions for each test box, black points
averages for each density; black curves give posterior predicted means and grey bands 95% credible
intervals.

Figure 3. Relationship between dispersal status and activity post-dispersal. White bars correspond to
observed frequencies, grey "eyes" to posterior predictions based on model intercepts and individual-
level random effects, i.e. averaging out the effects of sex, size and population density (black dots:
posterior means; segments: 95% credible intervals).

Figure 4. Relationship between dispersal status, activity post-dispersal, and mean predicted shell
mass, that is relative to the shell mass predicted for a snail of same shell height and sex (black dots:
posterior means; segments: 95% credible intervals).

Shell height

Peristome height

Shell width

1 mm