In Vivo Analysis of RNA Proximity Proteomes Using RiboPro

Xianzhi Lin[1][#], Kate Lawrenson[1,2][#]

[1] Women's Cancer Program at Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA.

[2] Center for Bioinformatics and Functional Genomics, Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA.

#Correspondence to: xianzhi.lin@cshs.org (XL) or kate.lawrenson@cshs.org (KL).

Functional and mechanistic annotation of uncharacterized long noncoding RNAs is challenging and often requires identification of interacting proteins. Here we report RiboPro, a flexible method that leverages the RNA-binding specificity of inactive Cas13b and proximity labeling activity of peroxidase APEX to permit rapid and unbiased discovery of target RNA binding proteins *in vivo*. RiboPro of poly(A)+ RNA reveals insights into poly (A)+ RNA nucleocytoplasmic transport, localization, turnover, and higher-order structure.

LncRNA transcripts do not encode protein, but they nonetheless play myriad roles in physiology and disease[1]. The spectrum of RNA binding proteins (RBPs) associated with a lncRNA transcript will largely dictate lncRNA activities[2]. RBPs control critical RNA activities including RNA editing, processing, stability, translation and transport[3]. Therefore, cataloguing the interacting proteins of lncRNAs is critical to understanding their function. Currently, functional and mechanistic characterization of lncRNAs is hampered by the limited set of research methods available[4,5]. We developed RiboPro (<u>Ribo</u>nucleic acid <u>pro</u>ximity protein labelling) method to identify proteins associated with a target RNA *in vivo* without cross-linking or tagging RNA with aptamer sequences – both of which can lead to false positive interactions not seen under normal physiologic conditions.

Inspired by the applications of type VI Cas13 members[6,7] and proximity labeling capacity of the engineered peroxidase APEX[8], we developed the RiboPro approach, based on a catalytically dead Cas13 (dCas13) fused to APEX (Fig. 1a). In the presence of a sequence-specific guide RNA (gRNA), the dCas13-APEX fusion protein is directed to a target RNA. APEX in the chimera will biotinylate RNA proximity proteins within a few nanometers of gRNA

target sequence[9]. The biotinylated proteins can then be readily enriched using streptavidin

beads and analyzed by mass spectrometry profiling or western blotting (Fig. 1a). We fused

catalytically dead Cas13b from *Prevotella sp. P5-125* (dPspCas13b)[6] to the engineered

soybean ascorbate peroxidase APEX2[8] with both Flag and HA tags (Fig. 1b). The expression

of the fusion protein dPspCas13b-Flag-APEX2-HA (from hereon is called the RiboPro protein)

was validated by western blotting using both anti-Flag and anti-HA antibodies after transfection

into HEK293T cells (Fig. 1c). The RiboPro protein is expressed in both the cytoplasm and the

nucleus (Fig. 1d). To verify the enzymatic activity of APEX2 in the chimera, HEK293T cells

transfected with the RiboPro plasmid were treated with biotin-phenol in the presence or

absence of H_2O_2. Biotinylated proteins were only detected when both biotin-phenol and H_2O_2

were added, indicating that the APEX2 retains its enzymatic activity in the RiboPro protein

(Fig. 1e).

We applied RiboPro to well-known *U1* small nuclear ribonucleoprotein (snRNP) to test

whether RiboPro would identify proteins known to directly bind to *U1*. Three guide RNAs

(gRNAs) were designed to target different regions of *U1* small nuclear RNA (snRNA) (U1-1,

U1-2, and U1-3) (Fig. 1f). Following co-transfection of *U1* gRNAs into HEK293T cells with wild-

type PspCas13b, expression of *U1* snRNA was significantly downregulated compared with

cells treated with a nontargeting control (NTC) gRNA, while the expression of *18S* rRNA was

not affected (Fig. 1g), indicating that all three *U1* gRNAs efficiently direct PspCas13b to the

target transcript. We then tested whether these *U1* gRNAs can direct the RiboPro protein to

bind *U1* snRNA. In RNA immunoprecipitation experiments, the RiboPro protein was efficiently

retrieved by anti-HA antibody but not isotype control IgG (Fig. 1h). *U1* snRNA was enriched

around 2-fold for *U1* gRNAs compared with the NTC gRNA (Fig. 1i), indicating that the

RiboPro protein was selectively enriched at *U1*. U1-70k, a known *U1* direct binding partner, was significantly enriched by RiboPro (Fig. 1j). *U1* proximity proteins were profiled using mass spectrometry with label-free quantification and in total, 8 out of 9 *U1* direct binding proteins were enriched by RiboPro (Fig. 1k and Supplementary Table 1). Together these data indicate that RiboPro is able to efficiently identify direct RNA-binding proteins of a target RNA of interest.

We next employed RiboPro to examine the RNA proximity proteome of poly (adenosine) (poly[A]) tails (Supplementary Fig.1a). Since oligomers of thirty nucleotides of poly (uridine) (poly[U]) are rare in the human transcriptome[10], poly(U) gRNA was used as control for poly(A) gRNA. A poly(A) gRNA or poly(U) gRNA was co-transfected into HEK293T cells with the RiboPro plasmid and then RiboPro was performed. Mass spectrometry profiling of two independent biological experiments uncovered 910 and 1295 proteins in the poly(A) tail RNA proximity proteome, and 676 and 1282 proteins in the poly(U) RNA proximity proteome, respectively (Fig. 2a). In comparison to the poly(U) pulldowns, 762 and 925 proteins were enriched in the two poly(A) pulldowns (Fig. 2a), 304 of which are shared by both replicates (label-free quantification fold change ≥ 2 in poly[A] compared with poly[U]) (Supplementary Fig. 1b). In contrast, only 106 and 56 proteins were enriched in poly(U) pulldowns relative to poly(A) pulldowns (Fig. 2a) with no proteins shared by both replicates when applying the same threshold (Supplementary Fig. 1b). When considering subunits from the same protein complex, there are 445 protein from replicate 1 and 452 proteins from replicate 2 were shared (Fig. 2b), indicating the two replicates are highly reproducible at the complex level (P=1, Binomial test). In contrast, in the poly(U) pulldowns, only 8 proteins from each replicate were shared (Fig. 2b), suggesting poor reproducibility between replicates in our negative control (P= 2.02×10^{-05},

Binomial test). Only 38 proteins overlapped between the poly(A) and poly(U) RNA proximity

proteome, indicating these sets of proteins are distinct (P= 1.90×10⁻¹⁷, Binomial test) (Fig. 2c).

Gene ontology (GO) analysis was performed on the 1347 proteins uniquely associated with

poly(A) tails (Supplementary Table 2). Thirteen of the top 20 enriched terms were RNA-related

and the majority were poly(A)+ RNA-related (Fig. 2d). Indeed, 360 proteins in poly(A) tail RNA

proximity proteins are poly(A)+ RNA binding proteins previously discovered in HEK293 cells[11]

(Fig. 2e) and 150 more poly(A)+ RNA binding proteins in HeLa or MCF7 cells[12,13]

(Supplementary Fig. 1c). We curated all known RBPs from literature (Supplementary Table 3,

see Methods and references therein) and found that 1230/1347 (91.31%) of the poly(A) tail

RNA proximity proteins identified by RiboPro are known RBPs (Fig. 2f) and 540/1230 (43.90%)

have previously been described as non-poly(A)-RBPs[14] (Supplementary Fig. 1d), suggesting

that more RBPs can associate with poly(A)+ RNA than previously appreciated. GO analysis of

117 novel poly(A) tail RNA binding proteins revealed by RiboPro suggest poly(A)+ RNA may

play a role in mitochondrial cristae formation, the Mediator complex and other novel processes

(Supplementary Fig. 1e).

   We next examined whether known categories of poly(A)-associated proteins are identified

among poly(A) tail RNA proximity proteome. Polyadenylation occurs co-transcriptionally and

requires cleavage and polyadenylation complex (CPA) that includes cleavage and

polyadenylation specificity factors (CPSFs), cleavage stimulating factors (CstFs), cleavage

factor I and II (CFI & CFII) and poly(A) polymerase[15]. CPA members for poly(A)+RNA including

CPSF1, CPSF3, CstF1 and CPSF7 are present in poly(A) tail RNA proximity proteome, while

factors specific for poly(A)- RNA such as SLBP and ZFP100 (ZNF473) are absent (Fig. 2g).

Among the 12 expressed known poly(A) binding proteins, RiboPro identifies 7 (58%) (Fig. 2g).

The poly(A) tail is part of the 3' untranslated region (UTR), and RiboPro identifies 30 out of 41

(73%) expressed known 3'UTR binding proteins (Fig. 2g). In addition, poly(A) RiboPro detects

3 out of 8 expressed known cap binding proteins (Fig. 2g), 15 out of 19 expressed known

5'UTR binding proteins (Fig. 2g), and 25 translation initiation factors (Fig. 2g), including EIF4G.

Finally, poly(A) tail RiboPro unearths 31 out of 33 small ribosomal subunits (RPSs) and 46 out

of 49 large ribosomal subunits (RPLs) (Fig. 2g). These data support that 5' and 3' of poly(A)+

mRNA form a closed-loop structure during translation[16] and provide novel evidence for the

importance of poly(A) tail in mRNA translation[17]. Interestingly, the poly(A) tail RNA proximity

proteome also enriches 13 tRNA ligases (Supplementary Table 2), suggesting poly(A) tails

may have a role in tRNA charging or tRNA ligases may have unknown function in regulating

poly(A).

Adding poly(A) tails by adenylation stabilizes RNA molecules and protects transcripts from

degradation[17], while deadenylation triggers decapping and decay[18]. Since the poly(A) gRNA is

30 nt in length, poly(A) RiboPro will identify proteins proximal to any poly(A) tail over 30

nucleotides in length (Supplementary Fig. 1a), which will include stable transcripts, transcripts

actively undergoing adenylation, and transcripts undergoing deadenylation. In addition to

proteins related to translation (Fig. 2g), poly(A) RiboPro uncovers dozens of proteins linked

with RNA decay, including deadenylase complex members[19], exosome components[20],

nonsense-mediated decay proteins[21], decapping factors, and proteins involved in AU-rich

element-containing mRNA degradation[22] (Fig. 2h). The decapping enzyme and 5'-3'

exonuclease are not enriched in the poly(A) tail RNA proximity proteome, suggesting that

mRNAs no longer form closed-loop structures once the decapping process is triggered by

deadenylation.

Poly(A) tails are also important for RNA nuclear export[23], and poly(A) RiboPro detects 10

proteins involved in RNA nuclear export (Fig. 2i), including XPO1 and NXF1. The observation

that two 5' end-associated nuclear components, NCBP1 and TREX complex[24], also associate

with poly(A) (Fig. 2i), suggests that poly(A)+ RNA may also form closed-loop structures during

nuclear export. Localization signals/zip code binding protein IGF2BP1 and 9 motor proteins

used in mRNA transport are also present in poly(A) tail RNA proximity proteome (Fig. 2i),

suggesting direct involvement of the poly(A) tail in RNA transport. Furthermore, poly(A)

associated proteins includes microfilaments (e.g. ACTG1 and ACTN4) and membrane proteins

(e.g. SLC3A2 and MPZL1) related to mRNA anchorage (Fig. 2i, supplementary Table 2),

suggesting that the poly(A) tail participates in multiple mechanisms for RNA localization[25,26].

Together, these data authenticate that poly(A) RiboPro recovers *bona fide* proteins associated

with poly(A) tails and also reveals putative novel poly(A) RBPs not detected by alternative

methods.

Data from both *U1* and poly(A) tail RNA proximity proteomic analysis using RiboPro

demonstrate that this method represents a valuable new tool for probing RNA binding proteins

in vivo that is expected to shed light on the function and mechanistic roles of lncRNA in diverse

biological systems.

Acknowledgements

This work was supported by the National Cancer Institute (K99/R00-CA184415 and R01-

CA207456 to K.L.), the Cedars-Sinai Samuel Oschin Comprehensive Cancer Institute (Cancer

Biology Grant 231433 to K.L.) and the Ovarian Cancer Research Alliance (Ann and Sol

Schreiber Mentored Investigator Award 458799 to X.L.). We thank Drs. Wei Yang and Bo Zhou

at the Cedars-Sinai Medical Center Biomarker Discovery Platform Core for label-free

quantitative mass spectrometry analysis. We also thank members of the Lawrenson lab for

stimulating discussions.

Competing interests

The authors declare no competing interests.

Methods

Plasmids and cloning. pC0046-EF1a-PspCas13b-NES-HIV was a gift from Dr. Feng Zhang (Addgene plasmid #

103862). pCMV-dPspCas13b-Flag-APEX2-HA was constructed by replacing ADAR2DD-delta-984-1090 in

pC0053-CMV-dPspCas13b-GS-ADAR2DD (E488Q)-delta-984-1090 (a gift from Feng Zhang, Addgene plasmid #

103869) with Flag-APEX2-HA subcloned from pcDNA3-APEX2-NES (a gift from Dr. Alice Ting, Addgene plasmid

49386) using the following primers: dPspCas13b_F:

5'TACCCATACGATGTTCCAGATTACGCTTAAGCGGCCGCTCGAGTC3', dPspCas13b_R:

5'GTCGTCATCCTTGTAGTCGGATCCCAGTGTCAGTCTTTCAAG3', FLAG_APEX2_HA_F:

5'GACTACAAGGATGACGACG3', FLAG_APEX2_HA_R:

5'TGGAACATCGTATGGGTACTGCAGGGCATCAGCAAAC3'.

PCR was performed using Q5 High-Fidelity DNA Polymerase (New England Biolabs, catalogue number M0491L).

PCR fragments were assembled using NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs,

catalogue number E2621S) according to manufacturer's instructions. DNA fragment was amplified from pC0043-

PspCas13b crRNA backbone (a gift from Dr. Feng Zhang, Addgene plasmid # 103854) using Cas13b_crRNA-

Forward: ggtgtttcgtcctttccaca and Cas13b_crRNA-Reverse: gttgtggaaggtccagtttt. Then the fragment was

assembled with the following oligos after annealing using NEBuilder HiFi DNA Assembly Master Mix.

Nontargeting control guide-Forward:

tgtggaaaggacgaaacaccgTTTTACAACGTCGTGACTGGGAAAACCCTGgttgtggaaggtccagtttt, Nontargeting control

guide-Reverse: aaaactggaccttccacaacCAGGGTTTTCCCAGTCACGACGTTGTAAAAcggtgtttcgtcctttccaca. U1 (1-

30) guide-Forward: tgtggaaaggacgaaacaccgATCATGGTATCTCCCCTGCCAGGTAAGTATgttgtggaaggtccagtttt,

U1 (1-30) guide-Reverse:

aaaactggaccttccacaacATACTTACCTGGCAGGGGAGATACCATGATcggtgtttcgtcctttccaca. U1 (101-130) guide-

Forward: tgtggaaaggacgaaacaccgCAAATTATGCAGTCGAGTTTCCCACATTTGgttgtggaaggtccagtttt, U1 (101-

130) guide-Reverse: aaaactggaccttccacaacCAAATGTGGGAAACTCGACTGCATAATTTGcggtgtttcgtcctttccaca.

U1(108-137) guide-Forward:

tgtggaaaggacgaaacaccgACTACCACAAATTATGCAGTCGAGTTTCCCgttgtggaaggtccagtttt, U1(108-137) guide-

Reverse: aaaactggaccttccacaacGGGAAACTCGACTGCATAATTTGTGGTAGTcggtgtttcgtcctttccaca. Poly (A)

guide-Forward: tgtggaaaggacgaaacaccgTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTgttgtggaaggtccagtttt, Poly (A)

guide-Reverse: aaaactggaccttccacaacAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAcggtgtttcgtcctttccaca. Poly

(U) guide-Forward: tgtggaaaggacgaaacaccgAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAgttgtggaaggtccagtttt.

Poly (U) guide-Reverse: aaaactggaccttccacaacTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTcggtgtttcgtcctttccaca.

The sequences of all constructs have been confirmed using Sanger sequencing.

Transfection and *in vivo* proximity biotinylation. HEK293T cells were seeded into 150 mm plates and were

transfected with 25 µg pCMV-dPspCas13b-Flag-APEX2-HA and 15 µg Cas13b gRNAs [NTC, U1-1, U1-2, U1-3,

poly (A), poly (U)] using Lipofectamine 3000 (Thermo Fisher Scientific, catalog number L3000015) while ~80%

confluency. HEK293T were incubated with 25 mL of DMEM media containing 25µL of 500 mM biotin-phenol (Iris

Biotech, catalog number LS-3500.1000) in DMSO (Sigma) for 30 min at 37 °C 24h after transfection. Cells were

then treated with 1 mM hydrogen peroxide (H_2O_2) (Sigma-Aldrich, catalog number H1009) for 1 min on a

horizontal shaker at room temperature. The labeling solution was aspirated, and cells were washed twice with 25

mL of quencher solution (10 mM sodium azide [Sigma-Aldrich, catalog number S2002-5G], 10 mM sodium

ascorbate [Sigma-Aldrich, catalog number PHR1279-1G], and 5 mM Trolox [Sigma-Aldrich, catalog number

238813-1G] in Dulbecco's phosphate buffed saline [DPBS, Thermo Fisher Scientific, catalog number 14040182]).

Cells were then washed three times with 15 mL of DPBS and were pelleted by centrifugation at 1,500×g for 5 min

at 4 °C. Cell pellets were snap frozen and stored at −80 °C.

Streptavidin enrichment of biotinylated proteins. Cell pellets from two 150 mm plates of transfected HEK293T

cells were lysed in 2 mL cell lysis buffer (10 mM HEPES, pH7.5 by KOH, 150 mM NaCl, 0.1% NP-40, 5 mM

EGTA, 5 mM Trolox, 10 mM Sodium ascorbate acid, 10 mM Sodium azide, 1 mM PMSF). Streptavidin magnetic

beads (Thermo Fisher Scientific, catalog number 88817) were washed twice with cell lysis buffer and 3.5 mg of

each whole cell lysate sample were incubated with 100 µL magnetic bead slurry with rotation for 2 h at room

temperature. After enrichment, the flowthrough was removed and beads were washed with 2×1 mL cell lysis

buffer, 1mL 1 M KCl, 1 mL 0.1 M Na_2CO_3, 1 mL of 2 M urea in 10 mM Tris-HCl (pH 8.0), and again with 2×1 mL

cell lysis buffer. These denaturing washes are important to disrupt protein-protein interactions and ensure

enrichment only of proteins directly biotinylated by APEX2. Biotinylated proteins were then eluted by boiling the

magnetic beads in 30 μL 4×Laemmli sample buffer (Bio-Rad, catalog number 1610747) supplemented with 20

mM DTT and 2 mM biotin.

Label-free quantitative mass spectrometry analysis

The proteins enriched by RiboPro were analyzed by label-free quantitative mass spectrometry analysis as

previously described[27] at Cedars-Sinai Medical Center Biomarker Discovery Platform Core.

Comparison of RBP databases

The proteins annotated as RBPs used in our study were retrieved from corresponding manuscripts [11–14,28–44]. A list

of human RBPs (GO_RNA Binding) were retrieved from QuickGO (https://www.ebi.ac.uk/QuickGO/) via searching

"RNA Binding" (GO:0003723) and select "*Homo sapiens*" under Taxon. Other categories of human RBPs were

retrieved in the same way by searching corresponding terms. The venn diagrams were generated using online

tool (http://bioinformatics.psb.ugent.be/webtools/Venn/) and pseudovenn diagrams were generated using R

package UpSetR.

GO analysis

GO analyses were performed using Metascape [45](http://metascape.org/gp/index.html#/main/step1).

Cellular fractionation. Six million HEK293T cells were treated with a plasma membrane lysis buffer (10 mM Tris-

HCl, pH 7.5, 0.15% NP-40, 150 mM NaCl) on ice for 4 min after homogenization by flicking. Lysates were loaded

onto a 24% sucrose cushion (24% RNase-free sucrose in plasma membrane lysis buffer) using large orifice tips,

and centrifuged at 15,000 × g for 10 min at 4°C. The supernatant (cytoplasmic fraction) was retained after

centrifugation, and the pellet (nuclear fraction) was washed with 1×PBS/1 mM EDTA and resuspended in 200 μL

of 1×PBS/1 mM EDTA. Fractionation efficiency was validated by Western blot using β-tubulin (Sigma Aldrich,

catalogue number T8328, 1:2,000) as a cytoplasmic marker and U1-70k (EMD Millipore, catalogue number 05-

1588, 1:1,000) as a nuclear marker.

RNA Immunoprecipitation (RIP). Twelve microliter Dynabeads Protein A (Thermo Fisher Scientific, catalog

number: 10001D) were washed with 200 μL HBS (150 mM NaCl, 10 mM HEPES, pH7.5 by KOH and incubated

with 2 μg anti-HA (Santa Cruz, catalogue number sc-7392) or 2 μg rabbit IgG isotype control (Thermo Fisher

Scientific, catalogue number 10500C) in the presence of 80 μL HBS buffer at room temperature for 1 h. Eight

million HEK293T cells were lysed with 800 μL cell lysis buffer (10 mM HEPES, pH7.5 by KOH, 150 mM NaCl,

0.1% NP-40, 5 mM EGTA, supplemented with 1× protease inhibitor cocktail and 1mM PMSF) at 4°C for 1 h. Cell

debris and insoluble proteins were removed by centrifugation at 4°C, 12,000 × g for 10 min, and the supernatants

were incubated with HA-conjugated Dynabeads at 4°C for 1 h. The Dynabeads were then washed 3 times with

wash buffer (150 mM NaCl, 10 mM HEPES, pH7.5 by KOH, 0.1% NP-40). Proteins were eluted with 22 μL

1×Laemmli sample buffer (Bio-Rad, catalog number: 1610747) by boiling for western blotting.

RNA extraction and RT-qPCR. Immunoprecipitated RNA associated with dPspCas13b-Flag-APEX2-HA was

extracted using TRIzol LS (Thermo Fisher Scientific, catalogue number: 10296028). M-MLV reverse transcriptase

(Promega, catalogue number: M5301) and random hexamers (Promega, catalogue number: C1181) were used

for reverse transcription. Gene expression was quantified by RT-qPCR using iQ SYBR Green supermix (Bio-Rad,

catalogue number: 170-8886). The relative gene expression was calculated using the $2^{-\Delta\Delta Ct}$ method and

normalized to GAPDH. Five nanograms cDNA was used for RT-qPCR analysis on CFX96 Touch Real-Time PCR

Detection System (Bio-Rad) using the following primer pairs: U1 snRNA-RT-Forward:

5'CCAGGGCGAGGCTTATCCATT3', U1 snRNA-RT-Reverse: 5'GCAGTCCCCCACTACCACAAAT3', 18S rRNA-

RT-Forward: 5'CAGCCACCCGAGATTGAGCA3', 18S rRNA-RT-Reverse: 5'TAGTAGCGACGGGCGTGTG3',

GAPDH-RT-Forward: 5'TGCCAAATATGATGACATCAAGAA3', GAPDH-RT-Reverse:

5'GGAGTGGGTGTCGCTGTTG3'.

Western blotting. Protein samples were run on 4-20% gradient precast protein gel (Bio-Rad, catalogue number:

456-1096) and then transferred onto PVDF membrane (Bio-Rad, catalogue number: 1704157). After one hour

blocking, membranes were incubated with anti-Flag (Santa Cruz, catalogue number sc-166384), anti-HA (Santa

Cruz, catalogue number sc-7392), anti-biotin (Santa Cruz, catalogue number sc-57636), or anti- β-actin (Santa

Cruz, catalogue number sc-47778, 1:2,000) at 4°C overnight. Membranes were washed three times with Tris-

buffered saline containing 0.5% Tween 20 (TBST) before incubating with HRP-conjugated secondary antibody at

room temperature for 2 h. Then the membranes were incubated briefly with ECL Western Blotting substrate

(Thermo Fisher Scientific, catalogue number: 32106) after three times wash with TBST. The membranes were

exposed to HyBlot Autoradiography Film (Denville Scientific, catalogue number: E3018).

Figure Legend

Figure 1. Proof of principle for RiboPro method

a. Schematic representation of RiboPro approach. A sequence-specific gRNA directs

dPspCas13-APEX to the target RNA and APEX in the chimera biotinylates proximal proteins in

the presence of biotin-phenol (BP) and H_2O_2. Biotinylated proteins are isolated by streptavidin

dynabeads and subsequently analyzed by mass spectrometry (MS) or western blotting

(WB). **b**. Diagram of dPspCas13b-Flag-APEX2-HA construct. **c**. Expression validation of

dPspCas13b-Flag-APEX2-HA by WB. **d**. dPspCas13b-Flag-APEX2-HA is expressed in both

cytoplasm and nucleus. **e**. Functional validation of dPspCas13b-Flag-APEX2-HA. HEK293T

cells transfected with dPspCas13b-Flag-APEX2-HA were treated with BP or BP+H_2O_2. Whole

cell lysates were blotted with anti-biotin antibody. β-actin was blotted as a loading control. **f**.

Designing of 3 gRNAs targeting *U1* snRNA. **g**. Verification of *U1* snRNA gRNAs. HEK293T

cells were transfected with PspCas13b and NTC gRNA or *U1* snRNA gRNAs (1:1 ratio). The

expression of *U1* snRNA or nontargeting *18S* rRNA was quantified by RT-qPCR. **h**. Western

blotting confirms successful pulldown of dPspCas13b-Flag-APEX2-HA by RIP. HEK293T cells

were transfected with dPspCas13b-Flag-APEX2-HA and NTC gRNA or *U1* snRNA gRNAs

(equal ratio). HA antibody or isotype control IgG were used to immunoprecipitate dPspCas13b-

Flag-APEX2-HA and total RNA was extracted and measured by RT-qPCR. **i**. RT-qPCR

measures the abundance of *U1* snRNA and *18S* rRNA from RIP. **j**. U1-70k was enriched by

specific gRNAs for *U1* snRNA. HEK293T cells transfected with pCMV-dPspCas13b-Flag-

APEX2-HA and NTC gRNA or *U1* snRNA gRNAs were treated with BP+H_2O_2. Whole cell

lysates (Input) or streptavidin enriched biotinylated proteins (RiboPro) were blotted. Short and

long exposures are shown for U1-70k. **k**. MS profiling reveals additional *U1* snRNA direct

binding proteins enriched by RiboPro.

Figure 2. RiboPro reveals human poly (A)+ tail RNA proximity proteome

a. Correlation of label-free quantification (LFQ) of proteins identified by RiboPro of poly (A) or

poly (U) tail from two independent replicates. R^2=0.24, P=2.12×10^{-5}, *Pearson*'s correlation

using shared candidates for both replicates. **b.** Shared proteins identified by two replicates. **c.**

Intersecting proteins between poly (A) and poly (U) protein proximitome. P= 1.90×10^{-17},

Binomial test. **d.** GO analysis of poly (A) tail RNA proximity proteome. Top 20 most

significantly enriched GO terms are shown. **e.** Comparison of poly (A) RiboPro with poly (A)

RNA binding proteins in HEK293 cells. **f.** Comparison of poly (A) RiboPro with all annotated

RBPs. **g.** Poly (A) RiboPro reveals known poly (A) RNA binding proteins for poly (A)+ mRNAs.

h. Poly (A) RiboPro uncovers proteins related to RNA turnover. **i.** Poly (A) RiboPro identifies

proteins related to RNA nuclear export, RNA transport, and RNA localization.

Supplementary Figure 1. Poly(A) tail RNA proximity proteome revealed by RiboPro

a. RiboPro of poly(A) tail RNA proximity proteome. In the presence of guide RNA, RiboPro

protein is directed to poly (A) tail ranging from 30 nt to 250 nt. Two hypothetical scenarios are

shown for poly (A)+ RNA, under which RiboPro should identify different categories of RNA

proximity proteins. In the unstructured scenario, RiboPro is likely to detect poly (A) RNA

binding proteins and 3'UTR binding proteins that bind proximal to poly (A) tail as the RNA

proximity proteome of poly (A) tail. In addition to these two categories of proteins, RiboPro will

likely identify cap-binding proteins, 5'UTR binding proteins that bind proximal to the cap in the

structured scenario. **b.** Intersected proteins between two biological replicates for both poly (A)

and poly (U) tail RNA proximity proteome. **c.** Pseudovenn diagram using UpSetR

demonstrates the intersections among poly (A) tail RNA proximity proteome, poly (A) RNA

binding proteomes in HEK293 cells, HeLa cells, and MCF7 cells, respectively. **d.** Pseudo-venn

diagram demonstrates the intersections among poly (A) tail RNA proximity proteome, total

RNA binding proteomes including both poly (A) RNA and non-poly (A) RNA binding proteins in

HEK293 cells, HeLa cells, and MCF7 cells, respectively. **e.** Enriched GO terms for unique poly

(A) tail RNA proximity proteins.

References

1. Wang, K. C. & Chang, H. Y. Molecular mechanisms of long noncoding RNAs. *Mol. Cell* **43**, 904–914

(2011).

2. Guttman, M. & Rinn, J. L. Modular regulatory principles of large non-coding RNAs. *Nature* **482**, 339–

46 (2012).

3. Glisovic, T., Bachorik, J. L., Yong, J. & Dreyfuss, G. RNA-binding proteins and post-transcriptional

gene regulation. *FEBS Lett.* **582**, 1977–1986 (2008).

4. Chu, C., Spitale, R. C. & Chang, H. Y. Technologies to probe functions and mechanisms of long

noncoding RNAs. *Nat. Struct. Mol. Biol.* **22**, 29–35 (2015).

5. Ramanathan, M., Porter, D. F. & Khavari, P. A. Methods to study RNA-protein interactions. *Nat.*

Methods **16**, 225–234 (2019).

6. Cox, D. B. T. *et al.* RNA editing with CRISPR-Cas13. *Science* **358**, 1019–1027 (2017).

7. Abudayyeh, O. O. *et al.* RNA targeting with CRISPR-Cas13. *Nature* **550**, 280–284 (2017).

8. Lam, S. S. *et al.* Directed evolution of APEX2 for electron microscopy and proximity labeling. *Nat.*

Methods **12**, 51–54 (2015).

9. Fazal, F. M. *et al.* Atlas of Subcellular RNA Localization Revealed by APEX-Seq. *Cell* (2019)

doi:10.1016/j.cell.2019.05.027.

10. Chang, H., Lim, J., Ha, M. & Kim, V. N. TAIL-seq: Genome-wide Determination of Poly(A) Tail Length

and 3′ End Modifications. *Molecular Cell* **53**, 1044–1052 (2014).

11. Baltz, A. G. *et al.* The mRNA-Bound Proteome and Its Global Occupancy Profile on Protein-Coding

Transcripts. *Molecular Cell* **46**, 674–690 (2012).

12. Milek, M. *et al.* DDX54 regulates transcriptome dynamics during DNA damage response. *Genome*

Res. **27**, 1344–1359 (2017).

13. Castello, A. *et al.* Insights into RNA biology from an atlas of mammalian mRNA-binding proteins.

Cell **149**, 1393–1406 (2012).

14. Trendel, J. *et al.* The Human RNA-Binding Proteome and Its Dynamics during Translational Arrest.

Cell **176**, 391-403.e19 (2019).

15. Shi, Y. & Manley, J. L. The end of the message: multiple protein–RNA interactions define the mRNA

polyadenylation site. *Genes Dev.* **29**, 889–897 (2015).

16. Vicens, Q., Kieft, J. S. & Rissland, O. S. Revisiting the Closed-Loop Model and the Nature of mRNA 5′

−3′ Communication. *Molecular Cell* **72**, 805–812 (2018).

17. Dreyfus, M. & Régnier, P. The Poly(A) Tail of mRNAs. *Cell* **111**, 611–613 (2002).

18. Norbury, C. J. Cytoplasmic RNA: a case of the tail wagging the dog. *Nat. Rev. Mol. Cell Biol.* **14**, 643–

653 (2013).

19. Collart, M. A. The Ccr4-Not complex is a key regulator of eukaryotic gene expression. *Wiley*

Interdiscip Rev RNA **7**, 438–454 (2016).

20. Chlebowski, A., Lubas, M., Jensen, T. H. & Dziembowski, A. RNA decay machines: The exosome.

Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms **1829**, 552–560 (2013).

21. Chang, Y.-F., Imam, J. S. & Wilkinson, M. F. The Nonsense-Mediated Decay RNA Surveillance

Pathway. *Annu. Rev. Biochem.* **76**, 51–74 (2007).

22. Laroia, G., Cuesta, R., Brewer, G. & Schneider, R. J. Control of mRNA decay by heat shock-ubiquitin-

proteasome pathway. *Science* **284**, 499–502 (1999).

23. Huang, Y. & Carmichael, G. G. Role of polyadenylation in nucleocytoplasmic transport of mRNA.

Mol. Cell. Biol. **16**, 1534–1542 (1996).

24. Cheng, H. *et al.* Human mRNA export machinery recruited to the 5' end of mRNA. *Cell* **127**, 1389–

1400 (2006).

25. Mofatteh, M. & Bullock, S. L. SnapShot: Subcellular mRNA Localization. *Cell* **169**, 178-178.e1 (2017).

26. Holt, C. E. & Bullock, S. L. Subcellular mRNA Localization in Animal Cells and Why It Matters. *Science*

326, 1212–1216 (2009).

27. Zhou, B. *et al.* Low-Background Acyl-Biotinyl Exchange Largely Eliminates the Coisolation of Non- *S* -

Acylated Proteins and Enables Deep *S* -Acylproteomic Analysis. *Anal. Chem.* **91**, 9858–9866 (2019).

28. Bao, X. *et al.* Capturing the interactome of newly transcribed RNA. *Nat. Methods* **15**, 213–220

(2018).

29. Beckmann, B. M. *et al.* The RNA-binding proteomes from yeast to man harbour conserved

enigmRBPs. *Nat Commun* **6**, 10127 (2015).

30. Brannan, K. W. *et al.* SONAR Discovers RNA-Binding Proteins from Analysis of Large-Scale Protein-

Protein Interactomes. *Molecular Cell* **64**, 282–293 (2016).

31. Castello, A. *et al.* Comprehensive Identification of RNA-Binding Domains in Human Cells. *Molecular*

Cell **63**, 696–710 (2016).

32. Caudron-Herger, M. *et al.* R-DeeP: Proteome-wide and Quantitative Identification of RNA-

Dependent Proteins by Density Gradient Ultracentrifugation. *Mol. Cell* **75**, 184-199.e10 (2019).

33. Cook, K. B., Kazan, H., Zuberi, K., Morris, Q. & Hughes, T. R. RBPDB: a database of RNA-binding

specificities. *Nucleic Acids Research* **39**, D301–D308 (2011).

34. Conrad, T. *et al.* Serial interactome capture of the human cell nucleus. *Nat Commun* **7**, 11212

(2016).

35. Gerstberger, S., Hafner, M. & Tuschl, T. A census of human RNA-binding proteins. *Nat. Rev. Genet.*

15, 829–845 (2014).

36. He, C. *et al.* High-Resolution Mapping of RNA-Binding Regions in the Nuclear Proteome of

Embryonic Stem Cells. *Molecular Cell* **64**, 416–430 (2016).

37. Kwon, S. C. *et al.* The RNA-binding protein repertoire of embryonic stem cells. *Nat. Struct. Mol.*

Biol. **20**, 1122–1130 (2013).

38. Mullari, M., Lyon, D., Jensen, L. J. & Nielsen, M. L. Specifying RNA-Binding Regions in Proteins by

Peptide Cross-Linking and Affinity Purification. *J. Proteome Res.* **16**, 2762–2772 (2017).

39. Perez-Perri, J. I. *et al.* Discovery of RNA-binding proteins and characterization of their dynamic

responses by enhanced RNA interactome capture. *Nat Commun* **9**, 4408 (2018).

40. Queiroz, R. M. L. *et al.* Comprehensive identification of RNA-protein interactions in any organism

using orthogonal organic phase separation (OOPS). *Nat. Biotechnol.* **37**, 169–178 (2019).

41. Ray, D. *et al.* A compendium of RNA-binding motifs for decoding gene regulation. *Nature* **499**, 172–

177 (2013).

42. Sundararaman, B. *et al.* Resources for the Comprehensive Discovery of Functional RNA Elements.

Molecular Cell **61**, 903–913 (2016).

43. Treiber, T. *et al.* A Compendium of RNA-Binding Proteins that Regulate MicroRNA Biogenesis.

Molecular Cell **66**, 270-284.e13 (2017).

44. Urdaneta, E. C. *et al.* Purification of cross-linked RNA-protein complexes by phenol-toluol

extraction. *Nat Commun* **10**, 990 (2019).

45. Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-level

datasets. *Nat Commun* **10**, 1523 (2019).