In vivo discovery of RNA proximal proteins in human cells via

proximity-dependent biotinylation

Xianzhi Lin[1,2]*, Marcos A. S. Fonseca[1,2], Rosario I. Corona[1,2], Kate Lawrenson[1,2,3]*

[1]Women's Cancer Research Program at Samuel Oschin Comprehensive Cancer Institute,

Cedars-Sinai Medical Center, Los Angeles, CA, USA.

[2]Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, Cedars-

Sinai Medical Center, Los Angeles, CA, USA.

[3]Center for Bioinformatics and Functional Genomics, Samuel Oschin Comprehensive

Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA.

*Correspondence: xianzhi.lin@cshs.org (X.L.) or kate.lawrenson@cshs.org (K.L.)

Abstract

RNA molecules function as messengers or noncoding adaptor molecules, structural components, and regulators of genome organization and gene expression. Their roles and regulation are mediated by other molecules they interact with, especially RNA binding proteins (RBPs). Here we report RNA proximity labeling (RPL), an RNA-centric method based on fusion of an endonuclease-deficient Type VI CRISPR-Cas protein (dCas13b) and engineered ascorbate peroxidase (APEX2) to discover *in vivo* target RNA proximal proteins (RPPs) through proximity-based biotinylation. *U1* RPPs enriched by proximity-based biotinylation included both *U1* snRNA canonical and noncanonical functions-related proteins. In addition, profiling of poly(A) tail proximal proteins uncovered expected categories of RBPs for poly(A) tails and also provided novel evidence for poly(A)$^+$ RNA 5'-3' proximity and expanded subcellular localizations. Our results suggest that RPL is a rapid approach for identifying both interacting and neighboring proteins associated with target RNA molecules in their native cellular contexts.

Keywords

Type VI CRISPR-Cas (Cas13), Engineered soybean ascorbate peroxidase APEX2, Proximity-dependent biotinylation, RNA proximity labeling (RPL), RNA proximal proteins (RPPs), RNA-centric method, RNA-protein interactions, poly(A) tail, RNA binding proteins (RBPs), *U1* snRNA interactors.

Introduction

RNA molecules include both messengers encoding proteins (mRNAs) and noncoding RNAs (ncRNAs) such as adaptor tRNAs and regulatory long noncoding RNAs (lncRNAs). Around 2% of the human genome encodes mRNAs (*International Human Genome Sequencing Consortium, 2001; Venter et al., 2001*), while the majority is pervasively transcribed into ncRNAs (*Berretta and Morillon, 2009; Djebali et al., 2012*), including lncRNAs that are widely considered as a large family of potential regulators (*Batista and Chang, 2013; Iyer et al., 2015; Yang et al., 2014*). However, only a small number of lncRNAs have been functionally and mechanistically studied and most remain uncharacterized (*Kopp and Mendell, 2018*).

The functions and regulation of RNA transcripts are mediated by other molecules they associate with, particularly RNA binding proteins (RBPs) that govern many critical RNA activities (*Dreyfuss et al., 2002; Glisovic et al., 2008; Hentze et al., 2018; Lunde et al., 2007*). Discovery of the interacting proteins for a given transcript plays pivotal role in unveiling its function and underlying mechanism. Currently, mechanistic study of lncRNAs is impeded by the shortage of RNA-centric tools and the limitations of existing methods (*Ci Chu et al., 2015; Ramanathan et al., 2019*). Antisense probe-based ChIRP (*C. Chu et al., 2015*) or RAP (*McHugh et al., 2015*) requires crosslinking *via* chemicals or UV light. However, chemicals such as formaldehyde also crosslink protein-protein interactions, which may lead to false-positive associations (*Panhale et al., 2019*). Since UV-crosslinking has very low efficiency, antisense probe-based purification methods usually require a large number of cells (~100-800 million) (*Lin et al., 2019; McHugh et al., 2015*), which may not be feasible for slow-growing model systems such as primary

cell cultures. Moreover, UV-crosslinking can induce RNA alterations like modifications

(*Wurtmann and Wolin, 2009*) that could change binding affinity of RNA to certain RBPs

(*Bernard et al., 2012*) and impair downstream protein analysis (*Urdaneta and*

Beckmann, 2019). An alternative approach, tagging of endogenous RNA requires

genetic manipulation and may interfere with endogenous RNA functions (*Laprade et al.,*

2020). Therefore, methods to discover endogenous RNA interacting proteins are needed.

In this study, we developed RPL (<u>R</u>NA <u>p</u>roximity <u>l</u>abelling) method to identify *in vivo*

target RNA proximal proteins (RPPs) without crosslinking or genetic manipulation. *U1*

RPPs recalled *U1* functional relevant proteins, while poly(A) tail RPPs recalled expected

categories of RBPs for poly(A) tails providing additional evidence for poly(A)$^+$ RNA 5'-3'

proximity and expanded subcellular localizations.

Results

Design and development of RPL, an RNA-centric method for screening RPPs

Inspired by the applications of RNA-targeting Type VI CRISPR-Cas systems (*Abudayyeh*

et al., 2017; Cox et al., 2017; Konermann et al., 2018; Yan et al., 2018) and proximity

labeling using engineered soybean ascorbate peroxidase (*Lam et al., 2015; Rhee et al.,*

2013) and biotin ligase (*Branon et al., 2018; Kim et al., 2016; Roux et al., 2012*), we

designed RPL, an RNA-centric approach based on a fusion protein of endonuclease-

deficient Cas13 (dCas13) and proximity labeling enzyme APEX2 (*Figure 1A*). The fusion

protein is directed to target RNA by a sequence-specific guide RNA (gRNA). In the

presence of hydrogen peroxide (H_2O_2), APEX2 in the fusion protein oxidizes substrate

biotin-phenol (BP) to short-lived biotin-phenoxyl radicals, which covalently react with

electron-rich amino acids (like tyrosine) on RPPs within a small radius (*Rhee et al., 2013*)

of the fusion protein (*Figure 1A*). The biotinylated RPPs, which may include target RNA

direct binding proteins, indirect binding proteins, and proximal proteins just present within

biotinylating radius, can be readily enriched using streptavidin beads and profiled by liquid

chromatography-tandem mass spectrometry (LC-MS/MS) (*Figure 1A*).

To construct the fusion protein, Cas13b was used for its high efficacy in RNA

knockdown with minimal off-target effect (*Cox et al., 2017*) and high specificity in RNA

labeling (*Yang et al., 2019*). For proximity labeling enzyme, we chose APEX2 for its fast

kinetics and high activity (*Lam et al., 2015*). Catalytically dead Cas13b from *Prevotella*

sp. P5-125 (dPspCas13b) (*Cox et al., 2017*) was fused to APEX2 with FLAG and HA

tags (*Figure 1B*). The expression of the fusion protein dCas13b-APEX2 (from hereon in

called the RPL protein) was confirmed by western blot using an anti-FLAG or anti-HA

Figure 1. Designing and developing the RPL method. (**A**). Schematic illustration of RPL workflow. A sequence-specific gRNA directs dCas13-APEX2 to target RNA and APEX2 in the fusion protein biotinylates target RNA proximal proteins (RPPs) *in vivo* in the presence of biotin-phenol and H_2O_2. Biotinylated RPPs are then enriched using streptavidin beads and analyzed by liquid chromatography-tandem mass spectrometry (LC-MS/MS). (**B**). Diagram of the fusion protein dPspCas13b-FLAG-APEX2-HA (dCas13b-APEX2, or the RPL protein) expression construct. (**C**). Expression validation of the RPL protein by western blot. HEK293T cells transfected with or without the RPL plasmid were harvested 24h-72h post transfection and whole cell lysates were blotted with an anti-FLAG or anti-HA antibody. (**D**). The RPL protein is expressed in both cytoplasm and nucleus. HEK293T cells transfected with the RPL plasmid for 24h were fractionated into cytoplasmic (Cy) and nuclear (Nu) fractions. Fractionation efficiency was evaluated by blotting cytoplasmic protein β-Tubulin and nuclear protein U1-70k. (**E**). Validation of enzymatic activity of APEX2 in the RPL protein. HEK293T cells transfected with the RPL plasmid were treated with different combinations of biotin-phenol (BP) and H_2O_2. Whole cell lysates were blotted with anti-biotin antibody. β-actin in (**C**) and (**E**) was used as loading control.

antibody (*Figure 1C*). The subcellular localization of the RPL protein was examined when

ectopically expressed in HEK293T cells. Efficient separation between cytoplasmic and

nuclear fractions was confirmed by blotting for cytoplasmic marker β-Tubulin and nuclear

marker U1-70k. The RPL protein was detected in both cytoplasm and nucleus (*Figure*

1D). To test if peroxidase activity of APEX2 is maintained in the RPL protein, HEK293T

cells were treated with different combinations of BP and H_2O_2 24h post transfection of the

RPL plasmid. The detection of biotinylated proteins requires both BP and H_2O_2, indicating

that APEX2 in the RPL protein retains peroxidase activity (*Figure 1E*). The results also

suggest that endogenous biotinylated proteins are rare in HEK293T cells and low level of

endogenous H_2O_2 (*Belousov et al., 2006; Huang and Sikes, 2014; Lyublinskaya and*

Antunes, 2019) could not trigger efficient biotinylation. These data suggest that the RPL

protein has peroxidase activity and can be applied to target both cytoplasmic and nuclear

transcripts.

Design and validation of gRNAs targeting *U1* snRNA

To test the approach, we asked whether *U1* snRNA proximal proteins (*U1* RPPs)

identified with the RPL protein include any known *U1* RBPs. The *U1* snRNA was selected

for three reasons: (1) its high abundance (*Gesteland, 1993*), (2) its structures in human

U1 small nuclear ribonucleoprotein (snRNP) and in spliceosome have been solved

(*Charenton et al., 2019; Pomeranz Krummel et al., 2009; Weber et al., 2010*), and (3)

its interacting proteins in *U1* snRNP (*Stark et al., 2001*) and in spliceosome (*Zhou et al.,*

2002) have been well documented.

Since Cas13b targets single-stranded RNA (*Cox et al., 2017; Smargon et al., 2017*),

three gRNAs (*U1*-1, *U1*-2, and *U1*-3) targeting *U1* single-stranded regions were designed

based on its structure in pre-B complex (*Charenton et al., 2019*) (*Figure 2A*). We first

tested if *U1* gRNAs direct wild-type PspCas13b to *U1* and cleave it by measuring *U1*

expression in HEK293T cells cotransfected with wild-type PspCas13b plasmid and

plasmid expressing *U1* gRNA or nontargeting control (NTC) gRNA at a 1:1 molar ratio.

The expression of *U1* was significantly lower in *U1* gRNA-transfected cells compared with

NTC gRNA-transfected cells (*Figure 2B*). The expression of a group of nontargets with

a wide range of abundance was not affected (*Figure 2B*), except *U2*, which may be

caused by Cas13b collateral activity (*Gootenberg et al., 2018*) since *U1* and *U2* are in

close contact during spliceosome assembly. The result indicated that *U1* gRNAs can

specifically direct PspCas13b to *U1*. We then tested if *U1* gRNAs deliver the RPL protein

to *U1* using RNA immunoprecipitation (RIP) experiment. Since the U6 promoter is slightly

stronger than CMV promoter in HEK293T cells (*Lebbink et al., 2011*), a 1:2 molar ratio

between the RPL plasmid (CMV promoter) and gRNA expressing plasmid (U6 promoter)

was used to avoid nonspecific targeting due to excess RPL protein. The RPL protein was

efficiently retrieved by anti-HA but not isotype control IgG (*Figure 2C*). Analysis of RNA

extracted from RIP experiment showed that anti-HA pulled down 5 times more RNA than

control (*Figure 2D*), certifying the RPL protein RNA binding activity. Although there is no

significant difference in the amount of RNA pulled down by the RPL protein with NTC or

U1 gRNAs (*Figure 2D*), *U1* gRNAs significantly enriched *U1* for ~2-3-fold compared with

NTC gRNA (*Figure 2E*). The fact that much more abundant *18S* was not enriched (*Figure*

2E) suggested that *U1* gRNAs are able to specifically direct the RPL protein to *U1*.

Fig. 2

Figure 2. Designing and validating gRNAs targeting *U1* snRNA. (**A**). Based on *U1* structure, three gRNAs
with spacers targeting *U1* nucleotides (nt) 1-30 (*U1*-1), 101-130 (*U1*-2), and 108-137 (*U1*-3) were designed.
Cartoon representation of *U1* (PDB ID: 6QX9, pre-B complex) is colored in black, 1-30 in green, 101-107
in red, 108-130 in magenta, and 131-137 in blue. (**B**). The expression of *U1* snRNA was significantly
downregulated in *U1* gRNA-transfected cells. HEK293T cells were cotransfected with plasmid expressing
wild-type PspCas13b and plasmid expressing *U1* or NTC gRNA (1:1 molar ratio). The expression of *U1* or
a group of nontargets was quantified by RT-qPCR and normalized to *GAPDH*. (**C**). Confirmation of pulldown
of the RPL protein by RIP using western blot. HEK293T cells were cotransfected with the RPL plasmid and
plasmid expressing *U1* or NTC gRNA (1:2 molar ratio). Anti-HA antibody or isotype control IgG were used
to immunoprecipitate the RPL protein. Clean-Blot IP detection reagent was used for blotting. (**D**). The
amount of total RNA extracted from RIP experiment. (**E**). *U1* gRNAs specifically directed the RPL protein
to *U1*. The expression of *U1* snRNA and nontarget *18S* rRNA was quantified by RT-qPCR and normalized
to *GAPDH*. Data shown in (**B**), (**D**), and (**E**) are mean ± SD from 5 independent experiments. ***$p<0.001$,
ns, not significant. Student's *t* test.

RPL-MS identified both *U1* canonical and noncanonical roles-related proteins

We next enriched *U1* RPPs using RPL with the same 1:2 molar ratio to avoid excess RPL protein that can cause nonspecific targeting and proximity labeling. *U1* has compact structure in pre-B complex (*Charenton et al., 2019*) (*Figure 2A*) and its size (less than ~10 nm in diameter) is much smaller than the biotinylating range of APEX2 (likely ~20-40 nm or larger in diameter) (*Fazal et al., 2019; Padrón et al., 2019; Rhee et al., 2013*), so we considered experiments using our three *U1* gRNAs as replicates. We analyzed streptavidin-enriched biotinylated proteins by LC-MS/MS (RPL-MS). Using label-free intensity-based absolute quantification (iBAQ) values to measure enrichment in *U1* gRNA relative to protein amounts in the NTC gRNA sample, RPL-MS identified 226 *U1* RPPs ($p < 0.05$ and $\log_2$ fold change [FC] > 2, false discovery rate [FDR] < 0.25, Benjamini-Hochberg method), including known *U1* direct RBPs (e.g. SNRNP70, also known as U1-70k) (*Stark et al., 2001*) and RBPs that likely interact with *U1* indirectly due to their function in the spliceosome (e.g. SNRPA1 and SNRPB2) (*Zhou et al., 2002*) (*Figure 3A, Table S1*). We verified the enrichment of U1-70k using western blot and found that it was enriched ~2-fold by all three *U1* gRNAs (*Figure 3B*), consistent with RPL-MS results (*Figure 3A*). Analysis of KEGG pathways enriched in the group of *U1* RPPs using STRING (*Szklarczyk et al., 2019*) showed that 'Spliceosome' is the most significantly enriched pathway (FDR < 10^{-8}) (*Figure 3C*). Indeed, *U1* RPPs included 99 splicing and related factors (*Cvitkovic and Jurica, 2013*), 56 proteins previously found by *U1* ChIRP-MS (*C. Chu et al., 2015*), and 58 proteins revealed by XLIP-MS using anti-U1A and/or anti-U1-70k antibody (*So et al., 2019*) (*Figure 3D*). In addition, the binding between *U1* and four *U1* RPPs was further supported by corresponding CLIP-Seq data as shown in

Fig. 3

Figure 3. *U1* RPPs identified by RPL-MS. (**A**). *U1* RPPs revealed by RPL-MS include known *U1* RBPs.

Volcano plot shows *U1*/NTC iBAQ ratio (fold change, FC) of protein quantification in *U1* gRNA cells

compared with NTC gRNA cells. RPL-MS enriched 226 *U1* RPPs (orange dots) that were statistically

significant ($p < 0.05$ and $\log_2$ FC > 2, FDR < 0.25, Benjamini-Hochberg method). Each dot represents the

average value from experiments using three *U1* gRNAs. Blue dots represent proteins from pre-B

spliceosome complex. (**B**). *U1* direct RBP U1-70k was enriched by RPL. HEK293T cells transfected with

the RPL plasmid and plasmid expressing *U1* gRNA or NTC gRNA were treated with BP and H_2O_2. Whole

cell lysates (Input) or streptavidin-enriched biotinylated proteins (RPL enriched) were blotted. Numbers

represent relative amount of U1-70k under the corresponding conditions from RPL enriched normalized to

input. (**C**). KEGG pathways significantly enriched by 226 *U1* RPPs using STRING. (**D**). Comparison of *U1*

RPPs, *U1* interactors identified by ChIRP-MS, *U1* interactors identified by XLIP-MS using anti-U1A and/or

anti-U1-70k antibody, and splicing & related proteins. Numbers listed below are total number of proteins

from each group. (**E**). List of 4 *U1* RPPs with CLIP-Seq data supporting their association of *U1* found in

ENCORI. (**F**). Summary of *U1* RPPs related to *U1* functions. (**G**). Inferred distances between *U1* or APEX2

in the RPL protein and those 6 *U1* RPPs present in pre-B complex shown in (**A**).

ENCORI (*Li et al., 2014*), including DDX3X that is not known to interact with *U1* in human

(*Deckert et al., 2006; Tarn and Chang, 2009*) (*Figure 3E*). These results together

validated that RPL can efficiently identify most known RBPs for *U1*.

U1 RPPs also included previously reported *U1* interactor RNA polymerase II (*Spiluttini*

et al., 2010; Yu and Reed, 2015) (*Figure 3F*), which is required for a noncanonical role

of *U1* in chromatin retention of ncRNAs (*Yin et al., 2020*). Moreover, *U1* RPL retrieved

proteins involved in chromatin remodeling, DNA modification, histone modification, and

transcription (*Table S1*), which could be regulated by chromatin-associated ncRNAs

(*Huang et al., 2020; Li and Fu, 2019*). The presence of GTF2F2 among the *U1* RPPs

may relate to a role for *U1* in regulation of transcription initiation (*Damgaard et al., 2008;*

Kwek et al., 2002) (*Figure 3F*). Interestingly, RPL-MS revealed nuclear import receptor

importin-β1 (KPNB1) (*Figure 3F*), which is required for *U1* nuclear import (*Palacios,*

1997). Six RBPs in pre-B complex (*Charenton et al., 2019*) were identified as *U1* RPPs

(*Figure 3A*). Their distances to *U1* snRNA in pre-B complex may provide insight to the

biotinylating range of the RPL protein. The inferred distances between APEX2 in the RPL

protein and those RBPs are all smaller than 12 nm and the average is 6.6 nm (*Figure*

3G), suggesting that APEX2 may biotinylate proteins within 12 nm. The inferred distances

between *U1* and associated RBPs range from 4.2 nm to 20.8 nm with an average of 12.5

nm (*Figure 3G*), suggesting that RPL can biotinylate proximal proteins within ~20 nm of

target RNA. These data indicated that RPL enables efficient identification of validated

RBPs associated with both canonical and noncanonical functions of *U1*.

RPL-MS recalled expected categories of proteins for poly(A) tails

To further test the generality of RPL method, we applied it to poly(A) tails, which are adenosines added to the 3' ends of the majority of eukaryotic mRNAs and many lncRNAs in the absence of template (*Derrien et al., 2012; Guttman et al., 2009; Tian, 2005; Yang et al., 2011*). Poly(A) tails play critical role in mRNA translation and stability (*Dreyfus and Régnier, 2002*) and their removal triggers mRNA decapping and decay (*Muhlrad et al., 1994; Norbury, 2013; Yamashita et al., 2005*). Although the 5' and 3' ends of pre-translational mRNAs (*Metkar et al., 2018*) and deadenylating mRNAs (*Chen and Shyu, 2011*) are distant (*Figure 4A, 5'-3' distance*), the physical distances between the two ends of diverse RNAs are incredibly close regardless of their length, type, species, or complexity (*Lai et al., 2018; Leija-Martínez et al., 2014*) (*Figure 4A, 5'-3' proximity*). As oligomers of 30 nt poly(U) are not found at the 3' of RNA (*Chang et al., 2014; Lim et al., 2018*) and rarely occur in the human transcriptome, poly(U)-targeting gRNA, or poly(U) gRNA, was used as negative control. The RPL plasmid was cotransfected with plasmid expressing poly(A) or poly(U) gRNA into HEK293T cells at a 1:2 molar ratio and then RPL was performed. Using label-free iBAQ values to measure enrichment in poly(A) gRNA relative to protein amounts in the poly(U) gRNA sample, RPL-MS enriched 786 proteins as poly(A) tail RPPs (Benjamini-Hochberg-adjusted $p < 0.05$ and $\log_2 FC > 2$) (*Figure 4B, Table S2*). Poly(A) tail RPPs included seven poly(A) binding proteins, fifteen 3'UTR binding proteins, ten 5'UTR binding proteins, and one cap binding protein (*Figure 4B, Table S3*), all of which are known to associate with poly(A) tails. Retrieval of proteins from both 5' and 3' ends by RPL within a small radius provided additional evidence for poly(A)[+]

RNA 5'-3' proximity.

Among poly(A) tail RPPs, at least 48% were RBPs interacting with poly(A)$^+$ RNA (*Baltz*

et al., 2012; Castello et al., 2012; Kwon et al., 2013; Milek et al., 2017) (*Figure 4C*). In

theory, poly(A) gRNA can direct the RPL protein to any transcripts with 30 nt-poly(A) tail

or longer (*Figure 4A*), including transcripts undergoing polyadenylation, readenylation,

deadenylation, or translation. We then interrogated poly(A) tail RPPs for other expected

classes of proteins, including factors involved in polyadenylation (*Shi and Manley, 2015*),

readenylation or deadenylation (*Yan, 2014*), and translation (*Dreyfus and Régnier,*

2002). Indeed, RPL-MS enriched five cleavage and polyadenylation factors for poly(A)$^+$

RNA (*Figure 4D*) but no such factors unique for poly(A)$^-$ RNA (e.g. SLBP and ZNF473)

(*Gilmartin, 2005*) (*Table S3*). Moreover, poly(A) tail RPPs included three exosome

proteins (*Chlebowski et al., 2013*), two deadenylase complex proteins (*Collart, 2016*),

as well as decapping factor EDC3 (*Mugridge et al., 2018*) (*Figure 4D, Table S3*).

Importantly, twenty translation initiation factors, fifteen translation elongation factors,

seventy ribosomal subunits, and eighteen tRNA ligases were identified by RPL-MS

(*Figure 4D, Table S3*), putatively supporting a model that poly(A) tail recruits translation

initiation factors to initiate translation at the 5' end like their viral counterparts (*Simon and*

Miller, 2013; Truniger et al., 2017). Moreover, RPL-MS revealed twelve proteins

involved in degradation of AU-rich element-containing mRNAs and 66 nonsense-

mediated decay proteins (including 58 ribosomal subunits) (*Chang et al., 2007; Laroia*

et al., 2002, 1999) (*Table S3*), further suggesting that RPL enables efficient discovery of

most relevant and validated RBPs proximal to poly(A) tails.

Fig. 4

Figure 4. RPL-MS revealed poly(A) tail RPPs in HEK293T cells. (**A**). Application of RPL to poly(A) tails. In

the presence of gRNA, the RPL protein (dCas13b-APEX2) is directed to poly(A) tails ranging from 30 nt up

to ~250 nt. In 5'-3' distance model, RPL will detect PABPs and 3'UTR binding proteins that bind proximal

to poly(A) tail within the biotinylating range. In 5'-3' proximity model, RPL will also identify cap-binding

proteins and 5'UTR binding proteins that bind proximal to the cap and lie within the biotinylating range. (**B**).

RPL-MS identified poly(A) tail RPPs. Volcano plot shows RPL-labeled proteins in HEK293T cells. For each

protein, the poly(A)/poly(U) iBAQ ratio reflects the enrichment of identified protein in poly(A) gRNA cells

compared with poly(U) gRNA transfected cells. RPL-MS identified 786 proteins (light purple dots) as

significantly enriched (Benjamini-Hochberg-adjusted $p < 0.05$ and $\log_2 FC > 2$). Each data point represents

the average value from biological triplicates. Red dots represent proteins belonging to PABPs, blue dots

for 3'UTR binding proteins, green dots for 5'UTR binding proteins, and black dot for cap binding protein.

(**C**). Venn diagram shows the comparison of poly(A) tail RPPs and RBPs associated with poly(A)+ RNA in

different cells. Numbers below each group represent the sizes of the protein cohort. (**D**). Summary of

expected categories of poly(A) tail RPPs that support 5'-3' proximity and the role of poly(A) tail in mRNA

translation. Each category of proteins points to a location/region of poly(A)$^+$ RNA where they most likely

associate with when identified by RPL. PABPs, poly(A) binding proteins; CPA, cleavage and

polyadenylation; TIFs, translation initiation factors; TEFs, translation elongation factors.

Localization analysis of poly(A) tail RPPs unveils expanded subcellular

localizations for poly(A)$^+$ RNA

Poly(A) tails are important for RNA nuclear export (*Huang and Carmichael, 1996*) *via*

the nuclear pore complex (NPC) (*Okamura et al., 2015*). This is further supported by the

presence of 90 mRNA processing factors, 20 mRNA nuclear export proteins, and 13 NPC

proteins in poly(A) tail RPPs (*Figure 5A*). It is not surprising that tRNAs and pre-miRNAs

nuclear export factors were also included (*Table S3*) since their precursors or primary

transcripts are also polyadenylated (*Cai, 2004; Kadaba et al., 2006*). Poly(A) tail RPPs

contained eight tRNA processing factors and five tRNA nuclear export factors (*Kruse et*

al., 2000), as well as three pri-miRNA processing factors and two pre-miRNA export

factors (*Bohnsack et al., 2004; Lund et al., 2004; Yi et al., 2003*) (*Table S3*), supporting

that their processing is coupled with export (*Kim, 2005; Köhler and Hurt, 2007*). Poly(A)

tail RPPs recovered 27 RBPs involved in mRNA transport (including zipcodes binding

protein IGF2BP1), 48 microtubule proteins, and 139 plasma membrane proteins that are

used by mRNAs to achieve different subcellular localizations (*Holt and Bullock, 2009;*

Mofatteh and Bullock, 2017) (*Figure 5A*), possibly suggesting a role for the poly(A) tail

in RNA subcellular localization.

Since unique localizations of RPPs reflect target RNA proximal localizations, we built

a putative subcellular localization map for poly(A)$^+$ RNA by comparing poly(A) tail RPPs

with proteins extracted from 22 subcellular compartments (*Figure 5B, Table S3*). The

results are generally consistent with previous reports that both mRNAs and ncRNAs have

multiple subcellular localizations (*Blower, 2013; Carlevaro-Fita and Johnson, 2019;*

Fazal et al., 2019; Wilk et al., 2016) and also support the presence of mRNAs in P-

bodies, stress granule, and the exosome (*Chlebowski et al., 2013; Decker and Parker,*

2012). Interestingly, RPL-MS also identified marker proteins of the endosome, lysosome,

proteasome, and Golgi apparatus, indicative of expanded subcellular localizations for

poly(A)$^+$ RNA (*Figure 5B, Table S3*). Discovery of lysosomal and proteasomal proteins

in poly(A) tail RPPs is compatible with the existence of RNA degradation pathway

'RNautophagy' in the lysosome (*Fujiwara et al., 2013*) and degradation function of

proteasomes for AU-rich element-containing mRNAs (*Laroia et al., 2002, 1999*). The

identification of endosomal proteins is in accordance with that late endosomes can be

used by mRNAs as a platform for translation (*Cioni et al., 2019*). Poly(A) tail RPPs

included Golgi marker *cis*-Golgi matrix protein GOLGA2 (*Munro, 2011*) (*Table S3*), which

has recently been annotated as an RBP by multiple groups (*Caudron-Herger et al., 2019;*

Queiroz et al., 2019; Trendel et al., 2019), suggesting that Golgi may be a novel

subcellular location for poly(A)$^+$ RNA. More experimental data are needed to determine

which specific transcripts are associated with GOLGA2 in the Golgi apparatus and the

biological significance of those interactions.

Figure 5. Poly(A) tail RPPs included proteins involved in subcellular localization of RNA. (**A**). RPL-MS

revealed proteins involved in RNA processing, nuclear export, transport, and subcellular localization for

poly(A) tail RPPs. (**B**). A putative subcellular localization map of poly(A)$^+$ RNA built upon subcellular

localization of poly(A) tail RPPs. Poly(A) tail RPPs were compared with proteins extracted from

corresponding GO terms. Numbers in brackets represent the size of each category of proteins. A full list of

proteins in each category can be found in *Table S3*.

Discussion

RPL: an RNA-centric approach for RPPs identification in living cells

We present an RNA-centric method, RPL, for discovering RPPs for transcripts of interest

and evaluate it in two distinct contexts–first interrogating a specific ncRNA target *U1* and

second surveying a heterogenous group of poly(A)$^+$ RNA in living cells. Both *U1* RPPs

and poly(A) tail RPPs demonstrated that RPL enables efficient discovery of functional

relevant RBPs for target transcripts. The recall of KPNB1 for *U1* nuclear import suggests

that RPL allows to detect transient and/or weak interacting proteins (*Branon et al., 2017;*

Roux et al., 2013). Compared with alternative methods, RPL needs no crosslinking or

sonication, requires far fewer cells (~20-40 million vs ~100-800 million) and involves no

genetic manipulation, which may interfere target RNA functions (*Laprade et al., 2020*).

The short pulse of labeling potentially permits RPL to be applied to study RNA-protein

dynamics. Recently, APEX2 has also been reported to biotinylate proximal nucleic acids

(*Fazal et al., 2019; Padrón et al., 2019; Y. Zhou et al., 2019*), suggesting that RPL could

be potentially applied to identify RNA and DNA in addition to proteins proximal to the

target RNA (together as 'RNA proximitome') within living cells.

During the preparation of our manuscript, similar strategies using different fusion

proteins of endonuclease-deficient Cas13 protein (dLwaCas13a, dPspCas13b, and

dRfxCas13d) and proximity labeling enzyme (APEX2, BioID2, BASU, and PafA) were

reported (*Han et al., 2020; Li et al., 2020; Yi et al., 2020; Zhang et al., 2020*).

Applications of these methods together with ours to both mRNAs and ncRNAs with wide

range of abundance (~10^2-10^6 copies/cell) demonstrate that these methods have broad

potential to identify functional relevant RBPs for diverse transcripts.

RPPs identified using the dCas13b-APEX2 are expected to include three types of

proteins theoretically: proteins that directly bind to target RNA, proteins that associated

with target RNA indirectly via protein-protein interactions, and proteins present within the

biotinylating range. Biological replicates are expected to help enrich the first two groups

of RBPs and reduce the third type as false positive candidates may not be enriched

repeatedly. In addition, an optimal molar ratio between the fusion protein and gRNA,

which enables efficient proximity-based biotinylation and prevents nonspecific labeling

due to excess fusion protein, is crucial for separating signal from noise. A validated set of

gRNAs that can specifically direct fusion protein to target RNA with low off-target activity

is another key factor. As complementarity between the gRNA spacer and targeted region

as well as local RNA accessibility are essential for RNA targeting (*Abudayyeh et al.,*

2017; Cox et al., 2017; Konermann et al., 2018; Smargon et al., 2017), general

principles for gRNA designing can provide critical help in choosing spacer sequence and

length for gRNA aiming at single-stranded region of target RNA (*Bandaru et al., 2020;*

Wessels et al., 2020).

Limitations and directions for improvement

High concentration of H_2O_2 (1 mM) used in the RPL method may cause oxidative stress

and necrosis to the cells (*Clément and Pervaiz, 2001*) and may preclude the application

of RPL to systems sensitive to oxidative stress and cell harm. Similar to other fusion

proteins, the RPL protein due to its large size (~130kDa) may pose steric hindrance to

access target RNA and increase the biotinylating range, which could reduce specificity

and limit the application to mapping RNA functional domains (*Quinn et al., 2014*) and

RNA-protein interactions at high resolution. Improvement could be achieved using smaller

Cas13 proteins by structure-guided truncations (*Zhang et al., 2018*). Alternatively, CIRTS

strategy could be applied to assemble a much smaller gRNA-dependent RNA proximity

labeling enzyme (*Rauch et al., 2019*).

Another limitation is that RPL and similar tools may not identify RBPs for a target RNA

as efficiently as antisense probe-based methods, as the RPL protein has to compete with

the RBPs bound to the target transcript (*Wessels et al., 2020*). The RPL protein can only

access single-stranded regions of target RNA (*Cox et al., 2017; Smargon et al., 2017*)

and only proteins with electron-rich amino acids like tyrosine exposed on the surface

within the biotinylating range have the opportunity to be labeled (*Rhee et al., 2013*). The

same limitation also applies to other proximity labeling enzymes including BioID and its

relatives and PafA, which all favor lysine as labeling substrate (*Liu et al., 2018;*

Samavarchi-Tehrani et al., 2020).

Perspectives

We anticipate that RPL and similar methods will be widely applied to characterize the

functions and regulation of diverse categories of RNA in multiple cell types and organisms.

Since both Cas13s and proximity labeling are very active research areas, further

optimization and refinements of RPL and similar methods are expected. Utilizations of

these tools together with protein-centric methods (*Licatalosi et al., 2008; Van Nostrand*

et al., 2016), annotation of RNA structure (*Spitale et al., 2015; Sun et al., 2019*) could

shed light on the molecular mechanisms of lncRNA functions, RNA-protein interactions,

RNA functional domain, and binding specificities for RBPs.

Acknowledgements

This work was supported by the Ovarian Cancer Research Alliance (Ann and Sol

Schreiber Mentored Investigator Award 458799 to X.L.) and the National Cancer Institute

(K99/R00-CA184415 and R01-CA207456 to K.L.), the Cedars-Sinai Samuel Oschin

Comprehensive Cancer Institute (Cancer Biology Grant 231433 to K.L.). We thank Drs.

Wei Yang and Bo Zhou at the Cedars-Sinai Medical Center Biomarker Discovery Platform

Core for label-free quantitative mass spectrometry analysis. We also thank anonymous

reviewers for their insightful comments and suggestions.

Author contributions

X.L conceived the project, designed and performed all experiments, and analyzed the

data. R.I.C. and M.A.S.F. analyzed the data. K.L. supervised the project. X.L. and K.L.

wrote the manuscript with input from all authors.

Competing interests

The authors declare no competing interests.

Materials and methods

Key Resources Table

Reagent or Resource	Source	Identifier
Antibodies		
U1-70k	EMD Millipore	Cat.# 05-1588, RRID: AB_11210916
β-tubulin	Sigma Aldrich	Cat.# T8328, RRID: AB_1844090
HA	Santa Cruz	Cat.# sc-7392, RRID: AB_627809
FLAG	Santa Cruz	Cat.# sc-166384, RRID: AB_2017592
Biotin	Santa Cruz	Cat.# sc-57636, RRID: AB_628778
β-actin	Santa Cruz	Cat.# sc-47778, RRID: AB_626632
IgG	Thermo Fisher Scientific	Cat.# 10500C, RRID: AB_2532981
Chemicals		
biotin-phenol	Iris Biotech	Cat.# LS-3500.1000, RRID: N/A
hydrogen peroxide (H_2O_2)	Sigma Aldrich	Cat.# H1009, RRID: N/A
sodium azide	Sigma Aldrich	Cat.# S2002-5G, RRID: N/A
sodium ascorbate	Sigma Aldrich	Cat.# PHR1279-1G, RRID: N/A
Trolox	Sigma Aldrich	Cat.# 238813-1G, RRID: N/A
Dynabeads Protein A	Thermo Fisher Scientific	Cat.#: 10001D, RRID: N/A
Streptavidin magnetic beads	Thermo Fisher Scientific	Cat.# 88817, RRID: N/A
Plasmids		
pC0046-EF1a-PspCas13b-NES-HIV	Addgene	RRID: Addgene_103862
pC0053-CMV-dPspCas13b-GS-ADAR2DD (E488Q)-delta-984-1090	Addgene	RRID: Addgene_103869
pCMV-dPspCas13b-FLAG-APEX2-HA	This study	N/A
pcDNA3-APEX2-NES	Addgene	RRID: Addgene_49386
PspCas13b NTC gRNA	Addgene	RRID: Addgene_103854
PspCas13b U1-1 gRNA	This study	N/A
PspCas13b U1-2 gRNA	This study	N/A
PspCas13b U1-3 gRNA	This study	N/A
PspCas13b poly(A) gRNA	This study	N/A
PspCas13b poly(U) gRNA	This study	N/A
Experimental Models: Cell lines		
Human HEK293T cells	ATCC	Cat.# CRL-3216, RRID: CVCL_0063
Software and Algorithms		
PyMOL	PyMOL	RRID: SCR_000305
Metascape	Metascape	RRID: SCR_016620
STRING	STRING	RRID: SCR_005223
Limma	Limma	RRID: SCR_010943

Plasmids and cloning. pC0046-EF1a-PspCas13b-NES-HIV was a gift from Dr. Feng Zhang (Addgene plasmid # 103862). pCMV-dPspCas13b-FLAG-APEX2-HA (RPL plasmid) was constructed by replacing ADAR2DD-delta-984-1090 in pC0053-CMV-dPspCas13b-GS-ADAR2DD (E488Q)-delta-984-1090 (a gift from Dr. Feng Zhang, Addgene plasmid # 103869) with FLAG-APEX2-HA subcloned from pcDNA3-

APEX2-NES (a gift from Dr. Alice Ting, Addgene plasmid # 49386) using the following primers:

dPspCas13b-For: 5'<u>TACCCATACGATGTTCCA</u>GATTACGCT<u>TAAGCGGCCGCTCGAGTC</u>3',

dPspCas13b-Rev: 5'<u>GTCGTCATCCTTGTAGTCGGATCCCAGTGTCAGTCTTTCAAG</u>3',

FLAG-APEX2-HA-For: 5'<u>GACTACAAGGATGACGACG</u>3',

FLAG-APEX2-HA-Rev: 5'<u>TGGAACATCGTATGGGTACTGCAGGGCATCAGCAAAC</u>3'.

PCR was performed using Q5 High-Fidelity DNA Polymerase (New England Biolabs, Cat.# M0491L). PCR

fragments were assembled using NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs, Cat.#

E2621S) according to manufacturer's instructions. The following spacer sequences were used to express

gRNAs using pC0043-PspCas13b crRNA backbone (a gift from Dr. Feng Zhang, Addgene plasmid #

103854):

NTC: ATGTCTTCCTGGGACGAAGACAA

$U1$-1_{1-30} : ATCATGGTATCTCCCCTGCCAGGTAAGTAT,

$U1$-$2_{101-130}$: CAAATTATGCAGTCGAGTTTCCCACATTTG,

$U1$-$3_{108-137}$: ACTACCACAAATTATGCAGTCGAGTTTCCC,

Poly(A): TTTTTTTTTTTTTTTTTTTTTTTTTTTTTT,

Poly(U) : AAAAAAAAAAAAAAAAAAAAAAAAAAAAAA.

The sequences of all constructs have been confirmed using Sanger sequencing.

Transfection and *in vivo* proximity dependent biotinylation. For validation of *U1* gRNAs in directing the

RPL protein to target *U1*, HEK293T cells were seeded into 12-well plates and were transfected with 1.5 µg

the RPL plasmid and 0.5 µg Cas13b gRNAs (NTC, *U1*-1, *U1*-2, *U1*-3) while ~80% confluency using

Lipofectamine 3000 (Thermo Fisher Scientific, Cat.# L3000015). For RIP experiments, HEK293T cells were

seeded into 6-well plates and were transfected with 2.5 µg the RPL plasmid and 1.5 µg Cas13b gRNAs

while ~80% confluency using Lipofectamine 3000. For proximity-dependent biotinylation, HEK293T cells

were seeded into 150 mm plate and were transfected with 25 µg the RPL plasmid and 15 µg Cas13b gRNAs

(NTC, *U1*-1, *U1*-2, *U1*-3, poly[A], poly[U]) while ~80% confluency using Lipofectamine 3000. HEK293T cells

were incubated with 25 mL of DMEM media containing 25 µL of 500 mM biotin-phenol (Iris Biotech, Cat.#

LS-3500.1000) in DMSO for 30 min at 37 °C 24h post transfection. Cells were then treated with 1 mM

hydrogen peroxide (H_2O_2) (Sigma Aldrich, Cat.# H1009) for 1 min on a horizontal shaker at room

temperature. The labeling solution was aspirated and cells were washed twice with 25 mL of quencher

solution (10 mM sodium azide [Sigma Aldrich, Cat.# S2002-5G], 10 mM sodium ascorbate [Sigma Aldrich,

Cat.# PHR1279-1G], and 5 mM Trolox [Sigma Aldrich, Cat.# 238813-1G] in DPBS (Thermo Fisher Scientific,

Cat.# 14040182). Cells were then washed three times with 15 mL of DPBS and were pelleted by

centrifugation at 1,500 g for 5 min at 4 °C. Cell pellets were snap frozen and stored at −80 °C.

Streptavidin enrichment of biotinylated proteins. Cell pellets from two 150 mm plates of transfected

HEK293T cells were lysed in 2 mL cell lysis buffer (10 mM HEPES, pH7.5 by KOH, 150 mM NaCl, 0.1%

NP-40, 5 mM EGTA, 5 mM Trolox, 10 mM Sodium ascorbate acid, 10 mM Sodium azide, 1 mM PMSF).

Streptavidin magnetic beads (Thermo Fisher Scientific, Cat.# 88817) were washed twice with cell lysis

buffer and 3.5 mg of each whole cell lysate sample were incubated with 100 µL magnetic bead slurry with

rotation for 2 h at room temperature. After enrichment, the flowthrough was removed and beads were

washed with 2 × 1 mL cell lysis buffer, 1mL 1 M KCl, 1 mL 0.1 M Na_2CO_3, 1 mL of 2 M urea in 10 mM Tris-

HCl (pH 8.0), and again with 2 × 1 mL cell lysis buffer. Biotinylated proteins were then eluted by boiling the

magnetic beads in 30 µL 4 × Laemmli sample buffer (Bio-Rad, Cat.# 1610747) supplemented with 20 mM

DTT and 2 mM biotin.

LC-MS/MS and label-free quantitative mass spectrometry proteomic analysis

The streptavidin-enriched proteins were profiled using label-free quantitative mass spectrometry as

previously described (B. Zhou et al., 2019) at Cedars-Sinai Medical Center Biomarker Discovery Platform

Core.

Data analysis for RNA proximal proteins

Data were first filtered to exclude non-human proteins and proteins that were detected in only one or none

of the *U1* replicates or poly(A) replicates. Then proteins detected with two or greater unique peptides were

subjected to log_2 transformation. Only the top gene name was kept from multiple candidates. Since *U1* has

compact structure in pre-B complex and its size is much smaller than the biotinylating range of APEX2,

experiments using *U1* gRNAs (*U1*-1, *U1*-2, *U1*-3) were considered as replicates to compare with

nontargeting controls (NTC1, partially [65%] targeting UBTF; NTC2, targeting poly[A]; NTC3, targeting

poly[U]). Moderated *t*-test with a paired design was used to compare the $\log_2$-transformed iBAQ values

between *U1* and NTC or between poly(A) and poly(U) using limma package (*Smyth, 2004*). *p* values were

adjust using the Benjamini-Hochberg (BH) method (*Benjamini and Hochberg, 1995*) for multiple

comparisons. Proteins with *p* < 0.05 were considered statistically significant. There are 226 *U1* RPPs with

p < 0.05, $\log_2$FC > 2, FDR < 0.25 and 786 poly(A) tail RPPs with BH-adjusted *p* < 0.05, $\log_2$FC > 2.

Comparison RPPs with different gene ontology (GO) terms

Lists of human proteins were retrieved (04/13/2020) from QuickGO (https://www.ebi.ac.uk/QuickGO/) *via*

searching corresponding GO terms and selecting '*Homo sapiens (9606)*' under Taxon, except P-bodies

and stress granule, which were both curated using data summarized from Wikipedia (04/24/2020)

(https://en.wikipedia.org/wiki/P-bodies, https://en.wikipedia.org/wiki/Stress_granule). The venn diagrams

were generated using online tools (http://bioinformatics.psb.ugent.be/webtools/Venn/).

Cellular fractionation. Cells were fractionated as previously described with slight modification (*Lin et al.,*

2019). Six million HEK293T cells were treated with PML buffer (10 mM Tris-HCl, pH 7.5, 0.15% NP-40, 150

mM NaCl) on ice for 4 min after homogenization by flicking. Lysates were loaded onto a 24% sucrose

cushion (24% RNase-free sucrose in PML buffer) using large orifice tips, and centrifuged at 15,000 × *g* for

10 min at 4°C. The supernatant (cytoplasmic fraction) was retained and the pellet (nuclear fraction) was

washed with 1 × PBS/1 mM EDTA and resuspended in 200 μL of 1 × PBS/1 mM EDTA. Fractionation

efficiency was validated by western blot using β-tubulin (Sigma Aldrich, Cat.# T8328, 1:2,000) as

cytoplasmic marker and U1-70k (EMD Millipore, Cat.# 05-1588, 1:1,000) as nuclear marker.

RNA Immunoprecipitation (RIP). RIP was performed as previously described with slight modification (*Lin*

et al., 2019). Twelve microliter Dynabeads Protein A (Thermo Fisher Scientific, Cat.#: 10001D) were

washed with 200 μL HBS (150 mM NaCl, 10 mM HEPES, pH7.5 by KOH) and incubated with 2 μg anti-HA

(Santa Cruz, Cat.# sc-7392) or 2 μg rabbit IgG isotype (Thermo Fisher Scientific, Cat.# 10500C) in the

presence of 80 µL HBS buffer at room temperature for 1 h. Eight million HEK293T cells were lysed with

800 µL cell lysis buffer (HBS, 0.1% NP-40, 5 mM EGTA, supplemented with 1 × protease inhibitor cocktail

[Roche, Cat.# 11873580001], 1 × PhosSTOP protease inhibitor cocktail [Roche, Cat.# 4906837001], 1 mM

PMSF [Sigma Aldrich, Cat.# 93482], and Superase-in [Ambion, Cat.# AM2696]) at 4°C for 1 h. Cell debris

and insoluble proteins were removed by centrifugation at 4°C, 12,000 *g* for 10 min, and the supernatants

were incubated with HA-conjugated or IgG-conjugated Dynabeads at 4°C for 1 h. The Dynabeads were

then washed 3 times with wash buffer (HBS, 0.1% NP-40) and aliquoted into two halves. Proteins

associated with half of the Dynabeads were eluted with 22 µL 4 × Laemmli sample buffer (Bio-Rad, Cat.#

1610747) by boiling at 95 °C for 5 min. RNA was extracted from the other half of Dynabeads using TRIzol

LS (Thermo Fisher Scientific, Cat.# 10296028).

RNA extraction and RT-qPCR. RNA associated with immunoprecipitated RPL fusion protein or RNA from

gRNA transfected cells were extracted using TRIzol LS. M-MLV reverse transcriptase (Promega, Cat.#

M5301) and random hexamers (Promega, Cat.# C1181) were used for reverse transcription. Gene

expression was quantified by RT-qPCR using iQ SYBR Green supermix (Bio-Rad, Cat.# 170-8886). The

relative gene expression was calculated using the $2^{-\Delta\Delta Ct}$ method and normalized to GAPDH. Five

nanograms cDNA was used for RT-qPCR analysis on CFX96 Touch Real-Time PCR Detection System

(Bio-Rad) using the following primer pairs:

U1-RT-For: 5'CCAGGGCGAGGCTTATCCATT3', U1-RT-Rev: 5'GCAGTCCCCCACTACCACAAAT3';

U2-RT-For: TTCTCGGCCTTTTGGCTAAG; U2-RT-Rev: CTCCCTGCTCCAAAAATCCA;

U6-RT-For: GCTTCGGCAGCACATATACTAAAAT; U6-RT-Rev: CGCTTCACGAATTTGCGTGTCAT;

5.8S-RT-For: GGTGGATCACTCGGCTCGT; 5.8S-RT-Rev: GCAAGTGCGTTCGAAGTGTC;

18S-RT-For: 5'CAGCCACCCGAGATTGAGCA3', 18S-RT-Rev: 5'TAGTAGCGACGGGCGTGTG3';

28S-RT-For: CCCAGTGCTCTGAATGTCAA; 28S-RT-Rev: AGTGGGAATCTCGTTCATCC;

GAPDH-RT-For: 5'TGCCAAATATGATGACATCAAGAA3',

GAPDH-RT-Rev: 5'GGAGTGGGTGTCGCTGTTG3'.

Western blot. Protein samples were run on 4-20% gradient precast protein gel (Bio-Rad, Cat.# 456-1096)

and transferred onto PVDF membrane (Bio-Rad, Cat.# 1704157). After 1 h blocking, membranes were

incubated with anti-FLAG (Santa Cruz, Cat.# sc-166384, 1:1,000), anti-HA (Santa Cruz, Cat.# sc-7392,

1:1,000), anti-biotin (Santa Cruz, Cat.# sc-57636, 1:1,000), or anti- β-actin (Santa Cruz, Cat.# sc-47778,

1:2,000) at 4°C overnight. Membranes were washed three times with Tris-buffered saline containing 0.5%

Tween 20 (TBST) before incubating with HRP-conjugated secondary antibody at room temperature for 2 h.

Then the membranes were incubated briefly with ECL Western Blotting Substrate (Thermo Fisher Scientific,

Cat.#: 32106) after three times wash with TBST. The membranes were exposed to HyBlot Autoradiography

Film (Denville Scientific, Cat.#: E3018).

Distance calculation. The distances between *U1* snRNA and *U1* RPPs identified by RPL in the pre-B

complex structure (PDB ID: 6QX9) were measured using PyMOL (*Schrodinger, 2020*). We used the

distance from *U1* snRNA (nucleotide 1) to the proximal residues of *U1* RPPs to estimate the actual distance

(D1). Since there is no structure available for PspCas13b, we used the structure of PbuCas13b (PDB ID:

6DTD) to infer the distance between *U1* RPPs and APEX2 in the RPL protein. Basically, the average

distances between gRNA (nucleotide 1, 12, and 23 of spacer) and the C-terminus of PbuCas13b, where

the APEX2 was fused to, were measured (D2). The inferred distances between APEX2 and RPPs were

then calculated as absolute value of the differences between D1 and D2.

Data availability. Raw images for western blots and raw mass spectrometry data for both *U1* RPPs and

poly(A) RPPs are included as supporting files.

References

Abudayyeh OO, Gootenberg JS, Essletzbichler P, Han S, Joung J, Belanto JJ, Verdine V, Cox DBT, Kellner MJ, Regev A, Lander ES, Voytas DF, Ting AY, Zhang F. 2017. RNA targeting with CRISPR-Cas13. *Nature* **550**:280–284. doi:10.1038/nature24049

Baltz AG, Munschauer M, Schwanhäusser B, Vasile A, Murakawa Y, Schueler M, Youngs N, Penfold-Brown D, Drew K, Milek M, Wyler E, Bonneau R, Selbach M, Dieterich C, Landthaler M. 2012. The mRNA-Bound Proteome and Its Global Occupancy Profile on Protein-Coding Transcripts. *Molecular Cell* **46**:674–690. doi:10.1016/j.molcel.2012.05.021

Bandaru S, Tsuji MH, Shimizu Y, Usami K, Lee S, Takei NK, Yoshitome K, Nishimura Y, Otsuki T, Ito T. 2020. Structure-based design of gRNA for Cas13. *Sci Rep* **10**:11610. doi:10.1038/s41598-020-68459-4

Batista PJ, Chang HY. 2013. Long Noncoding RNAs: Cellular Address Codes in Development and Disease. *Cell* **152**:1298–1307. doi:10.1016/j.cell.2013.02.012

Belousov VV, Fradkov AF, Lukyanov KA, Staroverov DB, Shakhbazov KS, Terskikh AV, Lukyanov S. 2006. Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nat Methods* **3**:281–286. doi:10.1038/nmeth866

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**:289–300. doi:10.1111/j.2517-6161.1995.tb02031.x

Bernard JJ, Cowing-Zitron C, Nakatsuji T, Muehleisen B, Muto J, Borkowski AW, Martinez L, Greidinger EL, Yu BD, Gallo RL. 2012. Ultraviolet radiation damages self noncoding RNA and is detected by TLR3. *Nat Med* **18**:1286–1290. doi:10.1038/nm.2861

Berretta J, Morillon A. 2009. Pervasive transcription constitutes a new level of eukaryotic genome regulation. *EMBO Rep* **10**:973–982. doi:10.1038/embor.2009.181

Blower MD. 2013. Molecular Insights into Intracellular RNA LocalizationInternational Review of Cell and Molecular Biology. Elsevier. pp. 1–39. doi:10.1016/B978-0-12-407699-0.00001-7

Bohnsack MT, Czaplinski K, Gorlich D. 2004. Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs. *RNA* **10**:185–191. doi:10.1261/rna.5167604

Branon T, Han S, Ting A. 2017. Beyond Immunoprecipitation: Exploring New Interaction Spaces with Proximity Biotinylation. *Biochemistry* **56**:3297–3298. doi:10.1021/acs.biochem.7b00466

Branon TC, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, Feldman JL, Perrimon N, Ting AY. 2018. Efficient proximity labeling in living cells and organisms with TurboID. *Nat Biotechnol* **36**:880–887. doi:10.1038/nbt.4201

Cai X. 2004. Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA* **10**:1957–1966. doi:10.1261/rna.7135204

Carlevaro-Fita J, Johnson R. 2019. Global Positioning System: Understanding Long Noncoding RNAs through Subcellular Localization. *Mol Cell* **73**:869–883. doi:10.1016/j.molcel.2019.02.008

Castello A, Fischer B, Eichelbaum K, Horos R, Beckmann BM, Strein C, Davey NE, Humphreys DT, Preiss T, Steinmetz LM, Krijgsveld J, Hentze MW. 2012. Insights into RNA biology from an atlas of mammalian mRNA-binding proteins. *Cell* **149**:1393–1406. doi:10.1016/j.cell.2012.04.031

Caudron-Herger M, Rusin SF, Adamo ME, Seiler J, Schmid VK, Barreau E, Kettenbach AN, Diederichs S. 2019. R-DeeP: Proteome-wide and Quantitative Identification of RNA-Dependent Proteins by Density Gradient Ultracentrifugation. *Mol Cell* **75**:184-199.e10. doi:10.1016/j.molcel.2019.04.018

Chang H, Lim J, Ha M, Kim VN. 2014. TAIL-seq: Genome-wide Determination of Poly(A) Tail Length and 3′ End Modifications. *Molecular Cell* **53**:1044–1052. doi:10.1016/j.molcel.2014.02.007

Chang Y-F, Imam JS, Wilkinson MF. 2007. The Nonsense-Mediated Decay RNA Surveillance Pathway. *Annu Rev Biochem* **76**:51–74. doi:10.1146/annurev.biochem.76.050106.093909

Charenton C, Wilkinson ME, Nagai K. 2019. Mechanism of 5' splice site transfer for human spliceosome activation. *Science* **364**:362–367. doi:10.1126/science.aax3289

Chen C-YA, Shyu A-B. 2011. Mechanisms of deadenylation-dependent decay: Mechanisms of deadenylation-dependent decay. *WIREs RNA* **2**:167–183. doi:10.1002/wrna.40

Chlebowski A, Lubas M, Jensen TH, Dziembowski A. 2013. RNA decay machines: The exosome. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms* **1829**:552–560. doi:10.1016/j.bbagrm.2013.01.006

Chu Ci, Spitale RC, Chang HY. 2015. Technologies to probe functions and mechanisms of long noncoding
RNAs. *Nat Struct Mol Biol* **22**:29–35. doi:10.1038/nsmb.2921
Chu C., Zhang QC, da Rocha ST, Flynn RA, Bharadwaj M, Calabrese JM, Magnuson T, Heard E, Chang
HY. 2015. Systematic discovery of Xist RNA binding proteins. *Cell* **161**:404–16.
doi:10.1016/j.cell.2015.03.025
Cioni J-M, Lin JQ, Holtermann AV, Koppers M, Jakobs MAH, Azizi A, Turner-Bridger B, Shigeoka T, Franze
K, Harris WA, Holt CE. 2019. Late Endosomes Act as mRNA Translation Platforms and Sustain
Mitochondria in Axons. *Cell* **176**:56-72.e15. doi:10.1016/j.cell.2018.11.030
Clément M-V, Pervaiz S. 2001. Intracellular superoxide and hydrogen peroxide concentrations: a critical
balance that determines survival or death. *Redox Report* **6**:211–214.
doi:10.1179/135100001101536346
Collart MA. 2016. The Ccr4-Not complex is a key regulator of eukaryotic gene expression. *Wiley Interdiscip*
Rev RNA **7**:438–454. doi:10.1002/wrna.1332
Cox DBT, Gootenberg JS, Abudayyeh OO, Franklin B, Kellner MJ, Joung J, Zhang F. 2017. RNA editing
with CRISPR-Cas13. *Science* **358**:1019–1027. doi:10.1126/science.aaq0180
Cvitkovic I, Jurica MS. 2013. Spliceosome database: a tool for tracking components of the spliceosome.
Nucleic Acids Res **41**:D132-141. doi:10.1093/nar/gks999
Damgaard CK, Kahns S, Lykke-Andersen S, Nielsen AL, Jensen TH, Kjems J. 2008. A 5′ Splice Site
Enhances the Recruitment of Basal Transcription Initiation Factors In Vivo. *Molecular Cell* **29**:271–
278. doi:10.1016/j.molcel.2007.11.035
Decker CJ, Parker R. 2012. P-Bodies and Stress Granules: Possible Roles in the Control of Translation
and mRNA Degradation. *Cold Spring Harbor Perspectives in Biology* **4**:a012286–a012286.
doi:10.1101/cshperspect.a012286
Deckert J, Hartmuth K, Boehringer D, Behzadnia N, Will CL, Kastner B, Stark H, Urlaub H, Lührmann R.
2006. Protein Composition and Electron Microscopy Structure of Affinity-Purified Human
Spliceosomal B Complexes Isolated under Physiological Conditions. *MCB* **26**:5528–5543.
doi:10.1128/MCB.00582-06
Derrien T, Johnson R, Bussotti G, Tanzer A, Djebali S, Tilgner H, Guernec G, Martin D, Merkel A, Knowles
DG, Lagarde J, Veeravalli L, Ruan X, Ruan Y, Lassmann T, Carninci P, Brown JB, Lipovich L,
Gonzalez JM, Thomas M, Davis CA, Shiekhattar R, Gingeras TR, Hubbard TJ, Notredame C,
Harrow J, Guigó R. 2012. The GENCODE v7 catalog of human long noncoding RNAs: analysis of
their gene structure, evolution, and expression. *Genome Res* **22**:1775–89.
doi:10.1101/gr.132159.111
Djebali S, Davis CA, Merkel A, Dobin A, Lassmann T, Mortazavi A, Tanzer A, Lagarde J, Lin W, Schlesinger
F, Xue C, Marinov GK, Khatun J, Williams BA, Zaleski C, Rozowsky J, Röder M, Kokocinski F,
Abdelhamid RF, Alioto T, Antoshechkin I, Baer MT, Bar NS, Batut P, Bell K, Bell I, Chakrabortty S,
Chen X, Chrast J, Curado J, Derrien T, Drenkow J, Dumais E, Dumais J, Duttagupta R, Falconnet
E, Fastuca M, Fejes-Toth K, Ferreira P, Foissac S, Fullwood MJ, Gao H, Gonzalez D, Gordon A,
Gunawardena H, Howald C, Jha S, Johnson R, Kapranov P, King B, Kingswood C, Luo OJ, Park
E, Persaud K, Preall JB, Ribeca P, Risk B, Robyr D, Sammeth M, Schaffer L, See LH, Shahab A,
Skancke J, Suzuki AM, Takahashi H, Tilgner H, Trout D, Walters N, Wang H, Wrobel J, Yu Y, Ruan
X, Hayashizaki Y, Harrow J, Gerstein M, Hubbard T, Reymond A, Antonarakis SE, Hannon G,
Giddings MC, Ruan Y, Wold B, Carninci P, Guigó R, Gingeras TR. 2012. Landscape of transcription
in human cells. *Nature* **489**:101–8. doi:10.1038/nature11233
Dreyfus M, Régnier P. 2002. The Poly(A) Tail of mRNAs. *Cell* **111**:611–613. doi:10.1016/S0092-
8674(02)01137-6
Dreyfuss G, Kim VN, Kataoka N. 2002. Messenger-RNA-binding proteins and the messages they carry.
Nat Rev Mol Cell Biol **3**:195–205. doi:10.1038/nrm760
ElMaghraby MF, Andersen PR, Pühringer F, Hohmann U, Meixner K, Lendl T, Tirian L, Brennecke J. 2019.
A Heterochromatin-Specific RNA Export Pathway Facilitates piRNA Production. *Cell* **178**:964-
979.e20. doi:10.1016/j.cell.2019.07.007
Engreitz JM, Sirokman K, McDonel P, Shishkin AA, Surka C, Russell P, Grossman SR, Chow AY, Guttman
M, Lander ES. 2014. RNA-RNA Interactions Enable Specific Targeting of Noncoding RNAs to
Nascent Pre-mRNAs and Chromatin Sites. *Cell* **159**:188–199. doi:10.1016/j.cell.2014.08.018
Fazal FM, Han S, Parker KR, Kaewsapsak P, Xu J, Boettiger AN, Chang HY, Ting AY. 2019. Atlas of
Subcellular RNA Localization Revealed by APEX-Seq. *Cell*. doi:10.1016/j.cell.2019.05.027

Fujiwara Y, Furuta A, Kikuchi H, Aizawa S, Hatanaka Y, Konya C, Uchida K, Yoshimura A, Tamai Y, Wada
K, Kabuta T. 2013. Discovery of a novel type of autophagy targeting RNA. *Autophagy* **9**:403–409.
doi:10.4161/auto.23002
Gesteland RF. 1993. The RNA world: the nature of modern RNA suggests a prebiotic RNA world. New
York: Cold Spring Harbor Laboratory Press.
Ghildiyal M, Zamore PD. 2009. Small silencing RNAs: an expanding universe. *Nat Rev Genet* **10**:94–108.
doi:10.1038/nrg2504
Gilmartin GM. 2005. Eukaryotic mRNA 3' processing: a common means to different ends. *Genes Dev*
19:2517–2521. doi:10.1101/gad.1378105
Glisovic T, Bachorik JL, Yong J, Dreyfuss G. 2008. RNA-binding proteins and post-transcriptional gene
regulation. *FEBS Lett* **582**:1977–1986. doi:10.1016/j.febslet.2008.03.004
Gootenberg JS, Abudayyeh OO, Kellner MJ, Joung J, Collins JJ, Zhang F. 2018. Multiplexed and portable
nucleic acid detection platform with Cas13, Cas12a, and Csm6. *Science* **360**:439–444.
doi:10.1126/science.aaq0179
Graindorge A, Pinheiro I, Nawrocka A, Mallory AC, Tsvetkov P, Gil N, Carolis C, Buchholz F, Ulitsky I,
Heard E, Taipale M, Shkumatava A. 2019. In-cell identification and measurement of RNA-protein
interactions. *Nat Commun* **10**:5317. doi:10.1038/s41467-019-13235-w
Guttman M, Amit I, Garber M, French C, Lin MF, Feldser D, Huarte M, Zuk O, Carey BW, Cassady JP,
Cabili MN, Jaenisch R, Mikkelsen TS, Jacks T, Hacohen N, Bernstein BE, Kellis M, Regev A, Rinn
JL, Lander ES. 2009. Chromatin signature reveals over a thousand highly conserved large non-
coding RNAs in mammals. *Nature* **458**:223–7. doi:10.1038/nature07672
Han S, Zhao BS, Myers SA, Carr SA, He C, Ting AY. 2020. RNA–protein interaction mapping via MS2- or
Cas13-based APEX targeting. *Proc Natl Acad Sci USA* 202006617. doi:10.1073/pnas.2006617117
Hentze MW, Castello A, Schwarzl T, Preiss T. 2018. A brave new world of RNA-binding proteins. *Nat Rev*
Mol Cell Biol **19**:327–341. doi:10.1038/nrm.2017.130
Holt CE, Bullock SL. 2009. Subcellular mRNA Localization in Animal Cells and Why It Matters. *Science*
326:1212–1216. doi:10.1126/science.1176488
Huang BK, Sikes HD. 2014. Quantifying intracellular hydrogen peroxide perturbations in terms of
concentration. *Redox Biology* **2**:955–962. doi:10.1016/j.redox.2014.08.001
Huang Y, Carmichael GG. 1996. Role of polyadenylation in nucleocytoplasmic transport of mRNA. *Mol Cell*
Biol **16**:1534–1542. doi:10.1128/MCB.16.4.1534
Huang Y, Guo Q, Ding X-P, Wang X. 2020. Mechanism of long noncoding RNAs as transcriptional
regulators in cancer. *RNA Biology* 1–13. doi:10.1080/15476286.2019.1710405
International Human Genome Sequencing Consortium. 2001. Initial sequencing and analysis of the human
genome. *Nature* **409**:860–921. doi:10.1038/35057062
Iyer MK, Niknafs YS, Malik R, Singhal U, Sahu A, Hosono Y, Barrette TR, Prensner JR, Evans JR, Zhao S,
Poliakov A, Cao X, Dhanasekaran SM, Wu Y-M, Robinson DR, Beer DG, Feng FY, Iyer HK,
Chinnaiyan AM. 2015. The landscape of long noncoding RNAs in the human transcriptome. *Nat*
Genet **47**:199–208. doi:10.1038/ng.3192
Kadaba S, Wang X, Anderson JT. 2006. Nuclear RNA surveillance in Saccharomyces cerevisiae: Trf4p-
dependent polyadenylation of nascent hypomethylated tRNA and an aberrant form of 5S rRNA.
RNA **12**:508–521. doi:10.1261/rna.2305406
Kim DI, Jensen SC, Noble KA, Kc B, Roux KH, Motamedchaboki K, Roux KJ. 2016. An improved smaller
biotin ligase for BioID proximity labeling. *MBoC* **27**:1188–1196. doi:10.1091/mbc.E15-12-0844
Kim VN. 2005. MicroRNA biogenesis: coordinated cropping and dicing. *Nat Rev Mol Cell Biol* **6**:376–385.
doi:10.1038/nrm1644
Kneuss E, Munafò M, Eastwood EL, Deumer U-S, Preall JB, Hannon GJ, Czech B. 2019. Specialization of
the Drosophila nuclear export family protein Nxf3 for piRNA precursor export. *Genes Dev* **33**:1208–
1220. doi:10.1101/gad.328690.119
Köhler A, Hurt E. 2007. Exporting RNA from the nucleus to the cytoplasm. *Nat Rev Mol Cell Biol* **8**:761–
773. doi:10.1038/nrm2255
Konermann S, Lotfy P, Brideau NJ, Oki J, Shokhirev MN, Hsu PD. 2018. Transcriptome Engineering with
RNA-Targeting Type VI-D CRISPR Effectors. *Cell* **173**:665-676.e14.
doi:10.1016/j.cell.2018.02.033
Kopp F, Mendell JT. 2018. Functional Classification and Experimental Dissection of Long Noncoding RNAs.
Cell **172**:393–407. doi:10.1016/j.cell.2018.01.011

Kruse C, Willkomm DK, Grünweller A, Vollbrandt T, Sommer S, Busch S, Pfeiffer T, Brinkmann J, Hartmann
RK, Müller PK. 2000. Export and transport of tRNA are coupled to a multi-protein complex. *Biochem*
J **346 Pt 1**:107–115.
Kwek KY, Murphy S, Furger A, Thomas B, O'Gorman W, Kimura H, Proudfoot NJ, Akoulitchev A. 2002. U1
snRNA associates with TFIIH and regulates transcriptional initiation. *Nat Struct Biol.*
doi:10.1038/nsb862
Kwon SC, Yi H, Eichelbaum K, Föhr S, Fischer B, You KT, Castello A, Krijgsveld J, Hentze MW, Kim VN.
2013. The RNA-binding protein repertoire of embryonic stem cells. *Nat Struct Mol Biol* **20**:1122–
1130. doi:10.1038/nsmb.2638
Lai W-JC, Kayedkhordeh M, Cornell EV, Farah E, Bellaousov S, Rietmeijer R, Salsi E, Mathews DH,
Ermolenko DN. 2018. mRNAs and lncRNAs intrinsically form secondary structures with short end-
to-end distances. *Nat Commun* **9**:4328. doi:10.1038/s41467-018-06792-z
Lam SS, Martell JD, Kamer KJ, Deerinck TJ, Ellisman MH, Mootha VK, Ting AY. 2015. Directed evolution
of APEX2 for electron microscopy and proximity labeling. *Nat Methods* **12**:51–54.
doi:10.1038/nmeth.3179
Laprade H, Querido E, Smith MJ, Guérit D, Crimmins H, Conomos D, Pourret E, Chartrand P, Sfeir A. 2020.
Single-Molecule Imaging of Telomerase RNA Reveals a Recruitment-Retention Model for
Telomere Elongation. *Molecular Cell* **79**:115-126.e6. doi:10.1016/j.molcel.2020.05.005
Laroia G, Cuesta R, Brewer G, Schneider RJ. 1999. Control of mRNA decay by heat shock-ubiquitin-
proteasome pathway. *Science* **284**:499–502. doi:10.1126/science.284.5413.499
Laroia G, Sarkar B, Schneider RJ. 2002. Ubiquitin-dependent mechanism regulates rapid turnover of AU-
rich cytokine mRNAs. *Proc Natl Acad Sci USA* **99**:1842–1846. doi:10.1073/pnas.042575699
Lebbink RJ, Lowe M, Chan T, Khine H, Wang X, McManus MT. 2011. Polymerase II promoter strength
determines efficacy of microRNA adapted shRNAs. *PLoS ONE* **6**:e26213.
doi:10.1371/journal.pone.0026213
Leija-Martínez N, Casas-Flores S, Cadena-Nava RD, Roca JA, Mendez-Cabañas JA, Gomez E, Ruiz-
Garcia J. 2014. The separation between the 5′-3′ ends in long RNA molecules is short and nearly
constant. *Nucleic Acids Research* **42**:13963–13968. doi:10.1093/nar/gku1249
Lemmens I, Jansen S, de Rouck S, de Smet A-S, Defever D, Neyts J, Dallmeier K, Tavernier J. 2020. The
Development of RNA-KISS, a Mammalian Three-Hybrid Method to Detect RNA–Protein
Interactions in Living Mammalian Cells. *J Proteome Res* acs.jproteome.0c00068.
doi:10.1021/acs.jproteome.0c00068
Li J-H, Liu S, Zhou H, Qu L-H, Yang J-H. 2014. starBase v2.0: decoding miRNA-ceRNA, miRNA-ncRNA
and protein–RNA interaction networks from large-scale CLIP-Seq data. *Nucl Acids Res* **42**:D92–
D97. doi:10.1093/nar/gkt1248
Li X, Fu X-D. 2019. Chromatin-associated RNAs as facilitators of functional genomic interactions. *Nat Rev*
Genet. doi:10.1038/s41576-019-0135-1
Li XZ, Roy CK, Dong X, Bolcun-Filas E, Wang J, Han BW, Xu J, Moore MJ, Schimenti JC, Weng Z, Zamore
PD. 2013. An ancient transcription factor initiates the burst of piRNA production during early
meiosis in mouse testes. *Mol Cell* **50**:67–81. doi:10.1016/j.molcel.2013.02.016
Li Y, Liu S, Cao L, Luo Y, Du H, Li S, You F. 2020. CBRPP: a new RNA-centric method to study RNA-
protein interactions (preprint). Molecular Biology. doi:10.1101/2020.04.09.033290
Licatalosi DD, Mele A, Fak JJ, Ule J, Kayikci M, Chi SW, Clark TA, Schweitzer AC, Blume JE, Wang X,
Darnell JC, Darnell RB. 2008. HITS-CLIP yields genome-wide insights into brain alternative RNA
processing. *Nature* **456**:464–469. doi:10.1038/nature07488
Lim J, Kim D, Lee Y, Ha M, Lee M, Yeo J, Chang H, Song J, Ahn K, Kim VN. 2018. Mixed tailing by TENT4A
and TENT4B shields mRNA from rapid deadenylation. *Science* **361**:701–704.
doi:10.1126/science.aam5794
Lin X, Spindler TJ, de Souza Fonseca MA, Corona RI, Seo J-H, Dezem FS, Li L, Lee JM, Long HW, Sellers
TA, Karlan BY, Noushmehr H, Freedman ML, Gayther SA, Lawrenson K. 2019. Super-Enhancer-
Associated LncRNA UCA1 Interacts Directly with AMOT to Activate YAP Target Genes in Epithelial
Ovarian Cancer. *iScience* **17**:242–255. doi:10.1016/j.isci.2019.06.025
Liu Q, Zheng J, Sun W, Huo Y, Zhang L, Hao P, Wang H, Zhuang M. 2018. A proximity-tagging system to
identify membrane protein–protein interactions. *Nat Methods* **15**:715–722. doi:10.1038/s41592-
018-0100-5

Lund E, Güttinger S, Calado A, Dahlberg JE, Kutay U. 2004. Nuclear export of microRNA precursors.
Science **303**:95–98. doi:10.1126/science.1090599

Lunde BM, Moore C, Varani G. 2007. RNA-binding proteins: modular design for efficient function. *Nat Rev*
Mol Cell Biol **8**:479–490. doi:10.1038/nrm2178

Lyublinskaya O, Antunes F. 2019. Measuring intracellular concentration of hydrogen peroxide with the use
of genetically encoded H_2O_2 biosensor HyPer. *Redox Biology* **24**:101200.
doi:10.1016/j.redox.2019.101200

Malone CD, Hannon GJ. 2009. Small RNAs as guardians of the genome. *Cell* **136**:656–668.
doi:10.1016/j.cell.2009.01.045

McHugh CA, Chen CK, Chow A, Surka CF, Tran C, McDonel P, Pandya-Jones A, Blanco M, Burghard C,
Moradian A, Sweredoski MJ, Shishkin AA, Su J, Lander ES, Hess S, Plath K, Guttman M. 2015.
The Xist lncRNA interacts directly with SHARP to silence transcription through HDAC3. *Nature*
521:232–6. doi:10.1038/nature14443

Metkar M, Ozadam H, Lajoie BR, Imakaev M, Mirny LA, Dekker J, Moore MJ. 2018. Higher-Order
Organization Principles of Pre-translational mRNPs. *Molecular Cell* **72**:715-726.e3.
doi:10.1016/j.molcel.2018.09.012

Milek M, Imami K, Mukherjee N, Bortoli FD, Zinnall U, Hazapis O, Trahan C, Oeffinger M, Heyd F, Ohler U,
Selbach M, Landthaler M. 2017. DDX54 regulates transcriptome dynamics during DNA damage
response. *Genome Res* **27**:1344–1359. doi:10.1101/gr.218438.116

Mofatteh M, Bullock SL. 2017. SnapShot: Subcellular mRNA Localization. *Cell* **169**:178-178.e1.
doi:10.1016/j.cell.2017.03.004

Mugridge JS, Coller J, Gross JD. 2018. Structural and molecular mechanisms for the control of eukaryotic
5′–3′ mRNA decay. *Nat Struct Mol Biol* **25**:1077–1085. doi:10.1038/s41594-018-0164-z

Muhlrad D, Decker CJ, Parker R. 1994. Deadenylation of the unstable mRNA encoded by the yeast MFA2
gene leads to decapping followed by 5'-->3' digestion of the transcript. *Genes Dev* **8**:855–866.
doi:10.1101/gad.8.7.855

Mukherjee J, Hermesh O, Eliscovich C, Nalpas N, Franz-Wachtel M, Maček B, Jansen R-P. 2019. β-Actin
mRNA interactome mapping by proximity biotinylation. *Proc Natl Acad Sci USA* **116**:12863–12872.
doi:10.1073/pnas.1820737116

Munro S. 2011. The Golgin Coiled-Coil Proteins of the Golgi Apparatus. *Cold Spring Harbor Perspectives*
in Biology **3**:a005256–a005256. doi:10.1101/cshperspect.a005256

Norbury CJ. 2013. Cytoplasmic RNA: a case of the tail wagging the dog. *Nat Rev Mol Cell Biol* **14**:643–653.
doi:10.1038/nrm3645

Okamura M, Inose H, Masuda S. 2015. RNA Export through the NPC in Eukaryotes. *Genes* **6**:124–149.
doi:10.3390/genes6010124

Padrón A, Iwasaki S, Ingolia NT. 2019. Proximity RNA Labeling by APEX-Seq Reveals the Organization of
Translation Initiation Complexes and Repressive RNA Granules. *Molecular Cell* **75**:875-887.e5.
doi:10.1016/j.molcel.2019.07.030

Palacios I. 1997. Nuclear import of U snRNPs requires importin beta. *The EMBO Journal* **16**:6783–6792.
doi:10.1093/emboj/16.22.6783

Panhale A, Richter FM, Ramírez F, Shvedunova M, Manke T, Mittler G, Akhtar A. 2019. CAPRI enables
comparison of evolutionarily conserved RNA interacting regions. *Nat Commun* **10**:2682.
doi:10.1038/s41467-019-10585-3

Pomeranz Krummel DA, Oubridge C, Leung AKW, Li J, Nagai K. 2009. Crystal structure of human
spliceosomal U1 snRNP at 5.5 Å resolution. *Nature* **458**:475–480. doi:10.1038/nature07851

Queiroz RML, Smith T, Villanueva E, Marti-Solano M, Monti M, Pizzinga M, Mirea D-M, Ramakrishna M,
Harvey RF, Dezi V, Thomas GH, Willis AE, Lilley KS. 2019. Comprehensive identification of RNA-
protein interactions in any organism using orthogonal organic phase separation (OOPS). *Nat*
Biotechnol **37**:169–178. doi:10.1038/s41587-018-0001-2

Quinn JJ, Ilik IA, Qu K, Georgiev P, Chu C, Akhtar A, Chang HY. 2014. Revealing long noncoding RNA
architecture and functions using domain-specific chromatin isolation by RNA purification. *Nat*
Biotechnol **32**:933–940. doi:10.1038/nbt.2943

Ramanathan M, Majzoub K, Rao DS, Neela PH, Zarnegar BJ, Mondal S, Roth JG, Gai H, Kovalski JR,
Siprashvili Z, Palmer TD, Carette JE, Khavari PA. 2018. RNA-protein interaction detection in living
cells. *Nat Methods* **15**:207–212. doi:10.1038/nmeth.4601

Ramanathan M, Porter DF, Khavari PA. 2019. Methods to study RNA-protein interactions. *Nat Methods* **16**:225–234. doi:10.1038/s41592-019-0330-1

Rauch S, He E, Srienc M, Zhou H, Zhang Z, Dickinson BC. 2019. Programmable RNA-Guided RNA Effector Proteins Built from Human Parts. *Cell* S0092867419306208. doi:10.1016/j.cell.2019.05.049

Rhee H-W, Zou P, Udeshi ND, Martell JD, Mootha VK, Carr SA, Ting AY. 2013. Proteomic Mapping of Mitochondria in Living Cells via Spatially Restricted Enzymatic Tagging. *Science* **339**:1328–1331. doi:10.1126/science.1230593

Roux KJ, Kim DI, Burke B. 2013. BioID: a screen for protein-protein interactions. *Curr Protoc Protein Sci* **74**:19.23.1-19.23.14. doi:10.1002/0471140864.ps1923s74

Roux KJ, Kim DI, Raida M, Burke B. 2012. A promiscuous biotin ligase fusion protein identifies proximal and interacting proteins in mammalian cells. *Journal of Cell Biology* **196**:801–810. doi:10.1083/jcb.201112098

Samavarchi-Tehrani P, Samson R, Gingras A-C. 2020. Proximity Dependent Biotinylation: Key Enzymes and Adaptation to Proteomics Approaches. *Mol Cell Proteomics* **19**:757–773. doi:10.1074/mcp.R120.001941

Schrodinger. 2015. The PyMOL Molecular Graphics System, Version 1.8.

Shi Y, Manley JL. 2015. The end of the message: multiple protein–RNA interactions define the mRNA polyadenylation site. *Genes Dev* **29**:889–897. doi:10.1101/gad.261974.115

Simon AE, Miller WA. 2013. 3' cap-independent translation enhancers of plant viruses. *Annu Rev Microbiol* **67**:21–42. doi:10.1146/annurev-micro-092412-155609

Smargon AA, Cox DBT, Pyzocha NK, Zheng K, Slaymaker IM, Gootenberg JS, Abudayyeh OA, Essletzbichler P, Shmakov S, Makarova KS, Koonin EV, Zhang F. 2017. Cas13b Is a Type VI-B CRISPR-Associated RNA-Guided RNase Differentially Regulated by Accessory Proteins Csx27 and Csx28. *Mol Cell* **65**:618-630.e7. doi:10.1016/j.molcel.2016.12.023

Smyth GK. 2004. Linear Models and Empirical Bayes Methods for Assessing Differential Expression in Microarray Experiments. *Statistical Applications in Genetics and Molecular Biology* **3**:1–25. doi:10.2202/1544-6115.1027

So BR, Di C, Cai Z, Venters CC, Guo J, Oh J-M, Arai C, Dreyfuss G. 2019. A Complex of U1 snRNP with Cleavage and Polyadenylation Factors Controls Telescripting, Regulating mRNA Transcription in Human Cells. *Molecular Cell* **76**:590-599.e4. doi:10.1016/j.molcel.2019.08.007

Spiluttini B, Gu B, Belagal P, Smirnova AS, Nguyen VT, Hebert C, Schmidt U, Bertrand E, Darzacq X, Bensaude O. 2010. Splicing-independent recruitment of U1 snRNP to a transcription unit in living cells. *Journal of Cell Science* **123**:2085–2093. doi:10.1242/jcs.061358

Spitale RC, Flynn RA, Zhang QC, Crisalli P, Lee B, Jung J-W, Kuchelmeister HY, Batista PJ, Torre EA, Kool ET, Chang HY. 2015. Structural imprints in vivo decode RNA regulatory mechanisms. *Nature* **519**:486–490. doi:10.1038/nature14263

Stark H, Dube P, Lührmann R, Kastner B. 2001. Arrangement of RNA and proteins in the spliceosomal U1 small nuclear ribonucleoprotein particle. *Nature* **409**:539–542. doi:10.1038/35054102

Sun L, Fazal FM, Li P, Broughton JP, Lee B, Tang L, Huang W, Kool ET, Chang HY, Zhang QC. 2019. RNA structure maps across mammalian cellular compartments. *Nat Struct Mol Biol* **26**:322–330. doi:10.1038/s41594-019-0200-7

Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, Simonovic M, Doncheva NT, Morris JH, Bork P, Jensen LJ, Mering C von. 2019. STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res* **47**:D607–D613. doi:10.1093/nar/gky1131

Tarn W-Y, Chang T-H. 2009. The current understanding of Ded1p/DDX3 homologs from yeast to human. *RNA Biology* **6**:17–20. doi:10.4161/rna.6.1.7440

Tian B. 2005. A large-scale analysis of mRNA polyadenylation of human and mouse genes. *Nucleic Acids Research* **33**:201–212. doi:10.1093/nar/gki158

Trendel J, Schwarzl T, Horos R, Prakash A, Bateman A, Hentze MW, Krijgsveld J. 2019. The Human RNA-Binding Proteome and Its Dynamics during Translational Arrest. *Cell* **176**:391-403.e19. doi:10.1016/j.cell.2018.11.004

Truniger V, Miras M, Aranda MA. 2017. Structural and Functional Diversity of Plant Virus 3'-Cap-Independent Translation Enhancers (3'-CITEs). *Front Plant Sci* **8**:2047. doi:10.3389/fpls.2017.02047

Urdaneta EC, Beckmann BM. 2019. Fast and unbiased purification of RNA-protein complexes after UV
cross-linking. *Methods* S1046202318304869. doi:10.1016/j.ymeth.2019.09.013

Van Nostrand EL, Pratt GA, Shishkin AA, Gelboin-Burkhart C, Fang MY, Sundararaman B, Blue SM,
Nguyen TB, Surka C, Elkins K, Stanton R, Rigo F, Guttman M, Yeo GW. 2016. Robust
transcriptome-wide discovery of RNA-binding protein binding sites with enhanced CLIP (eCLIP).
Nat Methods **13**:508–514. doi:10.1038/nmeth.3810

Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, Smith HO, Yandell M, Evans CA, Holt RA,
Gocayne JD, Amanatides P, Ballew RM, Huson DH, Wortman JR, Zhang Q, Kodira CD, Zheng XH,
Chen L, Skupski M, Subramanian G, Thomas PD, Zhang J, Gabor Miklos GL, Nelson C, Broder S,
Clark AG, Nadeau J, McKusick VA, Zinder N, Levine AJ, Roberts RJ, Simon M, Slayman C,
Hunkapiller M, Bolanos R, Delcher A, Dew I, Fasulo D, Flanigan M, Florea L, Halpern A,
Hannenhalli S, Kravitz S, Levy S, Mobarry C, Reinert K, Remington K, Abu-Threideh J, Beasley E,
Biddick K, Bonazzi V, Brandon R, Cargill M, Chandramouliswaran I, Charlab R, Chaturvedi K, Deng
Z, Francesco VD, Dunn P, Eilbeck K, Evangelista C, Gabrielian AE, Gan W, Ge W, Gong F, Gu Z,
Guan P, Heiman TJ, Higgins ME, Ji R-R, Ke Z, Ketchum KA, Lai Z, Lei Y, Li Z, Li J, Liang Y, Lin X,
Lu F, Merkulov GV, Milshina N, Moore HM, Naik AK, Narayan VA, Neelam B, Nusskern D, Rusch
DB, Salzberg S, Shao W, Shue B, Sun J, Wang ZY, Wang A, Wang X, Wang J, Wei M-H, Wides
R, Xiao C, Yan C, Yao A, Ye J, Zhan M, Zhang W, Zhang H, Zhao Q, Zheng L, Zhong F, Zhong W,
Zhu SC, Zhao S, Gilbert D, Baumhueter S, Spier G, Carter C, Cravchik A, Woodage T, Ali F, An H,
Awe A, Baldwin D, Baden H, Barnstead M, Barrow I, Beeson K, Busam D, Carver A, Center A,
Cheng ML, Curry L, Danaher S, Davenport L, Desilets R, Dietz S, Dodson K, Doup L, Ferriera S,
Garg N, Gluecksmann A, Hart B, Haynes J, Haynes C, Heiner C, Hladun S, Hostin D, Houck J,
Howland T, Ibegwam C, Johnson J, Kalush F, Kline L, Koduru S, Love A, Mann F, May D,
McCawley S, McIntosh T, McMullen I, Moy M, Moy L, Murphy B, Nelson K, Pfannkoch C, Pratts E,
Puri V, Qureshi H, Reardon M, Rodriguez R, Rogers Y-H, Romblad D, Ruhfel B, Scott R, Sitter C,
Smallwood M, Stewart E, Strong R, Suh E, Thomas R, Tint NN, Tse S, Vech C, Wang G, Wetter J,
Williams S, Williams M, Windsor S, Winn-Deen E, Wolfe K, Zaveri J, Zaveri K, Abril JF, Guigó R,
Campbell MJ, Sjolander KV, Karlak B, Kejariwal A, Mi H, Lazareva B, Hatton T, Narechania A,
Diemer K, Muruganujan A, Guo N, Sato S, Bafna V, Istrail S, Lippert R, Schwartz R, Walenz B,
Yooseph S, Allen D, Basu A, Baxendale J, Blick L, Caminha M, Carnes-Stine J, Caulk P, Chiang
Y-H, Coyne M, Dahlke C, Mays AD, Dombroski M, Donnelly M, Ely D, Esparham S, Fosler C, Gire
H, Glanowski S, Glasser K, Glodek A, Gorokhov M, Graham K, Gropman B, Harris M, Heil J,
Henderson S, Hoover J, Jennings D, Jordan C, Jordan J, Kasha J, Kagan L, Kraft C, Levitsky A,
Lewis M, Liu X, Lopez J, Ma D, Majoros W, McDaniel J, Murphy S, Newman M, Nguyen T, Nguyen
N, Nodell M, Pan S, Peck J, Peterson M, Rowe W, Sanders R, Scott J, Simpson M, Smith T,
Sprague A, Stockwell T, Turner R, Venter E, Wang M, Wen M, Wu D, Wu M, Xia A, Zandieh A,
Zhu X. 2001. The Sequence of the Human Genome. *Science* **291**:1304–1351.
doi:10.1126/science.1058040

Weber G, Trowitzsch S, Kastner B, Lührmann R, Wahl MC. 2010. Functional organization of the Sm core
in the crystal structure of human U1 snRNP. *EMBO J* **29**:4172–4184. doi:10.1038/emboj.2010.295

Wessels H-H, Méndez-Mancilla A, Guo X, Legut M, Daniloski Z, Sanjana NE. 2020. Massively parallel
Cas13 screens reveal principles for guide RNA design. *Nat Biotechnol*. doi:10.1038/s41587-020-
0456-9

Wilk R, Hu J, Blotsky D, Krause HM. 2016. Diverse and pervasive subcellular distributions for both coding
and long noncoding RNAs. *Genes Dev* **30**:594–609. doi:10.1101/gad.276931.115

Wurtmann EJ, Wolin SL. 2009. RNA under attack: cellular handling of RNA damage. *Crit Rev Biochem Mol*
Biol **44**:34–49. doi:10.1080/10409230802594043

Yamashita A, Chang T-C, Yamashita Y, Zhu W, Zhong Z, Chen C-YA, Shyu A-B. 2005. Concerted action
of poly(A) nucleases and decapping enzyme in mammalian mRNA turnover. *Nat Struct Mol Biol*
12:1054–1063. doi:10.1038/nsmb1016

Yan WX, Chong S, Zhang H, Makarova KS, Koonin EV, Cheng DR, Scott DA. 2018. Cas13d Is a Compact
RNA-Targeting Type VI CRISPR Effector Positively Modulated by a WYL-Domain-Containing
Accessory Protein. *Mol Cell* **70**:327-339.e5. doi:10.1016/j.molcel.2018.02.028

Yan Y-B. 2014. Deadenylation: enzymes, regulation, and functional implications: Regulation of
deadenylation. *WIREs RNA* **5**:421–443. doi:10.1002/wrna.1221

Yang L, Duff MO, Graveley BR, Carmichael GG, Chen L-L. 2011. Genomewide characterization of non-
polyadenylated RNAs. *Genome Biol* **12**:R16. doi:10.1186/gb-2011-12-2-r16

Yang L, Froberg JE, Lee JT. 2014. Long noncoding RNAs: fresh perspectives into the RNA world. *Trends*
in Biochemical Sciences **39**:35–43. doi:10.1016/j.tibs.2013.10.002

Yang L-Z, Wang Y, Li S-Q, Yao R-W, Luan P-F, Wu H, Carmichael GG, Chen L-L. 2019. Dynamic Imaging
of RNA in Living Cells by CRISPR-Cas13 Systems. *Molecular Cell* **76**:981-997.e7.
doi:10.1016/j.molcel.2019.10.024

Yi R, Qin Y, Macara IG, Cullen BR. 2003. Exportin-5 mediates the nuclear export of pre-microRNAs and
short hairpin RNAs. *Genes Dev* **17**:3011–3016. doi:10.1101/gad.1158803

Yi W, Li J, Zhu X, Wang X, Fan L, Sun W, Liao L, Zhang J, Li X, Ye J, Chen F, Taipale J, Chan KM, Zhang
L, Yan J. 2020. CRISPR-assisted detection of RNA–protein interactions in living cells. *Nat Methods*.
doi:10.1038/s41592-020-0866-0

Yin Y, Lu JY, Zhang X, Shao W, Xu Y, Li P, Hong Y, Cui L, Shan G, Tian B, Zhang QC, Shen X. 2020. U1
snRNP regulates chromatin retention of noncoding RNAs. *Nature* **580**:147–150.
doi:10.1038/s41586-020-2105-3

Yu Y, Reed R. 2015. FUS functions in coupling transcription to splicing by mediating an interaction between
RNAP II and U1 snRNP. *Proc Natl Acad Sci USA* **112**:8608–8613. doi:10.1073/pnas.1506282112

Zhang C, Konermann S, Brideau NJ, Lotfy P, Wu X, Novick SJ, Strutzenberg T, Griffin PR, Hsu PD, Lyumkis
D. 2018. Structural Basis for the RNA-Guided Ribonuclease Activity of CRISPR-Cas13d. *Cell*
175:212-223.e17. doi:10.1016/j.cell.2018.09.001

Zhang Z, Sun W, Shi T, Lu P, Zhuang M, Liu J-L. 2020. Capturing RNA–protein interaction via CRUIS.
Nucleic Acids Research gkaa143. doi:10.1093/nar/gkaa143

Zhou B, Wang Y, Yan Y, Mariscal J, Di Vizio D, Freeman MR, Yang W. 2019. Low-Background Acyl-Biotinyl
Exchange Largely Eliminates the Coisolation of Non- *S* -Acylated Proteins and Enables Deep *S* -
Acylproteomic Analysis. *Anal Chem* **91**:9858–9866. doi:10.1021/acs.analchem.9b01520

Zhou Y, Wang G, Wang P, Li Z, Yue T, Wang J, Zou P. 2019. Expanding APEX2 Substrates for Proximity-
Dependent Labeling of Nucleic Acids and Proteins in Living Cells. *Angew Chem Int Ed* **58**:11763–
11767. doi:10.1002/anie.201905949

Zhou Z, Licklider LJ, Gygi SP, Reed R. 2002. Comprehensive proteomic analysis of the human spliceosome.
Nature **419**:182–185. doi:10.1038/nature01031

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Raw western blot data

Figure 1C

Figure 1D

Figure 1E

Figure 3B

Whole cell lysate   Streptavidin enriched

Figure 2C