Differentiation potential of fish fins –Effective utilization of the fins as food

wastes–

Yusuke Tsuruwaka[1,2,3*], Eriko Shimada[1,4,5]

[1]Cellevolt, Niigata, Japan
[2]Institute for Advanced Biosciences, Keio University, Yamagata, Japan
[3]Marine Bioresource Exploration Research Team, Marine Biodiversity Research Program, Institute of
Biogeosciences, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Kanagawa, Japan
[4]Division of Applied Biosciences, Graduate School of Agriculture, Kyoto University, Kyoto, Japan
[5]Department of Pharmacology, University of California, Davis, Davis, California, USA.
12

13 ***Corresponding author:**
14 Yusuke Tsuruwaka
15 E-mail address: tsuru@ttck.keio.ac.jp
16 E-mail address: ytsuruwaka@gmail.com
17

18 Eriko Shimada
19 E-mail address: eshimada@alumni.ucdavis.edu
20

Abstract

Fish cells are largely affected by their culture media. Fibroblast-like cells obtained from the fish fins can differentiate to various kinds of cells such as skeletal muscle-like, neurofilaments and adipocytes. Our results suggest that the fins which are usually discarded as food wastes may practically applied to the clean meat technology.

Introduction

Introduction

In vitro meat production technology, also known as the cell culturing technology of 'clean meat' in the

laboratory, has gained more attention recently because the technology may contribute to sustainable food

production system. Currently, we face on food crisis with population growth, such as rapidly increasing

demand for livestock products. Raising livestock becomes an environmental stressor since it requires large

amount of natural resources and brings about 14.5% of total anthropogenic greenhouse gas emissions (Charles

et al., 2019; Gerber et al., 2013; Grossi et al., 2019). Therefore, establishment of sustainable food resource

production system for next generation is effective for environmental and ecological protection (Lynch &

Pierrehumbert, 2019; Tuomisto, 2019). In addition, it is expected that food wastes are reduced and recycled in

an effective manner (Lemaire & Limbourg, 2019).

Fish has been consumed increasingly worldwide, resulting in the enormous impacts on the ecosystem such as

overfishing. In the present study, we focus on the developmental technology which creates 'fish clean meat'

from fish cells for the first time in the world, in order to overcome the challenges in Sustainable Development

Goal (SDG) 14: Conserve and sustainably use the oceans, seas and marine resources for sustainable

development (FAO, 2015). We especially focused on fish fins and scales because 1) they are often discarded in

the meal and 2) collecting the scales and fins partially does not take a life in fish. We believe that technology of

creating 'aquatic clean meat' from fish brings potential benefits including environmental friendliness, animal

welfare and sustainability.

Fish have a high level of ability in regeneration. They are able to regenerate various body parts such as fins

which are equivalently a hand and foot in humans, hearts, neurons and so on. In general, regeneration occurs by

dedifferentiation and/or self-renewal with basal stem cells (Jopling et al., 2010; Kikuchi et al., 2010; Sada et al.,

2016; Sehring & Weidinger, 2020). Individual fish enables to regenerate their partially lost fins within a few

weeks. What about the fins cut from the body, on the other hand? The cultured fin tissues cut from fish body

shows explants around the tissues. However, the fin tissues have never been shaped to a complete fin as it was

in culture flask. Then, what ability or characteristics have those cells which gathered for regeneration retained?

If the cells have de-differentiated, what is the differentiation potential do they have?

In general, ES cells and the inner cell mass of early embryos as well as amphibian undifferentiated embryo

cells are cultured without adding specific stimulating factors when they are differentiated and cultured in vitro,

that is, serum, growth factors, etc. It is known to lead neural differentiation without external signals (Kamiya et

al., 2011). Thus, basal state of undifferentiated cells had a fate to neural induction. Shimada *et al.* (2013)

reported that the fins were derived from somatic mesoderm which has developed from somatic stem cells.

These results suggests the possibility that the fin cells in regeneration process was dedifferentiated and

become undifferentiated cells, they may become basal neural somatic stem cells. If differentiated fish cells have

the ability to differentiate into various types of cells such as muscle, fat, and fibers and nerves, they will be able

to transform into cultured 'aquatic' clean meat.

Materials and method

Animal

Live thread-sail filefish *Stephanolepis cirrhifer* was captured near Jogashima Island in Yokosuka City,

Kanagawa Prefecture, Japan. Specimens were maintained in aerated artificial seawater and its salinity and

water temperature were maintained at 34.0 ppt and 20±2°C, respectively.

Preparation of the cell line

The cell line of the thread-sail filefish *S. cirrhifer* was prepared the cells as follows. We obtained a 5 mm^2 cut

of fin tissue from the thread-sail filefish and placed it in 70% ethanol. The tissue was then washed three times

with phosphate buffered saline (PBS), then six times with penicillin and streptomycin (MP Biomedicals, Santa

Ana, CA) on ice. The dorsal fin tissue was cut into 1 mm squares in 0.25% Trypsin (MP Biomedicals)-0.02%

EDTA (MP Biomedicals) solution and incubated at room temperature for 20 min. The tissue was centrifuged at

1100 rpm and washed twice with Leibovitz's L-15 culture medium (Life Technologies, Carlsbad, CA). The

tissue was immersed in L-15 media containing 10% fetal bovine serum (FBS) (Biowest, Nuaillé, France) and

1% Zell Shield (Minerva Biolabs, Berlin, Germany), then seeded in a 25 cm^2 Collagen I coated flask (Thermo

Fisher Scientific, Waltham, MA), and cultured in an incubator at 25°C. After 24 h, we confirmed the outgrowth

surrounding the tissue. A couple of days later, the cells around the tissue were removed with TrypLE Express

(Life Technologies), and seeded into a new flask. The culture media was replaced every 3 d and the cells were

subcultured at 4.0×10^5 cells/ml five times to establish the cell line, named. TSF Cell morphology was observed

using an inverted microscope CKX41 (Olympus, Tokyo, Japan) connected to a digital camera

ARTCAM-300MI-WOM (Artray, Tokyo, Japan).

Induction of Differentiation

Fin cells from established *S. cirrhifer* cell line were evaluated their differentiation under various culture

conditions. 4.0×10^5 cells were seeded in L-15 media containing 10% FBS in a 25 cm^2 Collagen I coated flask

or Non-coated flask (Thermo Fisher Scientific). 24 h later, the cells were washed with PBS and treated with

different media. For those seeded in Collagen I coated flask, AIM V Medium (Thermo Fisher Scientific) + 10%

FBS, AIM V Medium (Thermo Fisher Scientific) only, L-15 + 10% heat-inactivated SeaGrow (EastCoast

Biologics, North Berwick, ME), KBM Neural Stem Cell medium (Kohjin Bio Co. Ltd., Saitama, Japan) + 1X

Neural Induction Supplement (Thermo Fisher Scientific). For the cells seeded in Non-coated flask, L-15 only.

Cell differentiation was photographed every 60 seconds and the time-lapse movies were created with Axio

Vision ver. 4.8 software (Carl Zeiss).

.

Lipid staining

The fin cells were fixed in 4% paraformaldehyde (Wako Pure Chemical Industries, Ltd., Osaka, Japan) in

phosphate-buffered saline (PBS) at RT for 1 h. The fixed cells were subjected to BODIPY® staining with

Adipocyte Fluorescent Staining kit (Cosmo Bio Co. Ltd., Tokyo, Japan) following to manufacture's protocol.

For Oil Red O staining, the fixed cells were treated with 60% isopropanol for 5 min. After the isopropanol was

removed, the cells were incubated with Oil Red O solution at RT for 25 min. The treated cells were observed

with the microscope as stated above.

Immunofluorescence

Differentiated cells were fixed in 4% paraformaldehyde (Wako Pure Chemical Industries, Ltd.) in PBS at RT

for 12 h. The cells were incubated with Milli-Mark™ FluoroPan Neuronal Marker (Mouse IgG conjugated with

Alexa 488) (Merck Millipore, Burlington, MA) as the manufacturer's instruction. The fluorescent cells were

observed with a Zeiss Axio Observer. D1 microscope (Carl Zeiss, Oberkochen, Germany). An AxioCam HRc

camera (Carl Zeiss) was used to photograph the images, and the images were analysed using AxioVision ver.

4.8 software (Carl Zeiss).

Scanning electron microscopy (SEM)

The adherent cells on the collagen I coated slide were fixed with 0.4 ml of 2.5 % glutaraldehyde in culture

medium in a 2 ml culture chamber overnight at 4 °C. The cells were postfixed with 2.0 % osmium tetroxide

dissolved in PBS for 2 hours at 4 °C, and then were dehydrated in a graded series of ethanol, and freeze dried in

a freeze drier VFD-21S (Vacuum device Inc., Ibaraki, Japan). They were coated with osmium using a POC-3

osmium plasma coater (Meiwafosis Co. Ltd., Tokyo, Japan) and observed in a field emission scanning electron

microscope JSM-6700F (JEOL, Ltd., Tokyo, Japan) operated at 5 kV.

Gas chromatography

Fatty acids in the fin, the fin cells and the differentiated cells were determined by gas chromatography. The

extract sample (1 μl) was injected into a gas chromatograph 7890A (Agilent Technologies Inc., Santa Clara,

CA) equipped with an Omega wax 320 30 m ×0.32 mm.

Results

Does the regeneration of fish fin tissue occur?

YES, the fin of *S. cirrhifer* began to regenerate soon after being cut. Basic structure of the fin ray was formed

within two weeks and recovered to an original state four weeks later (Fig.1). On the other hand, a piece of the

fin tissue which was cut apart from the fish body showed populated swarmer cells in the culture media (Fig. 2).

The cell was about 20-50 μm in size. Subculturing was performed subsequently, and stable fibroblast-like cells

were obtained at 5^{th} passage (Fig. 3a, Movie 1). We named the fibroblast-like cells as deSc (dedifferentiated

Stephanolepis cirrhifer). To verify a chromosomal mutation had not occurred in the deSc cells, Q-banding stain

analysis was performed and compared to wildtype *S. cirrhifer* which had been reported by Murofushi *et al*.

(1980). As a result, the number of chromosomes was 33 ($2n=30+X_1X_2Y$) in 96% of deSc cells, which was

identical to the wildtype *S. cirrhifer* (Supplementary Fig. 1). 4% of deSc cells showed 66 chromosomes since

they might be in the middle cell division with replicated DNA.

153 **Figure 1.** Regeneration of the fin of *S. cirrhifer*. upper; whole fish, lower; enlarged view of the cut fin, arrow;

154 the cut part, scale bar; 5 cm.

Figure 2. Swarmer calls from fin tissue of *S. cirrhifer*. Scale bar; 100 μm

Figure 3. deSc cells differentiated various cell types. a) Normal fibroblast-like cells, b) Skeletal muscle-like

cells, c) neural-like cells, d) neurofilaments. Upper; bright field image, scale bar; 50 μm. Lower; SEM image,

magnification; X1000 magnified, scale bar; 10 μm.

Do the fin cells differentiate?

YES, they do. They differentiate to many types of cells indeed! To explore an optimal culture condition for

the deSc cells, we had examined several culture media, serum, and extracellular matrix (ECM) and discovered

very interesting characteristics (Table 1). The cell changed their morphology variously based on the

combinations of those culture media, serum and ECM (Table 1, Fig. 3b, Movie 2). Within the various results,

we focused on the neural-like cells which differentiated own morphology under culturing without serum (L-15

medium only) in a non-coated flask (Fig. 3c, Movie 3).

Table 1. Differentiation of deSc cells under various combinations of culture medium, serum, flask and

additive.

Cell Morphology	Culture Medium	Serum	Flask	Additive
Fibloblast-like	L-15	FBS	Collagen I-coated	
Skeletal muscle-like	AIM V	FBS	Collagen I-coated	
Nerural-like	L-15	—	Non-coated	
Neurofilament	KBM	FBS	Collagen I-coated	Neural induction suppliment
Adipocyte	L-15	SeaGrow	Collagen I-coated	
Spheloid	L-15	FBS	Spheroid	
Cocoon	L-15	Horse	Non-coated	

Is there differentiation potency in deSc cells?

YES, there is. Basal state of ES cells in the absence of neural differentiation inhibitors such as serum and

transcription factor adopted neural fate (Kamiya et al., 2011). The present deSc cells which were cultured in the

same condition have also differentiated to neural-like cells within 24h. The result led us to speculate that the

deSc cells have potential of neural differentiate likewise if the cells had acquired pluripotency through

dedifferentiation process. To demonstrate the hypothesis, we first attempted to induce neural differentiation

directly with KBM Neural Stem Cell medium (Kohjin Bio Co., Ltd.) and Neural Induction Supplement

(Thermo Fisher Scientific). As the result, neurofilaments were formed with 465 μm maximum length, and

45.71 μm/h average elongation speed (Fig. 3d, Movie 4, Table 1). Neural immunofluorescence suggested that

those fin cells were virtually differentiated to the neural cells (Fig. 4). These results demonstrated the fin cells

possess the characteristic of practicable direct-differentiation only with culture medium components. We have

succeeded to induce the neural differentiation, which is a basal state of stem cells, by both the presence/absence

of serum and the direct differentiation. We then came up with a question, "What will happen to the

differentiation if serum is different?"

Figure 4. Immunofluorescence of neural-like cells. Scale bar; 50 μm.

What happens to the cell differentiation when the serum is changed?

Very interesting phenomena were observed. In human iPS cells, culturing with different mammalian sera is reported to affect cell proliferations, differentiations, gene expressions and a stability of transcriptomes (Gstraunthaler, 2003; Shahdadfaret al., 2005). We next examined cell differential morphology under culturing with different sera. First, we assessed salmon serum SeaGrow which was the same taxonomical group as the

deSc cells. Granule-like particles were appeared intracellularly five hours after the stimulation with the fish

serum. Another three hours later, the cell morphology changed to the round, larger in size, which were

adipocyte-like cells with 0.5-2.0 μm white droplets (Fig. 5a, Movie 5). In order to examine the white droplets

for details, the cells were stained with Oil Red O and BODYPY, respectively, and also analyzed with a gas

chromatography. The results showed that those white droplets were fat droplets (Supplementary Fig. 2).

Therefore, the replacement to salmon serum in the culture media resulted in the differentiation to adipocyte in

deSc cells.

Figure 5. deSc cells differentiated various cell types. a) upper adipocyte staining with oil red o, scale bar; 50

μm arrow SEM x1000 scale bar; 10 μm, b) upper; spheloid in SEM x200 Scale bar; 100 μm. upper; bright field

We next examined several mammalian sera other than FBS in the cell culture. With a rabbit and a sheep

serum, the deSc cells did not survive, or did not present stable and uniform differentiation states (data not

shown). However, we have witnessed an extraordinarily interesting phenomenon when examining a horse

serum.

We, organisms consist of various tissues such as bones, cartilages, muscles, skin, which are established on

living-scaffolds. In this point, the process of turning differentiated cells into a tissue body is significantly

important. Recently, a three-dimensional spheroid has been reported as an important *in vitro* model in terms of

similar functions and organization to biological tissues (Achilli et al., 2012). The deSc cells formed a spheroid

with three-dimensional culture, which mammalian cells did likewise. (Fig. 5b, Movie 6). The size of spheroids

were between 20 and 300 μm. Spheloids were formed by intaking their surrounding cells. Interestingly,

stimulating the cells with the horse serum showed colonies which looked like suspended cell aggregates with

adherent "feet" (Fig. 5c, Movie 7). We had named those colonies, "Cocoon." The cocoons "walked" around in

the culture flask at 38.46 μm/h average speed and also they moved largely every 3.5 h with absorbing or fusing

surrounding other cocoons (Movie 8). Furthermore, single deSc cells which had not formed the cocoon or had

separated from it showed the same behavior as the cocoons. Interestingly, 'the independent cells' which had not

constituted the colonies showed neither a cell division nor cell proliferation. The colonies showed various sizes

between 20 and 1000 μm diameter and the largest ones were able to be visualized macroscopically. Moreover,

our observations confirmed that the colonies were stable in culture at least three weeks, as seen in spheroids

culture too (Data not shown).

To date, we are not able to specify what characteristics the cocoons acquired and what they differentiated to.

However, they showed surprisingly dissimilar behaviors by acquiring the 3D structure with the legs compared

to the normal cells. The cell differentiations mentioned above; skeletal muscle-like, adipocyte, spheloid and

cocoons, were reversible processes by turning back to the basic culture condition; L-15 medium with 10% FBS

in collagen-I coated flask, except the neural differentiation. Dedifferentiation which observed in the

regenerating fin at an individual level was continuously observed at cellular levels too in fish. Moreover, the

differentiation processes were surprisingly simple since its triggers were the culture media, sera and ECM.

Discussion

Dedifferentiation mechanism is relatively a common phenomenon observed as a regeneration process in

organisms. Our present study reports for the first time in a cellular level that the fish cells sampled during the

regeneration process have a stem cell-like characteristic. We demonstrated that deSc cell differentiations were

induced by "simple stimulus" such as medium, serum and ECM, without a specialized technique like a gene

transfer. While the present studies focused on the thread-sail filefish *S. cirrhifer*, the differentiate phenomena in

two-dimension have been observed in other several fish cells, too, more or less with differentiation potential

(Suppremetary Fig. 3). To date, sophisficated biological technology, gene transfer, enables to create tissues

from various kinds of cells including iPS and ES cells (Takebe et al., 2019). Moreover, some research groups

succeeded to produce other cells directly from fibroblast without going through those stem cells (Wada et al.,

2013). Our studies suggested that the potential for 'simple and direct differentiation' is also commonly present

in fish as well as in mammals.

We believe that finding such potential in edible fish is extremely significant because it is interesting to utilize

the fins which are usually discarded as food wastes, effectively to the next-generation food processing

technologies, including clean meat. Especially in fish meals, taste and texture of the preferment parts in favorite

kinds of fishes will be able to be reproduced anytime without being affected by the seasonal catches. Moreover,

aquatic clean meat has a merit of unnecessity of microplastic pollutions.

Postscript

We would like to discuss the potential of fish cell research for its future development. Swann et al. (2014)

reported that mouse-derived thymic epithelial cells generated a fish-like thymus under coexpression of *Foxn4*

and *Foxn1*. Kherdjemil et al. (2016) and Nakamura et al. (2016) found that fish fin rays and fingers in

mammals emerge from the same group of embryonic cells. Fish is known to have high homology to humans

(Aparicio et al., 2002; Howe et al., 2013). These studies may suggest as follows; it is not impossible that cells

can be utilized universally beyond species by resetting cell functions with the reprogramming technology; in

other words, various types of mammalian cells can be created from fish cells. Moreover, our study shows the

cocoon, the 3-D structured cells, which was interesting phenomenon. The cocoons use their legs like tentacles,

which look like they have acquired a will, however, it is difficult to explain what the behavior and the legs are

for at present. Re-creating organisms from cultured cells is hard to imagine, but interesting.

Organisms have acquired a variety of lifestyle from ocean to land by changing their fins to legs through

evolution (Flammang et al., 2016). Fish fin is adopted and induced phenotypic changes in response to

environmental changes (Standen et al., 2014). We demonstrated that deSc cells, the fins in cellular level,

showed facile differentiation. These results and reports lead us an important factor – "serum," that is cells are

possible to make themselves change with the blood constituents. This is enormously interesting in terms of

discussing evolutionary aspect in biodiversity because differentiation is significant phenomenon in evolution.

At present, a mainstream approach to induce differentiation in scientific research is a DNA modification by

gene transfer. However, it is hard to imagine those modifications had occurred naturally in ancient times. Then,

what was involved with the differentiation other than the environmental gene alteration? What if a blood was

an inducer of the differentiation? Since organisms often contact with blood in nature through predation etc.,

can't we negate the possibility that the blood has been involve and served in biodiversity?

Finally, we came up with another question through our studies. Just as there are individual differences among

living things, cells also have characteristics individually. However, every cell differentiates almost identically

upon differentiation process. Could there be some cells that are difficult to differentiate? This is very

mysterious point. In recent years, it has been reported that pluripotent stem cells are inherently present in living

organisms in mesenchymal cells such as fibroblasts and bone marrow stromal cells in mammals (Kuroda et al.,

2010). What if there is a leader cell which plays a role ordering a command to the other cells even among cells

that look the same? When the leader cell is stimulated somehow, is a command issued so that the rest of all

cells in the flask begin to change? To support our hypothesis, a behavioral coupling mechanism between

neighboring epidermal stem cells was reported, that is, differentiation of a single stem cell was followed by the

division of local neighboring mouse cells (Mesa et al., 2018). Examining biological cells of not only mammals

but also various species will bring us greater knowledge of cell biological function in the future.

Funding

This research received no specific grant from any funding agency in public, commercial or not-for-profit

sectors.

Acknowledgement

We would like to thank Mr. Tomohisa Ogawa for technical support.

References

Achilli TM, Meyer J, Morgan JR. (2012). Advances in the formation, use and understanding of multi-cellular

spheroids. *Expert Opin Biol Ther*. **12**:1347-60.

Aparicio S, Chapman J, Stupka E, Putnam N, Chia JM, Dehal P, Christoffels A, Rash S, Hoon S, Smit A,

Gelpke MD, Roach J, Oh T, Ho IY, Wong M, Detter C, Verhoef F, Predki P, Tay A, Lucas S, Richardson P,

Smith SF, Clark MS, Edwards YJ, Doggett N, Zharkikh A, Tavtigian SV, Pruss D, Barnstead M, Evans C,

Baden H, Powell J, Glusman G, Rowen L, Hood L, Tan YH, Elgar G, Hawkins T, Venkatesh B, Rokhsar D,

Brenner S. (2002). Whole-genome shotgun assembly and analysis of the genome of Fugu rubripes. *Science*

297:1301-10.

Charles A, Kalikoski D, Macnaughton A. (2019). Addressing the climate change and poverty nexus: a

coordinated approach in the context of the 2030 agenda and the Paris agreement. Rome. FAO

Flammang BE, Suvarnaraksha A, Markiewicz J, Soares D. (2016). Tetrapod-like pelvic girdle in a walking

cavefish. *Scientific Reports* **6**:23711.

Gerber PJ, Steinfeld H, Henderson B, Mottet A, Opio C, Dijkman J, Falcucci A, Tempio G. (2013). Tackling

climate change through livestock: a global assessment of emissions and mitigation opportunities. Rome: FAO.

Available from http://www.fao.org/3/a-i3437e.pdf

Grossi G, Goglio P, Vitali A, Williams AG. (2019). Livestock and climate change: impact of livestock on

climate and mitigation strategies. *Animal Frontiers* **9**:69–76.

Gstraunthaler G. (2003). Alternatives to the Use of Fetal Bovine Serum: Serum-free Cell Culture. *ALTEX*.

20:275-81.

Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, Collins JE, Humphray S, McLaren K,

Matthews L, McLaren S, Sealy I, Caccamo M, Churcher C, Scott C, Barrett JC, Koch R, Rauch GJ, White S,

Chow W, Kilian B, Quintais LT, Guerra-Assunção JA, Zhou Y, Gu Y, Yen J, Vogel JH, Eyre T, Redmond S,

Banerjee R, Chi J, Fu B, Langley E, Maguire SF, Laird GK, Lloyd D, Kenyon E, Donaldson S, Sehra H,

Almeida-King J, Loveland J, Trevanion S, Jones M, Quail M, Willey D, Hunt A, Burton J, Sims S, McLay K,

Plumb B, Davis J, Clee C, Oliver K, Clark R, Riddle C, Elliot D, Threadgold G, Harden G, Ware D, Begum S,

Mortimore B, Kerry G, Heath P, Phillimore B, Tracey A, Corby N, Dunn M, Johnson C, Wood J, Clark S, Pelan

S, Griffiths G, Smith M, Glithero R, Howden P, Barker N, Lloyd C, Stevens C, Harley J, Holt K, Panagiotidis G,

Lovell J, Beasley H, Henderson C, Gordon D, Auger K, Wright D, Collins J, Raisen C, Dyer L, Leung K,

Robertson L, Ambridge K, Leongamornlert D, McGuire S, Gilderthorp R, Griffiths C, Manthravadi D, Nichol

S, Barker G, Whitehead S, Kay M, Brown J, Murnane C, Gray E, Humphries M, Sycamore N, Barker D,

Saunders D, Wallis J, Babbage A, Hammond S, Mashreghi-Mohammadi M, Barr L, Martin S, Wray P,

Ellington A, Matthews N, Ellwood M, Woodmansey R, Clark G, Cooper J, Tromans A, Grafham D, Skuce C,

Pandian R, Andrews R, Harrison E, Kimberley A, Garnett J, Fosker N, Hall R, Garner P, Kelly D, Bird C,

Palmer S, Gehring I, Berger A, Dooley CM, Ersan-Ürün Z, Eser C, Geiger H, Geisler M, Karotki L, Kirn A,

Konantz J, Konantz M, Oberländer M, Rudolph-Geiger S, Teucke M, Lanz C, Raddatz G, Osoegawa K, Zhu B,

Rapp A, Widaa S, Langford C, Yang F, Schuster SC, Carter NP, Harrow J, Ning Z, Herrero J, Searle SM,

Enright A, Geisler R, Plasterk RH, Lee C, Westerfield M, de Jong PJ, Zon LI, Postlethwait JH,

Nüsslein-Volhard C, Hubbard TJ, Roest Crollius H, Rogers J, Stemple DL. (2013). The zebrafish reference

genome sequence and its relationship to the human genome. Nature 496:498–503.

Jopling C, Sleep E, Raya M, Martí M, Raya A, Izpisúa Belmonte JC. (2010). Zebrafish heart regeneration

occurs by cardiomyocyte dedifferentiation and proliferation. *Nature* **464**:606-9.

Kamiya D, Banno S, Sasai N, Ohgushi M, Inomata H, Watanabe K, Kawada M, Yakura R, Kiyonari H, Nakao

K, Jakt LM, Nishikawa S, Sasai Y. (2011). Intrinsic transition of embryonic stem-cell differentiation into neural

progenitors. *Nature* **470**:503–509.

Kherdjemil Y, Lalonde RL, Sheth R, Dumouchel A, de Martino G, Pineault KM, Wellik DM, Stadler HS,

Akimenko MA, Kmita M. (2016). Evolution of Hoxa11 regulation in vertebrates is linked to the pentadactyl

state. *Nature* **539**:89–92.

Kikuchi K, Holdway JE, Werdich AA, Anderson RM, Fang Y, Egnaczyk GF, Evans T, Macrae CA, Stainier DY,

Poss KD. (2010). Primary contribution to zebrafish heart regeneration by gata4(+) cardiomyocytes. *Nature*

464:601-5.

Kuroda Y, Kitada M, Wakao S, Nishikawa K, Tanimura Y, Makinoshima H, Goda M, Akashi H, Inutsuka A,

Niwa A, Shigemoto T, Nabeshima Y, Nakahata T, Nabeshima Y, Fujiyoshi Y, Dezawa M. (2010). Unique

multipotent cells in adult human mesenchymal cell populations.*PNAS* **107**:8639-43.

Lemaire A, Limbourg S. (2019). How can food loss and waste management achieve sustainable development

goals? *Journal of Cleaner Production* **234**:1221–1234.

Lynch J, Pierrehumbert R. (2019). Climate impacts of cultured meat and beef cattle. *Front Sustain Food Syst*.

3:5.

Matsumoto T, Kano K, Kondo D, Fukuda N, Iribe Y, Tanaka N, Matsubara Y, Sakuma T, Satomi A, Otaki M,

Ryu J, Mugishima H. (2008). Mature adipocyte‐derived dedifferentiated fat cells exhibit multilineage

potential. *J. Cell Physiol*. **215**:210–222.

Mesa KR, Kawaguchi K, Cockburn K, Gonzalez D, Boucher J, Xin T, Klein AM, Greco V. (2018).

Homeostatic Epidermal Stem Cell Self-Renewal Is Driven by Local Differentiation. *Cell Stem Cell*

23:677-686.e4.

Murofushi M, Oikawa S, Nishikawa S, Yosida TH. (1980). Cytogenetical studies on fishes. III. Multiple sex

chromosome mechanism in the filefish, *Stephanolepis cirrhifer*. *The Japanese Journal of Genetics* **55**: 127-131.

Nakamura T, Gehrke AR, Lemberg J, Szymaszek J, Shubin NH. (2016). Digits and fin rays share common

developmental histories. *Nature* **537**:225-228.

Sada A, Jacob F, Leung E, Wang S, White BS, Shalloway D, Tumbar T. (2016). Defining the cellular lineage

hierarchy in the interfollicular epidermis of adult skin. *Nat Cell Biol*. **18**:619-31.

Sehring IM, Weidinger G. (2020). Recent advancements in understanding fin regeneration in zebrafish. Wiley

Interdiscip *Rev Dev Biol*. **9**:e367.

Shahdadfar A, Frønsdal K, Haug T, Reinholt FP, Brinchmann JE. (2005). In vitro expansion of human

mesenchymal stem cells: choice of serum is a determinant of cell proliferation, differentiation, gene expression,

and transcriptome stability. *Stem Cells* **23**:1357-66.

Shibata E, Ando K, Murase E, Kawakami A. Heterogeneous fates and dynamic rearrangement of regenerative

epidermis-derived cells during zebrafish fin regeneration. (2018). *Development* **145**: dev162016.

Shimada A, Kawanishi T, Kaneko T, Yoshihara H, Yano T, Inohaya K, Kinoshita M, Kamei Y, Tamura K,

Takeda H. (2013). Trunk exoskeleton in teleosts is mesodermal in origin. *Nat Commun*. **4**:1639.

Standen EM, Du TY, Larsson HC. (2014). Developmental plasticity and the origin of tetrapods. *Nature* **513**:

54-8.

Sugii S, Kida Y, Kawamura T, Suzuki J, Vassena R, Yin YQ, Lutz MK, Berggren WT, Izpisúa Belmonte JC,

Evans RM. (2010). Human and mouse adipose-derived cells support feeder-independent induction of

pluripotent stem cells. *PNAS* **107**: 3558-3563.

Swann JB, Weyn A, Nagakubo D, Bleul CC, Toyoda A, Happe C, Netuschil N, Hess I, Haas-Assenbaum A,

Taniguchi Y, Schorpp M, Boehm T. (2014). Conversion of the thymus into a bipotent lymphoid organ by

replacement of FOXN1 with its paralog, FOXN4. *Cell Reports* **8**:1184-97.

Tuomisto HL. (2019). The eco-friendly burger: Could cultured meat improve the environmental sustainability

of meat products? *EMBO Rep*. **20**: e47395.

Wada R, Muraoka N, Inagawa K, Yamakawa H, Miyamoto K, Sadahiro T, Umei T, Kaneda R, Suzuki T,

Kamiya K, Tohyama S, Yuasa S, Kokaji K, Aeba R, Yozu R, Yamagishi H, Kitamura T, Fukuda K, Ieda M.

(2013). Induction of human cardiomyocyte-like cells from fibroblasts by defined factors. *PNAS* **110**: 12667-72.

Zuk PA, Zhu M, Ashjian P, De Ugarte DA, Huang JI, Mizuno H, Alfonso ZC, Fraser JK, Benhaim P, Hedrick

MH. (2002). Human adipose tissue is a source of multipotent stem cells. *Molecular biology of the cell*, **13**:

4279–4295.

Supplemental Movie

Movie 1. Proliferation of deSc cells.

Movie 2. deSc normal cells differentiate to skeletal muscle-like cells.

Movie 3. deSc normal cells differentiate to neural-like cells.

Movie 4. deSc normal cells differentiate to neurofilaments.

Movie 5. deSc normal cells differentiate to adipocyte.

Movie 6. deSc normal cells differentiate to spheroids.

Movie 7. deSc normal cells differentiate to cocoons.

Movie 8. deSc normal cells differentiate to cocoons.

Supplemental Figure

Supplementary Figure 1. Chromosome of deSc cells, *Stephanolepis cirrhifer*, 2n=30+X_1X_2Y(33). Scale bar; 5

μm.

Supplementary Figure 2. Adipocyte staining with a) BODIPY and b) Oil-red O. Magnification x200, c) gas

chromatogram of normal deSc cells (upper) and adypocyte (lower); scale bar 50

Supplementary Figure 3. Differentiation of scorpion fish, *Sebastiscus marmoratu*s. a) normal, b) neural-like

cells under bright -field. Scale bar; 50 μm.