Corticospinal drive does not contribute to increased torque production in the presence of

residual force enhancement

Jasmin Frischholz[1], Brent J. Raiteri[1], Daniel Hahn[1,2]

Author contributions: JF, BJR and DH conceived and designed the research; JF performed

experiments; JF, BJR and DH analysed and interpreted the data; JF drafted the manuscript; JF

and DH prepared the figures; JF, BJR and DH edited and revised the manuscript.

[1] Human Movement Science, Faculty of Sport Science, Ruhr University Bochum, Germany

[2] School of Human Movement and Nutrition Sciences, University of Queensland, Brisbane,

Australia

Running head: Corticospinal drive and residual force enhancement

Corresponding author address:

Daniel Hahn

Human Movement Science – Faculty of Sport Science

Gesundheitscampus Nord 10

44801 Bochum – Germany

Tel.: +49 234 3227905

E-Mail: daniel.hahn@rub.de

Abstract

Following active muscle stretch, a muscle's force capacity is increased, which is known as

residual force enhancement (rFE). As earlier studies found modulations of cortical excitability

in the presence of rFE, this study aimed to test whether corticospinal drive contributes to rFE.

Fourteen participants performed submaximal plantar flexion stretch-hold and fixed-end

contractions at 30% of their maximal voluntary soleus muscle activity in a dynamometer.

During the steady state of the contractions, participants either received subthreshold or

suprathreshold transcranial magnetic stimulation (TMS) of their motor cortex while triceps

surae muscle responses to stimulation were obtained by electromyography (EMG) and net

plantar flexion torque was recorded. B-mode ultrasound imaging was used to confirm muscle

stretch during stretch-hold contractions in a subset of participants. Following stretch of the

plantar flexors, an average rFE of 7% and 11% was observed for contractions with

subthreshold and suprathreshold TMS, respectively. 41-46 milliseconds following

subthreshold TMS, triceps surae muscle activity was suppressed by 19-25%, but no difference

in suppression was found between contraction conditions. Similarly, the reduction in plantar

flexion torque following subthreshold TMS was not different between contraction conditions.

Motor evoked potentials, silent periods and superimposed twitches following suprathreshold

stimulations were also not different between contraction conditions. As stimulations of the

motor cortex by TMS did not result in any differences between stretch-hold and fixed-end

contractions, we conclude that corticospinal drive does not contribute to the increased torque

production in the presence of rFE following active muscle stretch.

New & Noteworthy

This study tested whether corticospinal drive contributes to the increased torque capacity in

the presence of rFE. Through subthreshold and suprathreshold TMS of the motor cortex,

triceps surae muscle activity was respectively supressed or increased in the presence of rFE

and during a reference contraction without rFE. As similar responses were observed between

contraction contractions, we conclude that corticospinal drive likely does not contribute to the

increased torque capacity in the presence of rFE.

Keywords: eccentric contraction, active muscle stretch, neural control, cortical excitability,

torque enhancement

Introduction

It is well known that stretch of an active muscle results in increased force production during the isometric steady state following stretch compared with the steady-state force produced at the same muscle length and activation level during a fixed-end contraction. This is referred to as residual force enhancement (rFE), which was initially investigated *in situ* (Abbott and Aubert, 1952) and in isolated muscle fibres (Edman et al., 1982). Edman et al. (1982) suggested that rFE after active muscle stretch results from non-uniformities in sarcomere lengths. Since then, a variety of studies have investigated the development of rFE and suggested additional potential underlying mechanisms. These suggestions include stretch-induced increases in the number of attached cross-bridges, an increase in the average cross-bridge force and/or the attachment of the second myosin head (for topical reviews, see Edman, 2012; Herzog, 2014; Rassier, 2012). Additionally, titin could potentially explain rFE by increasing its stiffness in the presence of calcium and by potentially interacting with actin (Freundt and Linke, 2019; Herzog, 2018; Nishikawa, 2016).

Complementary to *in vitro* research, several *in vivo* studies have investigated rFE during electrically-stimulated and voluntary contractions (for reviews, see Chapman et al., 2018; Seiberl et al., 2015). rFE during voluntary contractions has been observed for small muscles of the thumb (Lee and Herzog, 2002), for large muscles of the lower limb (Pinniger and Cresswell, 2007; Seiberl et al., 2010) and during multi-joint multi-muscle contractions of the lower limbs (Hahn et al., 2010). The potential relevance of rFE during human movement was demonstrated in studies that investigated rFE during submaximal voluntary contractions and at joint angle configurations that mimic those of human locomotion (Paternoster et al., 2016; Seiberl et al., 2013).

Besides rFE, several *in vivo* studies have documented an activation reduction (AR) during the steady state after active muscle stretch. In contrast to rFE, which is observed when muscle

activity is matched between stretch-hold (STR) and fixed-end reference (REF) contractions,

AR occurs when force or torque is matched between contraction conditions. AR refers to a

reduced muscle activity level needed to maintain a given force/torque following active muscle

stretch compared with fixed-end contractions at the same final muscle length. Several studies

have reported AR and concluded that neuromuscular efficiency after active muscle stretch is

increased, which might in part be explained by enhanced passive forces due to increased titin

stiffness (Altenburg et al., 2008; Jones et al., 2016, 2016; Mazara et al., 2018; Oskouei and

Herzog, 2005; Paquin and Power, 2018; Seiberl et al., 2012). However, Paquin and Power

(2018) also found a rightward shift in the EMG-torque relationship following active stretch

compared with fixed-end reference conditions, which indicates that rFE might alter the

neuromuscular activation strategy.

Despite the presence of rFE after active muscle stretch during submaximal and maximal

voluntary contractions, stretch contractions are often associated with neural inhibition

(Duchateau and Enoka, 2016). However, whether neural inhibition occurs during active

stretch under voluntary activation remains controversial (Hahn, 2018). The potentially unique

control of stretch contractions and the observation of AR following active stretch has

motivated a number of studies to investigate the neural control and/or neural modulations that

occur during the isometric steady state following active muscle stretch (Altenburg et al., 2008;

Contento et al., 2019; Hahn et al., 2012; Sypkes et al., 2018). For example, Altenburg et al.

(2008) examined single motor unit behaviour of the vastus lateralis muscle during AR after

active muscle stretch. These authors found similar discharge rates of VL motor units between

STR and REF conditions, which led them to conclude that a derecruitment of motor units

might have occurred during AR (Altenburg et al. 2008). Two further studies have used motor

evoked potentials (MEPs) and cervicomedullary motor evoked potentials (CMEPs) to

investigate modulations to the excitability of cortical and spinal sites in the presence of rFE

(Hahn et al., 2012; Sypkes et al., 2018). During submaximal dorsiflexion contractions, Sypkes

et al. (2018) found unchanged TA MEPs, but reduced TA CMEPs, following active muscle

stretch compared with fixed-end reference contractions. This is in line with the idea of

reduced motor unit recruitment (Altenburg et al. 2008) and might be caused by tension-related

afferent feedback as assessed by tendon-evoked inhibitory reflexes (Contento et al. 2019).

Contrary to these findings under submaximal contraction intensities, Hahn et al. (2012) found

larger MEPs, but unchanged CMEPs, following stretch of the plantar flexors during maximal

voluntary contractions. Both the results of Hahn et al. (2012) and Sypkes et al. (2018) could

be interpreted as an increase in cortical excitability in the presence of rFE. Further, enlarged

V-waves during rFE in the Hahn and colleagues (2012) study suggest a greater motoneuronal

output and/or an increase in stretch reflex excitability after active stretch (Hahn et al. 2012).

The above findings suggest that the underlying mechanisms of rFE during voluntary

contractions are not purely mechanical in nature. However, it remains unclear whether the

observed increase in cortical excitability contributes to the increased torque production in the

presence of rFE. Therefore, this study was designed to investigate whether corticospinal drive

contributes to increased torque production following active stretch of the plantar flexor

muscles during submaximal voluntary contractions. This was examined by inhibiting and

facilitating the motor cortex via transcranial magnetic stimulation (TMS) in the presence

(stretch-hold condition) and absence (fixed-end reference condition) of rFE.

Based on previous research (Hahn et al., 2012; Sypkes et al., 2018), we predicted that

inhibiting motor cortical neurons by subthreshold TMS (Davey et al., 1994; Petersen et al.,

2001; Zuur et al., 2010) would lead to a larger suppression in muscle activity (EMG) and

plantar flexion torque in the presence of rFE compared with fixed-end reference contractions

with a matched background muscle activity. Conversely, facilitation of cortical interneurons

and pyramidal neurons by suprathreshold TMS (Rossini et al., 2015; Rothwell, 1997) was

predicted to result in greater superimposed twitches in the presence of rFE compared with the

fixed-end reference contractions with a matched background muscle activity. If both

predictions hold, this would indicate that corticospinal drive contributes to the increased

torque production in the presence of rFE.

Materials and Methods

Participants

Fourteen recreationally-active participants (six women, 26.7 ± 5.3 yrs., 1.77 ± 0.11 m, and

74.0 ± 16.8 kg) voluntarily participated in this study after providing free written informed

consent. Participants were free of any neuromuscular disorders and injuries to their right

lower limb. The experimental protocol was approved by the local Ethics Committee of the

Faculty of Sport Science at Ruhr University Bochum, Germany.

Experimental set-up

During the experiment, the participants laid prone on a dynamometer bench with their upper

body supported by their forearms and their right foot tightly strapped onto the footplate

attachment of a dynamometer (IsoMed 2000, D&R Ferstl GmbH, GER). The axes of rotation

of the ankle joint and the dynamometer were aligned prior to testing and the foot was firmly

secured to the footplate using three straps (forefoot, ankle, heel) to minimize heel lift during

the plantar flexion contractions. Additionally, the participants' hips were secured to the

dynamometer bench with a belt. A computer monitor positioned in front of the participants

was used to provide visual feedback of their soleus muscle activity (moving 0.25-s root-mean-

square amplitude calculation) during the plantar flexion contractions. Net ankle joint torques

and ankle joint angles of the right leg were sampled at 1 kHz using a 16-bit Power-3 1401 and

Spike2 data collection system (v.8.01, Cambridge Electronics Design, Cambridge, UK)

Surface electromyography

Muscle activity and responses to TMS from soleus (SOL), medial gastrocnemius (MG) and

lateral gastrocnemius (LG) muscles were recorded via bipolar surface electromyography

(EMG). After skin preparation, self-adhesive surface electrodes (Ag/ AgCl, Kendall, ECG

Electrodes, 8 mm recording diameter) were positioned with an inter-electrode distance of 20

mm over the triceps surae muscle bellies following the recommendations of SENIAM

(Hermens et al., 1999). A reference electrode was placed over the right fibular head. Cables

were fixed to the skin with tape to prevent motion artefacts. EMG signals were amplified

1000 times by a multichannel analogue amplifier (AnEMG12, Bioelectronica, Turin, IT) and

band-pass filtered between 10 Hz and 4.4 kHz, prior to being sampled at 5 kHz. EMG and

torque/angle data were synchronised and sampled using the same analogue-to-digital

converter and software described above.

Transcranial Magnetic Stimulation (TMS)

TMS (MagPro Compact, MagVenture, Farum, Denmark) was used to either inhibit or

facilitate the motor cortical area of the right plantar flexor muscles in the left hemisphere,

slightly left of the vertex, via A double-cone coil (D-B80 Butterfly Coil, MagVenture, Farum,

Denmark). The coil had monophasic current running through its centre in an anterior posterior

direction. The vertex was marked on the scalp and defined as halfway between the left and

right processus zygomaticus ossis temporalis and halfway between the os nasale and the

external occipital protuberance. The vertex location helped to find the optimal location for

TMS of the motor cortical area innervating the right plantar flexor muscles (TMS hotspot),

which is generally defined as the position in which a single pulse stimulation evokes the

largest peak-to-peak MEP amplitude in the target muscle (Siebner and Ziemann, 2007). In

order to find the TMS hotspot, several stimuli were delivered while the TMS coil was slightly

left, in front or behind the vertex, while participants performed fixed-end plantar flexion (PF)

contractions at 30% of their maximal voluntary activity (MVA). Once the TMS hotspot was

located, the wings of the coil were marked on the scalp with a semi-permanent marker.

Subthreshold and suprathreshold intensities were then determined by either decreasing or

increasing the stimulator output. To achieve a suppression of soleus muscle activity,

stimulator output was reduced in small increments until the threshold for facilitation was

reached. This threshold was defined as the point when stimulation during 30% MVA resulted

in visible MEP responses in only 50% of ten consecutive trials (Petersen et al., 2001). For

facilitation, stimulator output was increased until MEPs were clearly visible in comparison to

the immediately preceding background EMG in at least five consecutive trials.

Ultrasound imaging

Muscle fascicle behaviour of the MG from three participants was examined during

familiarisation sessions using B-mode ultrasound imaging (LS128 CEXT-1Z, Telemed,

Vilnius, Lithuania) to ensure muscle fascicle stretch occurred during ankle rotation in the

stretch-hold condition. Ultrasound images were recorded using a flat, linear, 128-element

transducer (LV7.5/60/128Z-2, 6 MHz, 60 x 50 mm (width x depth)) operating at ~60 fps in

EchoWave II software (Telemed, Vilnius, Lithuania). The transducer was placed on the skin

over the MG mid-muscle belly in the longitudinal plane and rotated to obtain a clear image

with continuous aponeuroses and muscle fascicles. The position of the probe was maintained

and marked on the skin in order to ensure consistent placement during the stretch-hold

contractions.

Contraction conditions

The experiment involved participants performing submaximal plantar flexion contractions at

30% MVA. The background activity level was controlled throughout the contractions by

having participants visually match their SOL EMG amplitude (moving 0.25-s root-mean-

square (RMS) amplitude calculation) to a horizontal cursor on a screen. The conditions

involved fixed-end reference contractions (REF) at 20° dorsiflexion (DF) and active stretch-

hold contractions (STR) from 0°-20° DF (0° refers to the sole of the foot perpendicular to the

shank).

All contractions started with a 2-s linear ramp from 0-30% MVA. Following the ramp, the

isometric steady state during the fixed-end reference contraction was maintained for 13-s.

During the stretch-hold contractions, the ramps were followed by a 1-s isometric steady state

at the initial ankle joint angle (0° DF), before the plantar flexor muscles were actively

stretched to an ankle joint angle of 20° DF at an angular velocity of $30°s^{-1}$. Following active

stretch, participants maintained another isometric steady state for ~11-s, resulting in an

overall contraction duration of 15-s (Fig. 1).

Experimental protocol

Participants attended two sessions on two different days. In the first session, participants were

familiarised with the test protocol and trained to perform maximal voluntary fixed-end

contractions (MVC) and the submaximal contractions described above. Additionally,

participants were familiarised with TMS.

The second session consisted of the test protocol (Fig. 2). After a short warm-up (eight

submaximal PF contractions with increasing torque), at least two MVCs were performed to

determine 100% MVA. Participants were instructed to push their forefoot into the footplate as

hard as possible and to maintain the contractions, so that there was a clearly visible torque

plateau. MVC torque was calculated as the difference between peak torque during the

contraction and the mean baseline over 500 ms prior to the beginning of contraction. In order

to ensure that the MVCs were repeatable, peak-to-peak torques were required to be within a

5% range. 100% MVA was determined as the smoothed (moving 0.25-s RMS amplitude

calculation) SOL EMG amplitude at peak MVC torque from the best MVC. Three minutes of

rest was provided between MVCs to minimise fatigue.

Following the MVCs, the TMS hotspot and the subthreshold and suprathreshold TMS

intensities were determined during sustained fixed-end contractions at 30% MVA. Once

hotspot and stimulation intensities were determined, 100 subthreshold stimulations were

delivered for each contraction condition to study motor cortex inhibition. For this purpose,

contractions from each condition were separated into ten sets with ten TMS (1 Hz) during the

isometric steady state phase of the contractions at 20° DF starting 5-s after contraction onset

(Fig. 1). Contractions were randomised and a minimum of three minutes rest was provided

between each set. Facilitation of the motor cortex was investigated by providing 20

suprathreshold stimulations per contraction condition. Suprathreshold stimulations were

delivered at the same time points as the subthreshold stimulations, but participants performed

only two contractions per condition. The contractions with suprathreshold TMS were always

performed after all sets of the subthreshold stimulations were finished. During all

contractions, trials were excluded and repeated if participants could not maintain their soleus

EMG activity within 25-35% MVA.

Data analysis

Residual Force Enhancement (rFE)

rFE for the subthreshold and suprathreshold conditions was determined by calculating the

difference between the isometric steady-state torque during the STR and REF conditions. Net

PF torque was averaged across contractions for each contraction condition and mean rFE was

calculated during the isometric steady state at 20° DF, 500 ms after each TMS, for a time

window of 490 ms. Stimulations delivered before the isometric steady state of SOL muscle

activity were excluded.

TMS inhibition

Suppression of muscle activity (SOL, MG and LG) after subthreshold TMS was determined

via averaged rectified raw EMG signals from each muscle and contraction condition.

Waveform averages were calculated over a time window of 100 ms after stimulation and

again, stimulations delivered before the isometric steady state of SOL muscle activity were

excluded. The latency and duration of EMG amplitude suppression were determined using the

evaluation methods by Petersen et al. (2001) and Zuur et al. (2010). The onset of EMG

amplitude suppression was marked when the EMG amplitude first dropped under the

background EMG amplitude (mean value calculated 15–25 ms after stimulation) for at least 4-

ms. The offset of EMG amplitude suppression was calculated when the EMG amplitude rose

above the background EMG amplitude for 1 ms (Zuur et al., 2010) (Fig. 3). The mean EMG

amplitude between the onset and offset of suppression was calculated and compared with the

background EMG amplitude right before suppression.

Torque production following TMS-induced EMG amplitude suppression was analysed via

moving correlations (50 ms time intervals) between the torque of the REF and STR

conditions. A reduction in torque due to TMS-induced EMG amplitude suppression was

identified when torque data of REF and STR contractions was correlated ($r \geq 0.7$). This data

analysis was based on the random fluctuations in torque steadiness that should result in non-

correlated torque signals. However, when torque was affected by the TMS-induced EMG

amplitude suppression in a similar manner, positive correlations were expected. Once the time

window of torque reduction was identified, the torque offset between the REF and STR

conditions was removed by subtracting the mean torque difference between REF and STR

over the first 10 ms after the onset of torque reduction. Finally, the mean torque during the

period of TMS-induced torque reduction (i.e. during the times where $r \geq 0.7$) was calculated

for REF and STR and compared between contraction conditions.

TMS facilitation

MEPs of SOL, MG and LG following suprathreshold TMS were calculated as peak-to-peak

amplitudes from the raw EMG signals and averaged across contractions for each muscle and

contraction condition (Lewis et al., 2014) (Fig. 4). Additionally, the silent period (SP)

duration was analysed for both contraction conditions as the time from stimulation to the end

of the SP. The end of the SP was defined as the time when the EMG signal crossed the

threefold standard deviation (SD) of the SP raw EMG following the MEP for the first time

(Fig. 4). SD during the SP was determined as the smallest SD of the raw EMG over a moving

30ms window. Superimposed torque twitches following suprathreshold TMS were calculated

separately as peak-to-peak amplitudes and averaged for each contraction condition.

Statistical analysis

All data were tested for normality using the Shapiro-Wilk test. Paired t-tests were used to test

for significant differences (alpha level of 0.05) in triceps surae muscle activity, net plantar

flexion torque, EMG amplitude suppression, torque reduction, MEP, SP and superimposed

twitch amplitude between REF and STR conditions. Statistical analysis was performed using

SPSS software (v22, IBM, New York, USA). Values are presented as mean ± SD in the text.

Results

Five and four participants were excluded from further analysis as they did not show rFE

following the subthreshold (n=9) and suprathreshold (n=10) stimulations, respectively.

Contraction conditions

During both contraction conditions, participants managed to maintain a constant level of

muscle activity (~30 % MVA). During the isometric steady state, EMG of SOL, MG and LG

did not differ between STR and REF conditions during contractions with subthreshold TMS

(SOL: $P = 0.308$, MG: $P = 0.091$, LG: $P = 0.144$; Fig. 5) or suprathreshold TMS (SOL: $P =$

0.528, MG: $P = 0.490$, LG: $P = 0.349$). Ultrasound imaging visually confirmed that the

muscle-tendon unit stretch during the 20° dynamometer rotation of the stretch-hold

contraction resulted in fascicle stretch of the MG muscle at 30% MVA (n=3).

Contractions with subthreshold TMS

The isometric steady-state net plantar flexion torque in the STR condition exceeded the time-

matched steady-state torque in the REF condition (STR: 105.9 ± 34.5 Nm; REF: 99.0 ± 33.4

Nm, $P = 0.002$), which resulted in mean rFE of 7.3 ± 4.2% (Fig. 6A).

Subthreshold stimulation of the motor cortex led to significant ($P < 0.001$) EMG amplitude

suppression for SOL, MG and LG without any difference between STR and REF conditions

(SOL: $P = 0.483$; MG: $P = 0.378$; LG: $P = 0.399$). The mean EMG amplitude suppression

ranged between 19-25% of the background EMG amplitude, it occurred 41-46 ms after

stimulation, and it lasted 10-15 ms in total (Table 1). The reduction in net plantar flexor

torque following EMG amplitude suppression did not differ ($P = 0.228$) between STR and

REF conditions.

Contractions with suprathreshold TMS

The isometric steady-state net plantar flexion torque in the STR condition exceeded the time-

matched steady-state torque in the REF condition (STR: 89.5 ± 26.0 Nm; REF: 82.2 ± 27.2

Nm, $P = 0.002$), which resulted in mean rFE of 10.8 ± 10.0 % (Fig. 6B). For all muscles

analysed (SOL, MG, LG), MEP and SP after suprathreshold TMS did not differ between STR

and REF conditions (MEP: SOL: $P = 0.304$, MG: $P = 0.594$, LG: $P = 0.453$; SP: SOL: $P =$

0.320, MG: $P = 0.283$, LG: $P = 0.265$; Table 2). TMS-evoked superimposed twitches

following suprathreshold stimulations were 9.1 ± 3.6 Nm (STR) and 8.6 ± 3.5 Nm (REF), and

did not differ between STR and REF conditions ($P = 0.259$).

Discussion

The aim of this study was to determine whether corticospinal drive contributes to the

increased torque production in the presence of rFE following active stretch of the plantar

flexor muscles. To achieve this, reductions in EMG activity and torque production following

inhibition of motor cortical neurons, and increases in MEP amplitudes, SP durations and

superimposed twitch torque amplitudes following facilitation of motor cortical neurons were

compared between submaximal stretch-hold and fixed-end contractions.

Participants managed to keep muscle activity constant (30% MVA) throughout the fixed-end

reference contractions and during the isometric steady state of the stretch-hold contractions.

At this matched background EMG, mean rFE magnitudes of 7% and 11% were observed

during the steady state in STR compared with REF, which is in line with former studies

(Oskouei and Herzog, 2005; Paternoster et al., 2016; Pinniger and Cresswell, 2007; Seiberl et

al., 2013). Inhibition of motor cortical neurons by subthreshold TMS caused reductions in

EMG activity and net plantar flexion torque, however the reductions did not differ

significantly between contraction conditions. Similarly, MEP amplitudes, SP durations and

superimposed twitch torque amplitudes evoked by suprathreshold TMS were not significantly

different between STR and REF conditions.

Following subthreshold TMS, we found a suppression of triceps surae muscle activity by 19-

24% relative to the background activity. This is similar to the ~15% EMG amplitude

suppression in SOL induced by subthreshold TMS during walking and jumping (Petersen et

al., 2001; Zuur et al., 2010), but smaller compared with the 50% suppression in hand and arm

muscles during voluntary fixed-end contractions (Davey et al., 1994). Also, the latency and

the duration of the observed EMG amplitude suppression was similar to values reported

previously (Petersen et al., 2001; Zuur et al., 2010). Importantly, the TMS-induced EMG

amplitude suppression showed no difference between STR and REF conditions, which also

resulted in the same magnitudes of torque reduction for both contraction conditions. Contrary

to our first prediction, this indicates that corticospinal drive does not contribute to the
increased torque production following active muscle stretch.

Our data also does not support our second prediction that suprathreshold TMS elicits larger
superimposed twitch torque amplitudes in the presence of rFE compared with fixed-end
reference contractions (twitch amplitude of ~9 Nm for both contraction types). This further
supports the interpretation of the subthreshold TMS-induced EMG amplitude suppression that
corticospinal drive does not contribute to the increased torque production following active
muscle stretch. As the facilitation of the motor cortex with a given suprathreshold stimulation
intensity elicited comparable twitch torques for both contraction conditions, despite steady-
state torques after active stretch being enhanced because of rFE, our findings indicate that rFE
is a mechanical property of muscle (Freundt and Linke, 2019; Herzog, 2018; Nishikawa,
2016).

A limitation of the current study was that we only performed TMS of the motor cortex to
assess corticospinal excitability, which does not allow us to distinguish between cortical and
spinal aspects of corticospinal modulation. The unchanged MEP amplitudes and SP durations
that we found in the presence of rFE compared with the reference contractions is in line with
the findings of Sypkes et al. (2018) and indicate that the overall corticospinal excitability was
unaffected. However, unchanged MEPs do not exclude potential spinal modulations and the
smaller CMEP amplitudes in the presence of rFE observed by Sypkes et al. (2018) could be
interpreted as reduced spinal motoneuron excitability, which might be due to tension-related
afferent feedback from Golgi tendon organs (Contento et al. 2019). Accordingly, we assume
that the unchanged MEP amplitudes in our study reflect an increased cortical excitability,
which did not contribute to the increased torque production following active muscle stretch.

In conclusion, we found that subthreshold and suprathreshold TMS of motor cortical neurons
affected muscle activity and torque production, but the changes did not differ between stretch-

hold and fixed-end reference contractions. Contrary to our predictions, a contribution of

corticospinal drive to the increased torque production following active muscle stretch could

not be confirmed. However, it remains unknown whether and how the spinal modulations

observed by others (Altenburg et al. 2008, Sypkes et al. 2018, Contento et al. 2019) affect

torque production in the presence of rFE.

Disclosures

The authors report no conflicts of interest.

Endnote

At the request of the authors, readers are herein alerted to the fact that additional materials

related to this manuscript may be found at https://figshare.com/articles/Frischholz_et_al_data

_xlsx/11919828. These materials are not a part of this manuscript and have not undergone

peer review by the American Physiological Society (APS). APS and the journal editors take

no responsibility for these materials, for the website address, or for any links to or from it.

[Data currently stored, but not publicly available. Data will be formally published, including

details to accepted manuscript, once paper accepted.]

References

Abbott BC, Aubert XM. The force exerted by active striated muscle during and after change

of length. *The Journal of Physiology* 117: 77–86, 1952.

Altenburg TM, Ruiter CJ de, **Verdijk PWL, van Mechelen W, Haan A** de. Vastus

lateralis surface and single motor unit EMG following submaximal shortening and

lengthening contractions. *Applied Physiology, Nutrition, and Metabolism* 33: 1086–1095,

2008.

Chapman N, Whitting J, Broadbent S, Crowley-McHattan Z, Meir R. Residual Force

Enhancement in Humans. A Systematic Review. *Journal of Applied Biomechanics* 34: 240–

248, 2018.

Contento VS, Dalton BH, Power GA. The Inhibitory Tendon-Evoked Reflex Is Increased in

the Torque-Enhanced State Following Active Lengthening Compared to a Purely Isometric

Contraction. *Brain Sciences* 10, 2019.

Davey NJ, Romaiguère P, Maskill DW, Ellaway PH. Suppression of voluntary motor

activity revealed using transcranial magnetic stimulation of the motor cortex in man. *The*

Journal of Physiology 477: 223–235, 1994.

Duchateau J, Enoka RM. Neural control of lengthening contractions. *The Journal of*

Experimental Biology 219: 197–204, 2016.

Edman KAP. Residual force enhancement after stretch in striated muscle. A consequence of

increased myofilament overlap? *The Journal of Physiology* 590: 1339–1345, 2012.

Edman KAP, Elzinga G, Noble MIM. Residual Force Enhancement after Stretch of

Contracting Frog Single Muscle Fibres. *The Journal of General Physiology* 80: 769–784,

1982.

Freundt JK, Linke WA. Titin as a force-generating muscle protein under regulatory control.

Journal of Applied Physiology 126: 1474–1482, 2019.

Hahn D. Stretching the limits of maximal voluntary eccentric force production in vivo.

Journal of Sport and Health Science 7: 275–281, 2018.

Hahn D, Hoffman BW, Carroll TJ, Cresswell AG. Cortical and spinal excitability during
and after lengthening contractions of the human plantar flexor muscles performed with
maximal voluntary effort. *PloS one* 7: e49907, 2012.

Hahn D, Seiberl W, Schmidt S, Schweizer K, Schwirtz A. Evidence of residual force
enhancement for multi-joint leg extension. *Journal of Biomechanics* 43: 1503–1508, 2010.

Herzog W. Mechanisms of enhanced force production in lengthening (eccentric) muscle
contractions. *Journal of Applied Physiology* 116: 1407–1417, 2014.

Herzog W. The multiple roles of titin in muscle contraction and force production. *Biophysical*
Reviews 10: 1187–1199, 2018.

Jones AA, Power GA, Herzog W. History dependence of the electromyogram. Implications
for isometric steady-state EMG parameters following a lengthening or shortening
contraction. *Journal of Electromyography and Kinesiology* 27: 30–38, 2016.

Joumaa V, Herzog W. Energy cost of force production is reduced after active stretch in
skinned muscle fibres. *Journal of Biomechanics* 46: 1135–1139, 2013.

Mazara N, Hess AJ, Chen J, Power GA. Activation reduction following an eccentric
contraction impairs torque steadiness in the isometric steady-state. *Journal of Sport and*
Health Science 7: 310–317, 2018.

Nishikawa K. Eccentric contraction. Unraveling mechanisms of force enhancement and
energy conservation. *The Journal of Experimental Biology* 219: 189–196, 2016.

Oskouei AE, Herzog W. Observations on force enhancement in submaximal voluntary
contractions of human adductor pollicis muscle. *Journal of Applied Physiology* 98: 2087–
2095, 2005.

Paquin J, Power GA. History dependence of the EMG-torque relationship. *Journal of*
Electromyography and Kinesiology 41: 109–115, 2018.

Paternoster FK, Seiberl W, Hahn D, Schwirtz A. Residual force enhancement during

multi-joint leg extensions at joint- angle configurations close to natural human motion.

Journal of Biomechanics 49: 773–779, 2016.

Petersen NT, Butler JE, Marchand-Pauvert V, Fisher R, Ledebt A, Pyndt HS, Hansen

NL, Nielsen JB. Suppression of EMG activity by transcranial magnetic stimulation in

human subjects during walking. *The Journal of Physiology* 537: 651–656, 2001.

Pinniger GJ, Cresswell AG. Residual force enhancement after lengthening is present during

submaximal plantar flexion and dorsiflexion actions in humans. *Journal of Applied*

Physiology 102: 18–25, 2007.

Rassier DE. The mechanisms of the residual force enhancement after stretch of skeletal

muscle. Non-uniformity in half-sarcomeres and stiffness of titin. *Proceedings. Biological*

Sciences 279: 2705–2713, 2012.

Rossini PM, Burke D, Chen R, Cohen LG, Daskalakis Z, Di Iorio R, Di Lazzaro V,

Ferreri F, Fitzgerald PB, George MS, Hallett M, Lefaucheur JP, Langguth B,

Matsumoto H, Miniussi C, Nitsche MA, Pascual-Leone A, Paulus W, Rossi S,

Rothwell JC, Siebner HR, Ugawa Y, Walsh V, Ziemann U. Non-invasive electrical and

magnetic stimulation of the brain, spinal cord, roots and peripheral nerves. Basic principles

and procedures for routine clinical and research application. An updated report from an

I.F.C.N. Committee. *Clinical Neurophysiology* 126: 1071–1107, 2015.

Rothwell JC. Techniques and mechanisms of action of transcranial stimulation of the human

motor cortex. *Journal of Neuroscience Methods* 74: 113–122, 1997.

Seiberl W, Hahn D, Herzog W, Schwirtz A. Feedback controlled force enhancement and

activation reduction of voluntarily activated quadriceps femoris during sub-maximal muscle

action. *Journal of Electromyography and Kinesiology* 22: 117–123, 2012.

Seiberl W, Hahn D, Kreuzpointner F, Schwirtz A, Gastmann U. Force Enhancement of

Quadriceps Femoris in Vivo and Its Dependence on Stretch-Induced Muscle Architectural

Changes. *Journal of Applied Biomechanics* 26: 256–264, 2010.

Seiberl W, Paternoster F, Achatz F, Schwirtz A, Hahn D. On the relevance of residual

force enhancement for everyday human movement. *Journal of Biomechanics* 46: 1996–

2001, 2013.

Seiberl W, Power GA, Hahn D. Residual force enhancement in humans. Current evidence

and unresolved issues. *Journal of Electromyography and Kinesiology* 25: 571–580, 2015.

Sypkes CT, Kozlowski BJ, Grant J, Bent LR, McNeil CJ, Power GA. The influence of

residual force enhancement on spinal and supraspinal excitability. *PeerJ* 6: e5421, 2018.

Zuur AT, Lundbye-Jensen J, Leukel C, Taube W, Grey MJ, Gollhofer A, Nielsen JB,

Gruber M. Contribution of afferent feedback and descending drive to human hopping. *The*

Journal of Physiology 588: 799–807, 2010.

Figure legends

Figure 1. Example data from the active stretch-hold (STR, dark blue) and fixed-end reference

(REF, light blue) contraction conditions. A: The traces show soleus (SOL) EMG (moving

0.25-s RMS amplitudes). Transcranial magnetic stimulation (TMS, vertical black lines) was

delivered at 1 Hz from 5-s after contraction onset (marked as time zero). In case the first

stimulation was delivered before SOL EMG reached the target level, the stimulation was

excluded from analysis. B: Traces show the corresponding crank arm angles.

Figure 2. A: Schematic diagram of the experimental protocol. The protocol started and ended

with maximal voluntary contractions (MVC, dark grey). Transcranial magnetic stimulation

(TMS) hotspot and sub-/suprathreshold intensities (light grey) were then determined before

data collection. Twenty contractions with subthreshold TMS (dark blue) and 4 contractions

with suprathreshold TMS (light blue) were separated into three sets, with 5 min rest between

each set. B: Schematic description of contractions within sets (blue bars) and delivery of TMS

(same procedure for sub- and suprathreshold TMS). Vertical black lines indicate the timing of

TMS during the submaximal contractions. Each set during subthreshold TMS consisted of 10

contractions of 15-s duration, followed by a 3 min rest.

Figure 3. Example data of rectified and averaged soleus (SOL) EMG from contractions with

subthreshold transcranial magnetic stimulation (TMS). The horizontal dotted line represents

the averaged background EMG amplitude. Vertical dashed lines indicate onset and offset of

EMG amplitude suppression following TMS. Time zero indicates the time of stimulation.

Figure 4. Representative example of motor evoked potentials (MEPs) and silent periods (SPs)

from raw soleus (SOL) EMG signals elicited by suprathreshold transcranial magnetic

stimulation (TMS). The light blue traces show the single MEPs (n = 12) and the dark blue

trace shows the average of all SOL MEPs from one participant. The horizontal dashed black

line indicates the duration of the silent period (SP) and the vertical black line indicates the

peak-to-peak amplitude. Time zero indicates the time of stimulation.

Figure 5. Mean EMG amplitudes from the triceps surae muscles following subthreshold

transcranial magnetic stimulation for individual participants (blue symbols and grey lines) and

the group average (black horizontal lines) during active stretch-hold (STR) and fixed-end

reference (REF) contractions.

Figure 6. A: Individual mean net plantar flexion torque values 500-990 ms following

transcranial magnetic stimulation (TMS) with subthreshold intensity for active stretch-hold

(STR) and fixed-end reference (REF) contractions (blue symbols and grey lines). Black

horizontal lines represent the group means. B: Individual mean net plantar flexion torque

values 500-990-ms following TMS with suprathreshold intensity for active stretch-hold (STR)

and fixed-end reference (REF) contractions (blue symbols and grey lines). Black horizontal

lines represent the group means. *indicates that STR and REF were significantly different.

Table 1. Mean values ± SD of EMG amplitude suppression following subthreshold TMS from soleus (SOL), medial gastrocnemius (MG) and lateral gastrocnemius (LG) muscles during active stretch-hold (STR) and fixed-end reference (REF) contractions.

	SOL	MG	LG
EMG suppression REF [%]	23.5 ± 6.5	22.9 ± 8.4	19.4 ± 2.3
EMG suppression STR [%]	24.1 ± 5.1	23.7 ± 6.9	20.9 ± 6.4
Latency REF [ms]	44 ± 4	42 ±5	44 ± 5
Latency STR [ms]	44 ± 3	43 ± 6	46 ± 5
Duration REF [ms]	13 ± 4	13 ± 5	12 ± 5
Duration STR [ms]	13 ± 5	12 ± 4	10 ± 4

Table 2. Mean values ± SD of motor evoked potentials (MEP) and silent periods (SP) after suprathreshold TMS from soleus (SOL), medial gastrocnemius (MG) and lateral gastrocnemius (LG) muscles during active stretch-hold (STR) and fixed-end reference (REF) contractions.

	SOL	MG	LG
MEP REF [V]	1.30 ± 0.78	2.54 ± 2.67	1.19 ± 0.46
MEP STR [V]	1.37 ± 0.72	2.70 ± 2.97	1.14 ± 0.50
SP REF [ms]	115 ± 31	112 ± 22	118 ± 29
SP STR [ms]	116 ± 31	115 ± 23	123 ± 34

A

MVC

TMS hotspot & intensity

Data collection (subthreshold TMS)

Data collection (suprathreshold TMS)

100% MVC

30% MVC

~ 20 min   ~ 35 min   5 min   ~ 35 min   ~ 15 min

B