Genomic diversity of *Escherichia coli* isolates from non-human primates in the Gambia

Ebenezer Foster-Nyarko[1,2], Nabil-Fareed Alikhan[1], Anuradha Ravi[1], Gaëtan Thilliez[1],

Nicholas Thomson[1], David Baker[1], Gemma Kay[1], Jennifer D. Cramer[3], Justin O'Grady[1],

Martin Antonio[2,4], Mark J. Pallen[1,5†]

[1] Quadram Institute Bioscience, Norwich Research Park, Norwich, Norfolk, United Kingdom

[2] Medical Research Council Unit the Gambia at the London School of Hygiene and Tropical

Medicine, Atlantic Boulevard Road, Fajara, The Gambia

[3] American Public University System, Charles Town, WV, USA

[4] Microbiology and Infection Unit, Warwick Medical School, University of Warwick,

Coventry, United Kingdom

[5] School of Veterinary Medicine, University of Surrey, Guildford, Surrey, United Kingdom.

[†]Correspondence: Professor Mark Pallen, Quadram Institute Bioscience, Norwich Research

Park, Norwich, Norfolk, United Kingdom

Email: Mark.Pallen@quadram.ac.uk

Abstract

Increasing contact between humans and non-human primates provides an opportunity for the transfer of potential pathogens or antimicrobial resistance between host species. We have investigated genomic diversity, and antimicrobial resistance in *Escherichia coli* isolates from four species of non-human primate in the Gambia: *Papio papio* (n=22), *Chlorocebus sabaeus* (n=14), *Piliocolobus badius* (n=6) and *Erythrocebus patas* (n=1). We performed Illumina whole-genome sequencing on 101 isolates from 43 stools, followed by nanopore long-read sequencing on eleven isolates. We identified 43 sequence types (STs) by the Achtman scheme (ten of which are novel), spanning five of the eight known phylogroups of *E. coli*. The majority of simian isolates belong to phylogroup B2—characterised by strains that cause human extraintestinal infections—and encode factors associated with extraintestinal disease. A subset of the B2 strains (ST73, ST681 and ST127) carry the *pks* genomic island, which encodes colibactin, a genotoxin associated with colorectal cancer. We found little antimicrobial resistance and only one example of multi-drug resistance among the simian isolates. Hierarchical clustering showed that simian isolates from ST442 and ST349 are closely related to isolates recovered from human clinical cases (differences in 50 and seven alleles respectively), suggesting recent exchange between the two host species. Conversely, simian isolates from ST73, ST681 and ST127 were distinct from human isolates, while five simian isolates belong to unique core-genome ST complexes—indicating novel diversity specific to the primate niche. Our results are of public health importance, considering the increasing contact between humans and wild non-human primates.

Keywords

Non-human primates, *Escherichia coli,* phylogenomic diversity, Extraintestinal pathogenic *E. coli*.

Impact statement

Little is known about the population structure, virulence potential and the burden of antimicrobial resistance among *Escherichia coli* from wild non-human primates, despite increased exposure to humans through the fragmentation of natural habitats. Previous studies, primarily involving captive animals, have highlighted the potential for bacterial exchange between non-human primates and humans living nearby, including strains associated with intestinal pathology. Using multiple-colony sampling and whole-genome sequencing, we investigated the strain distribution and population structure of *E. coli* from wild non-human primates from the Gambia. Our results indicate that these monkeys harbour strains that can cause extraintestinal infections in humans. We document the transmission of virulent *E. coli* strains between monkeys of the same species sharing a common habitat and evidence of recent interaction between strains from humans and wild non-human primates. Also, we present complete genome assemblies for five novel sequence types of *E. coli*.

Author notes

All supporting data, code and protocols have been provided within the article or through supplementary data files. Nine supplementary figures and six supplementary files are available with the online version of this article.

Abbreviations

ExPEC, Extraintestinal pathogenic *Escherichia coli*; ST, Sequence type; AMR, Antimicrobial resistance; MLST, Multi-locus sequence typing; VFDB, Virulence factors database; SNP, single nucleotide polymorphism; SPRI, Solid phase reversible immobilisation.

Data summary

The raw sequences and polished assemblies from this study are available in the National

Center for Biotechnology Information (NCBI) Short Read Archive, under the BioProject

accession number PRJNA604701. The full list and characteristics of these strains and other

reference strains used in the analyses are presented in Table 1 and Supplementary Files 1-4

(available with the online version of this article).

Introduction

Escherichia coli is a highly versatile species, capable of adapting to a wide range of

ecological niches and colonising a diverse range of hosts (1, 2). In humans, *E. coli* colonises

the gastrointestinal tract as a commensal, as well as causing intestinal and extraintestinal

infection (2). *E. coli* is also capable of colonising the gut in non-human primates (3-5), where

data from captive animals suggest that gut isolates are dominated by phylogroups B1 and A,

which, in humans, encompass commensals as well as strains associated with intestinal

pathology (6-9). *E. coli* strains encoding colibactin, or cytotoxic necrotising factor 1 have

been isolated from healthy laboratory rhesus macaques (4, 10), while enteropathogenic *E.

coli* strains can—in the laboratory—cause colitis in marmosets (11), rhesus macaques

infected with simian immunodeficiency virus (12) and cotton-top tamarins (13).

There are two potential explanations for the co-occurrence of *E. coli* in humans and non-

human primates. Some bacterial lineages may have been passed on through vertical

transmission within the same host species for long periods, perhaps even arising from

ancestral bacteria that colonised the guts of the most recent common ancestors of humans and

non-human primate species (14-16). In such a scenario, isolates from non-human primates

would be expected to be novel and distinct from the diversity seen in humans. However, there

is also clearly potential for horizontal transfer of strains from one host species to another

(17).

The exchange of bacteria between humans and human-habituated animals, particularly

non-human primates, is of interest in light of the fragmentation of natural habitats globally

(18-28). We have seen that wild non-human primates in the Gambia are frequently exposed

to humans through tourism, deforestation and urbanisation. In Uganda, PCR-based studies

have suggested transmission of *E. coli* between humans, non-human primates and livestock

(26-28). Thus, wild non-human primates may constitute a reservoir for the zoonotic spread of

E. coli strains associated with virulence and antimicrobial resistance to humans.

Alternatively, humans might provide a reservoir of strains with the potential for

anthroponotic spread to animals—or transmission might occur in both directions (29).

We do not know how many different lineages can co-exist within the same non-human

primate host. Such information may help us contextualise the potential risks associated with

transmission of bacterial strains between humans and non-human primates. In humans, up to

eleven serotypes could be sampled from picking eleven colonies from individual stool

samples (30).

To address these issues, we have exploited whole-genome sequencing to explore the

colonisation patterns, population structure and phylogenomic diversity of *E. coli* in wild non-

human primates from rural and urban Gambia.

Methods

Study population and sample collection

In June 2017, wild non-human primates were sampled from six sampling sites in the Gambia:

Abuko Nature Reserve (riparian forest), Bijilo Forest Park (coastal fenced woodland),

Kartong village (mangrove swamp), Kiang West National park (dry-broad-leaf forest),

Makasutu Cultural Forest (ecotourism woodland) and River Gambia National park (riparian

forest) (Figure 1). We sampled all four of the diurnal non-human primate species indigenous

to the Gambia. Monkeys in Abuko and Bijilo are frequently hand-fed by visiting tourists,

despite prohibiting guidelines (31).

Troops of monkeys were observed and followed. We collected a single freshly passed

formed stool specimen from 43 visibly healthy individuals (38 adults, 5 juveniles; 24

females, 11 males, 8 of undetermined sex), drawn from four species: *Erythrocebus patas*

(patas monkey), *Papio papio* (Guinea baboon), *Chlorocebus sabaeus* (green monkey) and

Piliocolobus badius (Western colobus monkey). Stool samples were immediately placed into

sterile falcon tubes, taking care to collect portions of stool material that had not touched the

ground, then placed on dry ice and stored at 80°C within 6 h. The sample processing flow is

summarised in Figure 2.

Microbiological processing

For the growth and isolation of *E. coli*, 0.1–0.2 g aliquots were taken from each stool sample

into 1.5 ml microcentrifuge tubes under aseptic conditions. To each tube, 1 ml of

physiological saline (0.85%) was added, and the saline-stool samples were vortexed for 2 min

at 4200 rpm. The homogenised samples were taken through four ten-fold serial dilutions and

a 100 µl aliquot from each dilution was spread on a plate of tryptone-bile-X-glucoronide agar

using the cross-hatching method. Plates were incubated at 37°C for 18–24 h in air. Colony

counts were performed for each serial dilution, counting translucent colonies with blue-green

pigmentation and entire margins as *E. coli*. Up to five colonies from each sample were sub-

cultured on MacConkey agar at 37°C for 18–24 h and then stored in 20% glycerol broth at -

80°C.

Genomic DNA extraction

A single colony from each subculture was picked into 1 ml Luria-Bertani broth and incubated

overnight at 37°C. Broth cultures were spun at 3500rpm for 2 min and lysed using lysozyme,

proteinase K, 10% SDS and RNase A in Tris EDTA buffer (pH 8.0). Suspensions were

placed on a thermomixer with vigorous shaking at 1600 rpm, first at 37°C for 25 min and

subsequently at 65°C for 15 min. DNA was extracted using solid-phase reversible

immobilisation magnetic beads (Becter Coulter Inc., Brea, CA, U.S.A.), precipitated with

ethanol, eluted in Tris-Cl and evaluated for protein and RNA contamination using A_{260}/A_{280}

and A_{260}/A_{230} ratios on the NanoDrop 2000 Spectrophotometer (Fisher Scientific,

Loughborough, UK). DNA concentrations were measured using the Qubit HS DNA assay

(Invitrogen, MA, USA). DNA was stored at -20°C.

Illumina sequencing

Whole-genome sequencing was carried out on the Illumina NextSeq 500 platform (Illumina,

San Diego, CA). We used a modified Nextera XT DNA protocol for the library preparation

as follows. The genomic DNA was normalised to 0.5 ng μl^{-1} with 10 mM Tris-HCl. Next, 0.9

µl of Tagment DNA buffer (Illumina Catalogue No. 15027866) was mixed with 0.09 µl of

Tagment DNA enzyme (Illumina Catalogue No. 15027865) and 2.01 µl of PCR-grade water

in a master-mix. Next, 3 µl of the master-mix was added to a chilled 96-well plate. To this, 2

µl of normalised DNA (1 ng total) was added, pipette-mixed and the reaction heated to 55°C

for 10 min on a PCR block. To each well, we added 11 µl of KAPA2G Robust PCR master-

mix (Sigma Catalogue No. KK5005), comprising 4 µl KAPA2G buffer, 0.4 µl dNTPs, 0.08

µl polymerase and 6.52 µl PCR-grade water, contained in the kit per sample. Next, 2 µl each

of P7 and P5 Nextera XT Index Kit v2 index primers (Illumina Catalogue numbers FC-131-

2001 to 2004) were added to each well. Finally, the 5 µl of Tagmentation mix was added and

mixed. The PCR was run as follows: 72°C for 3 min, 95°C for 1 min, 14 cycles of 95°C for

10 sec, 55°C for 20 sec and 72°C for 3 min. Following the PCR, the libraries were quantified

using the Quant-iT dsDNA Assay Kit, high sensitivity kit (Catalogue No. 10164582) and run

on a FLUOstar Optima plate reader. After quantification, libraries were pooled in equal

quantities. The final pool was double-SPRI size-selected between 0.5 and 0.7x bead volumes

using KAPA Pure Beads (Roche Catalogue No. 07983298001). We then quantified the final

pool on a Qubit 3.0 instrument (Invitrogen, MA, USA) and ran it on a high sensitivity D1000

ScreenTape (Agilent Catalogue No. 5067-5579) using the Agilent TapeStation 4200 to

calculate the final library pool molarity. The pooled library was run at a final concentration of

1.8 pM on an Illumina NextSeq500 instrument using a mid-output flow cell (NSQ® 500 Mid

Output KT v2 300 cycles; Illumina Catalogue No. FC-404-2003) following the Illumina

recommended denaturation and loading parameters, which included a 1% PhiX spike (PhiX

Control v3; Illumina Catalogue FC-110-3001). The data was uploaded to BaseSpace

(http://www.basespace.illumina.com) and then converted to FASTQ files.

Oxford nanopore sequencing

We used the rapid barcoding kit (Oxford Nanopore Catalogue No. SQK-RBK004) to prepare

libraries according to the manufacturer's instructions. We used 400 ng DNA for library

preparation and loaded 75 µl of the prepared library on an R9.4 MinION flow cell. The size

of the DNA fragments was assessed using the Agilent 2200 TapeStation (Agilent Catalogue

No. 5067-5579) before sequencing. The concentration of the final library pool was measured

using the Qubit high-sensitivity DNA assay (Invitrogen, MA, USA).

Genome assembly and phylogenetic analysis

Sequences were analysed on the Cloud Infrastructure for Microbial Bioinformatics (32).

Paired-end short-read sequences were concatenated, then quality-checked using FastQC

v0.11.7 (33). Reads were assembled using Shovill (https://github.com/tseemann/shovill) and

assemblies assessed using QUAST v 5.0.0, de6973bb (34). Draft bacterial genomes were

annotated using Prokka v 1.13 (35). Multi-locus sequence types were called from assemblies

according to the Achtman scheme using the mlst software (https://github.com/tseemann/mlst)

to scan alleles in PubMLST (https://pubmlst.org/) (36). To identify and assign new STs, we

used the ST search algorithm in EnteroBase, allowing for one allele mismatch (37). Snippy

v4.3.2 (https://github.com/tseemann/snippy) was used for variant calling and core genome

alignment, including references genome sequences representing the major phylogroups of *E.*

coli and *Escherichia fergusonii* as an outgroup (Supplementary File 1B). We used Gubbins

(Genealogies Unbiased By recomBinations In Nucleotide Sequences) to detect and remove

recombinant regions of the core genome alignment (38). RAxML v 8.2.4 (39) was used for

maximum-likelihood phylogenetic inference from this masked alignment based on a general

time-reversible nucleotide substitution model with 1,000 bootstrap replicates. The

phylogenetic tree was visualised using Mega v. 7.2 (40) and annotated using Adobe

Illustrator v 23.0.3 (Adobe Inc., San Jose, California). Pair-wise single nucleotide

polymorphism (SNP) distances between genomes were computed from the core-gene

alignment using snp-dists v0.6 (https://github.com/tseemann/snp-dists).

Population structure and analysis of gene content

Merged short reads were uploaded to EnteroBase (41) where we used the Hierarchical

Clustering (HierCC) algorithm to assign our genomes from non-human primates to HC1100

clusters, which in *E. coli* correspond roughly to the clonal complexes seen in seven-allele

MLST. Core genome MLST (cgMLST) profiles based on the typing of 2, 512 core loci for *E.*

coli facilitates single-linkage hierarchical clustering according to fixed core genome MLST

(cgMLST) allelic distances, based on cgMLST allelic differences. Thus, cgST HierCC

provides a robust approach to analyse population structures at multiple levels of resolution.

The identification of closely-related genomes using HierCC has been shown to be 89%

consistent between cgMLST and single-nucleotide polymorphisms (42). Neighbour-joining

trees were reconstructed with Ninja—a hierarchical clustering algorithm for inferring

phylogenies that is capable of scaling to inputs larger than 100,000 sequences (43).

ARIBA v2.12.1 (44) was used to search short reads against the Virulence Factors

Database (45) (VFDB-core) (virulence-associated genes), ResFinder (AMR) (46) and

PlasmidFinder (plasmid-associated genes) (47) databases (both ResFinder and

PlasmidFinder databases downloaded 29 October 2018). Percentage identity of ≥ 90% and

coverage of ≥ 70% of the respective gene length were taken as a positive result. Analyses

were performed on assemblies using ABRicate v 0.8.7

(https://github.com/tseemann/abricate). A heat map of detected virulence- and AMR-

associated genes was plotted on the phylogenetic tree using ggtree and phangorn in R studio

v 3.5.1. We searched EnteroBase for all *E. coli* strains isolated from humans in the Gambia

(n=128), downloaded the genomes and screened them for resistance genes using ABRicate v

0.9.8. Assembled genomes for isolates that clustered with our colibactin-encoding ST73,

ST127 and ST681 isolates were downloaded and screened for the colibactin operon using

ABRicate's VFDB database (accessed 28 July 2019). Assemblies reported to contain

colibactin genes were aligned against the colibactin-encoding *Escherichia coli* IHE3034

reference genome (NCBI Accession: GCA_000025745.1) using minimap2 2.13-r850. BAM

files were visualised in Artemis Release 17.0.1 (48) to confirm the presence of the *pks*

genomic island which encodes the colibactin operon.

Hybrid assembly and analysis of plasmids and phages

Base-called FASTQ files were concatenated into a single file and demultiplexed into

individual FASTQ files based on barcodes, using the qcat python command-line tool v 1.1.0

(https://github.com/nanoporetech/qcat). Hybrid assemblies of the Illumina and nanopore

reads were created with Unicycler (49). The quality and completion of the hybrid assemblies

were assessed with QUAST v 5.0.0, de6973bb and CheckM (34, 50). Hybrid assemblies were

interrogated using ABRicate PlasmidFinder and annotated using Prokka (35). Plasmid

sequences were visualised in Artemis using coordinates from ABRicate. Prophage

identification was carried out using the phage search tool, PHASTER (51).

Antimicrobial susceptibility

We determined the minimum inhibitory concentrations of amikacin, trimethoprim,

sulfamethoxazole, ciprofloxacin, cefotaxime and tetracycline for the isolates from non-human

primates using agar dilution (52). Two-fold serial dilutions of each antibiotic were performed

in molten Mueller-Hinton agar (Oxoid, Basingstoke, UK), from 32mg/L to 0.03 mg l^{-1} (512

mg l^{-1} to 0.03 mg l^{-1} for sulfamethoxazole), using *E. coli* NCTC 10418 as control. MICs were

performed in duplicate and interpreted using breakpoint tables from the European Committee

on Antimicrobial Susceptibility Testing v. 9.0, 2019 (http://www.eucast.org).

Results

Twenty-four of 43 samples (56%) showed growth indicative of *E. coli*, yielding a total of 106

colonies. The isolates were designated by the primate species and the site from which they

were sampled as follows: *Chlorocebus sabaeus*, 'Chlos'; *Papio papio*, 'Pap'; *Piliocolobus*

badius, 'Prob'; Abuko Nature Reserve, 'AN'; Bijilo Forest Park, 'BP'; Kartong village, 'K';

Kiang West National Park, 'KW'; Makasutu Cultural Forest, 'M'; and River Gambia

National Park, 'RG'. After genome sequencing, five isolates (PapRG-04, (n=1); PapRG-03

(n=1); ChlosRG-12 (n=1); ChlosAN-13 (n=1); ProbAN-19 (n=1)) were excluded due to low

depth of coverage (<20x), leaving 101 genomes for subsequent analysis (Table 1).

We recovered 43 seven-allele sequence types (ten of them novel), spanning five of the

eight known phylogroups of *E. coli* and comprising 38 core-genome MLST complexes

(Figure 3). The majority of strains belonged to phylogroup B2 (42/101, 42%), which

encompasses strains that cause extraintestinal infections in humans (ExPEC strains) (6-8).

Strains from phylogroup B2 carried colonisation and fitness factors associated with

extraintestinal disease in humans (Figure 3). A subset of the B2 strains (13/42, 31%),

belonging to STs 73, 681 and 127, carried the *pks* genomic island, which encodes the DNA

alkylating genotoxin, colibactin. Colibactin-encoding *E. coli* frequently cause colorectal

cancer, urosepsis, bacteraemia and prostatitis, and are highly associated with other virulence

factors such as siderophores and toxins (53-56).

Thirteen individuals were colonised by two or more STs and nine by two or more

phylogroups (Supplementary File 1A). Five colony picks from a single Guinea baboon

(PapRG-06) yielded five distinct STs, two of which are novel. Two green monkeys sampled

from Bijilo (ChlosBP-24 and ChlosBP-25) shared an identical ST73 genotype, while two

Guinea baboons from Abuko shared an ST226 strain—documenting transmission between

monkeys of the same species. Among the monkey isolates, we found several STs associated

with extraintestinal infections and/or AMR in humans: ST73, ST681, ST127, ST226, ST336,

ST349 (57-62).

In seventeen monkeys, we observed a cloud of closely related genotypes (seperated by 0-

5 SNPs, Table 2A) from each strain, suggesting evolution within the host after acqustion of

the strain. However, in two individuals, pair-wise SNP distances between genotypes from the

same ST were susbtantial enough (25 SNPs and 77 SNPs) to suggest multiple acquisitions of

each strain (Table 2B).

We identified the closest neighbours to all the recovered strains from our study (Table 3).

Our results suggest, in some cases, recent interactions between humans or livestock and non-

human primates. However, we also found a diversity of strains specific to the non-human

primate niche. Hierarchical clustering analysis revealed that simian isolates from ST442 and

ST349 (Achtman)— sequence types that are associated with virulence and AMR in humans

(49, 55)—were closely related to human clinical isolates, with differences of 50 alleles and

seven alleles in the core-genome MLST scheme respectively (Supplementary Figures 1-2).

Similarly, we found evidence of recent interaction between simian ST939 isolates and strains

from livestock (Supplementary Figure 3). Conversely, simian ST73, ST127 and ST681

isolates were genetically distinct from human isolates from these sequence types

(Supplementary Figures 4-6). The multi-drug resistant isolate PapAN-14-1 from ST349 was,

however, closely related to an environmental isolate recovered from water (Supplementary

Figure 7).

Five isolates were >1000 alleles away in the core-genome MLST scheme from anything in

EnteroBase (Supplementary Figures 8 & 9). Four of these were assigned to novel sequence

types in the seven-allele scheme (Achtman) (ST8550, ST8525, ST8532, ST8826), while one

belonged to ST1873, which has only two other representatives in EnteroBase: one from a

species of wild bird from Australia (*Sericornis frontalis*); the other from water. Besides,

ST8550, ST8525, ST8532, ST8826 belonged to novel HierCC 1100 groups (cgST

complexes), indicating that they were unrelated to any other publicly available *E. coli*

genomes.

We observed few antimicrobial resistance genes in our study population, compared to

what prevails in isolates from humans in the Gambia (Figure 4). Phenotypic resistance to

single agents was confirmed in ten isolates: to trimethoprim in a single isolate, to

sulfamethoxazole in four unrelated isolates and to tetracycline in four closely related isolates

from a single animal. A single ST2076 (Achtman) isolate (PapAN-14-1) belonging to the

ST349 lineage was resistant to trimethoprim, sulfamethoxazole and tetracycline. The

associated resistance genes were harboured on an IncFIB plasmid.

Eighty percent (81/101) of the study isolates harboured one or more plasmids. We

detected the following plasmid replicon types: IncF (various subtypes), IncB/K/O/Z, I1,

IncX4, IncY, Col plasmids (various subtypes) and plasmids related to p0111 (rep B)

(Supplementary File 2A). Long-read sequencing of six representative samples showed that

the IncFIB plasmids encoded acquired antibiotic resistance, fimbrial adhesins and colicins

(Supplementary File 2B). Also, the IncFIC/FII, ColRNAI, Col156 and IncB/O/K/Z plasmids

encoded fimbrial proteins and colicins. Besides, the IncX and Inc-I-Aplha encoded bundle

forming pili *bfp*B and the heat-stable enterotoxin protein *StbB* respectively.

We generated complete genome sequences of five novel sequence types of *E. coli*

(ST8525, ST8527, ST8532, ST8826, ST8827) within the seven-allele scheme (Achtman)

(Supplementary File 3A) (63). Although none of these new genomes encoded AMR genes,

ono of them (PapRG-04-4) contained an IncFIB plasmid encoding fimbrial proteins, and a

cryptic ColRNA plasmid. PHASTER identified thirteen intact prophages and four incomplete

phage remnants (Supplementary File 3B). Two pairs of genomes from Guinea baboons from

different parks shared common prophages: one pair carrying PHAGE_Entero_933W, the

other PHAGE-Entero_lambda.

336
337
Discussion

We have described the population structure of *E. coli* in diurnal non-human primates living in

rural and urban habitats from the Gambia. Although our sample size was relatively small, we

have recovered isolates that span the diversity previously described in humans and have also

identified ten new sequence types (five of them now with complete genome sequences). This

finding is significant, considering the vast number of *E. coli* genomes that have been

sequenced to date (9, 597 with MLST via sanger sequencing, and 127, 482 via WGS) (64).

Increasing contact between animal species facilitates the potential exchange of pathogens.

Accumulating data shows that ExPEC strains are frequently isolated from diseased

companion animals and livestock—highlighting the potential for zoonotic as well as

anthroponotic transmission (65-70). In a previous study, green monkeys from Bijilo Park

were found to carry lineages of *Staphylococcus aureus* thought to be acquired from humans

(31). Our analyses suggest similar exchange of *E. coli* strains between humans and wild non-

human primates. However, non-human primates also harbour *E. coli* genotypes that are

clinically important in humans, such as ST73, ST127 and ST681, yet are distinct from those

circulating in humans—probably reflecting lineages that have existed in this niche for long

periods.

We found that several monkeys were colonised with multiple STs, often encompassing

two or more phylotypes. Although colonisation with multiple serotypes of *E. coli* is common

in humans (30, 71) we were surprised to identify as many as five STs in a single baboon.

Sampling multiple colonies from single individuals also revealed within-host diversity arising

from microevolution. However, we also found evidence of acquisition in the same animal of

multiple lineages of the same sequence type, although it is unclear whether this reflects a

single transmission event involving more than one strain or serial transfers.

Antimicrobial resistance in wildlife is known to spread on plasmids through horizontal

gene transfer (72). Given the challenge of resolving large plasmids using short-read

sequences (73), we exploited long-read sequencing to document the contribution of plasmids

to the genomic diversity that we observed in our study population. Consistent with previous

reports (74), we found IncF plasmids which encoded antimicrobial resistance genes.

Virulence-encoding plasmids, particularly colicin-encoding and the F incompatibility group

ones, have long been associated with several pathotypes of *E. coli* (75). Consistent with this,

we found plasmids that contributed to the dissemination of virulence factors such as the heat-

stable enterotoxin protein *StbB*, colicins and fimbrial proteins.

This study could have been enhanced by sampling human populations living near those of

our non-human primates; however, we compensated for this limitation by leveraging the

wealth of genomes in publicly available databases. Besides, we did not sample nocturnal

monkeys due to logistic challenges; however, these have more limited contact with humans

than the diurnal species. Despite these limitations, however, this study provides insight into

the diversity and colonisation patterns of *E. coli* among non-human primates in the Gambia,

highlighting the impact of human continued encroachment on natural habitats and revealing

important phylogenomic relationships between strains from humans and non-human

primates.

References

1.   Sousa CP. The versatile strategies of *Escherichia coli* pathotypes: a mini review.

Journal of Venomous Animals and Toxins including Tropical Diseases. 2006;12:363-

73.

2.   Kaper JB, Nataro JP, Mobley HL. Pathogenic *Escherichia coli*. Nature Reviews

Microbiology. 2004;2(2):123-40.

3.   Carvalho VM, Irino K, Onuma D, Castro AFPd. Random amplification of

polymorphic DNA reveals clonal relationships among enteropathogenic *Escherichia*

coli isolated from non-human primates and humans. Brazilian Journal of Medical and

Biological Research. 2007;40:237-41.

4.   Martin HR, Taylor NS, Buckley EM, Marini RP, Patterson MM, Fox JG.

Characterization of cytotoxic necrotizing factor 1-producing *Escherichia coli* strains

from faeces of healthy macaques. Journal of Medical Microbiology.

2009;58(10):1354-8.

5.   enaillon O, Skurnik D, Picard B, Denamur E. The population genetics of commensal

Escherichia coli. Nature Reviews Microbiology. 2010;8(3):207-17.

6.   Beghain J, Bridier-Nahmias A, Le Nagard H, Denamur E, Clermont O.

ClermonTyping: an easy-to-use and accurate in silico method for *Escherichia* genus

strain phylotyping. Microbial Genomics. 2018;4(7):e000192.

7.   Escobar-Paramo P, Clermont O, Blanc-Potard AB, Bui H, Le Bouguenec C, Denamur

E. A specific genetic background is required for acquisition and expression of

virulence factors in *Escherichia coli*. Molecular Biology and Evolution.

2004;21(6):1085-94.

8.   Picard B, Garcia JS, Gouriou S, Duriez P, Brahimi N, Bingen E, et al. The link

between phylogeny and virulence in *Escherichia coli* extraintestinal infection.

Infection and Immunity. 1999;67(2):546-53.

9.   Clayton JB, Danzeisen JL, Trent AM, Murphy T, Johnson TJ. Longitudinal

characterization of *Escherichia coli* in healthy captive non-human primates. Frontiers

in Veterinary Science. 2014;1:24.

10.   Feng Y, Mannion A, Madden CM, Swennes AG, Townes C, Byrd C, et al. Cytotoxic

Escherichia coli strains encoding colibactin and cytotoxic necrotizing factor (CNF)

colonize laboratory macaques. Gut Pathogens. 2017;9:71.

11.   Thomson JA, Scheffler JJ. Hemorrhagic typhlocolitis associated with attaching and

effacing *Escherichia coli* in common marmosets. Laboratory Animal Science.

1996;46(3):275-9.

12.   Mansfield KG, Lin KC, Newman J, Schauer D, MacKey J, Lackner AA, et al.

Identification of enteropathogenic *Escherichia coli* in simian immunodeficiency

virus-infected infant and adult rhesus macaques. Journal of Clinical Microbiology.

2001;39(3):971-6.

13.   Mansfield KG, Lin KC, Xia D, Newman JV, Schauer DB, MacKey J, et al.

Enteropathogenic *Escherichia coli* and ulcerative colitis in cotton-top tamarins

(*Saguinus oedipus*). The Journal of Infectious Diseases. 2001;184(6):803-7.

14.   Ley RE, Lozupone CA, Hamady M, Knight R, Gordon JI. Worlds within worlds:

424   evolution of the vertebrate gut microbiota. Nature Reviews Microbiology.

425   2008;6(10):776-88.

426   15.   Muegge BD, Kuczynski J, Knights D, Clemente JC, González A, Fontana L, et al.

427   Diet drives convergence in gut microbiome functions across mammalian phylogeny

428   and within humans. Science (New York, NY). 2011;332(6032):970.

429   16.   Moeller AH, Caro-Quintero A, Mjungu D, Georgiev AV, Lonsdorf EV, Muller MN,

430   et al. Cospeciation of gut microbiota with hominids. Science (New York, NY).

431   2016;353(6297):380-2.

432   17.   Lozupone CA, Stombaugh J, Gonzalez A, Ackermann G, Wendel D, Vázquez-Baeza

433   Y, et al. Meta-analyses of studies of the human microbiota. Genome Research.

434   2013;23(10):1704-14.

435   18.   Iovine RdO, Dejuste C, Miranda F, Filoni C, Bueno MG, de Carvalho VM. Isolation

436   of *Escherichia coli* and *Salmonella* spp. from free-ranging wild animals. Brazillian

437   Journal of Microbiology. 2015;46(4):1257-63.

438   19.   Benavides JA, Godreuil S, Bodenham R, Ratiarison S, Devos C, Petretto M-O, et al.

439   No evidence for transmission of antibiotic-resistant *Escherichia coli* strains from

440   humans to wild western lowland gorillas in Lopé National Park, Gabon. Applied and

441   Environmental Microbiology. 2012;78(12):4281-7.

442   20.   Ryan SJ, Walsh PD. Consequences of non-intervention for infectious disease in

443   African great apes. PLOS one. 2011;6(12):e29030-e.

444   21.   Bublitz DC, Wright PC, Rasambainarivo FT, Arrigo-Nelson SJ, Bodager JR,

445   Gillespie TR. Pathogenic enterobacteria in lemurs associated with anthropogenic

446   disturbance. American Journal of Primatology. 2015;77(3):330-7.

22.   Calvignac-Spencer S, Leendertz SA, Gillespie TR, Leendertz FH. Wild great apes as

sentinels and sources of infectious disease. Clinical Microbiology and Infection.

2012;18(6):521-7.

23.   Daszak P, Cunningham AA, Hyatt AD. Anthropogenic environmental change and the

emergence of infectious diseases in wildlife. Acta Tropica. 2001;78(2):103-16.

24.   Weiss D, Wallace RM, Rwego IB, Gillespie TR, Chapman CA, Singer RS, et al.

Antibiotic-resistant *Escherichia coli* and class 1 integrons in humans, domestic

animals, and wild primates in rural Uganda. Applied and Environmental

Microbiology. 2018;84(21):e01632-18.

25.   Dobson A, Foufopoulos J. Emerging infectious pathogens of wildlife. Philosophical

Transactions of the Royal Society of London. 2001;356(1411):1001-12.

26.   Goldberg TL, Gillespie TR, Rwego IB, Estoff EL, Chapman CA. Forest

fragmentation as cause of bacterial transmission among non-human primates, humans,

and livestock, Uganda. Emerging Infectious Diseases. 2008;14(9):1375-82.

27.   Goldberg TL, Gillespie TR, Rwego IB, Wheeler E, Estoff EL, Chapman CA. Patterns

of gastrointestinal bacterial exchange between chimpanzees and humans involved in

research and tourism in western Uganda. Biological Conservation. 2007;135(4):511-

7.

28.   Rwego IB, Isabirye-Basuta G, Gillespie TR, Goldberg TL. Gastrointestinal bacterial

transmission among humans, mountain gorillas, and livestock in Bwindi Impenetrable

National Park, Uganda. Conservation Biology. 2008;22(6):1600-7.

29.   Rwego IB, Isabirye-Basuta G, Gillespie TR, Goldberg TL. Gastrointestinal bacterial

transmission among humans, mountain gorillas, and livestock in Bwindi Impenetrable

National Park, Uganda. Conservation Biology. 2008;22(6):1600-7.

30.   Lidin-Janson G, Kaijser B, Lincoln K, Olling S, Wedel H. The homogeneity of the

faecal coliform flora of normal school-girls, characterized by serological and

biochemical properties. Medical Microbiology and Immunology. 1978;164(4):247-53.

31.   Senghore M, Bayliss SC, Kwambana-Adams BA, Foster-Nyarko E, Manneh J, Dione

M, et al. Transmission of *Staphylococcus aureus* from humans to green monkeys in

the Gambia as revealed by whole-genome sequencing. Applied and Environmental

Microbiology. 2016;82(19):5910-7.

32.   Connor TR, Loman NJ, Thompson S, Smith A, Southgate J, Poplawski R, et al.

CLIMB (the Cloud Infrastructure for Microbial Bioinformatics): an online resource

for the medical microbiology community. Microbial Genomics. 2016;2(9):e00008.

33.   Wingett SW, Andrews S. FastQ Screen: A tool for multi-genome mapping and quality

control. F1000Res. 2018;7:1338.

34.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for

genome assemblies. Bioinformatics (Oxford, England). 2013;29(8):1072-5.

35.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics (Oxford,

England). 2014;30(14):2068-9.

36.   Jolley KA, Maiden MCJ. BIGSdb: Scalable analysis of bacterial genome variation at

the population level. BMC Bioinformatics. 2010;11(1):595.

37.   A Achtman M, Wain J, Weill F-X, Nair S, Zhou Z, Sangal V, et al. Multilocus

sequence typing as a replacement for serotyping in *Salmonella enterica*. PLOS

Pathogens. 2012;8(6):e1002776.

38.   Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, et al. Rapid

phylogenetic analysis of large samples of recombinant bacterial whole genome

sequences using Gubbins. Nucleic Acids Research. 2015;43(3):e15.

39.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics (Oxford, England). 2014;30(9):1312-3.

497
40.   Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: Molecular

Evolutionary Genetics Analysis version 6.0. Molecular Biology and Evolution.

2013;30(12):2725-9.

41.   Zhou Z, Alikhan N-F, Mohamed K, Achtman M. The user's guide to comparative

genomics with EnteroBase. Three case studies: micro-clades within *Salmonella*

enterica serovar Agama, ancient and modern populations of *Yersinia pestis*, and core

genomic diversity of all *Escherichia*. BioRxiv. 2019:613554.

42.   Frentrup M, Zhou Z, Steglich M, Meier-Kolthoff JP, Göker M, Riedel T, et al. Global

genomic population structure of *Clostridioides difficile*. BioRxiv. 2019:727230.

43.   Wheeler TJ. Large-scale neighbor-joining with NINJA. In: Proceedings of the 9th

Workshop on Algorithms in Bioinformatics. 2009; 5724:375-389.

44.   Hunt M, Mather AE, Sánchez-Busó L, Page AJ, Parkhill J, Keane JA, et al. ARIBA:

rapid antimicrobial resistance genotyping directly from sequencing reads. Microbial

Genomics. 2017;3(10):e000131.

45.   Liu B, Zheng D, Jin Q, Chen L, Yang J. VFDB 2019: a comparative pathogenomic

platform with an interactive web interface. Nucleic Acids Research.

2019;47(D1):D687-D692.

46.   Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al.

Identification of acquired antimicrobial resistance genes. The Journal of

Antimicrobial Chemotherapy. 2012;67(11):2640-4.

47.   Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L, et al.

In silico detection and typing of plasmids using PlasmidFinder and plasmid

multilocus sequence typing. Antimicrobial Agents Chemother. 2014;58(7):3895-903.

48.   Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA. Artemis: an integrated

platform for visualization and analysis of high-throughput sequence-based

experimental data. Bioinformatics (Oxford, England). 2012;28(4):464-9.

49.   Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome

assemblies from short and long sequencing reads. PLOS Computational Biology.

2017;13(6):e1005595.

50.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing

the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. Genome Research. 2015;25(7):1043-55.

51.   Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better,

faster version of the PHAST phage search tool. Nucleic Acids Research.

2016;44(W1):W16-21.

52.   Wiegand I, Hilpert K, Hancock RE. Agar and broth dilution methods to determine the

minimal inhibitory concentration (MIC) of antimicrobial substances. Nature

Protocols. 2008;3(2):163-75.

53.   Krieger JN, Dobrindt U, Riley DE, Oswald E. Acute *Escherichia coli* prostatitis in

previously health young men: bacterial virulence factors, antimicrobial resistance, and

clinical outcomes. Urology. 2011;77(6):1420-5.

54.   Fais T, Delmas J, Barnich N, Bonnet R, Dalmasso G. Colibactin: More than a new

bacterial toxin. Toxins (Basel). 2018;10(4):151.

55.   Johnson JR, Johnston B, Kuskowski MA, Nougayrede JP, Oswald E. Molecular

epidemiology and phylogenetic distribution of the *Escherichia coli* pks genomic

island. Journal of Clinical Microbiology. 2008;46(12):3906-11.

56.   Micenkova L, Benova A, Frankovicova L, Bosak J, Vrba M, Sevcikova A, et al.

Human *Escherichia coli* isolates from hemocultures: Septicemia linked to urogenital

tract infections is caused by isolates harboring more virulence genes than bacteraemia

linked to other conditions. International Journal of Medical Microbiology.

2017;307(3):182-9.

57.   Ho W-S, Gan H-M, Yap K-P, Balan G, Yeo CC, Thong K-L. Genome sequence of

multidrug-resistant *Escherichia coli* EC302/04, isolated from a human tracheal

aspirate. Journal of Bacteriology. 2012;194(23):6691-2.

58.   Manges AR, Johnson JR. Reservoirs of extraintestinal pathogenic *Escherichia coli*.

Microbiology Spectrum. 2015;3(5): UTI-0006-2012.

59.   Kamjumphol W, Wongboot W, Suebwongsa N, Kluabwang P, Chantaroj S, Okada K.

Draft genome sequence of a colistin-resistant *Escherichia coli* ST226: A clinical

strain harbouring an mcr-1 variant. Journal of Global Antimicrobial Resistance.

2019;16:168-9.

60.   Markovska R, Stoeva T, Boyanova L, Stankova P, Schneider I, Keuleyan E, et al.

Multicentre investigation of carbapenemase-producing *Klebsiella pneumoniae* and

Escherichia coli in Bulgarian hospitals – Interregional spread of ST11 NDM-1-

producing *K. pneumoniae*. Infection, Genetics and Evolution. 2019;69:61-7.

61.   Salinas L, Cárdenas P, Johnson TJ, Vasco K, Graham J, Trueba G. Diverse

Commensal *Escherichia coli* clones and plasmids disseminate antimicrobial

resistance genes in domestic animals and children in a semirural community in

Ecuador. mSphere. 2019;4(3):e00316-19.

62.   Manges AR, Geum HM, Guo A, Edens TJ, Fibke CD, Pitout JDD. Global

Extraintestinal Pathogenic *Escherichia coli* (ExPEC) Lineages. Clinical Microbiology

Reviews. 2019;32(3):e00135-18.

63.   Wirth T, Falush D, Lan R, Colles F, Mensa P, Wieler LH, et al. Sex and virulence in

Escherichia coli: an evolutionary perspective. Molecular Microbiology.

2006;60(5):1136-51.

64.   Zhou Z, Alikhan N-F, Mohamed K, Achtman M. The user's guide to comparative

genomics with EnteroBase. Three case studies: micro-clades within *Salmonella*

enterica serovar Agama, ancient and modern populations of *Yersinia pestis*, and core

genomic diversity of all *Escherichia*. BioRxiv. 2019:613554.

65.   Zogg AL, Zurfluh K, Schmitt S, Nuesch-Inderbinen M, Stephan R. Antimicrobial

resistance, multilocus sequence types and virulence profiles of ESBL producing and

non-ESBL producing uropathogenic *Escherichia coli* isolated from cats and dogs in

Switzerland. Veterinary Microbiology. 2018;216:79-84.

66.   Johnson JR, Clabots C, Kuskowski MA. Multiple-host sharing, long-term persistence,

and virulence of *Escherichia coli* clones from human and animal household members.

Journal of Clinical Microbiology. 2008;46(12):4078-82.

584
67.   Johnson JR, Owens K, Gajewski A, Clabots C. *Escherichia coli* colonization patterns

among human household members and pets, with attention to acute urinary tract

infection. Journal of Infectious Disease. 2008;197(2):218-24.

68.   Johnson JR, Miller S, Johnston B, Clabots C, Debroy C. Sharing of *Escherichia coli*

sequence type ST131 and other multidrug-resistant and urovirulent *E. coli* strains

among dogs and cats within a household. Journal of Clinical Microbiology.

2009;47(11):3721-5.

69. Achtman M, Heuzenroeder M, Kusecek B, Ochman H, Caugant D, Selander RK, et

al. Clonal analysis of *Escherichia coli* O2:K1 isolated from diseased humans and

animals. Infection and Immunity. 1986;51(1):268-76.

70. Ewers C, Grobbel M, Stamm I, Kopp PA, Diehl I, Semmler T, et al. Emergence of

human pandemic O25:H4-ST131 CTX-M-15 extended-spectrum-β-lactamase-

producing *Escherichia coli* among companion animals. Journal of Antimicrobial

Chemotherapy. 2010;65(4):651-60.

71. Schlager TA, Hendley JO, Bell AL, Whittam TS. Clonal diversity of *Escherichia coli*

colonizing stools and urinary tracts of young girls. Infection and Immunity.

2002;70(3):1225-9.

72. Vittecoq M, Godreuil S, Prugnolle F, Durand P, Brazier L, Renaud N, et al.

Antimicrobial resistance in wildlife. Journal of Applied Ecology. 2016;53(2):519-29.

73. A rredondo-Alonso S, Willems RJ, van Schaik W, Schürch AC. On the

(im)possibility of reconstructing plasmids from whole-genome short-read sequencing

data. Microbial Genomics. 2017;3(10):e000128.

74. Ca Carattoli A. Resistance plasmid families in *Enterobacteriaceae*. Antimicrobial

Agents and Chemotherapy. 2009;53(6):2227.

75. Johnson TJ, Nolan LK. Pathogenomics of the virulence plasmids of *Escherichia coli*.

Microbiology and Molecular Biology Reviews. 2009;73(4):750-74.

611
612
Data bibliography

1. Foster-Nyarko, E. et al, NCBI BioProject PRJNA604701 (2020).

2. Forde, B. M., Ben Zakour, N. L., Stanton-Cook, M., Phan, M. D., Totsika, M. et al., 17

representative *E. coli* reference isolates (2014). NCBI accession numbers are provided in

Table 1B.

3.   Nougayrede J.P, Homburg S, Taieb F., Boury M., Brzuszkiewicz E., et al., *Escherichia*

coli induces DNA double-strand breaks in eukaryotic cells (2006). NCBI accession:

GCA_000025745.1.

Funding information

MP, EFN, NT, AR, GT, JO and GK were supported by the BBSRC Institute Strategic

Programme Microbes in the Food Chain BB/R012504/1 and its constituent projects

44414000A and 4408000A. NFA and DB were supported by the Quadram Institute

Bioscience BBSRC funded Core Capability Grant (project number BB/ CCG1860/1). The

funders had no role in the study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

Acknowledgements

We want to thank Dr Andrew Page and Dr Thanh Le-Viet for their thoughtful advice on the

long-read analysis. We also thank Dr Mark Webber for proofreading the manuscript and

giving constructive feedback.

Author contributions

Conceptualization, MA, MP; data curation, MP, NFA; formal analysis, EFN, analytical

support, GT; funding, MP and MA; sample collection, JDC; laboratory experiments, EFN,

DB; supervision, AR, NFA, GK, JO, MP, MA; manuscript preparation – original draft, EFN;

review and editing, NT, AR, JO, NFA, MP; review of final manuscript, all authors.

Conflicts of interest

The authors have no conflicts of interest to declare.

Ethical statement

645   No human nor animal experimentation is reported.

Table 1: Study isolates

Name	Source	Individual sampling number	Colony-pick	Sampling site	ST
PapRG-03-1	*Papio papio*	3	1	River Gambia national park	336
PapRG-03-2	*Papio papio*	3	2	River Gambia national park	336
PapRG-03-3	*Papio papio*	3	3	River Gambia national park	336
PapRG-03-4	*Papio papio*	3	4	River Gambia national park	336
PapRG-03-5	*Papio papio*	3	5	River Gambia national park	336
PapRG-04-1	*Papio papio*	4	1	River Gambia national park	1665
PapRG-04-2	*Papio papio*	4	2	River Gambia national park	1204
PapRG-04-4	*Papio papio*	4	3	Makasutu cultural forest	8826
PapRG-04-5	*Papio papio*	4	4	Makasutu cultural forest	1204
PapRG-05-2	*Papio papio*	5	1	Makasutu cultural forest	1431
PapRG-05-3	*Papio papio*	5	2	Makasutu cultural forest	99
PapRG-05-4	*Papio papio*	5	3	Makasutu cultural forest	6316
PapRG-05-5	*Papio papio*	5	4	Makasutu cultural forest	1431
PapRG-06-1	*Papio papio*	6	1	Makasutu cultural forest	4080
PapRG-06-2	*Papio papio*	6	2	Makasutu cultural forest	2521
PapRG-06-3	*Papio papio*	6	3	Makasutu cultural forest	8827
PapRG-06-4	*Papio papio*	6	4	Makasutu cultural forest	1204
PapRG-06-5	*Papio papio*	6	5	River Gambia national park	8525
ProbRG-07-1	*Piliocolobus badius*	7	1	River Gambia national park	73
ProbRG-07-2	*Piliocolobus badius*	7	2	River Gambia national park	73
ProbRG-07-3	*Piliocolobus badius*	7	3	River Gambia national park	73
ProbRG-07-4	*Piliocolobus badius*	7	4	River Gambia national park	73
ProbRG-07-5	*Piliocolobus badius*	7	5	River Gambia national park	73
ChlosRG-12-1	*Chlorocebus sabaeus*	12	1	River Gambia national park	8824
ChlosRG-12-2	*Chlorocebus sabaeus*	12	2	River Gambia national park	196
ChlosRG-12-3	*Chlorocebus sabaeus*	12	3	River Gambia national park	196
ChlosRG-12-5	*Chlorocebus sabaeus*	12	4	River Gambia national park	40
ChlosAN-13-1	*Chlorocebus sabaeus*	13	1	Abuko Nature Reserve	8526
ChlosAN-13-2	*Chlorocebus sabaeus*	13	2	Abuko Nature Reserve	8550
ChlosAN-13-4	*Chlorocebus sabaeus*	13	3	Abuko Nature Reserve	1973
ChlosAN-13-5	*Chlorocebus sabaeus*	13	4	Abuko Nature Reserve	1973
PapAN-14-1	*Papio papio*	14	1	Abuko Nature Reserve	2076
PapAN-14-2	*Papio papio*	14	2	Abuko Nature Reserve	939
PapAN-14-3	*Papio papio*	14	3	Abuko Nature Reserve	226
PapAN-14-4	*Papio papio*	14	4	Abuko Nature Reserve	226
PapAN-14-5	*Papio papio*	14	5	Abuko Nature Reserve	226
PapAN-15-1	*Papio papio*	15	1	Abuko Nature Reserve	226
PapAN-15-2	*Papio papio*	15	2	Abuko Nature Reserve	5073
PapAN-15-3	*Papio papio*	15	3	Abuko Nature Reserve	226
PapAN-15-4	*Papio papio*	15	4	Abuko Nature Reserve	126
PapAN-15-5	*Papio papio*	15	5	Abuko Nature Reserve	8823
ChlosAN-17-1	*Chlorocebus sabaeus*	17	1	Abuko Nature Reserve	681
ChlosAN-17-2	*Chlorocebus sabaeus*	17	2	Abuko Nature Reserve	362
ChlosAN-17-3	*Chlorocebus sabaeus*	17	3	Abuko Nature Reserve	681
ChlosAN-17-4	*Chlorocebus sabaeus*	17	4	Abuko Nature Reserve	681
ChlosAN-18-1	*Chlorocebus sabaeus*	18	1	Abuko Nature Reserve	681
ChlosAN-18-2	*Chlorocebus sabaeus*	18	2	Abuko Nature Reserve	681
ChlosAN-18-3	*Chlorocebus sabaeus*	18	3	Abuko Nature Reserve	681
ChlosAN-18-4	*Chlorocebus sabaeus*	18	4	Abuko Nature Reserve	681
ChlosAN-18-5	*Chlorocebus sabaeus*	18	5	Abuko Nature Reserve	349

ProbAN-19-2	*Piliocolobus badius*	19	1	Abuko Nature Reserve	8825
ChlosBP-21-1	*Chlorocebus sabaeus*	21	1	Bijilo forest park	677
ChlosBP-21-2	*Chlorocebus sabaeus*	21	2	Bijilo forest park	677
ChlosBP-21-3	*Chlorocebus sabaeus*	21	3	Bijilo forest park	677
ChlosBP-21-4	*Chlorocebus sabaeus*	21	4	Bijilo forest park	677
ChlosBP-21-5	*Chlorocebus sabaeus*	21	5	Bijilo forest park	677
ChlosBP-23-1	*Chlorocebus sabaeus*	23	2	Bijilo forest park	8527
ChlosBP-23-2	*Chlorocebus sabaeus*	23	3	Bijilo forest park	8527
ChlosBP-23-3	*Chlorocebus sabaeus*	23	4	Bijilo forest park	3306
ChlosBP-24-1	*Chlorocebus sabaeus*	24	1	Bijilo forest park	73
ChlosBP-24-2	*Chlorocebus sabaeus*	24	2	Bijilo forest park	73
ChlosBP-24-3	*Chlorocebus sabaeus*	24	3	Bijilo forest park	73
ChlosBP-24-4	*Chlorocebus sabaeus*	24	4	Bijilo forest park	73
ChlosBP-24-5	*Chlorocebus sabaeus*	24	5	Bijilo forest park	73
ChlosBP-25-1	*Chlorocebus sabaeus*	25	1	Bijilo forest park	73
ChlosBP-25-2	*Chlorocebus sabaeus*	25	2	Bijilo forest park	73
ChlosBP-25-3	*Chlorocebus sabaeus*	25	3	Bijilo forest park	73
ChlosBP-25-4	*Chlorocebus sabaeus*	25	4	Bijilo forest park	73
ChlosBP-25-5	*Chlorocebus sabaeus*	25	5	Bijilo forest park	73
ChlosM-29-1	*Chlorocebus sabaeus*	29	1	Makasutu cultural forest	1873
ChlosM-29-2	*Chlorocebus sabaeus*	29	2	Makasutu cultural forest	1873
PapM-31-1	*Papio papio*	31	1	Makasutu cultural forest	2800
PapM-31-2	*Papio papio*	31	2	Makasutu cultural forest	135
PapM-31-3	*Papio papio*	31	3	Makasutu cultural forest	5780
PapM-31-4	*Papio papio*	31	4	Makasutu cultural forest	1727
PapM-31-5	*Papio papio*	31	5	Makasutu cultural forest	5780
PapM-32-1	*Papio papio*	32	2	Makasutu cultural forest	8532
PapM-32-2	*Papio papio*	32	3	Makasutu cultural forest	212
PapM-32-3	*Papio papio*	32	4	Makasutu cultural forest	212
PapM-32-4	*Papio papio*	32	5	Makasutu cultural forest	212
PapM-32-5	*Papio papio*	32	6	Makasutu cultural forest	212
PapM-33-1	*Papio papio*	33	1	Makasutu cultural forest	8533
PapM-33-2	*Papio papio*	33	2	Makasutu cultural forest	8533
PapM-33-3	*Papio papio*	33	3	Makasutu cultural forest	8533
PapM-33-4	*Papio papio*	33	4	Makasutu cultural forest	38
PapM-33-5	*Papio papio*	33	5	Makasutu cultural forest	8533
PapM-34-1	*Papio papio*	34	1	Makasutu cultural forest	676
PapM-34-2	*Papio papio*	34	2	Makasutu cultural forest	676
PapM-34-3	*Papio papio*	34	3	Makasutu cultural forest	676
PapM-34-4	*Papio papio*	34	4	Makasutu cultural forest	676
PapM-36-1	*Papio papio*	36	1	Makasutu cultural forest	8535
PapM-36-2	*Papio papio*	36	2	Makasutu cultural forest	8535
PapKW-44-1	*Papio papio*	44	1	Kiang West national park	442
PapKW-44-2	*Papio papio*	44	2	Kiang West national park	442
PapKW-44-3	*Papio papio*	44	3	Kiang West national park	442
PapKW-44-4	*Papio papio*	44	4	Kiang West national park	442
ProbK-45-1	*Piliocolobus badius*	45	1	Kartong village	127
ProbK-45-2	*Piliocolobus badius*	45	2	Kartong village	127
ProbK-45-3	*Piliocolobus badius*	45	3	Kartong village	127
ProbK-45-4	*Piliocolobus badius*	45	4	Kartong village	127
ProbK-45-5	*Piliocolobus badius*	45	5	Kartong village	127

Table 2A: Within-host single nucleotide polymorphism diversity between multiple genomes of the same ST recovered from the same monkey

Sample ID	STs (colonies per ST)	Pair-wise SNP distances between multiple colonies of the same ST	Comment(s)
PapRG-03	336 (n=5)	0-2	
PapRG-04	1204 (n=2)	4	
PapRG-05	1431 (n=2)	0	
ProbRG-07	73 (n=5)	0-1	
ChlosRG-12	196 (n=2)	25	
PapAN-14	226 (n=3)	1	
PapAN-15	226 (n=2)	1	
ChlosAN-17	681 (n=3)	0-3	
ChlosAN-18	681 (n=4)	0	
ChlosBP-21	677 (n=4)	5	
ChlosBP-23	8527 (n=2)	0	
ChlosBP-24	73 (n=5)	0-5	
ChlosBP-25	73 (n=5)	0-79	Please see Table 2B
PapM-32	212 (n=4)	0	
PapM-33	8533 (n=4)	0-4	
PapM-34	676 (n=4)	0-1	
PapM-36	8535 (n=2)	0-1	
PapKW-44	442 (n=4)	1-2	
ProbK-45	127 (n=5)	0-4	

In individuals where multiple colonies yielded the same genotype (n=19), five had entirely identical genotypes, while we observed a cloud of closely related genetic variants (0-5 SNPs, Table 1) in twelve individuals. However, in two monkeys (highlighted with red boxes), pair-wise SNP comparisons suggested multiple infection events (See Table 2B).

Table 2B: Within-host diversity in green monkey 25 (ChlosBP-25)

Sample ID	Clone designation
ChlosBP-25	
ChlosBP-25-1	1
ChlosBP-25-2	2
ChlosBP-25-3	2
ChlosBP-25-4	2
ChlosBP-25-5	3

Pair-wise SNP distances between clones			
	Clone 1	Clone 2	Clone 3
Clone 1	0	12	79
Clone 2	12	0	67
Clone 3	79	67	0

Table 3: Genomic relationship between study isolates and publicly available *E. coli* genomes

7-allele ST	HC100 subgroups	Non-human primate host	Closest neighbours' source	Neighbours' country of isolation	Allelic distance
349	-	*Chlorocebus sabaeus* 18	Human (bloodstream infection)	Canada	7
2076	-	*Papio papio* 14	Environment (water)	Unknown	25
939	-	*Papio papio* 14	Livestock	US	40
442	-	*Papio papio* 44	Human	China	50
2800	-	*Papio papio* 31	Unknown	Vietnam	59
1973	-	*Chlorocebus sabaeus* 13	Unknown	Unknown	64
8533	-	*Papio papio* 33	Environment (water)	Unknown	69
6316	-	*Papio papio* 05	Human	Kenya	97
1727	-	*Papio papio* 34	Human	Kenya	98
676	-	*Papio papio* 34	Human (bloodstream infection)	UK	98
8823	-	*Papio papio* 15	Rodent (guinea pig)	Kenya	101
1431	-	*Papio papio* 05	Human	US	109
5073	-	*Papio papio* 15	Human	US	112
226	73641	*Papio papio* 14	Human	Tanzania	112
8827	-	*Papio papio* 06	Human	Unknown	122
1204	83197	*Papio papio* 04	Livestock	Japan	127
1204	83197	*Papio papio* 04	Livestock	Japan	130
677	-	*Chlorocebus sabaeus* 21	Human	US	132
40	-	*Chlorocebus sabaeus* 12	Human	UK	137
1204	83164	*Papio papio* 06	Livestock	Japan	173
99	-	*Papio papio* 05	Human	UK	180
362	-	*Chlorocebus sabaeus* 17	Food	Kenya	180
8825	-	*Piliocolobus badius* 19	Human	France	189
336	-	*Papio papio* 03	Poultry	Kenya	189
73	-	*Chlorocebus sabaeus* 24	Human	Sweden	189
196	-	*Chlorocebus sabaeus* 12	Human	Sweden	197
2521	-	*Papio papio* 06	Livestock	US	201
127		*Pioliocolobus badius* 45	Companion animal	US	229
681		*ChlosAN* 17	Human	Norway	251
38	-	*Papio papio* 33	human	UK	265
135	-	*Papio papio* 31	Poultry	US	281
8824	-	*Chlorocebus sabaeus* 12	Environmental*	US	296
226	100039	*Papio papio* 14	Human	Sri Lanka	318
8527	-	*Chlorocebus sabaeus* 23	Human	Kenya	323
8535	-	*Papio papio* 36	Environmental (soil)	US	368
1665	-	*Papio papio* 04	Livestock	UK	371
4080	-	*Papio papio* 06	Human	Denmark	507
8526	-	*Chlorocebus sabaeus* 13	Livestock	US	708
8532	-	*Papio papio* 32	Non-human primate	Gambia (PapM-31-3)	1102
8826	-	*Papio papio* 04	Livestock	Mozambique	1255
8525	-	*Papio papio* 06	Livestock/companion animal	Switzerland	1659
1873	-	*Chorocebus sabaeus* 29	Environment	US	1685
8550	-	*Chlorocebus sabaeus* 13	Unknown	Unknown	2006

*Source details unknown.

Isolates from humans were recovered from stools, except where indicated otherwise.

Figure legends

Figure 1. Study sites and distribution of study subjects.

Figure 2. Study sample-processing flow diagram.

Figure 3. A plot showing the maximum likelihood phylogeny of the study isolates overlaid with the prevalence of potential virulence genes among the study isolates. The tree was reconstructed based on non-repetitive core SNPs calculated against the *E. coli* K-12 reference strain (NCBI accession: NC_000913.3), using RAxML with 1000 bootstrap replicates. *E. coli* MG1655 was used as the reference and *E. fergusonii* as the outroot species. Recombinant regions were removed using Gubbins (Reference 38). The tip labels indicate the sample IDs, with the respective in silico Achtman sequence types (STs) and HC1100 (cgST complexes) are indicated next to the tip labels. Both the sample IDs and the STs (Achtman) are colour-coded to indicate the various phylogroups as indicated. Novel STs (Achtman) are indicated by an asterisk (*). *Escherichia fergusonii* and the *E. coli* reference genomes representing the major *E. coli* phylogroups are in black. Primate species are indicated in the strain names as follows: *Chlorocebus sabaeus*, 'Chlos'; *Papio papio*, 'Pap'; *Piliocolobus badius*, 'Prob'. The sampling sites are indicated as follows: BP, Bijilo forest park; KW, Kiang-West National park; RG, River Gambia National Park; M, Makasutu Cultural forest; AN, Abuko Nature reserve; K, Kartong village. Co-colonising seven-allele (Achtman) sequence types (STs) in single individuals are shown by the prefix of the strain names depicting the colony as 1, 2 up to 5. We do not show multiple colonies of the same Achtman ST recovered from a single individual. In such cases, only one representative is shown. Virulence genes are grouped according to their function, with genes encoding the colibactin genotoxin highlighted with a red box. The full names of virulence factors are provided in Supplementary file 5.

Figure 4: A bar graph comparing the prevalence of antimicrobial resistance genotypes in *E. coli* isolated from humans in the Gambia (n=128) as found in EnteroBase (Reference 41) to that found among the study isolates (n=101). The antimicrobial resistance genes detected were as follows: Aminoglycoside: *aph*(6)-Id, ant *aac*(3)-IIa, *ant*(3")-Ia, *aph*(3")-Ib, *aad*A1, *aad*A2; Beta-lactamase: *bla*OXA-1, *bla*TEM-1B, *bla*TEM-1B, *bla*TEM-1C, *bla*SHV-1; Trimethoprim: *dfr*A; Sulphonamide: *sul*1, *sul*2; Tetracycline: *tet*(A), *tet*(B), *tet*(34*), tet*(D); Macrolide, *mph*(A); Chloramphenicol*, cat*A1. Screening of resistance genes was carried out using ARIBA ResFinder (Reference 44) and confirmed by ABRicate (https://github.com/tseemann/abricate). A percentage identity of ≥ 90% and coverage of ≥ 70% of the respective gene length were taken as a positive result.

Supplementary Figure 1. A Ninja neighbour-joining tree showing the phylogenetic relationship between Achtman ST442 strains from this study and all other publicly available genomes that fell within the same HC1100 cluster (cgST complex). The locations of the isolates are displayed, with the genome count displayed in parenthesis. Branch lengths display the allelic distances separating genomes. Gambian strains are highlighted in red. The sub-tree (B) shows the closest relatives to the study strains, with the allelic distance separating them displayed with the arrow. Dotted lines represent long branches which have been shortened.

Supplementary Figure 2. A Ninja neighbour-joining tree showing the phylogenetic relationship between the ST349 (Achtman) strain from this study and all other publicly available genomes within the same HC1100 cluster (cgST complex). The legend shows the locations of the isolates, with genome counts displayed in parenthesis. Gambian strains are

highlighted in red. The study ST349 strain is separated from a clinical ST349 strain by only

seven alleles (<7 SNPs), as depicted in the subtree (B). Long branches are shortened

(indicated by dashes).

Supplementary Figure 3. A phylogenetic neighbour-joining tree reconstructed with the

study ST939 (Achtman) strain and all publicly available genomes that fell within the same

HC1100 cluster (cgST complex). The legend shows the locations of the isolates, with red

highlights around the nodes indicating the Gambian strains. The allelic distance between the

study strain and its nearest relative, a bovine ST939 strain, has been given, depicted by the

arrow. Dotted lines indicate shortened long branches.

Supplementary Figure 4. A Ninja neighbour-joining tree reconstructed with Achtman ST73

colibactin+ strains from this study and all other publicly available ST73 (Achtman) strains

that fell within the same HC1100 cluster (cgST complex) in EnteroBase (Reference 41). The

sources of the isolates are displayed, with Gambian strains highlighted in red. The Gambian

non-human primate strains are on separate long branches, although nested within clades

populated by human strains from other countries, suggestive of probably an ancient

transmission between the two hosts. The branch lengths for the Gambian strains are

displayed. Dotted lines represent long branches which have been shortened.

Supplementary Figure 5. A Ninja neighbour-joining tree showing the phylogenetic

relationship between ST127 strains from this study and other publicly available strains that

occur within the same HC1100 cluster (cgST complex). The sources of the isolates are

displayed in the legends, with Gambian strains highlighted in red. Branch lengths display the

allelic distances separating genomes. The sub-tree (B) shows the closest relatives to the study

strains, with the allelic distances separating them displayed with the arrow. Dotted lines represent long branches which have been shortened. Dotted lines represent long branches which have been shortened.

Supplementary Figure 6. A Ninja neighbour-joining tree showing the phylogenetic relationship between ST681 strains from this study and other publicly available strains that fell within the same HC1100 cluster (cgST complex). The study strains fell into two separate HC100 clusters, which are depicted in the two subtrees (B and C). The closest neighbours to both HC100 clusters are displayed, with the branch labels indicating the allelic distances between strains. The locations of the isolates are displayed for each tree, with Gambian strains highlighted in red. Dotted lines represent long branches which have been shortened.

Supplementary Figure 7. A phylogenetic tree showing the phylogenetic relationship between ST2076 strain (an MDR strain) and all other publicly available genomes that fell within the same HC1100 cluster (cgST complex). The legend shows the locations of the isolates, Gambian strains are highlighted in red. The subtree (B) shows the allelic distance between the study strain and its nearest relative, an ST2076 isolate recovered from water. Dotted lines indicate shortened long branches.

Supplementary Figure 8. A Ninja phylogenetic tree showing the closest neighbours of simian ST1873 strain—an environmental (soil) isolate belonging to ST83, separated from the study strain by 1659 alleles. The legends of both the main tree and the subtree show the locations of the isolates Gambian strains are highlighted in red. In the subtree (B), the closest neighbour to the simian ST1873 strain is also highlighted in red. Dotted lines are used to indicate shortened long branches.

Supplementary Figure 9. Ninja phylogenetic trees showing the closest neighbours to simian isolates belonging to novel sequence types (Achtman) ST8550 (A), ST8532 (B) and ST8525 (C), ST8826 (D). The allelic distances between these study isolates and their closest neighbours are >1100 alleles, and the closest neighbours belong to seven-allele STs which share less than five out of the seven MLST loci. Each genome (ST8550, ST8532, ST8525) belongs to a unique cgST complex (novel groups at HierCC 1100), indicative of novel diversity within the non-human primate niche.

Supplementary files

Supplementary File 1. A. Characteristics of the study population, displaying the primate species, their age and gender, and the *E. coli* sequence types (Achtman MLST STs) and phylotypes recovered from individual samples. Novel STs are designated by an asterisk (*).
B. Reference strains that were included in this study.

Supplementary File 2. A. Predicted plasmids from short-read sequences, using ARIBA PlasmidFinder (Reference 44).
B. A table indicating the virulence and (or) resistance genes located on representative plasmids that were sequenced by Oxford nanopore technology. The size of each plasmid and the functions of the respective genes encoded thereon are also indicated.

Supplementary File 3. A. A summary of the sequencing statistics of the novel sequence types derived from this study. **B.** Prophage types detected from long-read sequences using PHASTER (reference 51).

Supplementary File 4. A summary of the sequencing statistics of the study isolates.

Supplementary File 5. List of virulence factors detected using ARIBA VFDB (Reference 44).

Supplementary File 6. Pair-wise single nucleotide polymorphism distances calculated from the core genome alignment using snp-dists v0.6 (https://github.com/tseemann/snp-dists).

Bijilo National Park
Chlorocebus sabaeus (5)

Abuko Nature Park
Papio papio (2)
Chlorocebus sabaeus (5)
Piliocolobus badius (1)

River Gambia National Park
Papio papio (6)
Piliocolobus badius (4)
Erythrocebus patas (1)
Chlorocebus sabaeus (1)

Kartong village
Piliocolobus badius (1)

Makasutu Cultural Forest
Chlorocebus sabaeus (3)
Papio papio (7)

Kiang West National Park
Papio papio (7)