1   Title: Population structure and dispersal across small and large spatial scales in a direct
2   developing marine isopod

3   Authors: William S. Pearman[a*†], Sarah J. Wells[b*], Olin K. Silander[a], Nikki E. Freed[a], and
4   James Dale[a]

5
[a] School of Natural and Computational Sciences, Massey University, Auckland 0745, New
Zealand

[b] School of Environmental and Animal Sciences, Unitec Institute of Technology, Auckland
1025, New Zealand

*Contributed equally

[†] Corresponding author: wpearman1996@gmail.com

Abstract

Marine organisms generally exhibit one of two developmental modes: biphasic, with distinct adult and larval morphology, and direct development, in which larvae resemble adults. Developmental mode is thought to significantly influence dispersal, with direct developers expected to have much lower dispersal potential. However, in contrast to our relatively good understanding of dispersal and population connectivity for biphasic species, comparatively little is known about direct developers. In this study, we use a panel of 8,020 SNPs to investigate population structure and gene flow for a direct developing species, the New Zealand endemic marine isopod *Isocladus armatus*. On a small spatial scale (20 kms), gene flow between locations is extremely high and suggests an island model of migration. However, over larger spatial scales (600km), populations exhibit a clear pattern of isolation-by-distance. Because our sampling range is intersected by two well-known biogeographic barriers (the East Cape and the Cook Strait), our study provides an opportunity to understand how such barriers influence dispersal in direct developers. Our results indicate that *I. armatus* exhibits significant migration across these barriers, and suggests that ocean currents associated with these locations do not present a barrier to dispersal. Interestingly, we do find evidence of a north-south population genetic break occurring between Māhia and Wellington, two locations where there are no obvious biogeographic barriers between them. We conclude that developmental life history largely predicts dispersal in intertidal marine isopods. However, localised biogeographic processes can disrupt this expectation.

Introduction

Life history influences genetic population structure

A species' developmental life history has a significant effect on population connectivity and dispersal. For marine organisms, developmental life histories are generally classified in one of two ways. Direct developers have juveniles that resemble adults, lacking a pelagic larval phase, and are therefore often hypothesised to have limited dispersal ability relative to their biphasic counterparts. In contrast, biphasic species exhibit a pelagic larval stage have different morphologies from adults and generally disperse large distances via ocean currents (Cowen & Sponaugle, 2009; Puritz et al., 2017; Simpson et al., 2014). Consequently, direct-developing species tend to show greater population genetic structuring than biphasic species (Ayre et al., 2009; McMillan et al., 1992; Pelc et al., 2009; Waples, 1987). However, some species do not conform to this expected pattern (Ayre et al., 2009; Puritz et al., 2017; Shanks, 2009; Winston, 2012).

Dispersal (and thus population structure) is also affected by biogeographic barriers, and the effects of these barriers can themselves depend on the life history of the organism, thus life history and habitat availability jointly act to influence population structure (Ayre et al., 2009). However, habitat availability may be a more important determinant for species with reduced dispersal ability (i.e. direct developers). However, in some cases the lack of nearby suitable habitat appears to limit dispersal in biphasic species more than direct developers (Ayre et al., 2009). Indeed, cases where direct developers exhibit greater dispersal than predicted are not uncommon within intertidal organisms (Ayre et al., 2009; González-Wevar et al., 2018; Wells & Dale, 2018; Yoshino et al., 2018). Therefore, population connectivity cannot be entirely predicted by life history or the knowledge of biogeographic barriers. As a result, it is important to carefully quantify population structure and gene flow in direct developing marine species across a range of spatial scales and biogeographic contexts. This will help elucidate the complex relationships between life history, biogeography, and dispersal.

Population Genetics in *Isopoda*

Marine isopods offer a tractable model system for investigating the dispersal potential and population structure of direct-developing species. Many isopod species are abundant and easily sampled in intertidal zones across extensive geographic ranges. Nevertheless, the forces acting to maintain population genetic structure in marine isopods are not well understood. Some species exhibit genetic structure over small spatial scales, on the order of tens of kilometres or less (e.g *Idotea chelipes* (Jolly & Rogers, 2003), *Austridotea lacustris* (McGaughran et al., 2006), and *Jaera albifrons* (Piertney & Carvalho, 1994)). This is congruent with the hypothesis of reduced dispersal in direct developers, and may be

responsible for the widespread occurrence of multiple cryptic species of isopods (*Ligia* and

Tylos spp.) on the Southern Californian coastline (Hurtado et al., 2010, 2013; Markow &

Pfeiler, 2010).

In contrast, other species of isopods exhibit transoceanic distributions, indicative of

historically high dispersal rates. One example is *Sphaeroma terebrans*, which is distributed

across both the Atlantic and Indian Oceans. The apparently high dispersal rate and wide

distribution of this wood-boring isopod may be result of its reliance on "rafting" for dispersal

(a process in which individuals use large rafts of algae or other floating debris to disperse)

(Baratti et al., 2011). The contrast between *S. terebrans* and *Ligia* and *Tylos spp.* suggests

that life history (i.e. whether species are direct developers or not) may not always be a good

predictor of population structure in isopods.

Here, we investigate the New Zealand-wide population genetic structure of the intertidal

isopod *Isocladus armatus*. Endemic to New Zealand, this massively colour polymorphic

species is found in abundance on semi-sheltered rocky shorelines throughout the country

(Jansen, 1971). A previous study concluded that this direct-developing isopod species can

exhibit high dispersal rates because no significant population divergence was found between

two sites separated by 11km of coastline (Wells & Dale, 2018). In contrast, strong population

genetic structure was evident on a much larger scale (1,000kms). Therefore, it is unclear at

precisely what spatial scale this population structure begins to break down, and how gene

flow is affected by factors such as distance or the presence of biogeographic barriers.

In this study, we use genotyping-by-sequencing (GBS) to resolve population genomic

structure over a range of spatial scales. By quantifying population structure in this species,

we can identify the influence of geography (both distance and seascape features) on gene

flow and dispersal.

Methods

Sample Collection

We collected specimens of *Isocladus armatus* in June, 2018, from around the North Island,

New Zealand, from locations where *I. armatus* had previously been recorded (Hurley &

Jansen, 1977; *INaturalist.org*, n.d.). These sites were at Stanmore Bay, Browns Bay, Opito

Bay (Coromandel), Mt Maunganui, Māhia Peninsula, and Wellington (Fig. 1). At each site we

collected a minimum of 32 individuals, up to a maximum of 48, and attempted to sample

individuals as equally as possible amongst colour morph and sex. Where possible, we

collected specimens larger than 5mm in order to ensure enough DNA could be extracted for

sequencing (see Supplementary Material for details). The maximum distance between

individuals collected at any site did not exceed 30 m. Samples were stored at -80°C in 100%

ethanol until extraction. We have included previously collected samples from 2015 (see
(Wells & Dale, 2018) for sampling methods) to increase the number of sampled individuals
and the range of location. In addition, the temporal structure of this sampling may yield
insight into the short-term changes in allele frequencies within a population.

DNA Extraction

108   We extracted DNA following a modified Qiagen DNEasy Blood and Tissue protocol from
109   Wells and Dale (2018). Briefly, we used 178 µl of 0.5M EDTA and 22 µl of 20% SDS in each
110   extraction (rather than varying volumes by weight of tissue). Additionally, we eluted DNA
111   from the spin column three times, using 50 µl of nuclease-free water for each. We let the
112   eluent sit on the column for 15 minutes before centrifugation for one minute at 7,000 rcf.

Data Collection and Processing

DNA samples were processed by Diversity Arrays Technology (DArT) Ltd (Canberra,
Australia) using DArTseq, a genotyping-by-sequencing (GBS) approach. The methodology
behind this, including restriction enzyme choice are described in detail in Wells and Dale
(2018). DArT performs SNP calling using a proprietary pipeline. SNPs are only called if both
homozygous and heterozygous genotypes can be identified.

We analysed the dataset provided by diversityArrays together with the data from Wells and
Dale (2018). To ensure the datasets were compatible, we filtered each dataset separately
based on the conditions described below using the R packages dartR (Gruber et al., 2018)
and radiator (Gosselin, 2019). We then used only the loci shared across both datasets for
the remainder of the analyses.

We required SNPs to have a call rate ≥ 0.9 (where for at least 90% of individuals a genotype
was identified), a minor allele count of at least 3, observed heterozygosity > 0.5, minimum
depth of > 5X and maximum depth of 50X (excessively high coverage may suggest
duplicated genome elements which could confound further analyses). If we found multiple
SNPs on the same read, we removed the SNP with lowest replicability (based on the
number of technical replicates that resulted in the same allele being called). Finally, where
appropriate for further analysis, we removed SNPs outside of Hardy-Weinberg Equilibrium or
that we inferred as being under selection. Specific details of why these filters were
implemented and how are outlined in the Supplementary Material. One individual from
Browns Bay was excluded from all analyses, as this sample had a very high number of
SNPs missing across all loci (93%).

Figure 1. Map of sampling localities within New Zealand (coloured dots). The prevailing ocean currents and biogeographic breaks proposed by Shears et al (2008) are also indicated. Black dashed lines and unabbreviated labels indicate bioprovinces. Grey arrows indicate currents, abbreviated as: EAC (East Auckland Current), ECE (East Cape Eddy), ECC (East Cape Current), WE (Wairarapa Eddy), and SC (Southland Current).

Data Analysis

We calculated F-statistics using StAMPP (Pembleton et al., 2013). We used F_{st} as the primary measure of genetic differentiation because F_{st} remains relatively robust for biallelic markers such as SNPs (Whitlock, 2011).

We conducted principal component analyses (PCA) using the R package adegenet (Jombart & Ahmed, 2011). In order to understand the correspondence between the principal components and geography, we performed a Procrustes transformation of the first two principal components using MCMCpack in R (Martin et al., 2011). Procrustes transformations scale, stretch, and rotate the PCA in order to minimize the differences between two matrices (in this case, the difference between principal components and geographic coordinates).

We implemented clustering analyses using STRUCTURE 2.3.4 (Falush et al., 2003) performed with all populations (including the repeated samples of Stanmore Bay). For these

analyses, an admixture model was assumed. We ran the Markov Chain Monte Carlo
simulations with 100,000 iterations and a burn-in of 50,000. We conducted ten replicates of
each run, and varied K from 2 through 9. Final population inference was performed by
consolidating the results for each level of K in CLUMPP (Jakobsson & Rosenberg, 2007).
Additionally, we performed a separate STRUCTURE analysis on the Auckland populations
with the implementation of the locprior model at a K of 3, in order to test for fine-grain
population structure within Auckland. Due to concerns regarding the inferences made when
defining K, we chose to present a range of realistic values for K (Lawson et al., 2018;
Pritchard et al., 2010; Verity & Nichols, 2016).

We tested for Isolation-by-Distance by conducting a Mantel test using the R package vegan
(Oksanen et al., 2010). We used Slatkin's linearized F_{st} matrix (transformed using $1/1-F_{st}$
(Rousset, 1997)) and an overwater distance matrix for this test. We calculated overwater
distance using the marmap (Pante & Simon-Bouhet, 2013) and fossil (Vavrek, 2011) R
packages, finding the minimum distance between populations around the coast within a
depth range of 150 m.

Results

To examine population structure in *I. armatus* populations, we isolated 261 individuals
distributed across 8 populations across New Zealand (**Fig. 1**). We obtained DArTseq SNP
data for these 261 individuals, which identified 78,927 SNP loci as being polymorphic. After
stringent filtering (see **Methods** and **Supplementary Material**), 8,020 SNPs remained
(**Supplementary Table 2**).

We first used this filtered SNP data to test for population structure using F statistics (see
Methods). We found evidence of genetic structure (likely due to decreased rates of genetic
exchange) between populations that were more than 18 km apart (Browns Bay and
Stanmore Bay), with F_{st} values ranging from 0.06 to 0.25 (**Table 1**).

Table 1. Population F_{st} values. There is evidence of weak population structure within some of the Auckland populations (Hatfields Beach and Stanmore Bay with Browns Bay). However, population structure between all other populations is strong, with a maximal F_{st} of 0.25 between Kaikōura and the Māhia Peninsula. Unless otherwise indicated, all p-values are less than 0.0001.

	Hatfields Beach 2015	Stanmore Bay 2015	Stanmore Bay 2018	Browns Bay 2018	Opito Bay 2018	Mt Maunganui 2018	Māhia Peninsula 2018	Wellington 2018
Stanmore Bay 2015	0 (p=0.65)							
Stanmore Bay 2018	0.001 (p=0.02)	0 (p=0.79)						
Browns Bay 2018	0.002 (p=0.002)	0.002	0.002					
Opito Bay 2018	0.035	0.034	0.034	0.035				
Mt Maunganui 2018	0.067	0.069	0.067	0.068	0.032			
Māhia Peninsula 2018	0.092	0.090	0.091	0.091	0.070	0.063		
Wellington 2018	0.17	0.17	0.24	0.17	0.19	0.22	0.18	
Kaikōura 2018	0.18	0.18	0.19	0.18	0.20	0.23	0.25	0.12

We next quantified population structure using principal component analysis, which identifies the combinations of SNP loci that vary the most between individuals. We found that PC1 accounted for 19.5%, PC2 for 3.65%, and PC3 for 2.11% of the variance in SNP allele frequency (**Figs. 2A** and **2B**). PC1 primarily delineated the southern Wellington and Kaikōura populations from the other populations. PC2 primarily differentiated between the northern populations, and revealed three potential cases of migration between Mt Maunganui and the Māhia Peninsula (**Fig. 2A**). Finally, PC3 differentiated the southern populations, Wellington and Kaikoura. All Auckland populations clustered together, suggesting these individuals form one large population. We found no difference between the temporal samples from Stanmore Bay, indicating that allele frequency variation over short timescales is relatively stable.

Figure 2. Principal Component Analysis (PCA) indicates strong location-dependent population structure. PC1 (19.5% of the variance) largely differentiates the individuals in southern populations from those in the northern populations (i.e. Wellington and Kaikōura from the rest). PC2 (3.65% of the variance) differentiates between the northern populations. Finally, PC3 differentiates the populations within the southern group. Insets show the Auckland populations, and indicate minimal evidence for structure within them. Three potential recent migrant individuals are apparent, two from the Māhia population (dark green

204 points) to the Mt Maunganui population (pink points), and one from the Mt Maunganui
205 population to the Māhia population, visible in panel (B) within the dark green points.
206

A Procrustes transformation of the first two principal components of the PCA onto
geographical space revealed that the spacing of each genetic cluster was strikingly
concordant with the locations in geographical space, with the exception of Māhia and
Wellington (**Fig. 3**).

Figure 1. Procrustes transformation of PC1 vs PC2 onto geographical coordinates.
The transformation indicates a strong correspondence between sampling location and
genotype distance in principal component space. The arrows point to the location where a
population is actually found, while clusters indicate their location in Procrustes-transformed
space. The bottom arrow indicates the location of the Wellington population (purple) only.

As a complementary approach to the PCA in quantifying population structure, we
implemented a STRUCTURE analysis. This method implements a model-based clustering
approach, which probabilistically assigns individuals to one or more populations under an
admixture model. The admixture model assigns different proportions of an individual's
genome belonging to different ancestral populations (Pritchard et al., 2010). The
STRUCTURE analyses showed similar results to the PCA, with all individuals from the
Auckland locations consistently clustering together, while Kaikōura and Wellington also
always clustered together across all ranges of K (**Fig. 4**). Again we observed no fine-grain

structure within Auckland when sampling location was incorporated into the analysis

(through use of the locprior model) (**Suppl. Fig. 3**).

As we increased K, additional population genetic structure became apparent. When K=3,

Māhia individuals formed a distinct cluster exhibiting high admixture with both Mt. Maunganui

and Opito Bay. Increasing K further suggested genetic admixture between all adjacent

populations with the exception of Wellington and Māhia. With K=4, there was clear evidence

of two Māhia individuals with admixture profiles more similar to Mt. Maunganui individuals,

and one Mt. Maunganui individual with a profile similar to Māhia individuals (**Fig. 4**, bottom

two individuals in the Māhia sample and top individual in the Maunganui sample,

respectively). These correspond to the potential migrants identified in the PCA above.

Unexpectedly, with K=5 and K=6, there was evidence of admixture between the

northernmost populations (Auckland and Opito Bay) and the southernmost (Wellington and

Kaikōura). However, this may be an artefact of overfitting (Evanno et al., 2005).

Figure 4. Admixture plots based on Bayesian clustering analyses generated in
STRUCTURE. Each horizontal coloured line represents an individual from the locality
sampled (indicated on the left). Horizontal black lines designate the junction between
population samples. Results are based on 10 replicate runs for each K. At K=4 there are two
individuals from Māhia with Mt. Maunganui-like admixture profiles (bottom two bars in the
Māhia samples), and one individual from Mt. Maunganui with a Māhia-like admixture profile
(top bar in the Mt. Maunganui samples).

For all the analyses above (F_{st}, PCA, and STRUCTURE), there was clear evidence of

population structure. In addition, the Procrustes transformation showed high correspondence

between PCA distance and geographic distance (i.e. isolation-by-distance, IBD). In order to

further confirm this, we used a Mantel test. This test showed a significant positive correlation

between Slatkin's linearized F_{st} (used as a measure of population structure) and geographic

overwater distance (r = 0.83, P = 0.001; **Fig. 5**), again supporting the hypothesis of IBD. This

strong positive relationship remained even after excluding the southernmost populations

(Kaikōura and Wellington) (r = 0.92, P = 0.001). A molecular analysis of variance also

confirmed the existence of genetic structure between the southern populations (Wellington

and Kaikōura) and the northern populations, with the majority of between-population

variance being explained by this split (see **Supplementary Material**).

Figure 5. The positive correlation between overwater distance and Slatkin's linearized
F_{st} suggests a pattern of population genetic structure due to isolation-by-distance.
The black line indicates the slope of relationship for just the contrasts between the northern
populations (points in the lower left quadrant).

Discussion

Here we have quantified population genetic structure in a direct-developing marine

invertebrate across a wide range of spatial scales. By employing a GBS approach we aimed

to increase our power to detect even low levels of population structure.

We found minimal evidence of population structure of *Isocladus armatus* within the Auckland
region. Only the F_{st} analysis suggested that there is any population differentiation, between
Browns Bay and the other Auckland populations. These sites are separated by
approximately 20km, which may be the smallest distance at which population genetic
structure starts to occur in this species. Further resolution of this would require more fine-
scaled spatial sampling in this region. These results, indicating low levels of population
structure, are consistent with previous work in this species (Wells & Dale, 2018), as well as
some biphasically-developing marine invertebrates in the Hauraki Gulf surrounding
Auckland, such as the native New Zealand sea urchin *Evechinus chloroticus* (Nagel et al.,
2015) and the invasive tunicate *Styela clava* (Goldstien et al., 2010).

Over larger spatial scales, *I. armatus* exhibited strong patterns of isolation by distance (IBD).
The Procrustes-transformed PCA indicated that the first two principal components were
highly concordant with the geographic arrangement of populations. To test the hypothesis of
IBD more explicitly, we used a Mantel test, which indicated a strong and almost linear
correlation between geographic distance and genetic distance, indicative of a stepping stone
model of distribution. To try and estimate specific migration rates, we also performed a
BayesAss3 analysis (Mussman et al., 2019). However, the parameter estimates from this
analysis were internally inconsistent and, in some cases, had unbounded confidence
intervals (see **Supplementary Material**). This suggested that additional sampling would be
required to accurately estimate these parameters.

We found evidence for unusually high admixture between Mt. Maunganui and Māhia
Peninsula, despite their separation by a well-known biogeographic barrier, the East Cape.
The strongest evidence for this is the presence of three individuals that appeared to be
recent migrants between these populations. In one of these an individual from Māhia
population clustered with the Mt Maunganui in the PCA, while the opposite was the case for
the second individual. The third individual was found located between the associated PC
clusters. These three individuals also clustered with the other population, or were highly
admixed, in the STRUCTURE analysis, supporting the migrant hypothesis. The individual
with an intermediate genotype, as indicated by both the PCA and STRUCTURE analyses,
argues against these results being due to sample contamination.

The biogeographic break at the East Cape has been shown to affect population structure in
a range of species. This includes direct developers such as the anemone *Actinia tenebrosa*
and two species of amphipods (Stevens & Hogg, 2004; Veale & Lavery, 2012). Even
population structure in biphasic species with larval stages, such as the paua *Haliotis iris* (Will
et al., 2011), and the marine gastropod *Buccinulum vittatum* (Gemmell et al., 2018), are
affected by the East Cape. However, in contrast to these other direct developers, this break

does not appear to strongly affect *Isocladus armatus*, as F_{st} between Mt. Maunganui and

Māhia was no greater than F_{st} between Mt. Maunganui and the Auckland area. In addition,

the PCA indicated that the Mt. Maunganui population was almost exactly intermediate in

genetic similarity between the Māhia and Opito Bay populations, rather than being more

closely associated with the Opito Bay population, given their greater proximity. Finally, the

STRUCTURE analyses also showed admixture between Māhia and Mt. Maunganui across

all values of K.

Instead of the expected north-south genetic break at the East Cape, we found a strong

north-south break between Māhia and Wellington. This north-south break is congruent with

the placement of a proposed biogeographic region border in this area (Shears et al., 2008)

(**Fig. 1**). However, Shears et al. (2008) also proposed a biogeographic region border at East

Cape, which is inconsistent with our study. Indeed, our study contrasts with two previously

proposed hypotheses of strong genetic breaks associated with East Cape and the Cook

Strait. Instead, we observe high gene flow over the East Cape, and genetic clustering of

Wellington and Kaikōura populations.

One explanation for the strong genetic break may be the presence of cryptic species within *I.*

armatus, which would drive gene flow to near-zero levels. Within New Zealand, strong north-

south divergence between populations of the brooding brittle star, *Amphipolis squamata*, has

been suggested to be the result of cryptic speciation (Sponer & Roy, 2002). Cryptic species

have been frequently observed in isopods (Hurtado et al., 2016; Leese et al., 2008; Markow

& Pfeiler, 2010), and the degree of genetic divergence between the northern and southern

group in *I. armatus* is similar to that found between other cryptic species of isopods based on

mitochondrial DNA (Leese et al., 2008). The potential for cryptic species is further supported

by the observation of an individual from Browns Bay (which was excluded from all analyses)

that, despite appearing morphologically similar to *I. armatus*, lacked 93% of SNPs that were

present in other samples. While missing data is a common feature of reduced representation

datasets, excessively high missingness in GBS data has been associated with divisions

between species rather than populations (Tripp et al., 2017).

Conclusion

Isocladus armatus exhibits a surprisingly high amount of gene flow across small spatial

scales. However at distances greater than 20 km the level of population structure is

consistent with the expectation of reduced dispersal in direct developing species, and the

presence of IBD. Interestingly, the strongest genetic break we observed was between the

Māhia Peninsula and Wellington, with populations forming a clear northern and southern

grouping either side of this break. This was unexpected, as other well-known biogeographic
barriers – the East Cape and the Cook Strait – appeared to have little effect on population
genetic structure. Additional fine-scale sampling across this genetic break would help
determine whether differentiation across this break is a result of the existence of cryptic
species being present, or the result of a geophysical barrier that prevents dispersal in this
region.

References

Ayre, D. J., Minchinton, T. E., & Perrin, C. (2009). Does life history predict past and current

connectivity for rocky intertidal invertebrates across a marine biogeographic barrier?

Mol. Ecol., *18*(9), 1887–1903.

Baratti, M., Filippelli, M., & Messana, G. (2011). Complex genetic patterns in the mangrove

wood-borer Sphaeroma terebrans Bate, 1866 (Isopoda, Crustacea, Sphaeromatidae)

generated by shoreline topography and rafting dispersal. *J. Exp. Mar. Bio. Ecol.*,

398(1–2), 73–82.

Cowen, R. K., & Sponaugle, S. (2009). Larval Dispersal and Marine Population Connectivity.

Annual Review of Marine Science, *1*(1), 443–466.

https://doi.org/10.1146/annurev.marine.010908.163757

Dray, S., Dufour, A.-B., & Others. (2007). The ade4 package: Implementing the duality

diagram for ecologists. *J. Stat. Softw.*, *22*(4), 1–20.

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of

individuals using the software STRUCTURE: a simulation study. *Mol. Ecol.*, *14*(8),

2611–2620.

Excoffier, L., Smouse, P. E., & Quattro, J. M. (1992). Analysis of molecular variance inferred

from metric distances among DNA haplotypes: Application to human mitochondrial

DNA restriction data. *Genetics*, *131*(2), 479–491.

Falush, D., Stephens, M., & Pritchard, J. K. (2003). Inference of population structure using

multilocus genotype data: Linked loci and correlated allele frequencies. *Genetics*,

164(4), 1567–1587.

Foll, M., & Gaggiotti, O. (2008). A genome-scan method to identify selected loci appropriate

for both dominant and codominant markers: A Bayesian perspective. *Genetics*,

180(2), 977–993.

Gemmell, M. R., Trewick, S. A., Crampton, J. S., Vaux, F., Hills, S. F. K., Daly, E. E.,

Marshall, B. A., Beu, A. G., & Morgan-Richards, M. (2018). Genetic structure and

shell shape variation within a rocky shore whelk suggest both diverging and

constraining selection with gene flow. *Biol. J. Linn. Soc. Lond.*, *125*(4), 827–843.

Goldstien, S. J., Schiel, D. R., & Gemmell, N. J. (2010). Regional connectivity and coastal

expansion: Differentiating pre-border and post-border vectors for the invasive

tunicate Styela clava. *Mol. Ecol.*, *19*(5), 874–885.

González-Wevar, C. A., Segovia, N. I., Rosenfeld, S., Ojeda, J., Hüne, M., Naretto, J.,

Saucède, T., Brickle, P., Morley, S., Féral, J.-P., Spencer, H. G., & Poulin, E. (2018).

Unexpected absence of island endemics: Long-distance dispersal in higher latitude

sub-Antarctic Siphonaria (Gastropoda: Euthyneura) species. *Journal of*

Biogeography, *45*(4), 874–884. https://doi.org/10.1111/jbi.13174

Gosselin, T. (2019). *thierrygosselin/radiator: Official release*.

Gruber, B., Unmack, P. J., Berry, O. F., & Georges, A. (2018). dartr: An r package to

facilitate analysis of SNP data generated from reduced representation genome

sequencing. *Mol. Ecol. Resour.*, *18*(3), 691–699.

Hurley, D. E., & Jansen, K. P. (1977). The marine fauna of New Zealand: Family

Sphaeromatidae (Crustacea Isopoda: Flabellifera). *N.Z. Oceanogr. Inst. Mem.*, *63*,

1–80.

Hurtado, L. A., Lee, E. J., & Mateos, M. (2013). Contrasting Phylogeography of Sandy vs.

Rocky Supralittoral Isopods in the Megadiverse and Geologically Dynamic Gulf of

California and Adjacent Areas. *PLOS ONE*, *8*(7), e67827.

https://doi.org/10.1371/journal.pone.0067827

Hurtado, L. A., Mateos, M., Mattos, G., Liu, S., Haye, P. A., & Paiva, P. C. (2016). Multiple

transisthmian divergences, extensive cryptic diversity, occasional long-distance

395   dispersal, and biogeographic patterns in a marine coastal isopod with an amphi-

396   American distribution. *Ecology and Evolution*, *6*(21), 7794–7808.

397   https://doi.org/10.1002/ece3.2397

398 Hurtado, L. A., Mateos, M., & Santamaria, C. A. (2010). Phylogeography of supralittoral

399   rocky intertidal Ligia isopods in the Pacific region from Central California to Central

400   Mexico. *PLoS One*, *5*(7).

401 *INaturalist.org*. (n.d.). https://www.inaturalist.org/

402 Jakobsson, M., & Rosenberg, N. A. (2007). CLUMPP: a cluster matching and permutation

403   program for dealing with label switching and multimodality in analysis of population

404   structure. *Bioinformatics*, *23*(14), 1801–1806.

405   https://doi.org/10.1093/bioinformatics/btm233

406 Jansen, K. P. (1971). Ecological studies on intertidal New Zealand Sphaeromatidae

407   (Isopoda: Flabellifera). *Mar. Biol.*, *11*(3), 262–285.

408 Jolly, M. T., & Rogers, A. D. (2003). *Microgeographic generic variation of populations of*

409   *Idotea chelipes (Crustacea: Isopoda) in lagoons of the southern English coast*

410   *Microgeographic genetic variation of populations of Idotea chelipes (Crustacea:*

411   *Isopoda) in lagoons of the southern Eng. May 2014*.

412 Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: New tools for the analysis of genome-wide

413   SNP data. *Bioinformatics*, *27*(21), 3070–3071.

414 Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: An R package for genetic

415   analysis of populations with clonal, partially clonal, and/or sexual reproduction.

416   *PeerJ*, *2*, e281. https://doi.org/10.7717/peerj.281

417 Kimura, M., & Weiss, G. H. (1964). The Stepping Stone Model of Population Structure and

418   the Decrease of Genetic Correlation with Distance. *Genetics*, *49*(4), 561–576.

419 Lawson, D. J., van Dorp, L., & Falush, D. (2018). A tutorial on how not to over-interpret

420   STRUCTURE and ADMIXTURE bar plots. *Nature Communications*, *9*(1), 1–11.

421   https://doi.org/10.1038/s41467-018-05257-7

Leese, F., Kop, A., Wägele, J. W., & Held, C. (2008). Cryptic speciation in a benthic isopod

from Patagonian and Falkland Island waters and the impact of glaciations on its

population structure. *Front. Zool.*, *5*, 1–15.

Luikart, G., England, P. R., Tallmon, D., Jordan, S., & Taberlet, P. (2003). The power and

promise of population genomics: From genotyping to genome typing. *Nat. Rev.*

Genet., *4*(12), 981–994.

Markow, T. A., & Pfeiler, E. (2010). Mitochondrial DNA evidence for deep genetic

divergences in allopatric populations of the rocky intertidal isopod Ligia occidentalis

from the eastern Pacific. *Molecular Phylogenetics and Evolution*, *56*(1), 468–473.

https://doi.org/10.1016/j.ympev.2009.12.002

Martin, A., Quinn, K., & Park, J. H. (2011). MCMCpack: Markov Chain Monte Carlo in R.

Journal of Statistical Software, Articles, *42*(9), 1–21.

McGaughran, A., Hogg, I. D., Stevens, M. I., Lindsay Chadderton, W., & Winterbourn, M. J.

(2006). Genetic divergence of three freshwater isopod species from southern New

Zealand. *J. Biogeogr.*, *33*(1), 23–30.

McMillan, W. O., Raff, R. A., & Palumbi, S. R. (1992). Population Genetic Consequences of

Developmental Evolution in Sea Urchins (genus Heliocidaris). *Evolution*, *46*(5),

1299–1312. https://doi.org/10.1111/j.1558-5646.1992.tb01125.x

Morin, P. A., Leduc, R. G., Archer, F. I., Martien, K. K., Huebinger, R., Bickham, J. W., &

Taylor, B. L. (2009). Significant deviations from Hardy-Weinberg equilibrium caused

by low levels of microsatellite genotyping errors. *Mol. Ecol. Resour.*, *9*(2), 498–504.

https://doi.org/10.1111/j.1755-0998.2008.02502.x

Mussmann, S. M., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2019). BA3-SNPs:

Contemporary migration reconfigured in BayesAss for next-generation sequence

data. *Methods Ecol. Evol.*, *202*, 389.

Nagel, M. M., Sewell, M. A., & Lavery, S. D. (2015). Differences in population connectivity of

a benthic marine invertebrate Evechinus chloroticus (Echinodermata: Echinoidea)

across large and small spatial scales. *Conserv. Genet.*, *16*(4), 965–978.

https://doi.org/10.1007/s10592-015-0716-2

Pante, E., & Simon-Bouhet, B. (2013). marmap: A package for importing, plotting and

analyzing bathymetric and topographic data in R. *PLoS One*, *8*(9), e73051.

Pelc, R. A., Warner, R. R., & Gaines, S. D. (2009). Geographical patterns of genetic

structure in marine species with contrasting life histories. *Journal of Biogeography*,

36(10), 1881–1890. https://doi.org/10.1111/j.1365-2699.2009.02138.x

Pembleton, L. W., Cogan, N. O. I., & Forster, J. W. (2013). StAMPP: an R package for

calculation of genetic differentiation and structure of mixed-ploidy level populations.

Mol. Ecol. Resour., *13*(5), 946–952.

Piertney, S. B., & Carvalho, G. R. (1994). Microgeographic genetic differentiation in the

intertidal isopod Jaera albifrons Leach. I. Spatial distribution of allozyme variation.

Proc. R. Soc. Lond. B Biol. Sci.

Pritchard, J. K., Wen, W., & Falush, D. (2010). *Documentation for STRUCTURE software:*

Version 2.3.

http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.323.9675&rep=rep1&type=

pdf

Puritz, J. B., Keever, C. C., Addison, J. A., Barbosa, S. S., Byrne, M., Hart, M. W., Grosberg,

R. K., & Toonen, R. J. (2017). Life-history predicts past and present population

connectivity in two sympatric sea stars. *Ecol. Evol.*, *7*(11), 3916–3930.

Rousset, F. (1997). Genetic differentiation and estimation of gene flow from F-statistics

under isolation by distance. *Genetics*, *145*(4), 1219–1228.

Shanks, A. L. (2009). Pelagic larval duration and dispersal distance revisited. *The Biological*

Bulletin, *216*(3), 373–385. https://doi.org/10.1086/BBLv216n3p373

Shears, N. T., Smith, F., Babcock, R. C., Duffy, C. A. J., & Villouta, E. (2008). Evaluation of

biogeographic classification schemes for conservation planning: Application to New

Zealand's coastal marine environment. *Conserv. Biol.*, *22*(2), 467–481.

Simpson, S. D., Harrison, H. B., Claereboudt, M. R., & Planes, S. (2014). Long-Distance

Dispersal via Ocean Currents Connects Omani Clownfish Populations throughout

Entire Species Range. *PLOS ONE*, *9*(9), e107610.

https://doi.org/10.1371/journal.pone.0107610

Sponer, R., & Roy, M. S. (2002). Phylogeographic analysis of the brooding brittle star

Amphipholis squamata (Echinodermata) along the coast of New Zealand reveals

high cryptic genetic variation. *Evolution*, *56*(10), 1954–1967.

Stevens, M. I., & Hogg, I. D. (2004). Population genetic structure of New Zealand's endemic

corophiid amphipods: Evidence for allopatric speciation. *Biol. J. Linn. Soc. Lond.*,

81(1), 119–133.

Tripp, E. A., Tsai, Y.-H. E., Zhuang, Y., & Dexter, K. G. (2017). RADseq dataset with 90%

missing data fully resolves recent radiation of Petalidium (Acanthaceae) in the ultra-

arid deserts of Namibia. *Ecol. Evol.*, *7*(19), 7920–7936.

Van Wyngaarden, M., Snelgrove, P. V. R., DiBacco, C., Hamilton, L. C., Rodríguez-

Ezpeleta, N., Jeffery, N. W., Stanley, R. R. E., & Bradbury, I. R. (2017). Identifying

patterns of dispersal, connectivity and selection in the sea scallop, Placopecten

magellanicus, using RADseq-derived SNPs. *Evol. Appl.*, *10*(1), 102–117.

Vavrek, M. J. (2011). Fossil: Palaeoecological and palaeogeographical analysis tools.

Palaeontol. Electronica, *14*(1), 16.

Veale, A. J., & Lavery, S. D. (2012). The population genetic structure of the waratah

anemone (Actinia tenebrosa) around New Zealand. *N. Z. J. Mar. Freshwater Res.*,

46(4), 523–536.

Verity, R., & Nichols, R. A. (2016). Estimating the Number of Subpopulations (K) in

Structured Populations. *Genetics*, *203*(4), 1827–1839.

Waples, R. S. (1987). A Multispecies Approach to the Analysis of Gene Flow in Marine

Shore Fishes. *Evolution*, *41*(2), 385–400. JSTOR. https://doi.org/10.2307/2409146

Waples, R. S. (2015). Testing for Hardy-Weinberg proportions: Have we lost the plot? *J.*

Hered., *106*(1), 1–19. https://doi.org/10.1093/jhered/esu062

Wells, S. J., & Dale, J. (2018). Contrasting gene flow at different spatial scales revealed by

genotyping-by-sequencing in Isocladus armatus, a massively colour polymorphic

New Zealand marine isopod. *PeerJ*, *6*, e5462. https://doi.org/10.7717/peerj.5462

Whitlock, M. C. (2011). And D do not replace FST. *Mol. Ecol.*, *20*(6), 1083–1091.

Will, M., Hale, M. L., Schiel, D. R., & Gemmell, N. J. (2011). Low to moderate levels of

genetic differentiation detected across the distribution of the New Zealand abalone,

Haliotis iris. *Mar. Biol.*, *158*(6), 1417–1429.

Wilson, G. A., & Rannala, B. (2003). Bayesian inference of recent migration rates using

multilocus genotypes. *Genetics*, *163*(3), 1177–1191.

Winston, J. E. (2012). Dispersal in marine organisms without a pelagic larval phase.

Integrative and Comparative Biology, *52*(4), 447–457.

https://doi.org/10.1093/icb/ics040

Yoshino, H., Yamaji, F., & Ohsawa, T. A. (2018). Genetic structure and dispersal patterns in

Limnoria nagatai (Limnoriidae, Isopoda) dwelling in non-buoyant kelps, Eisenia

bicyclis and E. arborea, in Japan. *PLOS ONE, 13*(6), e0198451.

https://doi.org/10.1371/journal.pone.0198451

Supplementary Material

Supplementary Methods
Sample Collection
Isocladus armatus exhibits a wide range of colour morphs, which we refer to as striped,

variegated, white, unpatterned, or other (Wells and Dale 2018). During the collection of

individuals to quantify population structure, we collected at least 32 individuals from each

location (**Supplementary Table 1**). For cases in which more than 32 samples were

collected, we selected samples for DNA extraction and subsequent sequencing based on

the size and degree of colour morphotype (less ambiguous morphotypes were chosen

preferentially over the more ambiguous ones, **Supplementary Table 1**).

Supplementary Table 1. Number of samples from each location grouped by sex and colour morphotype. All sites are from 2018 except where explicitly stated.

Site	Sex	Striped	Unpatterned	Variegated	White	Other	Sum
Hatfields Beach 2015	Female	3	6	4	2	1	16
Hatfields 2015	Male	0	4	4	4	3	15
Stanmore Bay 2015	Female	4	4	4	4	0	16
Stanmore Bay 2015	Male	4	4	4	4	0	16
Stanmore Bay	Female	4	4	4	4	0	16
Stanmore Bay	Male	4	4	4	4	0	16
Browns Bay	Female	4	4	4	4	0	16
Browns Bay	Male	4	4	4	4	0	16
Māhia Peninsula	Female	0	6	4	3	0	13
Māhia Peninsula	Male	2	9	5	0	0	16
Mt Maunganui	Female	1	3	3	4	0	11
Mt Maunganui	Male	4	4	4	3	0	15
Opito Bay	Female	4	6	1	2	0	13
Opito Bay	Male	3	6	4	3	0	16
Wellington	Female	0	5	5	5	0	15
Wellington	Male	0	5	5	5	0	15
Kaikōura 2015	Female	1	7	4	3	0	15
Kaikōura 2015	Male	1	5	4	0	5	15

SNP Filtering
A SNP call rate of ≥ 0.9 was implemented. This ensures that for each locus, a SNP genotype

was identified in at least 90% of all individuals. Mean read depths of >5X and <50X were

also implemented. While high read depths are often desired in other scenarios, excessively

high coverage may suggest duplicated genome elements which could confound analyses.

On the other hand, low read depths (<5X) may reduce confidence in the accuracy of an

allele identification.

In cases where multiple SNPs were located on the same sequence read, we retained the

SNP that had the highest reproducibility (based on the proportion of technical replicates that

resulted in the same allele being called). In the event these were equal, then the SNP with

the highest amount variation across individuals was retained. Removing these SNPs

ensures that polymorphisms known to be tightly linked are removed, while removing the
SNP with the lowest variation ensures that the most informative loci are retained.

Finally, we removed loci that we inferred as being under strong selection. Non-neutral loci
are expected to exhibit different allele frequency spectra to neutral loci. Loci under strong
selection may bias inference of population demographics (Luikart et al., 2003). Loci under
selection were identified using BayeScan (Foll & Gaggiotti, 2008), with population as the
grouping factor. Loci exhibiting a q-value of ≤ 0.05 were excluded from any further analyses.

While it is common practice to filter loci based on being in Hardy-Weinberg equilibrium
(HWE) (Morin et al., 2009; Van Wyngaarden et al., 2017; Waples, 2015; Wells & Dale,
2018), this is not always the best practice. For example, STRUCTURE clusters individuals
into populations in Hardy-Weinberg equilibrium (Pritchard et al., 2010). By implementing a
filter on HWE, sites informative for population structure would be removed. The only
analyses for which a HWE filter was implemented were the F_{st} and the associated mantel
test. Loci out of HWE were first identified using the R package *pegas* with Bonferroni
correction for multiple testing, and removed using the dartR package.

Supplemental Analyses
Analysis of Molecular Variance
We conducted a hierarchical Analysis of Molecular Variance (AMOVA) using the R packages
ade4 (Dray et al., 2007) and *poppr* (Kamvar et al., 2014). This approach partitions the
variance in genetic data among and within each level of a predefined hierarchy (or example,
individuals, populations, or large geographically-defined groups) (Excoffier et al., 1992). We
tested the significance of the AMOVA by performing 1000 random permutations using the R
package *pegas*. We used a hierarchy (in ascending order) of sample, population, and then
by the observed north-south division found in our other analyses.

Migration Rates
We estimated migration rates between populations using BayesAss3-SNPs (Mussmann et
al., 2019; Wilson & Rannala, 2003). We used the BA3-Autotune software developed
alongside BayesAss3-SNPs to identify optimal parameters for the analysis (Mussmann et
al., 2019). We excluded the 2015 Stanmore Bay population for this analysis to avoid
duplicating a population for migration estimates.

We conducted this analysis using 1000 randomly sampled loci, and used the following
parameters from BA3-Autotune: allele-frequency mixing of 0.2125, migration mixing of
0.075, and inbreeding mixing of 0.075. We ran the analysis for 10 million iterations, with a
burn-in of 5 million iterations.

Supplementary Results

Supplementary Table 2 SNP filtering results and the amount of SNPs remaining after each filter. Datasets were first filtered separately and shared SNPs between the datasets were retained for the combined dataset. This was performed to minimise batch effects as a result of samples being sequenced at two different times. Additionally, the increased data output in the 2018 dataset (as a result of the inclusion of more individuals and populations) meant that almost twice as many SNPs were identified in this dataset, than in the 2015 dataset. By retaining shared SNPs the effect of this is minimized. Call rate refers to the proportion of individuals for which a genotype can be identified for the loci.

Condition	Number of SNPs remaining
Initial number	78,927
Present in Wells and Dale 2018 and the current study, filtered separately	
Call Rate > 0.8	57,348
Depth >3 and <50	54,123
Replicability > 0.9	52,800
Monomorphic Loci	51,370
Secondary loci	31,848
Shared between datasets (N=10,036 SNPs)	
Neutral loci	9,620
Call Rate > 0.9	8,334
Depth >3 and <50	8,334
Replicability > 0.95	8,334
Monomorphic loci	8,334
Secondary loci	8,334
Minor Allele Count > 3	8,037
Observed Heterozygosity < 0.5	8,020
Loci out of Hardy Weinberg	4,552

Supplementary Figure 1 Principal Component Analysis of Auckland populations, **coloured by morphotype**. No substructuring or stratifying effect of colouration is observed.

Supplementary Figure 2 - Admixture barplots generated through STRUCTURE and CLUMPP for a value of 3 for K. This analysis was done with the locprior model on just Auckland populations, and was used to identify fine grain structure.

In order to understand the levels at which genetic variation could be attributed, we performed an AMOVA. This analysis indicated some degree of population structure (**Supplementary Table 3**). The majority of genetic variation was within samples (56.2%), followed by regions (28.9%). Between population within region variance was just below 5%.

595 **Supplementary Table 3 AMOVA analysis shows population structure between**
596 **populations and regions based on genetic variation from 8,020 SNPs**. We defined
597 regions as the north group (all populations excluding Kaikōura and Wellington) and south
598 group (Kaikōura and Wellington).
599

Source of Variation	df	Percentage of Variation	p-value
Within samples	261	56.23%	<0.001
Between samples within morphotype	252	10.1%	<0.001
Between morphotype within populations	33	0.098%	0.2
Between populations within region	7	4.65%	<0.001
Between regions	1	28.91%	0.028

Estimation of Migration Rates
We tested for migration events using BayesAss3. This analysis suggested low migration

between most populations. These three groups correspond to 1) the Auckland based

populations, 2) the East Coast of North Island populations and, 3) the Southern populations,

similar to what was found in other analyses. These estimates indicated no migration in or out

of the Southern population, but some negligible migration into Wellington from Kaikōura (Fig

6 and Table 3).

Supplementary Table 4 BayesAss migration estimates (as proportion of migrant individuals in population). Values in bold are those used in the final analyses and used to create figure 6. Migration rates represent migration from the population on the top into the population on the left, i.e. the upper triangle represents northward migration, while the bottom triangle represents southward migration. Values in brackets represent standard deviations, as estimated by BayesAss. Heatmap colours are based on log10 transformed values.

	Hatfields Beach	Stanmore Bay	Browns Bay	Opito Bay	Mt Maunganui	Māhia Peninsula	Wellington	Kaikōura
Hatfields Beach	0.675 (0.008)	0.008 (0.008)	0.267 (0.021)	0.008 (0.008)	0.008 (0.008)	0.008 (0.008)	0.008 (0.008)	0.008 (0.008)
Stanmore Bay	0.009 (0.009)	0.676 (0.009)	0.258 (0.024)	0.009 (0.009)	0.009 (0.009)	0.01 (0.009)	0.01 (0.009)	0.009 (0.009)
Browns Bay	0.008 (0.008)	0.008 (0.008)	0.917 (0.023)	0.025 (0.014)	0.008 (0.008)	0.008 (0.008)	0.008 (0.008)	0.008 (0.008)
Opito Bay	0.009 (0.008)	0.009 (0.009)	0.009 (0.009)	0.925 (0.023)	0.014 (0.011)	0.009 (0.009)	0.009 (0.009)	0.009 (0.009)
Mt Maunganui	0.01 (0.009)	0.01 (0.009)	0.01 (0.009)	0.09 (0.027)	0.833 (0.029)	0.019 (0.013)	0.009 (0.009)	0.01 (0.009)
Māhia Peninsula	0.009 (0.009)	0.009 (0.009)	0.009 (0.009)	0.01 (0.009)	0.037 (0.017)	0.898 (0.025)	0.009 (0.009)	0.009 (0.009)
Wellington	0.009 (0.009)	0.009 (0.009)	0.009 (0.009)	0.009 (0.009)	0.009 (0.009)	0.009 (0.009)	0.926 (0.023)	0.011 (0.01)
Kaikōura	0.008 (0.008)	0.008 (0.008)	0.008 (0.008)	0.008 (0.008)	0.008 (0.008)	0.009 (0.008)	0.008 (0.008)	0.934 (0.021)

Supplementary Figure 6 Chord diagram of migration estimates between each population. These estimates represent the proportion of individuals in a population that are estimated to be migrants. The largest arc for each population indicates within population movement, such as a self-recruitment, while the smaller arcs are indicative of between population movement or migration. As a result, wider arcs indicate higher migration rates. Displayed migration rates are those for which the migration rate minus the standard deviation was greater than 0.005.

High migration within the Auckland populations was observed, with estimates of migration

rates approximately 0.25, this value represents the proportion of individuals within a

population that are estimated to be migrants from the source population. Further south,

among the populations along the East Coast of the North Island, migration estimates were

consistently higher among adjacent populations than between non-neighbouring

populations, suggestive of a stepping stone model of migration (Kimura & Weiss, 1964).

These migration rates were relatively low - generally between 0.01 and 0.02 (Supplementary

Table 3).

Supplementary Figure 7. Admixture plots based on Bayesian clustering analyses generated in STRUCTURE for K 7-9. Relatively little structure is observed, and a much higher amount of noise is observed as K increases. In all instances, Kaikoura and Wellington retain a high degree of separation from other peoples, and Māhia also retains some distinctiveness.