Title: Population structure and dispersal across small and large spatial scales in a direct

developing marine isopod

Authors: William S. Pearman[a]*[†], Sarah J. Wells[b]*, Olin K. Silander[a], Nikki E. Freed[a], and

James Dale[a]

[a] School of Natural and Computational Sciences, Massey University, Auckland 0745, New

Zealand

[b] School of Environmental and Animal Sciences, Unitec, Auckland 1025, New Zealand

*Contributed equally

[†] Corresponding author: wpearman1996@gmail.com

Abstract

Marine organisms generally develop in one of two ways: biphasic, with distinct adult and

larval morphology, and direct development, in which larvae look like adults. The mode of

development is thought to significantly influence dispersal, with direct developers having

much lower dispersal potential. While dispersal and population connectivity is relatively well

understood for biphasic species, comparatively little is known about direct developers. In this

study, we use a panel of 8,020 SNPs to investigate population structure and gene flow for a

direct developing species, the New Zealand endemic marine isopod *Isocladus armatus*. We

find evidence that on a small spatial scale (20 kms), gene flow between locations is

extremely high and suggestive of an island model of migration. However, over larger spatial

scales (600km), populations exhibit a strong pattern of isolation-by-distance. The

intersection of our sampling range by two well-known biogeographic barriers (the East Cape

and the Cook Strait) provides an opportunity to understand how such barriers influence

dispersal in direct developers. We find that *I. armatus* has high migration rates across these

barriers. However, we find evidence of a north-south population genetic break occurring

between Māhia and Wellington, although there are no obvious biogeographic barriers

between these locations. This study suggests that biogeographic barriers may affect

migration in direct development species in unexpected ways.

Introduction

Life history influences genetic population structure

A species' developmental life history has a significant effect on population connectivity and dispersal. For marine organisms, developmental life histories are generally classified in one of two ways: direct developers, in which larvae resemble adults and do not disperse large distances; and biphasic, in which larvae have different morphologies from adults and generally disperse large distances via ocean currents (Cowen & Sponaugle, 2009; Puritz et al., 2017; Simpson et al., 2014) during a pelagic larval stage. For this reason, direct-developing species tend to show greater population structure than biphasic species (Ayre et al., 2009; McMillan et al., 1992; Pelc et al., 2009; Waples, 1987). However, some species do not conform to this expected pattern (Ayre et al., 2009; Puritz et al., 2017; Shanks, 2009; Winston, 2012).

Dispersal (and thus population structure) is also affected by biogeographic barriers, and the effects of these barriers themselves depend on the life history of the organism. For example, off shore ocean currents may have large effects on biphasic species, but little effect on direct developers. Similarly, we would expect that direct developers with limited dispersal potential will be doubly limited if there is no suitable habitat within the range of their average dispersal distance. However, in some cases the lack of nearby suitable habitat does not obviously limit dispersal in direct developers, and but does appear to limits the dispersal of biphasic species (Ayre et al., 2009). This is contrary to predictions based on the expected relationship between life history and dispersal potential. These cases, where direct developers exhibit greater dispersal than otherwise predicted, are not uncommon within intertidal organisms (Ayre et al., 2009; González-Wevar et al., 2018; Wells & Dale, 2018; Yoshino et al., 2018), suggesting that population connectivity cannot be entirely predicted by life history or even the knowledge of biogeographic barriers. As a result, it is important to more carefully quantify population structure in direct developing marine species across a range of spatial scales and biogeographic contexts. This will help to clarify the complex relationships between life history, biogeography, and population structure.

Population Genetics in *Isopoda*

Marine isopods offer a highly tractable and attractive system for investigating the dispersal potential and population structure of direct-developing species. Many isopod species occur in large numbers in the easily sampled intertidal zone across extensive geographic ranges. However, the forces acting to maintain population genetic structure in marine isopods are not well understood. Some species exhibit genetic structure over small spatial scales, on the order of tens of kilometres. This is congruent with the hypothesis of reduced dispersal in

direct developers, and may be responsible for the widespread occurrence of multiple cryptic

species of isopods (*Ligia* and *Tylos spp.*) on the Southern California coastline and in Baja

(Markow & Pfeiler, 2010; Hurtado et al, 2010; Hurtado et al, 2013).

In contrast, other species of isopods exhibit transoceanic distributions, indicative of

historically high dispersal rates. One example is *Sphaeroma terebrans*, which is distributed

across both the Atlantic and Indian Oceans. The apparently high dispersal rate and wide

distribution of this wood-boring isopod may be result of its reliance on "rafting" for dispersal

(a process in which individuals use large rafts of algae or other floating debris to disperse)

(Baratti et al., 2011). The contrast between *Sphaeromona* and *Ligia* and *Tylos spp.* suggests

that for isopods, life history (i.e. whether species are direct developers or not) may not

always be a good predictor of population structure, and that behavioural aspects should also

be considered.

Here we investigate the New Zealand-wide population genetic structure of the intertidal

isopod *Isocladus armatus*. Endemic to New Zealand, this highly colour polymorphic species

is found in abundance on semi-sheltered rocky shorelines throughout the country. There is

evidence that this direct-developing isopod species can exhibit high dispersal rates (Wells &

Dale, 2018). In this case, the authors found no significant population divergence between

two sites separated by 11km of coastline. This is indicative of a highly mobile species. On a

much larger scale (1,000 kms), however, population structure was evident. It is unclear at

precisely what spatial scale this population genetic structure begins to break down, or

whether biogeographic barriers affect this.

To investigate this, we use genotyping-by-sequencing (GBS) to resolve population structure

over a range of spatial scales. By quantifying population genetic structure in this species, we

are able to identify the role of geography (both distance and seascape features) in

influencing gene flow and dispersal.

Methods

Sample Collection

We collected specimens of *Isocladus armatus* in June, 2018, from around the North Island,

New Zealand, from locations where *I. armatus* has previously been recorded (Hurley &

Jansen, 1977; *INaturalist.org*, n.d.). These sites were at Stanmore Bay, Browns Bay, Opito

Bay (Coromandel), Mt Maunganui, Māhia Peninsula, and Wellington (Fig. 1). At each site we

collected a minimum of 32 individuals, up to a maximum of 48, and attempted to sample

individuals as equally as possible amongst colour morph and sex. Where possible, we

collected specimens larger than 5mm in order to ensure sufficient DNA could be extracted

for sequencing (see supplementary methods for details). The maximum distance between

individuals collected at any site did not exceed 30 m. Samples were stored at -80°C in 100%

ethanol until extraction. We have included samples from Wells and Dale 2018 (see [ref] for

sampling methods) to increase the number of sampled individuals and the range of location.

In addition, the temporal structure of this sampling may yield insight into the short term

changes in allele frequencies within a population.

DNA Extraction

We extracted DNA following a modified Qiagen DNEasy Blood and Tissue protocol from

Wells and Dale (2018). Briefly, we used 178 µl of 0.5M EDTA and 22 µl of 20% SDS in each

extraction (rather than varying volumes by weight of tissue). Additionally, we eluted DNA

from the spin column three times, using 50 µl of nuclease-free water for each. We let the

eluent sit on the column for 15 minutes before centrifugation for one minute at 7,000 rcf.

Data Collection and Processing

DNA samples were processed by Diversity Arrays Technology (DArT) Ltd (Canberra,

Australia) using a genotyping-by-sequencing (GBS) approach. The methodology behind this,

including restriction enzyme choice are described in detail in Wells and Dale (2018). DaRT

performs SNP calling using a proprietary pipeline. SNPs are only called if both homozygous

and heterozygous genotypes can be identified.

We analysed the dataset provided by diversityArrays together with the data from Wells and

Dale (2018). To ensure the datasets were compatible, we filtered each dataset separately

based on the conditions described below using the R packages dartR (Gruber et al., 2018)

and radiator (Gosselin, 2019). We then used only the loci shared across both datasets for

the remainder of the analyses.

We required SNPs have a minimum call rate ≥ 0.9, a minor allele count of at least 3,

observed heterozygosity > 0.5, minimum depth of > 5X and maximum depth of 50X. If we

found multiple SNPs on the same read, then we removed the SNP with lowest replicability

(based on the number of technical replicates that resulted in the same allele being called).

Finally, where appropriate for further analysis, we removed SNPs outside of Hardy-Weinberg

Equilibrium or that we inferred as being under selection. Specific details of why these filters

were implemented and how are outlined in the supplementary methods. One individual was

excluded from all analyses (BBFA3), as this sample had a very high number of SNPs

missing across all loci (93%).

Figure 1. Map of sampling localities within New Zealand (coloured dots). The prevailing ocean currents and biogeographic breaks proposed by Shears et al (2008) are also indicated. Black dashed lines and unabbreviated labels indicate bioprovinces. Grey arrows indicate currents, abbreviated as: EAC (East Auckland Current), ECE (East Cape Eddy), ECC (East Cape Current), WE (Wairarapa Eddy), and SC (Southland Current).

Data Analysis

We calculated F-statistics using StAMPP (Pembleton et al., 2013). We used F_{st} as the

primary measure of genetic differentiation because F_{st} remains relatively robust for biallelic

markers such as SNPs (Whitlock, 2011).

We conducted principal component analyses (PCA) using the R package adegenet (Jombart

& Ahmed, 2011). In order to understand the correspondence between the principal

components and geography, we performed a Procrustes transformation of the first two

principal components using MCMCpack in R (Martin et al., 2011). Procrustes

transformations scale, stretch, and rotate the PCA in order to minimize the differences

between two matrices (in this case, the difference between principal components and

geographic coordinates).

We implemented STRUCTURE analyses using STRUCTURE 2.3.4 (Falush et al., 2003) and

performed with all populations (including the repeated samples of Stanmore Bay). For these

analyses, we assumed an admixture model. We ran the Markov Chain Monte Carlo

simulations with 100,000 iterations and a burn-in of 50,000. We conducted ten replicates of

each run, and varied K from 2 through 9. We performed final population inference by

consolidating the results for each level of K, in CLUMPP (Jakobsson & Rosenberg, 2007).

Additionally, we performed a separate STRUCTURE analysis on the Auckland populations

with the implementation of the locprior model at a K of 3, in order to test for fine-grain

population structure within Auckland. Due to concerns regarding the inferences made when

defining K, we chose to present a range of realistic values for K (Pritchard et al., 2010; Verity

& Nichols, 2016; Lawson van Dorp, Falush 2018).

We tested for Isolation-by-Distance by conducting a Mantel test using the R package vegan

(Oksanen et al., 2010). We used Slatkin's linearized F_{st} matrix (transformed using 1/1-F_{st}

(Rousset, 1997)) and an overwater distance matrix for this test. We calculated overwater

distance using the marmap (Pante & Simon-Bouhet, 2013) and fossil (Vavrek, 2011) R

packages, finding the minimum distance between populations around the coast within a

depth range of 150 m.

Results

To examine population structure in *I. armatus* populations, we isolated 261 individuals

distributed across 8 populations across New Zealand (**Fig. 1**). We obtained DaRT SNP data

for these 261 individuals, which identified 78,927 SNP loci as being polymorphic. After

stringent filtering (see **Methods** and **Supp. Methods**), 8,020 SNPs remained

(**Supplementary Table 2**).

We first used this filtered SNP data to test for population structure using F statistics (see

Methods). We found evidence for structure (likely due to decreased rates of genetic

exchange) only between populations at least 19 km apart (Browns Bay and Stanmore Bay),

with F_{st} values ranging from 0.06 to 0.25 (**Table 1**).

Table 1. Population F_{st} values. There is evidence of weak population structure within some of the
Auckland populations (Hatfields Beach and Stanmore Bay with Browns Bay). However, population
structure between all other populations is strong, with a maximal F_{st} of 0.25 between Kaikōura and the
Māhia Peninsula. Unless otherwise indicated, all p-values are less than 0.0001.

	Hatfields Beach 2015	Stanmore Bay 2015	Stanmore Bay 2018	Browns Bay 2018	Opito Bay 2018	Mt Maunganui 2018	Māhia Peninsula 2018	Wellington 2018
Stanmore Bay 2015	0 (p=0.65)							
Stanmore Bay 2018	0.001 (p=0.02)	0 (p=0.79)						
Browns Bay 2018	0.002 (p=0.002)	0.002	0.002					
Opito Bay 2018	0.035	0.034	0.034	0.035				
Mt Maunganui 2018	0.067	0.069	0.067	0.068	0.032			
Māhia Peninsula 2018	0.092	0.090	0.091	0.091	0.070	0.063		
Wellington 2018	0.17	0.17	0.24	0.17	0.19	0.22	0.18	
Kaikōura 2018	0.18	0.18	0.19	0.18	0.20	0.23	0.25	0.12

We next quantified population structure using principal component analysis, which identifies

the combinations of SNP loci that vary the most between individuals. We found that PC1

accounted for 19.5%, PC2 for 3.65%, and PC3 for 2.11% of the variance in SNP allele

frequency (**Figs. 2A** and **2B**). PC1 primarily delineated the southern Wellington and

Kaikōura populations from the other populations. PC2 primarily differentiated between the

northern populations, and revealed three potential cases of migration between Mt

Maunganui and the Māhia Peninsula (**Fig. 2A**). Finally, PC3 differentiated the southern

populations, Wellington and Kaikoura. All Auckland populations clustered together,

suggesting these individuals form one large population. We found no difference between the

temporal samples from Stanmore Bay, indicating that allele frequency variation over short

timescales is relatively stable.

Figure 2. Principal Component Analysis (PCA) indicates strong location-dependent population structure. PC1 (19.5% of the variance) largely differentiates the individuals in southern populations from those in the northern populations (i.e. Wellington and Kaikōura from the rest). PC2 (3.65% of the variance) differentiates between the northern populations. Finally, PC3 differentiates the populations within the southern group. Insets show the Auckland populations, and indicate minimal evidence for structure within within them. Three potential recent migrant individual are apparent, two from the Māhia population (green points) to the Mt Maunganui population (pink points), and one from the Mt Maunganui population to the Māhia population, visible in panel (B) within the green points.

A Procrustes transformation of the first two components of the PCA onto geographical space
revealed that the spacing of each genetic cluster was strikingly concordant with the locations
in geographical space, with the exception of Māhia and Wellington (**Fig. 3**).

Figure 1. Procrustes transformation of PC1 vs PC2 onto geographical coordinates.
The transformation indicates a strong correspondence between sampling location and
genotype distance in principal component space. The arrows point to the location where a
population is actually found, while clusters indicate their location in Procrustes-transformed
space. The bottom arrow indicates the location of the Wellington population (purple) only.

As a complementary approach to the PCA in quantifying population structure, we
implemented a STRUCTURE analysis. This method implements a model-based clustering
approach, which probabilistically assigns individuals to one or more populations under an
admixture model. The admixture model assigns different proportions of an individual's
genome belonging to different ancestral populations (Pritchard et al., 2010). The
STRUCTURE analyses showed similar results to the PCA, with all individuals from the
Auckland locations consistently clustering together, while Kaikōura and Wellington also
always clustered together across all ranges of K (**Fig. 4**). Again we observed no fine-grain
structure within Auckland when sampling location was incorporated into the analysis
(through use of the locprior model) (**Suppl. Fig. 3**).

As we increased K, additional population genetic structure (history) became apparent. When

K=3, Māhia individuals formed a distinct cluster exhibiting high admixture with both Mt.

Maunganui and Opito Bay. Increasing K further suggested genetic admixture between all

adjacent populations with the exception of Wellington and Māhia. With K=4, there was clear

evidence of two Māhia individuals with admixture profiles more similar to Mt. Maunganui

individuals, and one Mt. Maunganui individual with a profile similar to Māhia individuals (Fig.

5, bottom two individuals in the Māhia sample and top individual in the Maunganui sample,

respectively). These correspond to the potential migrants identified in the PCA above.

Unexpectedly, with K=5 and K=6, there was evidence of admixture between the

northernmost populations (Auckland and Opito Bay) and the southernmost (Wellington and

Kaikōura). However, this may be an artefact of overfitting (Evanno et al, 2005).

Figure 4. Admixture plots based on Bayesian clustering analyses generated in
STRUCTURE. Each horizontal coloured line represents an individual from the locality
sampled (indicated on the left). Horizontal black lines designate the junction between
population samples. Results are based on 10 replicate runs for each K. At K=4 there are two
individuals from Māhia with Mt. Maunganui-like admixture profiles (bottom two bars in the
Māhia samples), and one individual from Mt. Maunganui with a Māhia-like admixture profile
(top bar in the Mt. Maunganui samples).

For all the analyses above (F_{st}, PCA, and STRUCTURE), there was clear evidence of

population structure. In addition, the Procrustes transformation showed high correspondence

between PCA distance and geographic distance (i.e. isolation-by-distance, IBD). In order to

further confirm this, we used a Mantel test. This test showed a significant positive correlation

between Slatkin's linearized F_{st} (used as a measure of population structure) and geographic

overwater distance (r = 0.83, P = 0.001; **Fig. 5**), again supporting IBD. This strong positive

relationship remained even after excluding the southernmost populations (Kaikōura and

Wellington) (r = 0.92, P = 0.001), indicating a pattern of IBD across several spatial scales.

Figure 5. The positive correlation between overwater distance and Slatkin's linearized
F_{st}. suggests a pattern of population genetic structure due to isolation-by-distance.
The black line indicates the slope of relationship for just the contrasts between the northern
populations (points in the lower left quadrant).

Discussion

Here we have quantified population structure in a direct-developing marine invertebrate

across a wide range of spatial scales. By employing a GBS approach we aimed to increase

our power to detect even low levels of population structure.

We found minimal evidence of population structure of *Isocladus armatus* within the Auckland

region. Only the F_{st} analysis suggested that there is any population differentiation, between

the Browns Bay and other Auckland populations. These are separated by approximately

20km, which may be the smallest distance at which population genetic structure starts to

occur in this species. Further resolution of this would require more fine-scaled spatial

sampling in this region. These results, indicating low levels of population structure, are

consistent with previous work in this species (Wells and Dale, 2018), as well as biphasically

developing marine invertebrates in the Hauraki Gulf (offshore of the Auckland region), such

as the native New Zealand sea urchin *Evechinus chloroticus* (Nagel et al., 2015) and the

invasive tunicate *Styela clava* (Goldstien et al., 2010).

Over larger spatial scales, *I. armatus* exhibited strong patterns of isolation by distance (IBD).

The Procrustes-transformed PCA indicated that PC1 and PC2 were highly concordant with

the geographic arrangement of populations. To test the hypothesis of IBD more explicitly, we

used a Mantel test, which indicated a strong and almost linear correlation between

geographic distance and genetic distance, indicative of a stepping stone model of

distribution. To try and estimate specific migration rates, we also performed a BayesAss3

analysis (Mussman et al., 2019). However, the parameter estimates from this analysis were

internally inconsistent and in some cases had unbounded confidence intervals (see

Supplementary Methods). This suggested that additional sampling would be required to

accurately estimate these parameters.

We found evidence for unusually high admixture between Mt. Maunganui and Māhia

Peninsula, despite their separation by a well-known biogeographic barrier, the East Cape.

Perhaps the strongest evidence for this is the presence of three individuals that appeared to

be recent migrants between these populations. In one of these an individual from Māhia

population clustered with the Mt Maunganui in the PCA, while the opposite was the case for

the second individual. The third individual was found located between the associated PC

clusters. These three individuals also clustered with the other population, or were highly

admixed, in the STRUCTURE analysis, supporting the migrant hypothesis. The individual

with an intermediate genotype, as indicated by both the PCA and STRUCTURE analyses,

argues against these results being due to sample contamination.

The biogeographic break at the East Cape has been shown to affect population structure in

a range of species. This includes direct developers such as the anemone *Actinia tenebrosa*

and two species of amphipods (Stevens & Hogg, 2004; Veale & Lavery, 2012). Even

population structure in biphasic species with larval stages, such as the paua *Haliotis iris* (Will

et al., 2011), and the marine gastropod *Buccinulum vittatum* (Gemmell et al., 2018), are

affected by the East Cape. However, in contrast to these other direct developers, this break

does not appear to strongly affect *Isocladus armatus*, as F_{st} between Mt. Maunganui and

Māhia were no greater than the F_{st} between Mt. Maunganui and the Auckland area

populations. In addition, the PCA showed the Mt. Maunganui population as almost exactly

intermediate between the Māhia and Opito Bay populations, rather than being more closely

allied with the Opito Bay population. Finally, the STRUCTURE analyses also showed

admixture between Māhia and Maunganui across all values of K.

Instead of the expected north-south genetic break at the East Cape, we found a strong

north-south break between Māhia and Wellington. This north-south break is congruent with

the placement of a proposed biogeographic region border in this area (Shears et al., 2008)

(**Fig. 1**). However, Shears et al. also proposed a biogeographic region border at East Cape,

which is inconsistent with our study. Indeed, our study contrasts with two previously
proposed hypotheses of strong genetic breaks associated with East Cape and the Cook
Strait. Instead, we observe high gene flow over the East Cape, and genetic clustering of
Wellington and Kaikōura populations. This contrasts with what has been found for the
anemone *Actinia tenebrosa*, which has low gene flow over the East Cape (Veale & Lavery,
2012).

One explanation for the strong genetic break may be the presence of cryptic species within *I.*
armatus, which would drive gene flow to near-zero levels. Within New Zealand, strong north-
south divergence between populations of the brooding brittle star, *Amphipolis squamata*, has
been suggested to be the result of cryptic speciation (Sponer & Roy, 2002). Cryptic species
have been frequently observed in isopods (Hurtado et al., 2016; Leese et al., 2008; Markow
& Pfeiler, 2010), and the degree of genetic divergence between the northern and southern
group in *I. armatus* is similar to that found between other cryptic species of isopods based on
mitochondrial DNA (Leese et al., 2008). The potential for cryptic species is further supported
by the observation of an individual from Browns Bay (which was excluded from all analyses)
which, despite appearing morphologically similar to *I. armatus*, lacked 93% of SNPs that
were present in other samples. While missing data is a common feature of reduced
representation datasets, excessively high missingness in GBS data has been associated
with divisions between species rather than populations (Tripp et al., 2017).

Conclusion

Isocladus armatus exhibits a surprisingly high amount of gene flow across small spatial
scales. However at distances greater than 20 km the level of population structure is
consistent with the expectation of reduced dispersal in direct developing species, and the
presence of IBD. Interestingly, the strongest genetic break we observed was between the
Māhia Peninsula and Wellington, with populations forming a clear northern and southern
grouping either side of this break. This was unexpected, as other well-known biogeographic
barriers – the East Cape and the Cook Strait – appeared to have little effect on population
genetic structure. Additional fine-scale sampling across this genetic break would help in
determining whether differentiation across this break is a result of the existence of cryptic
species being present, or the result of a geophysical barrier that prevents dispersal in this
region.

References

Amos, W. (2010). Even small SNP clusters are non-randomly distributed: Is this evidence of mutational non-independence? *Proc. Biol. Sci.*, *277*(1686), 1443–1449.

Arranz Martinez, V. (2017). *Connectivity among marine communities: A multi-species approach to determining the major drivers of larval connection between populations of coastal species in New Zealand* [PhD Thesis].

Ayre, D. J., Minchinton, T. E., & Perrin, C. (2009). Does life history predict past and current connectivity for rocky intertidal invertebrates across a marine biogeographic barrier? *Mol. Ecol.*, *18*(9), 1887–1903.

Baratti, M., Filippelli, M., & Messana, G. (2011). Complex genetic patterns in the mangrove wood-borer Sphaeroma terebrans Bate, 1866 (Isopoda, Crustacea, Sphaeromatidae) generated by shoreline topography and rafting dispersal. *J. Exp. Mar. Bio. Ecol.*, *398*(1–2), 73–82.

Beerli, P., & Felsenstein, J. (2001). Maximum likelihood estimation of a migration matrix and effective population sizes in n subpopulations by using a coalescent approach. *Proceedings of the National Academy of Sciences*, *98*(8), 4563–4568. https://doi.org/10.1073/pnas.081068098

Cowen, R. K., & Sponaugle, S. (2009). Larval Dispersal and Marine Population Connectivity. *Annual Review of Marine Science*, *1*(1), 443–466. https://doi.org/10.1146/annurev.marine.010908.163757

Dray, S., Dufour, A.-B., & Others. (2007). The ade4 package: Implementing the duality diagram for ecologists. *J. Stat. Softw.*, *22*(4), 1–20.

Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. A., Kawamoto, K., Buckler, E. S., & Mitchell, S. E. (2011). A Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High Diversity Species. *PLoS One*, *6*(5), e19379.

Evanno, G., Regnaut, S., & Goudet, J. (2005) Detecting the number of clusters of individuals using the software structure: a simulation study. *Molecular Ecology*, 14(8), 2611-2620.

Excoffier, L., Smouse, P. E., & Quattro, J. M. (1992). Analysis of molecular variance inferred from metric distances among DNA haplotypes: Application to human mitochondrial DNA restriction data. *Genetics*, *131*(2), 479–491.

Falush, D., Stephens, M., & Pritchard, J. K. (2003). Inference of population structure using multilocus genotype data: Linked loci and correlated allele frequencies. *Genetics*, *164*(4), 1567–1587.

Fraser, C. I., Morrison, A. K., Hogg, A. M., Macaya, E. C., van Sebille, E., Ryan, P. G., Padovan, A., Jack, C., Valdivia, N., & Waters, J. M. (2018). Antarctica's ecological isolation will be broken by storm-driven dispersal and warming. *Nat. Clim. Chang.*, *8*(8), 704–708. https://doi.org/10.1038/s41558-018-0209-7

Gemmell, M. R., Trewick, S. A., Crampton, J. S., Vaux, F., Hills, S. F. K., Daly, E. E., Marshall, B. A., Beu, A. G., & Morgan-Richards, M. (2018). Genetic structure and shell shape variation within a rocky shore whelk suggest both diverging and constraining selection with gene flow. *Biol. J. Linn. Soc. Lond.*, *125*(4), 827–843.

Goldstien, S. J., Schiel, D. R., & Gemmell, N. J. (2010). Regional connectivity and coastal expansion: Differentiating pre-border and post-border vectors for the invasive tunicate Styela clava. *Mol. Ecol.*, *19*(5), 874–885.

González-Wevar, C. A., Segovia, N. I., Rosenfeld, S., Ojeda, J., Hüne, M., Naretto, J., Saucède, T., Brickle, P., Morley, S., Féral, J.-P., Spencer, H. G., & Poulin, E. (2018). Unexpected absence of island endemics: Long-distance dispersal in higher latitude sub-Antarctic Siphonaria (Gastropoda: Euthyneura) species. *Journal of Biogeography*, *45*(4), 874–884. https://doi.org/10.1111/jbi.13174

Gosselin, T. (2019). *thierrygosselin/radiator: Official release*.

Gruber, B., Unmack, P. J., Berry, O. F., & Georges, A. (2018). dartr: An r package to facilitate analysis of SNP data generated from reduced representation genome sequencing. *Mol. Ecol. Resour.*, *18*(3), 691–699.

Hurley, D. E., & Jansen, K. P. (1977). The marine fauna of New Zealand: Family Sphaeromatidae (Crustacea Isopoda: Flabellifera). *N.Z. Oceanogr. Inst. Mem.*, *63*, 1–80.

Hurtado, L. A., Mateos, M., Mattos, G., Liu, S., Haye, P. A., & Paiva, P. C. (2016). Multiple transisthmian divergences, extensive cryptic diversity, occasional long-distance dispersal, and biogeographic patterns in a marine coastal isopod with an amphi-American distribution. *Ecology and Evolution*, *6*(21), 7794–7808. https://doi.org/10.1002/ece3.2397

INaturalist.org. (n.d.). https://www.inaturalist.org/

Jakobsson, M., & Rosenberg, N. A. (2007). CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics*, *23*(14), 1801–1806. https://doi.org/10.1093/bioinformatics/btm233

Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: New tools for the analysis of genome-wide SNP data. *Bioinformatics*, *27*(21), 3070–3071.

Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: An R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*, *2*, e281. https://doi.org/10.7717/peerj.281

Keightley, P. D., Ness, R. W., Halligan, D. L., & Haddrill, P. R. (2014). Estimation of the spontaneous mutation rate per nucleotide site in a Drosophila melanogaster full-sib family. *Genetics*, *196*(1), 313–320.

Kilian, A., Wenzl, P., Huttner, E., Carling, J., Xia, L., Caig, V., Heller-uszynska, K., Jaccoud, D., Hopper, C., Aschenbrenner-kilian, M., Evers, M., Peng, K., Cayla, C., Hok, P., & Uszynski, G. (2012). Diversity Arrays Technology: A Generic Genome Profiling Technology on Open Platforms. In *Data Production and Analysis in Population Genomics* (Vol. 888, pp. 67–89).

Kimura, M., & Weiss, G. H. (1964). The Stepping Stone Model of Population Structure and the Decrease of Genetic Correlation with Distance. *Genetics*, *49*(4), 561–576.

Leese, F., Kop, A., Wägele, J. W., & Held, C. (2008). Cryptic speciation in a benthic isopod from Patagonian and Falkland Island waters and the impact of glaciations on its population structure. *Front. Zool.*, *5*, 1–15.

Markow, T. A., & Pfeiler, E. (2010). Mitochondrial DNA evidence for deep genetic divergences in allopatric populations of the rocky intertidal isopod Ligia occidentalis from the eastern Pacific. *Molecular Phylogenetics and Evolution*, *56*(1), 468–473. https://doi.org/10.1016/j.ympev.2009.12.002

Martin, A., Quinn, K., & Park, J. H. (2011). MCMCpack: Markov Chain Monte Carlo in R. *Journal of Statistical Software, Articles*, *42*(9), 1–21.

McGaughran, A., Hogg, I. D., Stevens, M. I., Lindsay Chadderton, W., & Winterbourn, M. J. (2006). Genetic divergence of three freshwater isopod species from southern New Zealand. *J. Biogeogr.*, *33*(1), 23–30.

McMillan, W. O., Raff, R. A., & Palumbi, S. R. (1992). Population Genetic Consequences of Developmental Evolution in Sea Urchins (genus Heliocidaris). *Evolution*, *46*(5), 1299–1312. https://doi.org/10.1111/j.1558-5646.1992.tb01125.x

Mussmann, S. M., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2019). BA3-SNPs: Contemporary migration reconfigured in BayesAss for next-generation sequence data. *Methods Ecol. Evol.*, *202*, 389.

Nagel, M. M., Sewell, M. A., & Lavery, S. D. (2015). Differences in population connectivity of a benthic marine invertebrate Evechinus chloroticus (Echinodermata: Echinoidea) across large and small spatial scales. *Conserv. Genet.*, *16*(4), 965–978. https://doi.org/10.1007/s10592-015-0716-2

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., O'hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., & Wagner, H. (2010). Vegan: Community ecology package. R package version 1.17-4. *Http://Cran. r-Project. Org>. Acesso Em*, *23*, 2010.

Pante, E., & Simon-Bouhet, B. (2013). marmap: A package for importing, plotting and
analyzing bathymetric and topographic data in R. *PLoS One*, *8*(9), e73051.
Pelc, R. A., Warner, R. R., & Gaines, S. D. (2009). Geographical patterns of genetic
structure in marine species with contrasting life histories. *J. Biogeogr.*, *36*(10), 1881–
1890.
Pembleton, L. W., Cogan, N. O. I., & Forster, J. W. (2013). StAMPP: an R package for
calculation of genetic differentiation and structure of mixed-ploidy level populations.
Mol. Ecol. Resour., *13*(5), 946–952.
Pritchard, J. K., Wen, W., & Falush, D. (2010). *Documentation for STRUCTURE software:*
Version 2.3.
http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.323.9675&rep=rep1&type=
pdf
Puebla, O., Bermingham, E., & Guichard, F. (2009). Estimating dispersal from genetic
isolation by distance in a coral reef fish (Hypoplectrus puella). *Ecology*, *90*(11),
3087–3098.
Puritz, J. B., Keever, C. C., Addison, J. A., Barbosa, S. S., Byrne, M., Hart, M. W., Grosberg,
R. K., & Toonen, R. J. (2017). Life-history predicts past and present population
connectivity in two sympatric sea stars. *Ecol. Evol.*, *7*(11), 3916–3930.
Rousset, F. (1997). Genetic differentiation and estimation of gene flow from F-statistics
under isolation by distance. *Genetics*, *145*(4), 1219–1228.
Santamaria, C. A., Mateos, M., & Hurtado, L. A. (2014). Diversification at the narrow sea-
land interface in the Caribbean: Phylogeography of endemic supralittoral Ligia
isopods. *Frontiers in Ecology and Evolution*, *2*.
https://doi.org/10.3389/fevo.2014.00042
Shanks, A. L. (2009). Pelagic larval duration and dispersal distance revisited. *The Biological*
Bulletin, *216*(3), 373–385. https://doi.org/10.1086/BBLv216n3p373
Shears, N. T., Smith, F., Babcock, R. C., Duffy, C. A. J., & Villouta, E. (2008). Evaluation of
biogeographic classification schemes for conservation planning: Application to New
Zealand's coastal marine environment. *Conserv. Biol.*, *22*(2), 467–481.
Simpson, S. D., Harrison, H. B., Claereboudt, M. R., & Planes, S. (2014). Long-Distance
Dispersal via Ocean Currents Connects Omani Clownfish Populations throughout
Entire Species Range. *PLOS ONE*, *9*(9), e107610.
https://doi.org/10.1371/journal.pone.0107610
Smouse, P. E., Long, J. C., & Sokal, R. R. (1986). Multiple Regression and Correlation
Extensions of the Mantel Test of Matrix Correspondence. *Systematic Zoology*, *35*(4),
627–632. JSTOR. https://doi.org/10.2307/2413122
Sponer, R., & Roy, M. S. (2002). Phylogeographic analysis of the brooding brittle star
Amphipholis squamata (Echinodermata) along the coast of New Zealand reveals
high cryptic genetic variation. *Evolution*, *56*(10), 1954–1967.
Stevens, M. I., & Hogg, I. D. (2004). Population genetic structure of New Zealand's endemic
corophiid amphipods: Evidence for allopatric speciation. *Biol. J. Linn. Soc. Lond.*,
81(1), 119–133.
Temunović, M., Franjić, J., Satovic, Z., Grgurev, M., Frascaria-Lacoste, N., & Fernández-
Manjarrés, J. F. (2012). Environmental heterogeneity explains the genetic structure
of Continental and Mediterranean populations of Fraxinus angustifolia Vahl. *PLoS*
One, *7*(8), e42764.
Teske, P., Papadopoulos, I., Zardi, G., McQuaid, C., Edkins, M., Griffiths, C., & Barker, N.
(2007). Implications of life history for genetic structure and migration rates of
southern African coastal invertebrates: Planktonic, abbreviated and direct
development. *Mar. Biol.*, *152*(3), 697–711.
Tripp, E. A., Tsai, Y.-H. E., Zhuang, Y., & Dexter, K. G. (2017). RADseq dataset with 90%
missing data fully resolves recent radiation of Petalidium (Acanthaceae) in the ultra-
arid deserts of Namibia. *Ecol. Evol.*, *7*(19), 7920–7936.

Underwood, J. N., Travers, M. J., & Gilmour, J. P. (2012). Subtle genetic structure reveals
restricted connectivity among populations of a coral reef fish inhabiting remote atolls.
Ecology and Evolution, *2*(3), 666–679. https://doi.org/10.1002/ece3.80
Van Wyngaarden, M., Snelgrove, P. V. R., DiBacco, C., Hamilton, L. C., Rodríguez-
Ezpeleta, N., Jeffery, N. W., Stanley, R. R. E., & Bradbury, I. R. (2017). Identifying
patterns of dispersal, connectivity and selection in the sea scallop, Placopecten
magellanicus, using RADseq-derived SNPs. *Evol. Appl.*, *10*(1), 102–117.
Vavrek, M. J. (2011). Fossil: Palaeoecological and palaeogeographical analysis tools.
Palaeontol. Electronica, *14*(1), 16.
Veale, A. J., & Lavery, S. D. (2012). The population genetic structure of the waratah
anemone (Actinia tenebrosa) around New Zealand. *N. Z. J. Mar. Freshwater Res.*,
46(4), 523–536.
Verity, R., & Nichols, R. A. (2016). Estimating the Number of Subpopulations (K) in
Structured Populations. *Genetics*, *203*(4), 1827–1839.
Waples, R. S. (1987). A Multispecies Approach to the Analysis of Gene Flow in Marine
Shore Fishes. *Evolution*, *41*(2), 385–400. JSTOR. https://doi.org/10.2307/2409146
Wells, S. J., & Dale, J. (2018). Contrasting gene flow at different spatial scales revealed by
genotyping-by-sequencing in Isocladus armatus, a massively colour polymorphic
New Zealand marine isopod. *PeerJ*, *6*, e5462. https://doi.org/10.7717/peerj.5462
Whitlock, M. C. (2011). And D do not replace FST. *Mol. Ecol.*, *20*(6), 1083–1091.
Will, M., Hale, M. L., Schiel, D. R., & Gemmell, N. J. (2011). Low to moderate levels of
genetic differentiation detected across the distribution of the New Zealand abalone,
Haliotis iris. *Mar. Biol.*, *158*(6), 1417–1429.
Wilson, G. A., & Rannala, B. (2003). Bayesian inference of recent migration rates using
multilocus genotypes. *Genetics*, *163*(3), 1177–1191.
Winston, J. E. (2012). Dispersal in marine organisms without a pelagic larval phase.
Integrative and Comparative Biology, *52*(4), 447–457.
https://doi.org/10.1093/icb/ics040
Yoshino, H., Yamaji, F., & Ohsawa, T. A. (2018). Genetic structure and dispersal patterns in
Limnoria nagatai (Limnoriidae, Isopoda) dwelling in non-buoyant kelps, Eisenia
bicyclis and E. arborea, in Japan. *PLOS ONE*, *13*(6), e0198451.
https://doi.org/10.1371/journal.pone.0198451