Running Title: Salt tolerance QTL derived from the Bangladeshi landrace *Horkuch*

Natural variation in growth and physiology under salt stress in rice: QTL mapping in a *Horkuch*

× IR29 mapping population at seedling and reproductive stages

Taslima Haque[1,2†], Sabrina M Elias[1,4,6†], Samsad Razzaque[1,2†], Sudip Biswas[1], Sumaiya Farah

Khan[1,5], G.M. Nurnabi Azad Jewel[1], Md. Sazzadur Rahman[3], Thomas E. Juenger[2] and Zeba I

Seraj[1*]

[1]Plant Biotechnology Lab, Department of Biochemistry and Molecular Biology, University of

Dhaka, Dhaka-1000, Bangladesh

[2]Department of Integrative Biology and Institute for Cellular and Molecular Biology, University

of Texas, Austin, Texas 78712, USA

[3]Plant Physiology Division, Bangladesh Rice Research Institute, Gazipur, Bangladesh

[4]Department of Agronomy and Horticulture, University of Nebraska, Lincoln, Nebraska 68583,

USA

[5]Department of Biochemistry and Molecular Biology, Jagannath University, Dhaka-1100,

Bangladesh

[6]School of Life Science, Independent University, Dhaka-1229, Bangladesh

†Contributed equally

*Correspondence:

Zeba I Seraj

zebai@du.ac.bd

Highlights:

We identified QTL for salt tolerance response for two different developmental stages of rice plant and detected a significant contribution of cytoplasm-nuclear genome interaction for few traits.

Abstract:

Salinity has a significant negative impact on the production of rice yield which will become severe due to recent climate changes. To cope with this increased soil salinity, we need to develop salt tolerant rice varieties that can maintaining higher yield. Rice landraces indigenous to the coastal region of Bangladesh can be a great resource to study the genetic basis of salt adaptation. In this study, we developed a reciprocal mapping population between a salt tolerant landrace *Horkuch* and a high yielding rice variety *IR29*. We applied a QTL analysis framework to identify genetic loci that contributes to salt adaptive responses for two different developmental stages of salinity treatment. We identified 14 QTL for 9 traits and found most QTL are unique to the specific developmental stage. Moreover, we discovered a significant effect of the cytoplasmic genome on the QTL model for some important traits such as leaf total potassium and filled grain number. This underscores the importance of considering cytoplasm-nuclear interaction for breeding programs. Along with that, we detected QTL co-localization for multiple traits that highlights the constraint of multiple QTL selection for breeding program. Overall, in this study we identified multiple QTL for different physiological and yield related traits for salinity treatment for two different developmental stages of the rice plant. We detected a significant contribution of cytoplasm-nuclear genome interaction for many traits. This study also suggests the selection constraint of donor alleles due to the presence of QTL co-localization.

Keywords: salinity, QTL, reciprocal cross, cytoplasm, crop breeding

Introduction

Rice production, (*Oryza sativa* L.), which feeds almost half of the world population, is under threat from global environmental changes such as increasing salinity, heat and drought (Seck et al., 2012; Ashikari and Ma, 2015). Among these abiotic stresses, salinity has already affected 45 million hectares of irrigated land worldwide and 1.5 million additional hectares are impacted each year (Munns and Tester, 2008). Bangladesh and other locales at or near sea level are particularly vulnerable to climate change-induced salinity. In Bangladesh, about 30% of the cultivable land along the coast is affected by salinity due to tidal flood during the wet season resulting in direct inundation by saline water, and upward or lateral movement of saline ground water during the dry season (Haque, 2006).

High yielding "elite" rice cultivars are especially susceptible to salinity stress. Recent studies have shown that production of high yielding rice varieties in Bangladesh will decrease by 15.6% in coastal districts where soil salinity is predicted to exceed 4 deciSiemens per meter (dSm^{-1}) by 2050 (Dasgupta *et al.*, 2014). However, the coastal belt of Bangladesh is enriched with many local rice landraces, among which a handful are adapted to high-to-moderate soil salinities. The rice landrace, *Pokkali*, has long been used as a salt tolerant landrace reference. Many other salt-tolerant landraces such as *Horkuch*, *Ashfal*, *Jatai* and *Balam* from the coastal region of southern Bangladesh have been identified and are currently grown by farmers in these salt-affected regions (Lisa *et al.*, 2004; Rahman *et al.*, 2016). Unfortunately, these landraces suffer from low yield, poor grain quality and longer duration to reach maturity and therefore cannot serve as good candidates for commercial crop varieties. However, studies of these adapted landraces to understand their salt tolerant mechanisms can open opportunities to incorporate desired traits to commercial rice varieties. Therefore, it is important for breeders to identify genetic variants of salt stress responses in these adapted landraces in order to design highly salt tolerant rice.

The effect of salinity on rice growth varies across various developmental stages (Lutts *et al.*, 1995). The rice plant is most sensitive to salinity at the early seedling stage and during panicle formation, whereas it is relatively tolerant during early germination, active tillering and maturity (Akbar *et al.*, 1972; Heenan *et al.*, 1988; Lutts *et al.*, 1995; Pearson and Bernstein, 1959; Singh and Flowers, 2010; Zeng and Shannon, 2000). During salt stress at the early seedling stage, there is a

significant decrease of dry matter as well as quantum yield of PSII and a significant increase of
sodium concentration in root, stem and shoot tissue (García Morales *et al.*, 2012). At the
reproductive stage, physiological studies under salinity stress show a significant decrease in
panicle weight, panicle length, primary branches per panicle, filled grains per panicle, total seeds
per panicle, total seed weight per panicle, 1000-seed weight and total seed weight per plant
(Abdullah *et al.*, 2001; Rao *et al.*, 2008). Moradi et al. (2003) have however shown that salinity
tolerance at the seedling and reproductive stages is only weakly associated. This emphasizes the
importance of discovering the contributing traits of these two very important growth stages of
rice.

The physiological basis of salt tolerance during the early seedling stage is well understood.
Munns and Tester (2008) and Roy (2014) have proposed several physiological mechanisms of
seedling tolerance such as sodium exclusion, compartmentalization of excessive sodium ions
(tissue tolerance) and shoot-ion independent tolerance for early stage tolerance. It has been
reported that *Pokkali* maintain lower shoot Na^+ accumulation and lower shoot Na^+/K^+ ratio under
high salinity compared to sensitive genotypes (Kavitha *et al.*, 2012; Sexcion *et al.*, 2009). The
enhancement of salinity tolerance by constitutive overexpression of the vacuolar Na^+/H^+
antiporter gene from *Pokkali* in transgenic rice plants suggest that this landrace may use a tissue
tolerance mechanism to lower shoot Na^+/K^+ ratio under high salinity (Amin *et al.*, 2016). Negrao
et al genotyped 392 rice accessions by EcoTILLING in order to understand allelic difference for
salt stress. They targeted five known genes that are involved in these different salt tolerant
mechanisms and assembled a set of accessions that represents all the haplotypes present in the
coding region of these five genes. The systematic study of phenotypes of this set suggest that
none of the main three mechanisms of tolerance is preferentially used over another (Inês *et al.*,
2015). Therefore, studies of different landraces can offer ways to understand individual salt
tolerance mechanisms in the rice plant (Lisa *et al.*, 2004; Yesmin *et al.*, 2014). However, the
mechanism associated with tolerance during the reproductive stage has been barely explored. As
mentioned earlier, high salinity in this stage can alter many traits associated with grain quality
and quantity, eventually decreasing yield significantly. Therefore, it is important to explore the
physiological response of the rice plant at both stages in order to obtain a superior variety from
breeding which can maintain salinity tolerance for both developmental stages.

The choice of female parents in breeding programs plays a critical role in the performance of crosses. Plants show evidence for complex nuclear-cytoplasmic interaction that may alter their phenotypes in both interspecific and intraspecific crosses. However, it still remains unclear to what extent these two components interact with each other and the role of environment in this interaction. Gregorio and Senadhira (1993) have studied the genetics of salinity tolerance on diallelic reciprocal crosses of nine different rice varieties and found significant reciprocal effects among crosses. The presence of maternal inheritance has also been reported for other abiotic stresses such as chilling response (Chung et al., 2003) and drought (Iida *et al.*, 2007). Therefore, in plant breeding programs where the aim is to produce stress tolerant high-yielding varieties, it is important to consider a specific cytoplasm and its interactions with nuclear donor alleles for determining the performance of plants under stress.

Quantitative trait loci (QTL) mapping has been implemented in many studies of rice to explore the genetic basis of traits involved in salinity stress for seedling stages, including salt injury/tolerance score, fresh and dry weight of shoot and root, Na^+ and K^+ content of shoot and root, and chlorophyll content (Cheng *et al.*, 2011; Lin *et al.*, 2004; Ren *et al.*, 2005; Sabouri *et al.*, 2009; Soltani *et al.*, 2016; Thomson *et al.*, 2010; Tian *et al.*, 2011; Wang *et al.*, 2012; Zheng *et al.*, 2015). However, very few studies have been conducted to understand the genetic basic of reproductive stage traits that are important for tolerance such as plant height, tiller number, panicle number, pollen fertility and yield (Hossain *et al.*, 2015; Zang *et al.*, 2008).

QTL co-localization has been reported for traits that are strongly correlated (Dechaine *et al.*, 2014). Many clustered, putatively pleiotropic QTL have been found that affect various life history and fitness characters, especially those that are related to yield, in rice, wheat, pea and rapeseed (Burstin *et al.*, 2007; Quarrie *et al.*, 2006; Shi *et al.*, 2009; Xue *et al.*, 2008). QTLs for two different traits can have the same/opposite sign of effects for both. If the genes involved help in coordination during multiple steps of development, then positive selection for one trait may have an outcome on several traits in the same positive direction, e.g. a pleiotropic same sign QTL for seed size and protein content in pea (Burstin *et al.*, 2007). Very strong opposite sign of effects for QTL has also been reported for plants including rice (Dechaine *et al.*, 2014; Xiao *et al.*, 1998) and

may represent tradeoffs. For breeding programs where breeders aim for QTL pyramiding for

multiple desired traits, opposite signed QTL for different traits may impose some constraints on

selecting co-localized QTL. Hence, it is beneficial to have a clear understanding about QTL co-

localization and perform careful selection of these genomic loci for pyramiding.

In this study we genotyped a reciprocal mapping population between the salt tolerant landrace,

Horkuch and a high yielding variety *IR29* by DArTseq technique (Akbari *et al.*, 2006). We

identified 14 QTL for 9 traits for salinity treatments at two different developmental stages of the

rice plant. One important finding of this study was to characterize the role of cytoplasm in a

plant's performance under salinity and implement an analysis incorporating this information to

estimate the effect of a QTL. Furthermore, in this QTL analysis framework, we applied a linear

mixed model to incorporate residual polygenic variation which is a better way to estimate QTL

effect for polygenic traits. In our previous study we applied Double digested Restriction

Associated DNA (ddRAD) technique to construct genetic map of this population where we failed

to map a substantial genetic space (Noor *et al.*, 2019). In this current study, with the aid of an

improved genetic map by DArTseq technique and a robust QTL analysis framework we were

able to identify additional QTLs with higher likelihood and tighter confidence interval. In

addition to that, we identified co-localized QTL within and across two different treatment stages,

which emphasizes the need for conditional selection of QTL in a breeding program in order to

combine survivability at the seedling stage and yield tolerance at reproductive stage. Taken

together, the findings of this study contribute to our understanding of the molecular mechanism

of salt tolerance for *Horkuch* and pave a way to introgress salinity tolerance into a commercial

cultivar that can maintain significant yields under stress.

Methods

Development of the reciprocally crossed populations

The rice cultivar *Horkuch* (IRGC 31804) and *IR29* (IRGC 30412) were acquired from the IRRI

Gene bank. Seeds were sown in a crossing block after breaking dormancy at 50°C for 5 days in

an oven during the wet season of 2011 (June-July) at IRRI, in the Philippines. Two reciprocally

crossed F_1 seeds were generated with *Horkuch* and *IR29* in October, 2011. F_1 plants were

initially confirmed by SSR marker (RM493) and both F_1s were advanced to the F_2 stage by

selfing. Our experimental approach centers on an $F_{2:3}$ design, whereby genotypes for mapping

are collected from F_2 individuals and phenotypes are obtained from a sampling of their $F_{2:3}$

progenies (Zhang et al, 2004). Approximately, 1200 $F_2$from both crosses (600 from each cross)

were advanced to F_3 at BRRI in Bangladesh. $F_{2:3}$ progenies derived from *Horkuch* × *IR29* are

referred to as *Horkuch*♀ and those from the reciprocal cross as *IR29*♀.

Physiological screening at the seedling stage

We randomly chose 204 $F_{2:3}$ families and screened for salinity stress-induced phenotypic traits at

the seedling stage (137 families from *IR29*♀ and 65 families from *Horkuch*♀ cross were chosen

for this screening). Plants were grown in hydroponic solution with 9 replicate siblings of each of

the 204 families (Yoshida, 1976). Two weeks after germination (four leaf stage), we started salt

stress from 6 dSm^{-1} through 12 dSm^{-1} salt stress (NaCl) with 2 dSm^{-1} increments per day. Then

after 2 weeks of salt stress, plants were scored using the SES approach (Standard Evaluation

System) (IRRI, 2002) which evaluates the leaf damage as well as the overall appearance of

plants. Here, a scale of 1-9 corresponds to ascending salt sensitivity. Physiological parameters

like stomatal conductance, chlorophyll content, potassium and sodium content, shoot and root

relative water content, dry weight, and length of the salt stressed plants were also measured (See

Supplementary Methods).

Physiological screening at the reproductive stage

Reproductive stage screening under salinity stress (NaCl stress at 8 dS/m applied at a time to

water in a bowl with the plants in soil in pots with holes) was performed for a subset of

Horkuch♀ and *IR29*♀ $F_{2:3}$ families in soil by following standard IRRI protocols (Gregorio *et al.*,

1997). For reproductive stage screening 140 families were chosen based on a selection of 70 F_2

families from each population. Our selection was based on the distribution of SES scores where

the lower tail (more tolerant families) was defined as SES scores from 3 to 5 and the upper tail

(sensitive) was defined as SES scores from 7 to 9. All families that were in the lower or upper

tail were selected along with 70 randomly chosen families from SES score in between 5 and 7.

The two F_0 parents were also included in our studies. A few families had poor germination and

were subsequently excluded from the reproductive screening experiment. In the end, 130

families were included in reproductive screening: 61 from *Horkuch*♀ and 69 from *IR29*♀

population. Stress was applied prior to panicle initiation at 30 days and continued until seeds

were mature. Five F_3 siblings from each F_2 individual were considered as replicates while

screening. Samples were randomized using an incomplete block design. Parameters that were

measured at this stage have been listed in Supplementary Methods.

DNA extraction, genotyping by DArTseq and linkage map construction

Genomic DNA of F_2 individuals and parents was extracted using the CTAB method (Doyle and

Doyle, 1990) from 1g of fresh leaf tissue after freezing in liquid nitrogen and grinding.

Genotyping was done by the DArTseq technique as described by Akbari et al (2006). For the

DArTseq method Nipponbare Genome from Phytozome (v9) was used as reference to determine

the physical position of each DArTseq clone. Analyses for linkage map construction were

completed with qtlTools (Lovell) and R/qtl (Broman *et al.*, 2003) packages. We filtered markers

that had more than 50% missing data and showed significant segregation distortion for chi-

square test (p-value < 0.001) from expected ratio (For a given locus,1:2:1 $\equiv$ homozygous parent

1 allele: heterozygous: homozygous for parent 2 allele). Similar markers were further removed

and marker order was obtained by the tspOrder function from TSPmap tool (Monroe *et al.*, 2017)

which applies a traveling salesperson problem solver to order markers. Detailed method for

constructing the linkage map is described in our Supplementary Methods. The final linkage map

was estimated by est.map function of R/qtl package with the Kosambi map function. To test for

the occurrence of cytoplasm-nuclear association we performed chi-square test of independence

on allele frequency for each locus grouped by cytoplasm and significant association was

determined by adjusting p-values using FDR method (FDR threshold =0.1).

QTL analysis

Genotype probabilities were calculated at a 1 cM step interval using the calc.genoprob function.

QTL mapping was executed using the Haley-Knott regression algorithm (Broman *et al.*, 2003) of

the R/qtl2 (Broman *et al.*, 2019; Lovell)(Broman *et al.*, 2019) package where we fit the LOCO

(Leave One Chromosome Out) model for each trait and included kinship as a covariate. The

LOCO model utilizes the kinship matrix to reduce background polygenetic variation except for the chromosome that is being tested for QTL mapping. We also tested cytoplasm as an additive and interactive covariate in the QTL model using likelihood ratio tests and retained factors in the models when significant. To test whether our selection of a test cohort imposed significant population structure for the reproductive stage treatment we assigned each F_2 family into one of the following categories: tolerant, intermediate and sensitive. We incorporated selection cohort as a covariate while building QTL model for traits at reproductive stage. However, we did not find any significant effect of selection cohort on QTL models and therefore did not include selection cohort as a covariate for further analysis. Significance thresholds for QTL were determined for each trait by 1000 permutations (alpha=0.05) and QTL peaks that passed the threshold were considered for further analysis. Permutations were stratified by cytoplasm for the QTL models where cytoplasm was considered as a covariate. We also evaluated the normality of the QTL model residuals and found this assumption has been violated of for trait TK. Confidence intervals (1.5 LOD drop) for each QTL were calculated using the *lodind* function of the R/qtl package (Broman *et al.*, 2003) expanding it to a true marker on both sides of the QTL. Codes for QTL analysis are available in GitHub repository (Haque, 2019).

Identification of candidate genes within QTL confidence interval and Gene ontology enrichment analysis

In order to identify candidate gene models within a given QTL interval, we integrated the genetic and physical map based on the marker order of genetic map. We first pulled out the genetic markers flanking a given QTL confidence interval and their basepair positions to define the physical interval on the genome for that QTL. Gene models in these physical QTL intervals were retrieved using the structural gene annotation of the rice Nipponbare reference genome from Phytozome 9. We used the Gene ontology (GO) annotated for each gene model of this reference (Phytozome 9) for GO enrichment analysis. We then tested for the enrichment of GO terms for each QTL interval using the classical Fisher's exact test available in the topGO (Alexa and Rahnenfuhrer, 2019) package in R.

RESULTS

Phenotypic traits vary between cross direction in both developmental stages:

We used a bi-directional $F_{2:3}$ experimental design to examine the effect of salinity on various
growth, yield and physiological parameters of rice as well as the role of cytoplasm on these
traits. Rice is most susceptible to salinity during seedling and reproductive growth stages.
Therefore, we focused on phenotypes at these two stages that can potentially mediate stress
during salinity treatment. For the seedling stage, we scored traits that were related to survival,
photosynthesis and mineral elements in leaves including Standard Evaluation Score (SES), Total
Chlorophyll Content (Tchlr), Shoot Length (SL), Root length (RL), Total Sodium (TNa), Total
Potassium (TK) and Potassium by Sodium (K/Na). For the reproductive stage treatment, we
focused on yield-related parameters including Panicle Exsertion (PE), Total Tiller Number (TT),
Effective Tiller Number (ET), Filled Grain Weight (FGW), Filled Grain Number (FGN) and
Spikelet Fertility (SF).

We found striking difference between the two parents for many of our measured traits across
both stages of salt treatment (Table 1). As reported in previous studies, we have found that
Horkuch is more tolerant to salinity (measured by SES score) compared to *IR29* which is highly
sensitive for salt at the seedling stage (p-value < 0.01). For reproductive stage salinity treatment,
these two parents show significantly different responses for ET, SF, FGW, DF and HI where the
Horkuch parent had higher number of ET and also higher SF, FGW and HI.

It has been reported that complex cytoplasm-nuclear interaction can alter plant phenotypes
(Joseph *et al.*, 2013a). Therefore, we aimed to test whether cytoplasm has a role on the
performance of this mapping population under salinity stress. We found many traits were
significantly different as a result of cytoplasmic background (Table 1). Tchlr, TNa, TK and K/Na
were found to differ significantly among the Horkuch and IR29 cytoplasms at the seedling stage
treatment. For reproductive stage treatment PE, TT, ET, FGN were found to differ significantly
among the two parental cytoplasms. This observation indicates that cytoplasm can explain a
significant amount of variation in the population.

The generation of extreme phenotypes in a crossing population, or transgressive segregation, has
been reported in many plants which may be due to the effect of complementary genes, over-
dominance or epistasis (Rieseberg *et al.*, 1999). Here, we further investigated the role of cytoplasm
on the transgressive segregation of these traits. In this experiment we found that SES, SRWC,
TChlr, TK, FGN, FGW clearly segregate transgressively (Table 1). Interestingly, TK, TChlr,

FGN and FGW showed significant differences in the two reciprocal crosses with respect to the

segregation pattern. The distribution of TK showed a strong bimodal pattern with very little

overlap of distribution between cytoplasmic backgrounds (Figure 1). This observation is

indicative of an important role of cytoplasm on transgressive segregation for this population.

To understand the partitioning of genetic variation we used principal component analysis of 130

families which had complete observations of all traits for both the stages of treatment. The first

two PC axes comprise the majority of genetic trait variation in this population (31.3%) (Figure

2). This indicated substantial genetic correlation among these traits in this population.

Interestingly, principal component one separates individuals into two groups almost exclusively

depending on their respective cytoplasm. This observation further supports a significant role of

cytoplasm on the performance of a plant under different treatment stages. We also found

moderate correlation of some traits between two different stages of treatment: PH showed

positive correlation with SL and yield related traits such as FGN, FGW, SF, HI and PE are

negatively correlated with TK (Figure 2). This correlation suggests some possible shared

mechanisms of salinity responses that has some trade-off between two different treatment stages.

Linkage map construction

In our previous study we constructed a linkage map for this reciprocal mapping population by

applying ddRAD technique. Unfortunately, we had high genotyping error and inflated

segregation distortion for the two parental alleles for a given locus. Therefore, we failed to

capture the genome-wide linkage map which resulted in the removal of the entire chromosome 5

for QTL mapping. In this study, we genotyped this reciprocal mapping population using

DArTSeq (Diversity Array Technology coupled with Sequencing) technique in order to improve

genotyping error by adding higher sequence coverage. DArTSeq can generate low to moderate

density SNP information with high coverage and low cost. This method uses restriction enzymes

to reduce the complexity of the genome and has been optimized for various plant species to

achieve best complexity reduction. We used this platform for our mapping population to generate

genotype information at ~10 thousand loci that are well-distributed in the rice genome. For

analyses, we filtered all the DArTseq SNP loci to obtain polymorphic homozygous SNPs for the

parents and retained only loci that had minimum 50% representation in the population. With

these filter criteria we obtained 2,230 high quality SNPs for this mapping population. Among

these loci, 956 markers that showed significant segregation distortion by chi-square test [P value < 1e-2] were removed. More distorted loci were skewed towards the *Horkuch* parent than the *IR29* parent. 739 markers that were similar were dropped using a minimum recombination fraction threshold of 0.03. Markers were reordered by the concorde program as determined by the tspOrder function (Monroe *et al.*, 2017). The final map was constructed by "kosambi" map function after dropping 36 markers which had very different orders between their genetic map and physical position in genome. The final map had 499 markers with a map size of 2004.8 cM and average distance between markers of 4.1 cM (Supplementary Figure 1A). The maximum gap of 22.3 cM was found in chromosome 5. Chromosome 7 had the fewest markers (22 markers). Supplementary Figure 1B presents the concordance of genetic map with the physical map of rice genome. As mentioned in earlier, we detected significant association of cytoplasm for multiple traits in both stages of treatment therefore we aimed to test (Chi-square test of independence, see Method section) for cytoplasm-nuclear association for each marker in this genetic map. We detected 10 loci that showed significant cytoplasm-nuclear association [FDR < 0.1] and these are mostly clustered in chromosomes 2, 3 and 7. This association suggested the presence of cytoplasm-nuclear linkage disequilibrium in this mapping population resulting from selection at the reproductive stage during the development of the mapping population. Overall, we constructed a linkage map with moderate marker density that was closely aligned with the physical map of rice genome.

Significant QTL at both growth stages

QTL for seedling stage treatment

We measured eight traits that reflect the survival performance of rice seedlings under salinity stress. We found six QTL for three traits: SL, RL and TK (Table 2, Figure 3). We found three significant QTL for SL occurring at qSL.1@183 (reporting a QTL for SL located at chromosome 1 at 183 cM), qSL.3@218, and qSL.5@160. For these QTL, the positive alleles were from *Horkuch* parent and we detected no significant cytoplasmic effects or interactions. The QTL at qSL.1@183 had a large effect corresponding to a ~3.5 cm increase in seedling length and was localized to 3.5 Mbp at the end of chromosome 1 (~38.4 Mbp). The second QTL, qSL.5@160, had a small confidence interval of ~ 1 Mbp but the effect size was moderate. The third QTL at qSL.5@160 had a very larger confidence interval (~ 9 Mbp) with small effect size.

We identified one large effect QTL for RL at qRL.2@167 with a confidence interval of ~ 4 Mb.

Here, the *Horkuch* parent contributed the positive allele. For TK, cytoplasm was a significant

covariate that interacted with two QTL that were detected at qTK.2@45 and qTK.3@203. In

addition, the QTL qTK.2@45 is co-localized with the cluster of genetic loci that showed

significant cytoplasm-nuclear association. The other QTL for TK, qTK.3@204 did not overlap

with the association cluster in chromosome 3 but resided in close proximity. This evidence

suggests a possible role of cytoplasmic-nuclear interaction for this trait.

QTL for reproductive stage treatment

Very little work has been published on salinity stress during rice reproductive stages. For this

stage, our primary focus was on yield responses of the plants under salt stress. We found 8 QTL

for 6 traits at reproductive stage salinity treatment (Table 2, Figure 3). A major effect QTL for

FGN was found on Chromosome 10 at 58.5 cM (qFGN.10@58.5). Here, the *IR29* parent

contributed the positive allele. We found significant cytoplasm-nuclear interaction for QTL

model where the *IR29* allele had positive effects only for *IR29*♀ (Figure 4). However, this QTL

had a very wide confidence interval of ~6 Mbp. We found two co-localized QTL in this same

region including a QTL for FGW and another for SF at qFGW.10@58.5 and qSF.10@59

respectively. These two QTL also had significant interactions with cytoplasm in their

corresponding QTL models. *IR29* contributed the positive allele for both qFGN.10@58.5 and

qFGW.10@58.5. For FGW, as like FGN, the *IR29* allele had a positive effect only for *IR29*♀ but

for SF this allele not only had positive effect for *IR29*♀ but also had a negative effect for

Horkuch♀. We also found a QTL for HI at qHI.10@104 for which the positive allele was from

IR29. However, this model only had an additive effect of cytoplasm.

Another QTL, qET.7@97 cM was detected for ET where the *Horkuch* parent contributed the

positive allele. Similar with the other grain related traits, cytoplasm contributed significantly to

this QTL model. However, the positive allele from *Horkuch* performed better in *IR29*♀. We also

found three PH QTL occurring at qPH.1@215, qPH.3@211 and qPH.5@144, for which the

positive alleles were from the *Horkuch* parent. The first two QTL models showed only additive

contribution of cytoplasm but the third showed also an interactive effect of cytoplasm. Overall,

we found a hotspot of QTL on chromosome 10 for multiple parameters related to yield. The

significant correlation of these traits may reflect related metrics of yield performance in rice.

QTL co-localization

In this study, we found that some QTL intervals of various traits overlapped, therefore we annotated these overlapping intervals as QTL clusters. We detected a co-localized QTL at 1@175:220 cM [QTL Cluster 1 (QC 1)] affecting PH and SL from two different stages of treatment. These traits represent the vigor of plants at the two different growth stages and had significant positive correlation (Pearson correlation coefficient = 0.54). In parallel with this co-relation, these two QTL had positive alleles from the *Horkuch* parent which indicate the same sign of effects for these QTL. Another wide co-localized cluster was found at 3@194:273 cM [QTL Cluster 2 (QC 2)] impacting SL, PH and TK with the positive allele from the *Horkuch* parent. However, TK showed no significant correlation with PH which was causal for the co-localization for QC1. A third QTL Cluster 3 (QC 3) was found at 5@144:170 for SL and PH where the positive alleles were from *Horkuch* parent. The fourth QTL Cluster (QC 4) at 10@58:107 was found for four yield related traits including FGN, FGW, SF and HI for which the positive alleles were from the *IR29* parent.

GO enrichment analysis of candidate genes within QTL confidence intervals

To understand the molecular mechanism of salt tolerance, we further investigated the function of candidate genes that were located within the QTL confidence intervals (Supplementary File 1).We applied GO enrichment analysis on the candidate gene lists against the genome-wide background frequency of GOs. The results for GO enrichment analysis have been provided in Supplementary File 2. For the seedling stage, QTL at qSL.1@183 showed significant enrichment of GO terms such as pollination, protein lipidation, lipoprotein and liposaccharide metabolic process, various transport and DNA-directed RNA polymerase complexes. Another QTL for SL at qSL.3@218 was significantly enriched with the GO terms protein transport and localization, amide and lipid transport, cytoskeleton and actin binding. The third QTL for SL, qSL.5@160 had significant enriched GO terms for chromatin assembly, nucleosome organization, DNA packaging, ion transport, amide and peptide transport. QTL qRL.2@167 showed significant enrichment of GO terms such as carboxylic, dicarboxylic and C4-dicarboxylate transport, malate transport, anion transport and sexual reproduction. Two QTL for TK, qTK2@45 and qTK3@204, were both enriched with the GO terms oxidoreductase activity, various

transmembrane transporter activity, potassium ion transmembrane transporter and cation
transmembrane transporter.

For traits at reproductive stage, the QTL qPH.1@215 was significantly enriched with the GO
terms anion and potassium ion transmembrane transport, divalent metal ion transport,
pollination, reproduction process, endoplasmic reticulum and organelle sub-compartment.
Another PH QTL, qPH.3@211 showed enrichment for GO terms such as cellular nitrogen
compound metabolism, organic acid transport, mitochondrial membrane and protein complex.
The third QTL, qPH.5@144 also showed significant enrichment for nitrogen compound
metabolic process. This QTL was also enriched for various mitochondria and cytoplasm related
GO terms. The QTL, qET.7@97 was enriched with cytoplasm and mitochondrial membrane part
and chlorophyll metabolic process. QTL for FGW, FGN, SF and HI share the same QTL
intervals therefore the gene models within this interval were identical. This interval was enriched
with cell wall macromolecule catabolic process, amino sugar and glycan metabolic process,
protein localization to organelles and mitochondrial transport. The significant enrichment of
mitochondrial and organelle related GO terms for some QTL confidence intervals suggests a
possible explanation for the significant cytoplasm and cytoplasmic-nuclear interactions detected
in our study.

GO enrichment analysis of differentially expressed genes of this mapping population

In our previous gene expression studies (Razzaque *et al.*, 2019; Razzaque *et al.*, 2017), we
investigated the association of gene expression differences with salt stress on this reciprocal
mapping population for both seedling and reproductive stage salinity treatment. To understand
the molecular mechanism of cytoplasm-treatment interaction, we further tested for GO
enrichment of genes that showed significant cytoplasm*treatment in our previous studies
(Supplementary File 3). From our seedling stage salinity treatment study, the expression profile
of the reciprocal mapping population for shoot and root tissues for 24 hours and 72 hours after
salt treatment was available. Differentially expressed genes (DEGs) for cytoplasm*treatment
were significantly enriched with processes such as chromatin assembly, amide biosynthesis,
various cation transports, photosynthesis, cellular response to stress and energy coupled proton
transport. Cellular components were enriched with GO terms involved in various organelles,
mitochondrial matrix and photosystems. GO terms such as sodium ion transmembrane activity

and various sodium symporters were enriched for molecular function. DEGs for root tissues were
enriched with biological processes such as response to oxidative stress and transports of malate,
dicarboxylic acid and copper ion. The only significant enriched GO term for the cellular
component was for extracellular activity. For molecular functions, DEGs were enriched with GO
terms such as oxidoreductase, hydrolase activity and voltage gated potassium channel activity.

For the reproductive stage of salinity treatment, the expression profile of shoot and root tissues at
72 hours after salinity treatment was available. The number of DEGs that showed significant
interaction between cytoplasm and treatment was very low (5 significant DEGs) for root tissue
therefore we only tested enrichment for DEGs in shoot tissue. DEGs for shoot tissues for
reproductive stage were enriched with biological processes such as different lipid metabolic
processes, tRNA processing and various ion transporters. Similar to shoot DEGs at seedling
stage, significant enriched GO terms for the cellular component were involved with
mitochondria and organelles. For molecular function, significant enriched GO terms showed
phosphatase and oxidoreductase activity, sodium ion transporter activity and various sodium
symporter activity including organic acid:sodium symporter, bile acid:sodium symporter activity.
These were also enriched for DEGs for the seedling stage shoot tissue. Many of these symporters
are located in the mitochondrial membrane or organelle lumen. This is one of the reasons why
we had many significantly enriched GO terms for the cellular component of mitochondria.

For QTL models that had significant cytoplasmic effect, 1473 annotated genes are present in the
respective confidence intervals. Among the genes that are present in the QTL confidence
interval, 188 showed significant cytoplasm*treatment interaction in our previous expression
studies (Supplementary File 4). In order to identify their association for cellular component we
further tested for enrichment of GOs for the same. We found these genes to be significantly
enriched with GO terms such as mitochondrial proton-transporting ATP-synthesis complex,
mitochondrial protein-complex and mitochondrial membrane.

DISCUSSION

In this study, we explored the responses of rice to salinity stress at two different growth stages
with a reciprocal mapping population. Among the 14 QTL that we reported, 8 QTL models

showed significant effect of cytoplasm. This finding underlines the importance of considering
both organelle and nuclear genome for complex traits such as salinity tolerance.

Cytoplasmic background may play an important role in trait genetic architecture by itself or
through complex interactions with the nuclear genome. (Joseph *et al.*, 2013b; Lovell *et al.*, 2015;
Moison *et al.*, 2010; Tang *et al.*, 2013). Gregorio and Senadhira (1993) reported significant
reciprocal effects among crosses for salinity response in their study of nine different rice
varieties and suggested using susceptible plants as male parent for hybridization programs. In
order to identify the best candidates for QTL pyramiding by breeders, it is essential to estimate
single QTL effect for the trait of interest. Hence, it is important to test for the random effect of
covariates such as cytoplasm in a QTL model and estimate its size. In this way, the causality,
contribution and combinations of cytoplasm and nuclear-donor alleles of QTL can be defined.
Moreover, including cytoplasm as covariate in QTL mapping, can increase the ability to detect
unrelated QTL peaks. Considering all these aspects, in this study we employed a QTL modeling
framework where the cytoplasm-nuclear interaction was also considered as a contributor to
phenotypic variance. For TK, FGN, FGW the additive effect of cytoplasm was significantly large
compared to the effect of a single QTL (Figure 4). Overall, we found significant contribution of
cytoplasm for traits related to yield, such as FGN, FGW and SF as well as one important trait for
the seedling stage TK. Identification of causal impacts of cytoplasm will help to define the best
combination of cytoplasm and nuclear-donor materials and will underscore the selection trade-
off for multiple desired traits. For instance, on the one hand, we found the positive nuclear allele
of *IR29* had its effect only in *IR29*♀ for the QTL model of yield related traits; while on the other
hand we had the strong positive effect of *Horkuch* cytoplasm for the QTL model of TK at
seedling stage treatment. The latter trait is a highly desired one for breeding salt tolerant
varieties. Hence, estimating the contributions of cytoplasm for multiple traits can help
understand the performance trade-off in breeding program for QTL pyramiding.

Cytoplasmic genome can influence the interaction of alleles from nucleus and cytoplasm and can
favor the evolutionary co-adaptation of high-fitness. In the current study we found a significant
association of cytoplasm for some traits and therefore further tested for non-random interaction
of alleles for nucleus and cytoplasm. We found that the QTL qTK.2@45 was a hotspot of
cytoplasm-nuclear interaction on chromosome 2. Similarly, qPH.3@211 was another similar

hotspot on chromosome 3. Both of these QTL models showed significant effect for cytoplasm.
For qTK2@45, the effect of cytoplasm was mostly additive where *Horkuch*♀ contributed large
positive effect. On the contrary, for qPH3@211, cytoplasm had an interactive effect. *Horkuch*
nuclear allele had a positive effect on PH but the effect was even higher for *IR29*♀
(Supplementary Figure 3). Taken together, this suggests a significant interaction of nuclear
alleles with the cytoplasmic genome. This further supports the fact that selection of female plant
plays an important role for the performance of a breeding population and while pyramiding QTL,
the conditional selection of cytoplasm may have some trade-off on a hybrid plant's performance.
We detected significant cytoplasm-nuclear linkage of a few markers that overlapped with some
QTL intervals. Therefore, careful consideration is needed in order to select these loci for QTL
pyramiding.

One important finding in this study is that we have detected multiple co-localized QTL within
and among the two different stages of salinity treatment. This finding emphasizes the possible
constraints during selection of QTL in a breeding program. Here we identified four QTL clusters
where multiple trait QTL co-localized. Co-localized QTL can impose constraints on selection for
QTL pyramiding. As an example, we found that QTL cluster 1 had a positive effect for the
Horkuch parental allele for PH and SL. However, a taller plant is not the desired plant
architecture for a breeding program for high-yielding rice varieties since this will lead to over-
investment of energy in vegetative growth and potential lodging. On the other hand, Leon et al.
(2015) reported that percent of shoot length reduction under saline treatment is highly co-related
to saline sensitivity. This conditional relationship between traits results in some possible trade-
offs between favorable and undesirable traits. The same logic is applicable for QTL cluster 2
where traits (SL, TK and PH) for these clusters are positively correlated but increased PH is not
desirable for any breeding program. On the other hand, for QTL cluster 4, all the yield related
such as FGN, FGW, SF and HI could be combined where the *IR29* parent contributes all the
positive alleles. This finding underscores the importance of studying the performance of a plant
for different developmental stage. In addition to that, we need to consider the fact that selection
on multiple traits may not be orthogonal due to the complex mechanisms of salt adaptation.

To understand the molecular mechanism of salt response and the effect of cytoplasm for salt
tolerance we tested for enrichment of GO functions for genes within QTL confidence intervals.

Both the QTL intervals for TK were enriched with various transmembrane transporter activity,

and potassium ion transmembrane transporter. K^+ is involved in numerous metabolic process in

plants and excess Na^+ interferes with the K^+ homeostasis during salinity stress. To maintain the

cellular homeostasis of K^+ various potassium transmembrane transporters have been reported

that showed increase salt tolerance in various glycophytes (Tester and Davenport, 2003). For the

reproductive stage, we found most of the QTL intervals for PH, ET, FGW, FGN, SF and HI were

enriched with mitochondria, cytoplasm and organelle related GOs. This supported the

observation that these QTL models also showed significant interaction with cytoplasm. Thus a

possible interaction of the cytoplasm genome with nuclear alleles present in the region of QTL

confidence intervals is likely. Additionally, enrichment analysis of DEGs (significant for

cytoplasm*treatment model) from our previous studies (Razzaque *et al.*, 2019; Razzaque *et al.*,

2017)   on this mapping population were enriched with GO terms such as organelle, thylakoid,

mitochondria, photosynthesis, cation transmembrane transporter and various sodium symporter

activities. Salt stress inhibits photosynthesis of plants but how this affects the ionic balance of

chloroplasts has not been studied much until recently. Bose et al. (2017) has proposed some

candidate transporters that are involved for the movement of sodium, potassium and chloride

across chloroplast membrane in glycophytes and halophytes and explained how these

transporters may regulate photosynthesis in chloroplast. These candidate symporters include bile

acid: sodium symporter and cation transmembrane transporter which have possible role in

maintaining chloroplast ion homeostasis. From our gene expression studies of

cytoplasm*treatment DEGs, enrichment of symporter GOs that are localized in mitochondria and

organelles suggest a possible role of mitochondria and chloroplast during salinity stress and

tolerance or sensitivity to it. This evidence also suggests a plausible explanation why we found

cytoplasm as a covariate in QTL models for this study. These are likely candidates for future

functional genomic studies of salinity tolerance in rice.

In this QTL analysis framework, we applied linear mixed model which can handle cytoplasm

and alleles as fixed effect predictors. This model can also consider residual polygenic variation

as random effect using a kinship matrix. Here we implemented DArtSeq technique which can

genotype a moderate number of SNPs that are well-dispersed in rice genome and aimed to select

SNPs close to gene space of the rice genome. In our previous study, we had generated a genetic

map on this mapping population by ddRAD technique which failed to capture a significant space

of genetic map due to erroneous genotyping and high rate of missing SNP calls (Noor *et al.*,
2019). In this current study, we implemented a robust QTL analysis framework on this improved
genetic map and we were able to detect three QTLs for SL and one RL at seedling stage salinity
treatment which we could not detect in our earlier study (Supplementary Table 1). For
reproductive stage salinity treatment, we were able to detect additional five QTL for PH, ET and
SF. We have also detected one big effect QTL for FGN and FGW in a different chromosome in
this current study due to the fact that in our previous study we failed to capture markers at that
region. In addition to that, this framework provided QTL with higher likelihood and tighter
confidence interval and provided better estimation of effect size of each QTL for a given trait.
Therefore, this additional detected QTL with high LOD scores and tighter confidence intervals
may contribute significantly for the improvement of salt tolerant high yielding rice variety
development.

CONCLUSION

In this study, we aimed to identify genetic loci for salinity tolerance of a rice landrace, *Horkuch,*
at two sensitive developmental stages. We found 14 QTL for 9 traits under salinity treatment.
We detected some overlap in the genomic regions affecting traits across developmental stages.
One chief finding of this study was the significant contribution of cytoplasm on many traits and
eventually the effect on their corresponding QTL model. Enrichment analyses suggest that the
observed cytoplasmic effect could be causally related to plastid symporter activity and their
interaction with nuclear genes. Collectively, this study helped to understand the genetic basis of
salt tolerant mechanism of a local rice landrace *Horkuch*. Moreover, careful implementation of
pyramiding of QTLs that were detected in this study can pave a way to generate high yielding
salt tolerant rice varieties.

SUPPLEMENTARY DATA

Supplementary_Material: Detailed supplementary methods, tables and figures

Supplementary_File_1: List of genes in QTL confidence interval where one sheet represents
one single QTL

Supplementary_File_2: List of significant GO terms for genes in QTL confidence interval where one sheet represents one single QTL

Supplementary_File_3: List of significant GO terms of significant DEGs for cytoplasm*treatment interaction model of seedling shoot tissue, seedling root tissue and reproductive shoot tissue

Supplementary_File_4: List of significant GO terms for the common genes between all candidate genes of QTL confidence interval and significant DEGs for cytoplasm*treatment interaction model of seedling shoot tissue, seedling root tissue and reproductive shoot tissue

CONTRIBUTIONS

Z.I.S. and S.M.E. designed the experiment. M.S.R. did reciprocal crossing. S.M.E., S.R., S.F. K., S.B. and G.M.N.A.J grew plant and collected phenotypes. S.M.E. and S.R. isolated DNA for DArTseq. T.H and S.M.E did genotype calling from raw data. T. H. did the modeling for QTL and other statistical analysis. T.E.J provided his feedback for statistical analysis. T.H. wrote the manuscript. T.E.J. and Z.I.S. provided their significant feedbacks for writing. S.M.E. and S.R. revised the manuscript.

ACKNOWLEDGEMENT

We thanks to Nazrul Islam and other technicians at BRRI and Dhaka University who took care for the plant materials. We would like to thank the members of the Plant Biotechnology Lab and Juenger Lab who provided their feedbacks on this research project. This work was supported by the PEER science grant to Z.I.S. by the National Science Foundation (Project 2-004) and USAID (agreement number AID-OAA-A-11-00012). The grant was hosted by an existing NSF grant from the Plant Genome Research Program (IOS-0922457) to T.E.J. S.M.E. was supported through a PhD fellowship grant from Monsanto's Beachell-Borlaug International Scholars Program (MBBISP). We would like to thank Dr. Harkamal Walia for supervising S.M.E for her MBBISP. We are thankful to Dr. Adam Price for recommending us DArtseq for genotyping and also supervising S.F.K. for her Split Commonwealth Fellowship. We would also like to thank Dr. Abdelbagi M. Ismail to allow us to use IRRI facilities for raising reciprocal population.

REFERENCES

Abdullah Z, Khan MA, Flowers TJ. 2001. Causes of Sterility in Seed Set of Rice under
Salinity Stress. Journal of Agronomy and Crop Science **187**, 25-32.
Akbar M, Yabuno T, Nakao S. 1972. Breeding for Saline-resistant Varieties of Rice : I.
Variability for Salt Tolerance among Some Rice Varietles. Japanese Journal of Breeding **22**,
277-284.
Akbari M, Wenzl P, Caig V, Carling J, Xia L, Yang S, Uszynski G, Mohler V, Lehmensiek
A, Kuchel H, Hayden MJ, Howes N, Sharp P, Vaughan P, Rathmell B, Huttner E, Kilian A.
2006. Diversity arrays technology (DArT) for high-throughput profiling of the hexaploid wheat
genome. Theoretical and Applied Genetics **113**, 1409-1420.
Alexa A, Rahnenfuhrer J. 2019. topGO: Enrichment Analysis for Gene Ontology.
Amin USM, Biswas S, Elias SM, Razzaque S, Haque T, Malo R, Seraj ZI. 2016. Enhanced
Salt Tolerance Conferred by the Complete 2.3 kb cDNA of the Rice Vacuolar Na+/H+
Antiporter Gene Compared to 1.9 kb Coding Region with 5′ UTR in Transgenic Lines of Rice.
Frontiers in Plant Science **7**.
Bose J, Munns R, Shabala S, Gilliham M, Pogson B, Tyerman SD. 2017. Chloroplast
function and ion regulation in plants growing on saline soils: lessons from halophytes. Journal of
Experimental Botany **68**, 3129-3143.
Broman KW, Gatti DM, Simecek P, Furlotte NA, Prins P, Sen Ś, Yandell BS, Churchill
GA. 2019. R/qtl2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data
and Multiparent Populations. Genetics **211**, 495-502.
Broman KW, Wu H, Sen Ś, Churchill GA. 2003. R/qtl: QTL mapping in experimental crosses.
Bioinformatics **19**, 889-890.
Burstin J, Marget P, Huart M, Moessner A, Mangin B, Duchene C, Desprez B, Munier-
Jolain N, Duc G. 2007. Developmental Genes Have Pleiotropic Effects on Plant Morphology
and Source Capacity, Eventually Impacting on Seed Protein Content and Productivity in Pea.
Plant Physiology **144**, 768-781.
Cheng L, Wang Y, Meng L, Hu X, Cui Y, Sun Y, Zhu L, Ali J, Xu J, Li Z. 2011.
Identification of salt-tolerant QTLs with strong genetic background effect using two sets of
reciprocal introgression lines in rice. Genome **55**, 45-55.
Dasgupta S, Hossain MM, Huq M, Wheeler D. 2014. Climate Change, Soil Salinity, and the
Economics of High-Yield Rice Production in Coastal Bangladesh. Policy Research working
paper **no. WPS 7140**.
De Leon TB, Linscombe S, Gregorio G, Subudhi PK. 2015. Genetic variation in Southern
USA rice genotypes for seedling salinity tolerance. Frontiers in Plant Science **6**, 374.
Dechaine JM, Brock MT, Weinig C. 2014. QTL architecture of reproductive fitness characters
in Brassica rapa. BMC Plant Biology **14**, 1-12.
Doyle JJ, Doyle JL. 1990. Isolation of plant DNA from fresh tissue. Focus **12**, 13-15.
García Morales S, Trejo-Téllez LI, Gómez Merino FC, Caldana C, Espinosa-Victoria D,
Herrera Cabrera BE. 2012. Growth, photosynthetic activity, and potassium and sodium
concentration in rice plants under salt stress. Acta Scientiarum. Agronomy **34**, 317-324.
Gregorio GB, Senadhira D. 1993. Genetic analysis of salinity tolerance in rice (Oryza sativa
L.). Theoretical and Applied Genetics **86**, 333-338.
Gregorio GB, Senadhira D, Mendoza RD. 1997. Screening rice for salinity tolerance.
International Rice Research Institute discussion paper series.
Haque SA. 2006. Salinity Problems and Crop Production in Coastal Regions of Bangladesh.
Pakistan Journal of Botany **38**, 1359-1365.

Haque T. 2019.

Heenan D, Lewin L, McCaffery D. 1988. Salinity tolerance in rice varieties at different growth
stages. Australian Journal of Experimental Agriculture **28**, 343-349.

Hossain H, Rahman MA, Alam MS, Singh RK. 2015. Mapping of Quantitative Trait Loci
Associated with Reproductive-Stage Salt Tolerance in Rice. Journal of Agronomy and Crop
Science **201**, 17-31.

Iida S, Yamada A, Amano M, Ishii J, Kadono Y, Kosuge K. 2007. Inherited maternal effects
on the drought tolerance of a natural hybrid aquatic plant, Potamogeton anguillanus. Journal of
Plant Research **120**, 473-481.

Inês PS, Sónia N, Margarida OM, D. PM. 2015. Comprehensive phenotypic analysis of rice
(Oryza sativa) response to salinity stress. Physiologia Plantarum **155**, 43-54.

IRRI. 2002. Standard evaluation system for rice.

Joseph B, Corwin JA, Li B, Atwell S, Kliebenstein DJ. 2013a. Cytoplasmic genetic variation
and extensive cytonuclear interactions influence natural variation in the metabolome. eLife **2**,
e00776.

Joseph B, Corwin JA, Züst T, Li B, Iravani M, Schaepman-Strub G, Turnbull LA,
Kliebenstein DJ. 2013b. Hierarchical Nuclear and Cytoplasmic Genetic Architectures for Plant
Growth and Defense within Arabidopsis. The Plant Cell **25**, 1929-1945.

Kavitha PG, Miller AJ, Mathew MK, Maathuis FJM. 2012. Rice cultivars with differing salt
tolerance contain similar cation channels in their root cells. Journal of Experimental Botany **63**,
3289-3296.

Lin HX, Zhu MZ, Yano M, Gao JP, Liang ZW, Su WA, Hu XH, Ren ZH, Chao DY. 2004.
QTLs for Na+ and K+ uptake of the shoots and roots controlling rice salt tolerance. Theoretical
and Applied Genetics **108**, 253-260.

Lisa LA, Seraj ZI, Fazle Elahi CM, Das KC, Biswas K, Islam MR, Salam MA, Gomosta
AR. 2004. Genetic variation in microsatellite DNA, physiology and morphology of coastal saline
rice (Oryza sativa L.) landraces of Bangladesh. Plant and Soil **263**, 213-228.

Lovell JT. qtlTools.

Lovell JT, Mullen JL, Lowry DB, Awole K, Richards JH, Sen S, Verslues PE, Juenger TE,
McKay JK. 2015. Exploiting Differential Gene Expression and Epistasis to Discover Candidate
Genes for Drought-Associated QTLs in Arabidopsis thaliana. The Plant Cell **27**, 969-983.

Lutts S, Kinet JM, Bouharmont J. 1995. Changes in plant response to NaCl during
development of rice (Oryza sativa L.) varieties differing in salinity resistance. Journal of
Experimental Botany **46**, 1843-1852.

Moison M, Roux F, Quadrado M, Duval R, Ekovich M, Lê D-H, Verzaux M, Budar F.
2010. Cytoplasmic phylogeny and evidence of cyto-nuclear co-adaptation in Arabidopsis
thaliana. The Plant Journal **63**, 728-738.

Monroe JG, Allen ZA, Tanger P, Mullen JL, Lovell JT, Moyers BT, Whitley D, McKay JK.
2017. TSPmap, a tool making use of traveling salesperson problem solvers in the efficient and
accurate construction of high-density genetic linkage maps. BioData Mining **10**, 38.

Moradi F, Ismail A, Gregorio G, Egdane J. 2003. Salinity tolerance of rice during
reproductive development and association with tolerance at the seedling stage. Indian Journal of
Plant Physiology **8**.

Munns R, Tester M. 2008. Mechanisms of Salinity Tolerance. Annual Review of Plant Biology
59, 651-681.

Noor AUZ, Nurnabi Azad Jewel GM, Haque T, Elias SM, Biswas S, Rahman MS, Seraj ZI.
2019. Validation of QTLs in Bangladeshi rice landrace Horkuch responsible for salt tolerance in
seedling stage and maturation. Acta Physiologiae Plantarum **41**, 173.

Pearson GA, Bernstein L. 1959. Salinity effects at several growth stages of rice. Agronomy
Journal **51**, 654–657.

Quarrie S, Pekic Quarrie S, Radosevic R, Rancic D, Kaminska A, Barnes J, Leverington
M, Ceoloni C, Dodig D. 2006. Dissecting a wheat QTL for yield present in a range of
environments: from the QTL to candidate genes. Journal of Experimental Botany **57**, 2627-2637.

Rahman MA, Thomson MJ, Shah-E-Alam M, de Ocampo M, Egdane J, Ismail AM. 2016.
Exploring novel genetic sources of salinity tolerance in rice through molecular and physiological
characterization. Annals of Botany **117**, 1083-1097.

Rao PS, Mishra B, Gupta SR, Rathore A. 2008. Reproductive stage tolerance to salinity and
alkalinity stresses in rice genotypes. Plant Breeding **127**, 256-261.

Razzaque S, Elias SM, Haque T, Biswas S, Jewel GMNA, Rahman S, Weng X, Ismail AM,
Walia H, Juenger TE, Seraj ZI. 2019. Gene Expression analysis associated with salt stress in a
reciprocally crossed rice population. Scientific Reports **9**, 8249.

Razzaque S, Haque T, Elias SM, Rahman MS, Biswas S, Schwartz S, Ismail AM, Walia H,
Juenger TE, Seraj ZI. 2017. Reproductive stage physiological and transcriptional responses to
salinity stress in reciprocal populations derived from tolerant (Horkuch) and susceptible (IR29)
rice. Scientific Reports **7**, 46138.

Ren Z-H, Gao J-P, Li L-G, Cai X-L, Huang W, Chao D-Y, Zhu M-Z, Wang Z-Y, Luan S,
Lin H-X. 2005. A rice quantitative trait locus for salt tolerance encodes a sodium transporter.
Nature Genetics **37**, 1141.

Rieseberg LH, Archer MA, Wayne RK. 1999. Transgressive segregation, adaptation and
speciation. Heredity **83**, 363-372.

Roy SJ, Negrão S, Tester M. 2014. Salt resistant crop plants. Current Opinion in Biotechnology
26, 115-124.

Sabouri H, Rezai AM, Moumeni A, Kavousi A, Katouzi M, Sabouri A. 2009. QTLs mapping
of physiological traits related to salt tolerance in young rice seedlings. Biologia Plantarum **53**,
657-662.

Sexcion FSH, Egdane JA, Ismail AM, Dionisio-Sese ML. 2009. Morpho-physiologicall traits
associated with tolerance of salinity during seedling stage in rice (*Oryza sativa* L.). Phil J Crop
Sci **34**, 27-37.

Shi J, Li R, Qiu D, Jiang C, Long Y, Morgan C, Bancroft I, Zhao J, Meng J. 2009.
Unraveling the Complex Trait of Crop Yield With Quantitative Trait Loci Mapping in Brassica
napus. Genetics **182**, 851-861.

Singh RK, Flowers TJ. 2010. Physiology and Molecular Biology of the Effects of Salinity on
Rice. *Handbook of Plant and Crop Stress,Third Edition*: CRC Press, 899-939.

Soltani A, Kumar A, Mergoum M, Pirseyedi SM, Hegstad JB, Mazaheri M, Kianian SF.
2016. Novel nuclear-cytoplasmic interaction in wheat (Triticum aestivum) induces vigorous
plants. Functional & Integrative Genomics **16**, 171-182.

Tang Z, Yang Z, Hu Z, Zhang D, Lu X, Jia B, Deng D, Xu C. 2013. Cytonuclear epistatic
quantitative trait locus mapping for plant height and ear height in maize. Molecular Breeding **31**,
1-14.

Tester M, Davenport R. 2003. Na+ Tolerance and Na+ Transport in Higher Plants. Annals of
Botany **91**, 503-527.

Thomson MJ, de Ocampo M, Egdane J, Rahman MA, Sajise AG, Adorada DL, Tumimbang-Raiz E, Blumwald E, Seraj ZI, Singh RK, Gregorio GB, Ismail AM. 2010. Characterizing the Saltol Quantitative Trait Locus for Salinity Tolerance in Rice. Rice **3**, 148-160.

Tian L, Tan L, Liu F, Cai H, Sun C. 2011. Identification of quantitative trait loci associated with salt tolerance at seedling stage from Oryza rufipogon. Journal of Genetics and Genomics **38**, 593-601.

Wang Z, Cheng J, Chen Z, Huang J, Bao Y, Wang J, Zhang H. 2012. Identification of QTLs with main, epistatic and QTL × environment interaction effects for salt tolerance in rice seedlings under different salinity conditions. Theoretical and Applied Genetics **125**, 807-815.

Xiao J, Li J, Grandillo S, Ahn SN, Yuan L, Tanksley SD, McCouch SR. 1998. Identification of Trait-Improving Quantitative Trait Loci Alleles From a Wild Rice Relative, Oryza rufipogon. Genetics **150**, 899-909.

Xue W, Xing Y, Weng X, Zhao Y, Tang W, Wang L, Zhou H, Yu S, Xu C, Li X, Zhang Q. 2008. Natural variation in Ghd7 is an important regulator of heading date and yield potential in rice. Nat Genet **40**, 761-767.

Yesmin N, Elias SM, Rahman MS, Haque T, Mahbub Hasan AKM, Seraj ZI. 2014. Unique Genotypic Differences Discovered among Indigenous Bangladeshi Rice Landraces. International Journal of Genomics **2014**, 11.

Zang J, Sun Y, Wang Y, Yang J, Li F, Zhou Y, Zhu L, Jessica R, Mohammadhosein F, Xu J, Li Z. 2008. Dissection of genetic overlap of salt tolerance QTLs at the seedling and tillering stages using backcross introgression lines in rice. Science in China Series C: Life Sciences **51**, 583-591.

Zeng L, Shannon MC. 2000. Salinity Effects on Seedling Growth and Yield Components of Rice. **40**, 996-1003.

Zheng H, Wang J, Zhao H, Liu H, Sun J, Guo L, Zou D. 2015. Genetic structure, linkage disequilibrium and association mapping of salt tolerance in japonica rice germplasm at the seedling stage. Molecular Breeding **35**, 152.

785   Table1: Descriptive statistics of phenotypes measured at seedling and reproductive stages under salinity stress.

	Phenotypes	Abbreviation	Population		Parent	
			Mean	SD	*Horkuch*	*IR29*
Seedling Stage	Shoot Relative Water Content (%)	SRWC	70.06	5.07	67.72	62.43
	Standard Evaluation System (number score)	SES	5.83	0.99	**5.60**	**7.80**
	Shoot Length (cm)	SL	38.09	4.64	**41.78**	**21.35**
	Root Length (cm)	RL	10.71	2.15	**11.51**	**7.60**
	Total Chlorophyll ***(mg chl per gram fresh weight)	Tchlr	5.04	0.92	**5.31**	**3.38**
	Total Sodium *** (mmol/g dry wt)	TNa	3.07	0.77	**3.08**	**6.23**
	Total Potassium *** (mmol/g dry wt)	TK	0.38	0.08	**0.38**	**0.21**
	Potassium by Sodium (ration) **	K/Na	0.14	0.04	**0.13**	**0.04**
Reproductive Stage	Plant Height (cm)	PH	104.5	17.61	**129.81**	**75.50**
	Panicle Exsertion (%) ***	PE	99.24	1.42	100.00	98.81
	Total Tiller*** (number)	TT	4.66	1.16	5.07	4.36
	Effective Tiller (number)	ET	3.43	0.8	**3.87**	**3.23**
	Filled Grain Number (number) ***	FGN	140.87	92.96	156.29	149.85
	Filled Grain Weight (gm) ***	FGW	2.85	1.95	**3.91**	**2.00**
	Spikelet Fertility (%) ***	SF	48.63	15.93	**49.4**	**45.08**
	Days to Flower (day) ***	DF	72.08	9.83	**101.93**	**66.60**
	Harvest Index***(ratio)	HI	0.26	0.12	0.23	0.32

787   Asterisks indicates significant effect of cytoplasm (*** for $P < 0.001$, ** for $P < 0.01$ and * for $P < 0.05$). For the
788   parent's column, trait values in bold character indicate significant difference between the group means of two
789   parents (p-value < 0.05)

793   \

Table 2: QTL models

Phenotypes	QTL model	Chr	Position	LOD	Lower CI	Upper CI	Positive Allele
SL	qSL.1@183	1	183.00	13.42	175.11	190.57	Horkuch
SL	qSL.3@218	3	218.00	4.82	212.12	236.45	Horkuch
SL	qSL.5@160	5	160.43	3.83	102.86	170.27	Horkuch
RL	qRL.2@167	2	167.00	10.67	161.14	176.67	Horkuch
TK	qTK.2@45* Cyto	2	45.00	6.11	24.05	66.99	Horkuch†
TK	qTK.3@204* Cyto	3	203.78	6.46	194.44	209.29	Horkuch†
PH	qPH.1@215	1	215.00	5.59	175.11	222.54	Horkuch
PH	qPH.3@211	3	211.02	5.15	203.78	272.38	Horkuch
PH	qPH.5@144 * Cyto	5	144.00	6.64	124.73	170.27	Horkuch†
ET	qET.7@97 * Cyto	7	97.00	5.82	85.83	104.41	Horkuch
FGN	qFGN.10@58 * Cyto	10	58.48	7.72	50.30	107.07	IR29
FGW	qFGW.10@58 * Cyto	10	58.48	9.13	50.30	107.07	IR29
SF	qSF.10@59 * Cyto	10	59.00	7.71	50.30	107.07	IR29
HI	qHI.10@104+ Cyto	10	103.75	8.48	50.30	107.07	IR29

796   Each QTL model was built by linear mixed model using kinship matrix as a covariate. *cytoplasm denotes

797   interaction whereas (+) sign denotes only additive cytoplasmic effect in the QTL model. † denotes to QTL that has

798   both main and interaction effect therefore only considering main effect direction can be misleading

Figure 1: Frequency distribution of traits showing transgressive segregation in the F_2 population

and the individual subsets of cross directions. Blue and orange histograms indicate samples from

Horkuch♀ and *IR29*♀ cytoplasm respectively. Curves in blue and orange indicate distribution

plots of *Horkuch*♀ and *IR29*♀ cytoplasm respectively and dotted curve in black indicates the

distribution plot of total population. Parental values are marked by a dotted vertical line where

blue indicates *Horkuch* and orange indicates *IR29*.

Figure 2: PCA on trait correlations in the F_2 mapping population. Each point represents the

genetic means of each F_2 family whereas the shape of point indicates the cytoplasm (cross

direction). Direction of variation for axis 1 and 2 of each trait has been plotted as arrow and are

color labelled depending on two different treatment stages: green indicates Seedling stage

treatment and red indicates reproductive stage treatment. Labels of traits are printed close to the

arrow-head. Small insert-plot at the bottom-right shows the correlations of traits where brown

color shows positive correlation and light-blue indicates negative correlation. Traits labelled with

green color indicates seedling stage ones and red indicates reproductive stage traits.

Figure 3: Illustration of QTL across chromosomes. QTL are denoted as a point and 1.5 LOD

drop confidence intervals extended to a true marker is indicted by the bar for each QTL. Peaks of

the QTL were marked as black line in QTL intervals. QTL from same trait are marked with

same color. Line width represents the magnitude of LOD score. Genomic regions that showed

significant association with cytoplasm are marked here with black line segment.

Figure 4: Interaction plots of allelic effect of QTL and cytoplasm on different traits from two

different treatment stages. Blue line shows plants with *Horkuch* cytoplasm whereas orange line

indicates plants with *IR29* cytoplasm. Alleles are plotted on x-axis where AA, AB and BB

indicate homozygous *Horkuch*, heterozygous of *Horkuch*/*IR29* and homozygous *IR29*

respectively. Allelic means +/- SE are reported. Representative QTL effects for SL and PH are

presented in the upper panel and exhibit no significant interaction with cytoplasm. The third plot

from the left on upper panel demonstrates significant additive effects of the maternal cytoplasms

on TK. In the bottom panel, plot two and three from the left demonstrate significant interaction

of QTL alleles with cytoplasm for traits such FGW, ET.