Defining Strawberry Uniformity using 3D Imaging and Genetic Mapping

Bo Li[1,2#*], Helen M. Cockerton[1*], Abigail W. Johnson[1], Amanda Karlström[1], Eleftheria Stavridou[1], Greg Deakin[1], Richard J. Harrison[1,3]

[1]NIAB EMR, East Malling, Kent, ME19 6BJ, United Kingdom
[2]University of the West of England, Bristol, BS16 1QY, United Kingdom
[3]NIAB, Huntingdon Road, Cambridge, CB3 0LE, United Kingdom

Bo2.li@uwe.ac.uk
Helen.cockerton@emr.ac.uk
Abi.johnson@emr.ac.uk
Amanda.karlstrom@emr.ac.uk
Eleftheria.stavridou@emr.ac.uk
Greg.deakin@emr.ac.uk
Richard.harrison@niab.com

Corresponding author: bo2.li@uwe.ac.uk. Tel: +44(0)1173281117

* Authors contributed equally.

Abstract

Strawberry uniformity is a complex trait, influenced by multiple genetic and environmental components. To complicate matters further, the phenotypic assessment of strawberry uniformity is confounded by the difficulty of quantifying geometric parameters 'by eye' and variation between assessors. An in-depth genetic analysis of strawberry uniformity has not been undertaken to date, due to the lack of accurate and objective data. Nonetheless, uniformity remains one of the most important fruit quality selection criteria for the development of a new variety. In this study, a 3D-imaging approach was developed to characterise berry uniformity. We show that circularity of the maximum circumference had the closest predictive relationship with the manual uniformity score. Combining five or six automated metrics provided the best predictive model, indicating that human assessment of uniformity is highly complex. Furthermore, visual assessment of strawberry fruit quality in a multi-parental QTL mapping population has allowed the identification of genetic components controlling uniformity. A "regular shape" QTL was identified and found to be associated with three uniformity metrics. The QTL was present across a wide array of germplasm, indicating a strong candidate for marker-assisted breeding. A greater understanding of berry uniformity has been achieved through the study of the relative impact of automated metrics on human perceived uniformity. Furthermore, the comprehensive definition of strawberry uniformity using 3D imaging tools has allowed precision phenotyping, which has improved the accuracy of trait quantification. This tool has allowed us to illustrate the use of advanced image analysis towards the breeding of greater uniformity in strawberry.

Key Words

3D imaging, uniformity, achene, carpel-viability, breeding, fruit development

Introduction

Strawberries (*Fragaria x ananassa*) are not true fruits. The red fleshy pseudocarp of a strawberry is formed from a swollen flower base or receptacle. The true fruits are, in fact, the achenes which develop from a whorl of carpels and together form an aggregate-accessory fruit. The viability of both carpels and pollen play an important role in the resulting uniformity of berries [1]. Carpel position, density and viability dictate the shape, size and uniformity of a strawberry. Indeed, strawberry breeders have selected for high carpel densities in order to produce larger fruits [1]. Simple, classical studies

which remove all or part of the achenes from undeveloped pseudocarps has led to a
cessation in the auxin "swelling signal" in the area beneath each achene and thus
uneven fruit development [2]. In a similar fashion to achene removal, uneven pollination
of the carpels, or absence of achene development, are the main causes of uneven
pseudocarps [3]. Uneven successful pollination can be caused by damage to flowers
through high temperature, frost or precipitation [1]. A late frost in spring could lead to
carpel and other damage, resulting not only in malformation but also complete lack of
strawberry development [1]. Strawberry flowers have a variable proportion of viable
carpels and anthers between flower orders, both within a plant and also between
different cultivars [4,5]. Indeed, primary fruit are more likely to be malformed due to the
relatively lower quantities of viable anthers and pollen [6,7].
85
In spite of the environmental factors known to influence uniformity, literature has shown
that strawberry uniformity still has a large genetic component and can be improved
through breeding [8,9]. Indeed, where breeders have selected for increased uniformity
within and among berries, improvements in uniformity were observed over time [8].
Cultivars have been shown to differ in their susceptibility to misshapen fruit, indicating a
significant genetic component controlling uniformity [1,8,10]. For example, 'Florida Elyana'
is susceptible to rain damage, disrupting carpel development and thus misshapen fruit
leading to lower market value [9], similarly 'Camerosa' has been noted as a cultivar
which is particularly susceptible to misshapen fruit with ~4% of yields lost as a result of
misshapes [10,11]. By contrast, 'Florida Radiance' has high marketable yields and does
not exhibit a high proportion of misshapen fruits [9]. Breeders can influence the proportion
of uniform strawberries through selecting- be it directly or indirectly- for 1) even
allocation of viable carpels across the receptacle within the flower 2) ready access to
pollen within flowers and 3) high fertility of carpels ensuring even successful pollination.
100
Strawberry is an important fruit crop with a global market revenue of 21,171 million
USD in 2015 [12]. Producing visually appealing strawberry fruit is one of the primary
objectives in a strawberry breeding program [13]. Shape uniformity is an essential trait of
strawberry fruits due to the direct association with product quality and value [14].
Increasing the uniformity of berries can increase the proportion of marketable fruit as
berry irregularity is one of the primary imperfections leading to culling and reduced
marketable yield [8].
108
As there is no well-defined strawberry phenotyping guidelines for fruit uniformity, the
current system at NIAB EMR relies on visual assessments, which are subjective and
laborious. Unlike morphological traits such as length, volume and colour, which can be
accurately measured manually in a low-throughput manner, uniformity assessment is

extremely subjective. As there is no quantitative method of generating phenotypic data
for uniformity, the genetic determinants of strawberry uniformity are still unknown.

Computer vision has shown great potential to quantify external fruit quality and 2D
imaging has been successfully implemented to measure the shape and size of fruits
such as strawberries [15], apples [16], watermelon [17], cherries [18] and mangos [19]. Basic
shape traits such as length, width, aspect ratio and volume, and more sophisticated
traits such as elliptic Fourier descriptors [20] have been quantified and used to describe
variation in fruit quality. 3D imaging has been successfully used for phenotyping the
crop canopy [21,22] and root architecture [23,24], and a 3D strawberry phenotyping platform
has been explored in our previous study [25]. With the 3D point cloud reconstructed based
on the Structure from Motion (SfM) method [26], basic size-related parameters have been
measured in three dimensions allowing volume estimation with high accuracy [27].
Compared with shape and size evaluation, uniformity is a multi-dimensional trait,
therefore it is not possible to quantify through 2D image analysis with a single viewing
angle. The application of 3D image analysis for phenotyping the external qualities of
fruit has not been sufficiently explored, and the basic, previously characterised, shape-
and size-related parameters are not adequate for understanding uniformity.

Here the application of a 3D phenotyping platform allows us to investigate the genetic
basis of strawberry uniformity. The 3D image analysis software leverages the previously
developed platform [25] in order to define eight new external variables and investigate
their importance on manual uniformity assessment. This method was applied to a multi-
parental strawberry mapping population in order to quantify the genetic components
underpinning strawberry uniformity.

139 Materials and methods

Plant material and experimental set-up
A multi-parental strawberry population was generated through crossing 26 diverse
cultivars and breeding lines to create a population of 416 genotypes made up of 26
families each containing 16 individuals (Suppl. Figure 1). Progenitors were selected to
represent diversity across multiple fruit quality traits. Twelve replicate runner plants
were pinned down from each genotype into 9 cm square pots containing compost.
Clonal plants were separated from parental plants and then placed in cold storage (-2
$^{\circ}$C) until the start of the experiment. Plants were potted into 2 L pots containing coir and
fertigated at 1kg L^{-1} (rate: 10 seconds every 45 min) using Vitex Vitafeed (N:P:K,
176:36:255). Blocks were horizontal intersections across the polytunnels. Due to the
large scale of the experiment, replicate blocks were set up at three week intervals. A
Natupol Koppert bumble bee hive was added into each polytunnel to assist even

pollination. Strawberries were picked when ripe into egg boxes. Boxes were labeled
with QR codes to assist tracking of genotypes. Strawberry uniformity was scored on a
scale from 1 (irregular) to 9 (uniform) with extensive training provided for all assessors.
Strawberry shape was allocated into 9 categories: globose, globose-conic, conic, long-
conic, bi-conic, conic-wedge, wedge, square and miscellaneous. Manual uniformity
scores were recorded in the field book app [28], the QR scanning feature allowed quick
access to the correct entry form.
159
Genotyping and Linkage map
DNA was extracted for each genotype from unopened leaflets using the Qiagen DNeasy
plant mini extraction kit. Genotyping was conducted using the Axiom® IStraw35 384HT
array [29] (i35k). Crosslink was used to generate linkage maps- a program developed
specifically for polyploid plant species [30]. The map orders from 5 populations were
combined to make the consensus map as detailed in the study of Harrison et al [31].
Fragaria × ananassa chromosome number is denoted by 1-7 and the sub-genome
number is represented by A-D as specified in [32].
168
3D reconstruction
The 3D imaging platform was a modified version of that developed by He et al. [25].
Strawberry fruit were placed in the middle of a turntable, on a dark blue holder made by
polymeric foam (38 mm x 19 mm x 19 mm; height, length and width). Unlike the
previous study, a webcam (Logitech C920, Newark, CA, USA) was fitted at a height of
30 cm and horizontal distance of 25 cm away from the sample. QR codes on containers
were scanned through the webcams allowing tracking of berries and automated labeling
of image files. The imaging rig was placed inside a photography studio tent with
constant LED illumination. The turntable rotated at a frequency of 50 seconds per full
turn, and an image was captured every second. Six imaging platforms allowed
concurrent imaging of replicate berries. The 3D reconstruction was implemented with
Agisoft Photoscan (Agisoft, LLC, St. Petersburg, Russia), and in order to increase the
accuracy and processing speed, all images were pre-processed by cropping to a
smaller size (400 x 600 pixels). Background subtraction was achieved through arbitrary
colour thresholding. The image processing software for webcam control and automated
image pre-processing were written in C++ utilising the OpenCV Library [25,33].
185
Data processing pipeline of phenotypic traits extraction
Point cloud preprocessing
In the preprocessing stage (Fig. 1), each point cloud model was converted from the
colour space of RGB (Red, Green and Blue) to HSV (Hue, Saturation and Value).
Arbitrary thresholding on the hue channel was used to remove the noise introduced in
the reconstruction stage. The clean point cloud was translated to the origin of the 3D

coordinate system based on the distance between the moment of the point cloud and
the origin. By calculating the eigenvector associated with the largest eigenvalue of the
coordinates of points, a rotation matrix could be derived to represent the main
orientation of the point cloud, which can be used to rotate the point cloud with the main
orientation aligned with the z-axis. After rotation, the arbitrary threshold was applied
again on the hue channel in order to segment the strawberry body and blue holder from
the whole point cloud. The height of the holder was obtained by calculating the
difference between the maximum and minimum values of the holder point cloud on the
z-axis. As the original coordinate system generated by Structure from Motion (SfM)
method has an arbitrary scale, each point cloud model needed to be standardised by
the height of the holder, so that the sizes of all point clouds are comparable.

Original point cloud　　　　　Preprocessed point cloud

203
Figure 1. Point cloud pre-processing for strawberry body extraction, translation to origin
of xyz coordinate system and size standardisation.

Uniformity-related traits measurements
Eight uniformity-related traits were calculated from the point cloud data of strawberry
body after preprocessing. These are:

Coefficient of variation (CV) of side view areas (CV_A) and the ratio between
maximum and minimum side view area (Max_A/Min_A)
All side views should be identical in a perfectly uniform strawberry. In order to eliminate
the heterogeneity introduced from the calyx and the holder, only the points within the
middle 50% of the body height of each point cloud were retained for analysis (Fig. 2). In
order to understand the heterogeneity of different side views of a point cloud, each point
cloud was rotated along the z-axis by 3.6° for 99 rotations, and after each rotation, the

side view of the point cloud was projected onto the x-z plane in 2D (labelled in white). A convex hull was fitted to each projected image and the contour area was calculated. For area metrics, two traits were obtained; the CV of side view areas (CV_A) and the ratio between the maximum and minimum area (Max_A/Min_A). An ideal uniform strawberry will have a value of zero for CV_A and one for Max_A/Min_A.

CV of principal orientations (CV_D)

The major eigenvector indicating the main orientation was calculated by principal component analysis (PCA) for all 100 side view projected images, and the heterogeneity of the orientations of the projected images was quantified by calculating the CV of angles of the main orientation. Like CV_A, a perfectly uniform strawberry will have a value of zero for CV_D.

Figure 2. Side view of strawberry body for the CV measurement of the area and principal orientations. Convex hulls are outlined in blue, and blue and green arrows indicate the principal orientations (a). Extraction of example slice images horizontal to x-

235 y plane at the height of 20%, 40%, 60% and 80% of the total height. A minimum
236 bounding box is fitted to each slice image (b). Sixteen patches of points labelled in
237 different colours for curvature estimation (c).
238

Aspect ratio of the minimum bounding box (L/W)
A lateral slice image was obtained by identifying the intersection between the plane in
parallel with the x-y plane and point cloud (Fig. 2b). Based on the values on the z-axis,
100 evenly spaced slice images were obtained. The slice image with the largest contour
was obtained by calculating the contour area of the convex hulls for all slice images.
The main orientation of the contour was indicated by the major eigenvector of the PCA
and the minimum bounding box was fitted to the slice images. The ratio between the
length and width of the largest bounding box was derived and the ratio should be one
for a perfectly uniform fruit.
248

Circularity of the maximum circumference (CIR)
250 Visually, the circularities of the contours in horizontal slice images are high if the
251 strawberry is uniform. Circularity (CIR) was calculated as previously described [34]:
252

$$C = \frac{4\pi A}{p^2}$$

254 Where A and p are the area and perimeter of the convex hull respectively. For each
255 point cloud, the circularity was calculated for the slice image with largest contour area.
256

Straightness of centre axis (STR)
258 The coordinates of the centroids for each horizontal slice image can be located by
259 calculating the moment of the contour. The centroids can be connected as a straight
260 line for a uniform strawberry. The centroids were calculated for all the slice images
261 within the middle 80% of the body height, and the straightness of the central axis was
262 characterised by:
263

$$STR = \frac{\sum_{i=1}^{N-1} d_i}{D}$$

265 Where N (N = 80) is the number of slice images used for the analysis, d_i is the
266 Euclidean distance between neighbouring slice images, and D is the Euclidean distance
267 between the centroids of the top and bottom slice images.
268

CV of curvature and the ratio between maximum and minimum curvatures (Max_C/Min_C)
The principal curvature can be calculated for each point in the point cloud, which
describes how much the curve deviates from a straight line at this point. It can be
imagined that the 3D curve can be sliced orthogonally around the direction of normal in

to 2D curve, and the maximum curvature k_1 and minimum curvatures k_2 are the two
principal curvatures for the 3D curve [35]. The average curvature k, which is defined as
the mean value of the magnitudes of principal curvatures in the two main directions was
applied to quantify the curvature for a given point.
278
As the curvature measurement is sensitive to noise, the point cloud surface of
strawberry body was first smoothed by using Moving Least Squares (MLS) method [36],
which could reconstruct a smooth surface from the noisy point cloud. Sixteen patches of
the points were selected evenly from the points forming the largest slice in parallel with
x-y plane (Fig. 2c). For each patch, the first half largest curvatures were averaged and
used to represent the curvature of the patch. With the curvatures of all 16 patches, the
CV of curvature (CV_C) and the ratio between maximum and minimum curvatures
(Max_C/Min_C) were calculated.
287
288
Statistical analysis
Ordinal regression
Statistical analysis was performed using R (version 3.5.1) and the Genstat statistical
package (Version 13.0, VSN International Ltd. England). Differences in uniformity traits
within each shape groups were distinguished using ANOVA and Tukey post-hoc test.
Pearson coefficients of correlation were calculated between all proposed uniformity-
related traits. As the group labels are ordinal dependent variables, ordinal regression
was used to evaluate the performances of all traits [37]. Model fit was ascertained by
using selection criterion values based on the Akaike Information Criterion (AIC) and the
Bayesian Information Criterion (BIC). In general, a better model fit generates lower
values for both AIC and BIC [38]. In order to identify the optimal variable combination
related to manual assessment, stepwise AIC and BIC methods were applied [39]. The
most significant variable was identified by comparing the criterion values of all models.
Other variables were added successively and retained if the model fit was improved.
303
304 **Genetic Analysis**
305 The best linear unbiased estimate (BLUE) was calculated for all genotypes in order to
306 correct for the influence of assessor, data and block. Linear mixed-effects models were
307 generated for each phenotypic trait with and without covariates. Grand scores for each
308 genotype were calculated using mixed models to account for significant covariates.
309
Composite interval mapping
Multiparental QTL mapping was conducted in R using package "mppR" [40]. A
permutation test determined the significance threshold [41]. A two-step QTL analysis was
implemented: the selection of cofactors was achieved through Simple Interval Mapping

(SIM) proceeded by a multi-QTL model search using composite interval mapping (CIM)
[42,43]. As a multi-parent population CIM works on parent relationships. Therefore the
'CPEM0162' x 'Rumba' cross was removed as it is not directly related through the
parental cultivar network. All other crosses were interrelated and formed a single
network (Sup. Figure 1).
319

Results

Characterisation of uniformity-related traits
All the uniformity-related traits were calculated based on the point cloud, the mean
values and the standard errors for each visual uniformity class are presented in Figure
3. ANOVA results showed that significant differences were observed between uniformity
classes for all traits ($p < 0.001$). The Pearson's linear correlation coefficients were
calculated between all traits, and strong correlations were found amongst
Max_A/Min_A, L/W and CIR (Table 1).
328

Ordinal regression models were constructed for all variables and each variable
independently. L/W was not significant due to the high correlation with other variables
and CIR showed the best model fit with the lowest AIC and BIC values (Table 2). New
variables were added sequentially to the model until no further improvement of the
criterion value was observed. The AIC and BIC based stepwise selection methods
showed inconsistent results (Table 3). The AIC based method showed the optimal
criterion value with all variables except L/W and Max_C/Min_C, but BIC based method
showed that STR could not improve the model fit.
337

The influence of Shape on uniformity
The shape of a strawberry influences the uniformity trait score. Bi-conic strawberries
were seen to have high uniformity based on the area overlap measures (CV_A &
Max_A/Min_A), L/W and CIR scores indicating bi-conic strawberries have consistently
circular horizontal transects at the mid point. Whereas for curvature uniformity
measures (CV_C & Max_C/Min_C) globose fruit are the most uniform and
miscellaneous fruit the least (Data not shown). Both the manual uniformity score and
CIR could discriminate miscellaneous shapes from the other shape categories (Figure
4).
347

QTL identification
A total of 28 QTL were found to be associated with uniformity traits (Table 4). Of which
25 were detected in more than one progenitor (Figure 5). Five focal SNP's, on
chromosome 2B and 5D were found to represent more than one trait (Table 4, Figure
6). The same focal SNP AX.166521303 was identified as important region in

Max_A/Min_A, CV_A and CIR uniformity traits. Global adjusted R^2 values for linear
models were between 5.07 and 32.15 indicating the proportion of variation explained by
identified QTLs (Table 5). All uniformity traits apart from CV_A were significantly
affected by date of picking. CV_D had the largest broad sense heritability score of 38.4
(Table 5).

Figure 3. Mean value and standard error of calculated uniformity-related traits by the
newly developed 3D image analysis software against defined uniformity scale based on
manual assessment.

363
364 **Figure 4.** CIR scores for each manually classified strawberry shape category. Letters
365 denote significant differences between categories. Error bars are standard errors of the
366 mean.

367
368 **Table 1**. Pearson's linear correlation coefficients among all uniformity-related traits. All
369 values are significant at $p < 0.05$ level.

	CV_A	Max_A/Min_A	CV_D	L/W	CIR	STR	CV_C	Max_C/Min_C
CV_A	1.00							
Max_A/Min_A	0.61	1.00						
CV_D	0.13	0.29	1.00					
L/W	0.54	0.90	0.27	1.00				
CIR	-0.48	-0.85	-0.32	-0.85	1.00			
STR	0.12	0.26	0.06	0.26	-0.29	1.00		
CV_C	0.21	0.24	0.07	0.23	-0.25	0.15	1.00	

| Max_C/Min_C | 0.17 | 0.20 | 0.15 | 0.17 | -0.27 | 0.10 | 0.57 | 1.00 |

Table 2. Summary of individual ordinal models and variable significance of ordinal model with all variables, towards prediction of manual assessment uniformity scores. * p < 0.05, ** p < 0.01, *** p < 0.001.

Model	LogLik	AIC	BIC	Signif. codes
CIR	-1691.82	3401.65	3445.35	***
Max_A/Min_A	-1707.44	3432.89	3476.85	***
L/W	-1742.98	3503.96	3547.66	
CV_A	-1790.55	3599.09	3642.79	***
CV_C	-1824.20	3666.41	3710.11	***
Max_C/Min_C	-1839.63	3697.25	3740.95	*
CV_D	-1857.94	3733.87	3777.57	***
STR	-1861.85	3741.70	3785.39	*

Table 3. Model comparison values for uniformity metrics, towards prediction of manual assessment uniformity scores based on AIC and BIC.

Model	1	2	3	4	5	6	7	8
CIR	x	x	x	x	x	x	x	x
Max_A/Min_A		x	x	x	x	x	x	x
L/W			x					
CV_A				x	x	x	x	x
CV_C					x	x	x	x
Max_C/Min_C						x		
CV_D							x	x
STR								x

AIC	3401.65	3374.40	3374.29	3362.57	3302.01	3299.53	3291.87	3288.40
BIC	3445.35	3422.95	3427.70	3415.97	3360.28	3362.65	3354.99	3356.38

383 **Supplementary Table 4.** Focal SNPs representing strawberry uniformity QTL. The
384 position of QTL is reported in Mb as scaled to the vesca version 4 genome

Marker Names	Chromosome	Pos Mb	$\log^{10} p$ value	R^2	Trait
AX.89804099	3B	96.3	3.29	6.9	Visual Uniformity
AX.123357183	2D	63.1	4.07	3.81	Max_C/Min_C
AX.166513757	4B	239.6	3.03	5.05	Max_C/Min_C
AX.166523206	4D	272.0	3.89	4.03	Max_C/Min_C
AX.166514922	5D	1.8	3.02	9.9	Max_C/Min_C
AX.89886024	5D	36.8	3.32	2.87	Max_C/Min_C
AX.166525020	6B	34.4	3.49	7.59	Max_C/Min_C
AX.166526040	7B	47.5	3.07	2.71	Max_C/Min_C
AX.166521293	2B	60.7	3.74	6.41	CV_C
AX.166519032	2C	183.5	3.04	5.07	STR
AX.166521303	**2B**	**63.1**	**3.29**	**9.41**	**CIR**
AX.89791395	4B	130.4	3.17	2.71	CIR
AX.166509340	4C	241.7	3.00	2.37	CIR
AX.89890707	5C	212.2	3.11	1.86	CIR
AX.166526395	7C	228.6	3.81	3.95	CIR
AX.89873861	1A	139.7	3.73	5.27	L/W
AX.123357666	3B	303.6	3.35	3.42	L/W
AX.89829301	4D	182.3	3.03	2.06	L/W

AX.166524494	5B	103.6	3.25	3.83	L/W
AX.89791880	**5D**	**123.7**	**3.52**	**1.46**	**L/W**
AX.89792701	5D	182.0	3.61	10.64	L/W
AX.166503621	2C	266.2	3.89	2.19	CV_D
AX.166504095	3A	149.5	4.10	2.8	CV_D
AX.166525754	6C	74.7	3.25	3.58	CV_D
AX.166521303	**2B**	**63.1**	**4.55**	**15.08**	Max_A/Min_A
AX.89791880	**5D**	**123.7**	**4.22**	**3.11**	Max_A/Min_A
AX.166506309	5D	150.8	3.16	2.48	Max_A/Min_A
AX.166523943	5D	233.9	3.42	4.47	Max_A/Min_A
AX.166521303	**2B**	**63.1**	**3.86**	**6.28**	**CV_A**
AX.89788547	5D	40.9	3.61	5.8	CV_A

386
387 **Table 5.** Broad sense heritability scores for each automated uniformity trait, the
388 influence of block and date of assessment on the trait measured. The number of QTL
389 and the coefficient of determination associated with combined QTL.
390

Trait	Heritability	Significance of Block	Significance of Date	GxE	Number of QTL	R2	R2 adj
STR	21.70	***	***	***	1	5.07	4.63
CIR	22.00	***	***	***	5	21.2	19.3
CV_D	38.40	***	***	***	3	9.35	8.05
CV_A	19.95	NS	NS	***	2	12.13	11.29
CV_C	24.63	***	***	***	1	6.41	5.97
L/W	19.20	***	**	***	6	24.77	22.59
Max_C/Min_C	16.70	NS	*	NS	7	32.15	29.85

	22.70	**	***	***	4	23.1	21.62
Max_A/Min_A	22.70	**	***	***	4	23.1	21.62
Uniformity	8.00	NS	***	***	1	6.9	6.46

391
392
393

	Man Uni	CV_A		Max_A/Min_A				CV_D			L:W						CIR					STR	CV_C	Max_C/Min_C						
	AX.89800099	AX.166521303	AX.89788547	AX.166521303	AX.165906309	AX.89791880	AX.165923943	AX.166503621	AX.166504095	AX.166525754	AX.89873861	AX.123357666	AX.89829301	AX.165924494	AX.89791880	AX.89792701	AX.166521303	AX.89791395	AX.165909340	AX.89890707	AX.166526395	AX.166519032	AX.166521293	AX.123357183	AX.166513757	AX.166523206	AX.1665134922	AX.89886024	AX.165525020	AX.165526040
FA2311	1.26						-0.68														-0.49	1.01						NA	1.04	
FA2907			-0.72				-0.68			-1.56								-0.52	-0.84										-0.75	
FA0123							-0.68	NA												-0.49	-0.74									-0.80
FA0193					0.65		-0.68											-0.52	-0.84										1.04	
FA0248					0.65		-0.68														-0.74	1.01								
FA0307	1.26		-0.72				-0.68		1.18		1.11			-0.90	1.37						-0.49					0.70			-0.75	-0.40
FA0676							-0.68	2.32			2.23								-0.84	-0.49	-0.74	1.01				0.70			1.04	-0.40
FA0677			NA				-0.68		NA				-0.90					-0.52				1.01								
FA0909					0.65		-1.35														-0.49	1.01			-0.86					
FA1111																					-0.49					0.70			2.07	-0.40
FA1211	1.26																	-0.52	-0.84		-0.49					0.70			2.07	-0.40
FA1217		-1.03	-0.72	-1.73	0.65		-1.35				1.11					-2.66	1.35	-0.52									-1.16	-0.86		-0.40
FA2654											1.11									-0.49	-0.74									
FA6969	1.26	-1.03	-0.72	-1.73			-0.68				1.11	1.80				-2.66	1.35	-0.52				1.01	-1.16							-0.40
FA0508	1.26		-1.44				-0.68													-0.49	-1.47								-0.75	-0.40
FA9521			-1.44	-1.73			-0.68	NA					1.80					-1.04		-0.99	-1.47				-0.86	0.70			-0.75	-0.40
FA2907		-1.03		-1.73			-0.68						1.80				1.35	-0.52			-1.16					0.70				-0.40
FA1204	1.26				0.65		-1.35															1.01				2.47				
FA0406		-1.03	-0.72	-1.73	0.65		-1.35			-1.56							1.35				-0.49	1.01	-1.16			0.70				
FA1711			-0.72		0.65	1.25				-1.56					NA	0.82				-0.49	-0.74				-0.86	2.47	1.72		1.04	
FA0908					0.65		-1.35											NA	-0.84		-0.74	1.01								
FA1805							NA											-0.52			-0.74									
FA8752		-1.03	-0.72	-1.73	NA		-0.68			-1.56	NA					NA	1.35		-0.84	-0.49		NA	-1.16			0.70		NA	1.04	NA
M.Centenary		-1.03	-0.72	-1.73	NA	1.25	-0.68			NA						0.82	1.35	-0.52	-0.84	-0.49	-0.74	NA	-1.16			0.70		NA	1.04	NA
FA0407			-0.72		NA			2.32					1.80			NA		-0.52		-0.49		NA			-0.86			NA	NA	NA
Vibrant					NA											NA		-0.52				NA						NA	NA	NA

Figure 5. Effect sizes associated with each QTL in each of the 26 progenitors; blue colour is associated with lower uniformity and red colour is associated with higher uniformity.

400
401
402 **Figure 6.** Location of QTL on the octoploid consensus map scaled to the *Fragaria*
403 *vesca* 'version four' genome. Horizontal grey lines represent istraw 35k axiom array
404 markers
405

406
407 **Supplementary Figure 1.** A network of crosses conducted to generate the
408 multiparental mapping population used in this study. Cultivars are represented by
409 circles, families are represented by lines.
410

411
412 (a)

413
414 (b)
415

416
417 (c)
418 **Supplementary Figure 2.** Manhattan plots of **(**a) Max_A/Min_A, (b) L/W and (c) CIR.
419
420

Discussion

We report for the first time a robust method to measure strawberry uniformity and apply
this technique to generate genetic markers for uniformity traits. Several studies have
attempted to quantify strawberry fruit shape using 2D images with neural networks [44],
3D imaging [45] and by machine learning[15]. However, none of these studies investigated
berry uniformity. Unlike the aforementioned studies, who measure a relatively small
number of genotypes intensively, we have implemented a high throughput imaging
platform across a large population to facilitate genetic analysis of the trait. Although
strawberry shape has received greater attention in the literature, berry uniformity is a
more important trait for a breeder to improve (Personal communication, Abigail
Johnson).
432

In current strawberry breeding practice, there is no widely accepted criteria for quantifying uniformity due to the difficulty of defining a multidimensional trait. Here, the manual strawberry uniformity scale has been designed by NIAB EMR breeders. As such, the absence of a straightforward definition, has meant that it has not been possible to study the genetic components controlling strawberry uniformity in the past. To overcome this, we have used 3D image analysis to define the parameters underlying a breeder's perception of strawberry uniformity. The original 3D strawberry phenotyping system [25] could accurately measure basic size-related traits. In this study, the point cloud analysis software was further developed to quantify strawberry uniformity through eight proposed metrics. By comparing with the manual scale, the image processing pipeline has demonstrated an objective method of characterising strawberry uniformity components.

Quantifying berry uniformity

Circularity of the maximum circumference (CIR) of strawberries showed the best predictive ability for manual uniformity scores based on the ordinal regression model fit, when studying individual variables alone. A completely misshapen fruit with a severely undulating fruit surface will score a value of 1 for manual assessments, and these completely misshapen fruits were the easiest category to identify by eye, as they were clearly distinct from regular shapes. A low CIR value appears to represent the undulating misshapen and "miscellaneous" fruit (Figure 4 & 6). Miscellaneous berries are the most undesirable fruit shape category therefore it is highly beneficial to select against them. When multiple traits are combined to describe uniformity, the best fitting model required the combination of CIR, CV_A and Max_A/Min_A, CV_D and CV_C. The five factors required for optimal model construction indicate that there are multiple uniformity components influencing the manual uniformity score.

Misshapen fruit QTL

One of the QTL represented by the focal marker AX.166521303 on chromosome 2B was found to be associated with CIR, this QTL was also associated with CV_A and Max_A/Min_A, each of which were found in the best fitting model used to describe the manual uniformity score. The focal SNP AX.166521303 was found to be present and significant in six progenitors and had an effect size of 9.14% on CIR. Therefore, this marker is a good candidate for marker assisted breeding in selection against completely mis-shapen and irregular strawberries. Furthermore, this work has highlighted a region of interest for further study to pinpoint the causative allele associated with reduced uniformity. Dissecting the contribution of genetic and environmental components believed to underpin strawberry uniformity; susceptibility to heat stress, carpel and pollen viability, achene position, size and distribution[1] may help to further elucidate the mechanism of uniformity segregating in the multiparental population.

Uniformity trait selection

The trait L/W shows little improvement on the overall combined trait model fit due to the
high correlation with other traits including Max_A/Min_A and CIR, but it was still a good
predictor of uniformity based on the model fit when studying individual variables alone.
AIC and BIC based stepwise feature selection showed disagreement on the selection of
the STR parameter. The difference between calculating AIC and BIC is that AIC does
not account for the sample size, so when sample number is large, BIC applies larger
penalty for complex models and leads to a simpler model [46]. However, this study does
not aim to identify the optimal feature combination to develop prediction model related
to manual uniformity evaluation, but develop a new image based quantification to
replace the manual scale, because the ground-truth data are subjective and as such
any large bias can reduce the robustness of model development. Moreover, the manual
scale cannot be considered a comprehensive assessment as the parameter STR
cannot be visually evaluated by eye. However, it must be said that if a trait cannot be
detected by the human eye, then it is not a valuable trait for a strawberry breeder to
select upon.
489

Limitations of the system

The 3D point cloud analysis software is independent of the imaging acquisition system,
and the uniformity-related traits can be extracted automatically in a high-throughput
manner. However, the imaging collection throughput was 50 seconds per fruit and the
3D reconstruction has to be performed separately, which limits use to pre breeding
experiments as opposed to use as a breeders tool. Due to the occlusion from the
viewing angles, the strawberry nose cannot be fully reconstructed especially for globose
shaped fruit, which decreases the accuracy of STR measurements and also limits the
study on automated shape classification. To increase the throughput and accuracy of
3D phenotyping, it is necessary to further develop the hardware with multiple cameras
to allow more viewing angles or a structured light based imaging system with a robotic
arm, and also integrate the hardware driver with 3D reconstruction software. The
current point cloud image analysis software was able to characterise many key external
traits which are important for strawberry breeding, however, the measurement of other
parameters such as achene density must be investigated through an improved
phenotyping platform in future studies.
506

Genetic Control of Fruit Quality

508 Papers detailing strawberry fruit quality QTL report genetic alleles associated with
509 multiple fruit quality traits including fresh weight, metabolites, external colour and
510 firmness [29,47,48], however, there are currently no papers which report QTL associated
511 with strawberry uniformity. Here, we provide a phenotyping platform which has
512

facilitated the assessment of the genetic components underlying strawberry uniformity
for the first time. The use of a multi-parental population has allowed the study of a
diverse set of germplasm and has ensured that resulting QTL to have a greater
relevance for breeders when compared to alleles identified in bi-parental studies.
Overall, 25 of the QTL were found to have an effect on uniformity in more than one of
the 26 progenitors indicating that there has been limited linkage decay between the
causal allele and marker, and that the relationships have been maintained across
generations. Furthermore, the QTL on chromosome 2B was observed three times
across different uniformity traits, such traits are only partially correlated and thus
describe discrete components, as such this allele can be seen to play a major role in
uniformity.
524
Genetic control of strawberry fruit shape
Unlike uniformity the mechanism controlling fruit shape has been studied extensively in
the wild strawberry; *Fragaria vesca* and may act as a surrogate model for the cultivated
octoploid strawberry *Fragaria* x *ananassa*. In *F. vesca*, fruit shape is primarily controlled
by phytohormones [49,50]. Auxin increases the width of fruit and by contrast gibberellic
acid (GA) increases the length of a strawberry whereas Abscisic acid (ABA) down
regulates both Auxin and GA and thus reduces fruit expansion [49,50]. GA deficient Vesca
mutants were found to have a "short" or globose berry shape, which, through the
application of GA to the berry, could be restored to result in a "long" or long conic fruit
shape [50].
535
It is clear that breeders wish to select for greater berry uniformity however the
confounding relationship between shape and uniformity must also be considered. For
example, square, wedge and wedge-conic strawberries may have high 2D symmetry
but not 3D symmetry. UK breeders primarily aim to select for conic or long conic fruit
whereas globose, square, wedge and miscellaneous berries are classified as
undesirable and biconic, globose-conic and conic-wedge fruit are seen as acceptable
shapes (Personal Communication, Abigail Johnson). Here we provide an objective
measure (CIR) that can be used to discriminate the least desirable berries -
miscellaneous or misshapen berries and select for regular fruit shapes.
545
Heritability of Uniformity
Broadsense heritability scores were between 16.70 and 38.40 for automated uniformity
metrics indicating a greater genetic component than that associated with manual
uniformity (8.00). These values indicate the proportion of variation segregating in the
study population, however improvement in the heritability may also be caused, in part,
by more accurate phenotypic measurements. In particular, high heritability was
observed for CV_D which indicates the angle of a strawberry related to whorl of carpels

(Figure 4) is under strong genetic control. Date of picking was seen to have a significant
impact on all uniformity metrics apart from CV_D which had a large genetic component.
The high significance of date indicates the developmental environmental conditions has
a significant impact on strawberry uniformity. Extreme temperatures were observed
during the experiment which may have caused the significance of date. All traits apart
from CV_C showed a significant genotype by environment interaction indicating that
genotypes were responding differently to heat stress. Misshapen fruit have been found
to have a greater proportion of small underdeveloped achenes following exposure
extreme temperatures during embryo development [51,52].
562
Here we provide a comprehensive dissection of the traits underlying strawberry uniformity and
show that the visual perception of a strawberry can be represented by 5 metrics. The generation
of an objective measure of uniformity has allowed the assessments of genetic components in a
multi-parental breeding population. We show uniformity has a strong genetic component that
can be improved by breeding and identify genetic components controlling uniformity that are
present across a wide array of germplasm.
569
570

Availability of data, materials and methods

572 The software developed and datasets generated and analysed during the current study
573 are available from the corresponding author on reasonable request.

Acknowledgements

The authors acknowledge project partners Soloberry, Sainsburys, Botanicoir and
Agrovista for their involvement and support of the project. The authors acknowledge
Robert Vickerstaff for generating the octoploid consensus map as part of other projects
and Dr Beatrice Denoyes, INRA and Dr Amparo Monfort, CRAG for granting the use of
their informative markers in the production of the strawberry consensus linkage map.
This work was supported by grants from the Biotechnology and Biological Sciences
Research Council (BBSRC) BB/M01200X/2, BB/P005039/1 and Innovate UK project
101914.

584
585

Authors' contributions

BL – Analysed imaging data
AJ, HMC, ES, BL, RJH - Conceived and designed experiments
HMC – Conducted quantitative genetics analysis
AJ, HMC, AK - Performed experiments
AK - Performed genotyping

GD - Ordinal regression analysis

BL, HMC & RJH wrote the manuscript with contributions from all authors.

BL and HMC equally contributed to the manuscript.

Conflict of interests

On behalf of all authors, the corresponding author states that there is no conflict of
interests regarding the publication of this work.

List of Abbreviations

CIM- composite interval mapping
CIR - Circularity
CV_A - Coefficient of Variation of side view areas
CV_C - Coefficient of Variation of curvatures
CV_D - Coefficient of Variation of principal orientations
i35k - Istraw35 Affymetrix chip
L/W - Aspect ratio of the minimum bounding box
Max_A/Min_A - ratio between maximum and minimum side view areas
Max_C/Min_C - ratio between maximum and minimum curvatures
MedR - Median number of roots
QTL- Quantitative Trait Loci
QR - Quick Response
RGB- Red Green Blue
SNP - Single Nucleotide Polymorphism
STR- Straightness of centre axis

619 Reference

1	Carew JG, Morretini M, Battey NH. Misshapen fruits in strawberry. *Small Fruits*
Review. 2003; **2**: 37–50.

2	Nitsch JP. Growth and morphogenesis of the strawberry as related to Auxin.
American Journal of Botany. 1950; **37**: 211.

3	Kronenberg HG, Braak JP, Zeilinga AE. Poor fruit setting in strawberries. II.
Euphytica. 1959; **8**: 245–251.

4	Thompson PA. Environmental effects on pollination and receptacle development in
the strawberry. *Journal of Horticultural Science*. 1971; **46**: 1–12.

5   Darrow GM. The importance of sex in the strawberry. *Journal of Heredity*. 1925; **16**:
193–204.

6   Darrow GM. The strawberry. *The Strawberry* 1966.

7   Gilbert C, Breen PJ. Low pollen production as a cause of fruit malformation in
strawberry. *J Am Soc Hortic Sci* 1987.

8   Whitaker VM, Hasing T, Chandler CK, Plotto A, Baldwin E. Historical trends in
strawberry fruit quality revealed by a Trial of University of Florida Cultivars and
Advanced Selections. *HortScience*. 2011; **46**: 553–557.

9   Chandler CK, Santos BM, Peres NA, Jouquand C, Plotto A. 'Florida Elyana'
strawberry. *HortScience*. 2009; **44**: 1775–1776.

10 Zanin DS, Fagherazzi AF, dos Santos AM, Martins R, Kretzschmar AA, Rufato L.
Agronomic performance of cultivars and advanced selections of strawberry in the
South Plateau of Santa Catarina State. *Revista Ceres*. 2019; **66**: 159–167.

11 Ariza MT, Soria C, Medina-Mínguez JJ, Martínez-Ferri E. Incidence of misshapen
fruits in strawberry plants grown under tunnels is affected by cultivar, planting date,
pollination, and low temperatures. *HortScience*. 2012; **47**: 1569–1573.

12 Xiong Y, From PJ, Isler V. Design and evaluation of a novel cable-driven gripper
with perception capabilities for strawberry picking robots. *2018 IEEE International
Conference on Robotics and Automation (ICRA)*. 2018.
doi:10.1109/icra.2018.8460705.

13 Faedi W, Mourgues F, Rosati C. Strawberry breeding and varieties: situation and
perspectives. *Acta Horticulturae*. 2002; 51–59.

14 Nielsen JA, Lovell PH. Value of morphological characters for cultivar identification
in strawberry (Fragariaxananassa). *New Zealand Journal of Crop and Horticultural
Science*. 2000; **28**: 89–96.

15 Ishikawa T, Hayashi A, Nagamatsu S *et al.* Classification of strawberry fruit shape
by machine learning. *ISPRS - International Archives of the Photogrammetry,
Remote Sensing and Spatial Information Sciences*. 2018; **XLII-2**: 463–470.

16 Mir JI, Ahmed N, Singh DB *et al.* Diversity evaluation of fruit quality of apple (Malus
× domestica Borkh.) germplasm through cluster and principal component analysis.
Indian Journal of Plant Physiology. 2017; **22**: 221–226.

17 Akodagali J, Balaji S. Computer Vision and Image Analysis based techniques for
automatic characterization of fruits a review. *International Journal of Computer
Applications*. 2012; **50**: 6–12.

18 Beyer M, Hahn R, Peschel S, Harz M, Knoche M. Analysing fruit shape in sweet

cherry (Prunus avium L.). *Scientia Horticulturae*. 2002; **96**: 139–150.

19   Naik S, Patel B, Pandey R. Shape, size and maturity features extraction with fuzzy
classifier for non-destructive mango (Mangifera Indica L., cv. Kesar) grading. 2015
IEEE Technological Innovation in ICT for Agriculture and Rural Development
(TIAR). 2015. doi:10.1109/tiar.2015.7358522.

20   Hiraoka Y, Kuramoto N. Identification of Rhus succedanea L. cultivars using Elliptic
Fourier Descriptors based on fruit shape. *Silvae Genetica*. 2004; **53**: 221–226.

21   Paproki A, Sirault X, Berry S, Furbank R, Fripp J. A novel mesh processing based
technique for 3D plant analysis. *BMC Plant Biol* 2012; **12**: 63.

22   Coupel-Ledru A, Pallas B, Delalande M *et al.* Multi-scale high-throughput
phenotyping of apple architectural and functional traits in orchard reveals genotypic
variability under contrasted watering regimes. *Hortic Res* 2019; **6**: 52.

23   Topp CN, Iyer-Pascuzzi AS, Anderson JT *et al.* 3D phenotyping and quantitative
trait locus mapping identify core regions of the rice genome controlling root
architecture. *Proc Natl Acad Sci U S A* 2013; **110**: E1695–704.

24   Cockerton, H.M., Li, B., Stavridou, E., Johnson, A., Karlström, A., Armitage, A.D.,
Martinez-Crucis, A., Arjona, L.G. and Harrison, R.J. Genetic and phenotypic
associations between root architecture, arbuscular mycorrhizal fungi colonisation
and low phosphate tolerance in strawberry (Fragaria × ananassa).
doi:10.21203/rs.2.18300/v1.

25   He JQ, Harrison RJ, Li B. A novel 3D imaging system for strawberry phenotyping.
Plant Methods 2017; **13**: 93.

26   Westoby MJ, Brasington J, Glasser NF, Hambrey MJ, Reynolds JM. 'Structure-
from-Motion' photogrammetry: A low-cost, effective tool for geoscience
applications. *Geomorphology*. 2012; **179**: 300–314.

27   Kovacs L, Eder M, Hollweck R *et al.* Comparison between breast volume
measurement using 3D surface imaging and classical techniques. *Breast* 2007; **16**:
137–145.

28   Rife TW, Poland JA. Field Book: An Open-Source Application for field data
collection on Android. *Crop Science*. 2014; **54**: 1624.

29   Verma S, Zurn JD, Salinas N *et al.* Clarifying sub-genomic positions of QTLs for
flowering habit and fruit quality in U.S. strawberry (Fragaria×ananassa) breeding
populations using pedigree-based QTL analysis. *Horticulture Research*. 2017; **4**.
doi:10.1038/hortres.2017.62.

30   Vickerstaff RJ, Harrison RJ. Crosslink: A fast, scriptable genetic mapper for
outcrossing species. *BioRxiv*, doi:10.1101/135277.

31   Cockerton HM, Vickerstaff RJ, Karlström A *et al.* Identification of powdery mildew
700   resistance QTL in strawberry (Fragaria × ananassa). *Theor Appl Genet* 2018; **131**:
701   1995–2007.

702   32   van Dijk T, Pagliarani G, Pikunova A *et al.* Genomic rearrangements and
703   signatures of breeding in the allo-octoploid strawberry as revealed through an allele
704   dose based SSR linkage map. *BMC Plant Biol* 2014; **14**: 55.

705   33   Laganiere R. OpenCV 3 Computer Vision Application Programming Cookbook.
706   Packt Publishing Ltd, 2017.

707   34   Podczeck F, Rahman SR, Newton JM. Evaluation of a standardised procedure to
708   assess the shape of pellets using image analysis. *Int J Pharm* 1999; **192**: 123–138.

709   35   Taubin G. Estimating the tensor of curvature of a surface from a polyhedral
710   approximation. *Proceedings of IEEE International Conference on Computer Vision*
711   1995. doi:10.1109/iccv.1995.466840.

712   36   Lancaster P, Salkauskas K. Surfaces generated by moving least squares methods.
713   *Mathematics of Computation*. 1981; **37**: 141–141.

714   37   Gutierrez PA, Perez-Ortiz M, Sanchez-Monedero J, Fernandez-Navarro F, Hervas-
715   Martinez C. Ordinal Regression methods: survey and experimental study. *IEEE*
716   *Transactions on Knowledge and Data Engineering*. 2016; **28**: 127–146.

717   38   Chakrabarti A, Ghosh JK. AIC, BIC and recent advances in model selection.
718   Philosophy of Statistics. 2011; 583–605.

719   39   Yamashita T, Yamashita K, Kamimura R. A stepwise AIC method for variable
720   selection in linear regression. Communications in Statistics - *Theory and Methods*.
721   2007; **36**: 2395–2403.

722   40   Garin V, Wimmer V, Mezmouk S, Malosetti M, van Eeuwijk F. How do the type of
723   QTL effect and the form of the residual term influence QTL detection in multi-parent
724   populations? A case study in the maize EU-NAM population. *Theor Appl Genet*
725   2017; **130**: 1753–1764.

726   41   Churchill GA, Doerge RW. Empirical threshold values for quantitative trait mapping.
727   *Genetics* 1994; **138**: 963–971.

728   42   Zeng ZB. Theoretical basis for separation of multiple linked gene effects in mapping
729   quantitative trait loci. *Proc Natl Acad Sci U S A* 1993; **90**: 10972–10976.

730   43   Zeng Z-B. Precision mapping of quantitative trait loci. *Genetics* 1994; **136**: 1457–
731   1468.

732   44   Oo LM, Aung NZ. A simple and efficient method for automatic strawberry shape
733   and size estimation and classification. *Biosystems Engineering*. 2018; **170**: 96–107.

45   Kochi N, Tanabata T, Hayashi A *et al.* A 3D shape-measuring system for assessing
strawberry fruits. *International Journal of Automation Technology.* 2018; **12**: 395–
404.

46   Aho K, Derryberry D, Peterson T. Model selection for ecologists: the worldviews of
AIC and BIC. Ecology. 2014; **95**: 631–636.

47   Castro P, Lewers KS. Identification of quantitative trait loci (QTL) for fruit-quality
traits and number of weeks of flowering in the cultivated strawberry. *Molecular*
Breeding. 2016; **36**. doi:10.1007/s11032-016-0559-7.

48   Lerceteau-Köhler E, Moing A, Guérin G *et al.* Genetic dissection of fruit quality
traits in the octoploid cultivated strawberry highlights the role of homoeo-QTL in
their control. *Theor Appl Genet* 2012; **124**: 1059–1077.

49   Liao X, Li M, Liu B *et al.* Interlinked regulatory loops of ABA catabolism and
biosynthesis coordinate fruit growth and ripening in woodland strawberry. *Proc Natl*
Acad Sci U S A 2018; **115**: E11542–E11550.

50   Wang S-M, Wei-jia LI, Yue-xue LIU, He LI, Yue MA, Zhang Z-H. Comparative
transcriptome analysis of shortened fruit mutant in woodland strawberry (Fragaria
vesca) using RNA-Seq. *Journal of Integrative Agriculture.* 2017; **16**: 828–844.

51   Ariza MT, Soria C, Medina JJ, Martínez-Ferri E. Fruit misshapen in strawberry
cultivars (Fragaria×ananassa) is related to achenes functionality. *Annals of Applied*
Biology. 2011; **158**: 130–138.

52   Pipattanawong R, Yamane K, Fujishige N, Bang S-W, Yamaki Y. Effects of high
temperature on pollen quality, ovule fertilization and development of embryo and
achene in 'Tochiotome' strawberry. *Journal of the Japanese Society for*
Horticultural Science. 2009; **78**: 300–306.