Title

Switching secretory pathway direction for organelle acquisition in plants

Authors

Takehiko Kanazawa[1,2], Hatsune Morinaka[3], Kazuo Ebine[1,2], Takashi L. Shimada[4],

Sakiko Ishida[5], Naoki Minamino[1], Katsushi Yamaguchi[6], Shuji Shigenobu[6], Takayuki

Kohchi[5], Akihiko Nakano[7], Takashi Ueda[1,2*]

Affiliations

[1]Division of Cellular Dynamics, National Institute for Basic Biology, Nishigonaka 38,

Myodaiji, Okazaki, Aichi, 444-8585 Japan

[2]The Department of Basic Biology, SOKENDAI (The Graduate University for

Advanced Studies), Nishigonaka 38, Myodaiji, Okazaki, Aichi, 444-8585 Japan

[3]Department of Biological Sciences, Graduate School of Sciences, The University of

Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

[4]Department of Applied Biological Chemistry, Graduate School of Horticulture, Chiba

University, 648 Matsudo, Matsudo, Chiba, Japan

[5]Graduate School of Biostudies, Kyoto University, Kitashirakawa-oiwake-cho,

Sakyo-ku, Kyoto 606-8502, Japan

[6]Functional Genomics Facility, National Institute for Basic Biology (NIBB), Okazaki,

Aichi 444-8585, Japan

[7]Live Cell Super-Resolution Imaging Research Team, RIKEN Center for Advanced

Photonics, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

*email: tueda@nibb.ac.jp

Abstract

Eukaryotic cells acquired novel organelles during evolution through mechanisms that remain largely obscure. The existence of the unique oil body compartment is a synapomorphy of liverworts that represents lineage-specific acquisition of this organelle during evolution, although its origin, biogenesis, and physiological function are yet unknown. We found that two Syntaxin 1 paralogs in the liverwort, *Marchantia polymorpha*, are distinctly targeted to forming cell plates and the oil body, suggesting these structures share some developmental similarity. Oil body formation is under the regulation of an ERF/AP2-type transcription factor and loss of the oil body increased *M. polymorpha* herbivory. These findings highlight a common strategy for the acquisition of organelles with distinct functions in plants, via periodical switching in secretion direction depending on cellular phase transition.

Main text:

Eukaryotic cells originated from prokaryotes by expanding their endomembrane network during evolution, with the last eukaryotic common ancestor (LECA) likely possessing a complex set of organelles[1]. New membrane trafficking pathways were added to the LECA endomembrane network, some of which were secondarily lost in a lineage-specific manner, resulting in divergent and organism-specific membrane trafficking systems and organelle compositions of extant eukaryotes[2-4]. It remains mostly unknown how organelles were acquired during evolution. We report that the liverwort-specific oil body was acquired by the cyclic switching of secretory pathway direction regulated by a single transcription factor MpERF13. A pair of Syntaxin 1 homologs in the liverwort

(*Marchantia polymorpha*) were distinctly targeted to the forming cell plate and oil body. Clathrin-coated vesicles formed at the oil body membrane and plasma membrane proteins were targeted to either the oil body or plasma membranes depending on distinct cellular phases of the hypothetical oil body cycle, indicating a shared characteristic between the plasma and oil body membranes. We identified a key transcription factor regulating oil body formation and/or the oil body cycle, and found that oil bodies confer resistance against herbivores. These findings highlight a common strategy for the acquisition of plant-specific organelles/cell structures with distinct functions during evolution.

The liverwort oil body was first described in 1834[5]. It contains bioactive compounds such as sesquiterpenoids and cyclic bisbibenzyl compounds and is not related to the oil body that stores neutral lipids in storage organs like seeds and fruits (i.e. lipid body or oleosome). About 90% of liverwort species have this organelle; however, its origin, biogenesis, and physiological function remain unclear with controversial origins proposed from microscopic observations[6-9]. Through the systematic analysis of SNARE proteins in the liverwort, *Marchantia polymorpha* (hereafter referred to as Marchantia), we identified an oil body-resident protein, MpSYP12B[10], which is a homolog of animal Syntaxin 1 that acts in the final step of exocytosis. Functions of SYP1 members to which MpSYP12B belongs dramatically diversified during plant evolution, suggesting expanded secretory trafficking systems in plants. Arabidopsis SYP1 consists of three subgroups, SYP11, SYP12, and SYP13. The SYP13 group mediates constitutive secretion, whereas the other groups are involved in plant-specific higher ordered functions; for example, KNOLLE/SYP111 is specifically

Fig. 1 | MpSYP12A is important for cell plate formation in *M. polymorpha* thallus cells.
(a, c, and e) Single confocal images of Marchantia thallus cells expressing mCitrine-MpSYP12A (a), mCitrine-MpSYP13A (c), and mCitrine-MpSYP13B (e) stained with FM4-64. (b, d, and f) Line graphs showing relative fluorescence intensities from mCitrine and FM4-64 along the white arrows in panels a (b), c (d), and e (f). (g) Single confocal images of thallus cells of Tak-1 (left) and a putative knock-out mutant (right) stained with FM4-64. (h and i) Accumulation of PM-resident MpSYP13 proteins driven by the Mp*SYP12A* promoter at forming cell plates. Single confocal images of FM4-64 staining in dividing thallus cells expressing mCitrine-MpSYP13A (h) or mCitrine-MpSYP13B (i). Expression of the Mp*CYCB1* promoter results in accumulation of MpSYP13A at the forming cell plate (j). A single confocal image of FM4-64-stained dividing thallus cells expressing mCitrine-MpSYP13A are shown with asterisks indicate forming cell plates. Green, magenta, and blue pseudo-colours indicate fluorescence from mCitrine, FM4-64, and chlorophyll, respectively. Bars = 10 μm.

involved in membrane fusion at forming cell plates, and PEN1/SYP121 is required for

intact penetration resistance against powdery mildew fungi and regulation of the ion

channel activity[11-16]. For more insight into the functional diversification of SYP1

members in Marchantia and the mechanism of oil body biogenesis, we further

characterized four SYP1 members in this organism[10,17].

Our phylogenetic analysis suggested that MpSYP1 members belong to two

distinct subgroups: the SYP13 group (MpSYP13A and 13B) and the SYP11/12 group

(MpSYP12A and 12B) that contains SYP11 and SYP12 groups in seed plants[18] (Fig.

S1a). MpSYP12A, 13A and 13B were ubiquitously expressed and localized to the

plasma membrane (PM) in thallus tissues, whereas MpSYP12B exhibited specific

expression in a subpopulation of thallus cells (Fig S1b-e). Using some ubiquitously

expressed mCitrine-MpSYP1 protein fusion constructs driven by their own regulatory

elements, we detected specific localization of MpSYP12A at forming cell plates

associated with the phragmoplast that rapidly stained with FM4-64 (Figs. 1a-f, S1f, and

S1g). This localization suggested that MpSYP12A could be a functional counterpart of

KNOLLE in Arabidopsis, which was further verified genetically.

90   We tried to generate a complete Mp*syp12a* knockout mutant by genome editing

(Fig. S1h) but were not successful, probably because it is an essential gene. However,

we succeeded in isolating chimeric plants comprising mutated and wild-type cells, in

which we frequently observed enlarged cells with cell wall stabs, suggesting that

MpSYP12A is required for cell plate formation during cytokinesis similar to the

function of KNOLLE (Fig. 1g). The *KNOLLE* promoter is sufficient to target

non-cytokinesis-specific SYP132 to forming cell plates[19]. Similarly, mCitrine-tagged

MpSYP13A and 13B were localized to cell plates and the PM when expressed under the

Mp*SYP12A* promoter (Fig. 1h, i). Furthermore, a similar localization was also observed

when mCitrine-MpSYP13 was expressed by the Mp*CYCB1* promoter (Figs. 1j and S1i),

indicating that the cell cycle-dependent transcriptional regulation is also critical for cell

plate targeting of SYP1 proteins in Marchantia. We also found that clathrin light chain

(CLC) tagged with mCitrine localized to the PM and *trans*-Golgi network in

non-dividing thallus cells, as well as on forming cell plates in Marchantia as has been

reported in other plants (Fig. S1j, k)[20-22]. These results strongly suggested that

fundamental mechanisms of cell plate formation were conserved during land plant

evolution.

Distinct from the other MpSYP1 members, MpSYP12B exhibits specific

expression in oil body cells and localizes to the oil body membrane[10]. Oil body cells

expressing a *pro*Mp*SYP12B:*2×YFP construct indicated that MpSYP12B distributed

Fig. 2 | The oil body membrane shares common properties with the plasma membrane. (a) A five-day-old thallus expressing 2×Citrine (YFP) under the regulation of the Mp*SYP12B* promoter observed with a light sheet microscope. (b) A single confocal image of an oil body cell expressing mCitrine-MpSYP12B under its own regulatory elements. The bright field image was merged with the fluorescent image. (c and d) Immunoelectron micrographs of oil body cells in thalli of Tak-1 with (c) or without (d) mCitrine-MpSYP12B expression. Cyt, cytosol, and OB, oil body. (e) A single confocal image of an oil body cell expressing GFP-HDEL (an ER marker, magenta) and mCitrine-MpSYP12B (green). (f) An electron micrograph of the oil body cell. The arrowhead and arrow indicate a clathrin-coated vesicle and a clathrin-coated pit, respectively. The magnified image of the boxed region is shown in the inset. (g) A single confocal image of an oil body cell expressing MpCLC1-Citrine. Arrowheads indicate clathrin-coated vesicles/pits forming at the oil body membrane. The bright field image is also shown in the right panel. (h) A single confocal image of oil body cells expressing mCitrine-MpPIP2 under its own regulatory elements. (i) A single confocal image of an oil body cell expressing sec-mRFP driven by the Mp*EF1α* promoter. The bright field image was merged with the fluorescent image. The blue pseudo-colour indicates chlorophyll autofluorescence. Bars = 500 μm in (a), 10 μm in (b), (e), and (g - i), and 200 nm in (c), (d), and (f).

110 around meristematic regions in young thalli, which was observed using light-sheet

111 microscopy (Fig. 2a; Supplementary movie S1). These results suggest that oil body

112 formation occurs around meristematic regions and into thalli during the growth of

113 thallus tissues. mCitrine-MpSYP12B expressed under its own regulatory elements

114 localized to the oil body membrane with a faint signal on the PM (Fig. 2b), which was

115 also confirmed by immuno-EM analysis using an anti-GFP antibody (Fig. 2c, d). We

116 then tested whether organelle markers for the endoplasmic reticulum (MpSEC20,

117 MpUSE1A, MpSEC22, mRFP-HDEL, and GFP-HDEL), Golgi apparatus (MpGOS11

118 and MpSFT1), *trans*-Golgi network (MpSYP6A and MpSYP4), and tonoplast

119 (MpSYP2 and MpVAMP71) were targeted to the oil body membrane, none of which

120 were detected (Figs. 2e and S2b, c). Using transmission electron microscopy, we found

121 that clathrin-coated vesicles formed from the oil body membrane, and emergence and

Fig. 3 | The oil body is formed by directional switching of the secretory pathway. (a and b) Single confocal images of thallus cells including oil body cells expressing mCitrine-MpSYP13A (a) or mCitrine-MpSYP13B (b) under their own regulatory elements. (c - e) Single confocal images of thallus cells including oil body cells expressing mCitrine-MpSYP13A (c), mCitrine-MpSYP13B (d), or mCitrine-MpPIP2 (e) under the Mp*SYP12B* promoter. Bright field images are also shown in the right panel for each fluorescent image. (f and g) Single confocal images of thallus cells including oil body cells expressing sec-mRFP under the Mp*SYP12B* promoter (f) or the Mp*SYP13B* promoter (g). Bright field images are merged with fluorescent images. (h - j) Single confocal images of oil body cells at distinct developmental stages expressing mCitrine-MpSYP12B and mRFP-MpSYP13B. Green, magenta, and blue pseudo-colours indicate from mCitrine, mRFP, and chlorophyll signals, respectively. Bars = 10 μm. (k) A scatter diagram of the areas of oil body cells on the X axis and the areas of oil bodies on the Y axis. R^2 = 0.9547. (l and m) Schemes of the oil body cycle hypothesis. Newly synthesized PM proteins and secreted cargos are targeted to the oil body membrane in the oil body phase, and then to the PM and extracellular space during the PM phase (l). Expression levels of MpSYP12B and MpSYP13B periodically oscillate depending on the cell phases; whereas, MpSYP13A is constitutively expressed, resulting in dual localization at the oil body membrane and the PM (m).

disappearance of clathrin-positive foci at the oil body membrane was also observed in

transgenic plants expressing Citrine-tagged MpCLC (Fig. 2g; Supplementary movie

S2).

Given that clathrin-mediated endocytosis occurs at the PM and SYP1 members

generally function on the PM, the oil body membrane could share similar characteristics

with the PM. This hypothesis was supported by dual localization of the PM proteins,

PM-type aquaporin MpPIP2 (Fig. 2h) and PM-resident SNARE MpSYP13A (Fig. 3a)

driven by their own regulatory elements. The plasma membrane-like nature of the oil

body membrane and the existence of the SYP1 member that is homologous to KNOLLE

suggested that the oil body could be formed by the fusion of secretory vesicles similar

to the cell plate. Thus, the luminal space of the oil body should be topologically

equivalent to the extracellular space, which was tested by expressing a general secretion

marker sec-mRFP, which is composed of the signal peptide for ER translocation and

mRFP under the constitutive Mp*EF1α* promoter. In non-oil body cells, mRFP

fluorescence was detected only in the extracellular space (Fig. S2d). However, mRFP

accumulated in the oil body in addition to the extracellular space in oil body cells,

demonstrating equivalent topology between the lumen of the oil body and extracellular

space (Fig. 2i). Unlike MpSYP12A and KNOLLE, MpSYP12B loss of function did not

result in a detectable abnormality in oil body formation and transport of MpSYP13A or

sec-mRFP (Fig. S3), probably reflecting a functional redundancy between MpSYP12B

and 13A similar to the partial functional redundancy of KNOLLE and SYP132[23].

Distinct from MpSYP13A with the dual localization to the plasma and oil body

membranes, the close homolog MpSYP13B was only localized to the PM in oil body

cells (Fig. 3a, b). Surprisingly, both of these proteins were targeted predominantly to the

oil body membrane when expressed under the regulation of the Mp*SYP12B* promoter

(Fig. 3c, d). This effect was not restricted to SYP1 homologs; the PM-resident protein

MpPIP2 and a general secretion cargo sec-mRFP were also targeted almost exclusively

to the oil body when driven by the Mp*SYP12B* promoter (Fig. 3e, f). These results

indicated that the secretory pathway is redirected inward and vesicles fuse with each

other to form the oil body during the phase in which the Mp*SYP12B* promoter is active.

In contrast, when Mp*SYP13B* promoter is active, the secretory pathway is directed to

the PM and extracellular space, which was further confirmed by accumulation of

sec-mRFP predominantly in the extracellular space when the construct was driven by

the Mp*SYP13B* promoter (Fig. 3g). Other organelle markers did not change their

localization even when expressed under the Mp*SYP12B* promoter (Fig. S2b, c). These

results indicated that the switching of directions of the secretory pathway in oil body

cells is regulated at the transcription level. The switching of secretory directions should

take place periodically and repeatedly through oil body cell development, because we

never observed oil body membrane localization for MpSYP13B at any developmental

stages of oil body cells (Fig. 3h-j), and increase in the size of the oil body cell along

with the increase in the oil body size was observed (Fig. 3k). Based on these findings,

we propose the "oil body cycle hypothesis", which states that Marchantia oil body cells

cycle between two distinct cellular phases: the "PM phase" in which the secretory

pathway is oriented to the PM and extracellular space, and the "oil body phase" when

the secretory pathway is oriented to the oil body, with the phase transition under the

regulation of a transcriptional regulatory system (Fig. 3m, l).

To investigate the regulatory system of oil body formation and oil body cycle,

we screened mutants defective in oil body formation (Fig. S4a). From 48,825 T-DNA

insertion lines, we identified a mutant (Mp*erf13*^{GOF}) with an increased number of oil

bodies. The mutant gemmae contained 334.4 ± 85.6 oil bodies (mean $\pm$ SD), whereas

wild-type gemmae possessed 51.9 ± 8.7 oil bodies under our experimental conditions

(Fig. 4a-e). The T-DNA was inserted 2674-bp upstream of the start codon of

Mapoly0060s0052.1/Mp*ERF13*, which encodes a putative ERF/AP2 transcription factor

in the subgroup containing DREB1A and TINY in Arabidopsis (Fig. S4b). Moreover,

Mp*ERF13* and Mp*SYP12B* transcripts accumulated in this mutant compared to wild

type as detected by RNA-sequencing (RNA-Seq) and quantitative RT-PCR analyses

(Fig. S5b). We then generated knockout mutants in which the Mp*ERF13* gene was

deleted by genome editing (Fig. S4c). Two independent mutants (Mp*erf13-1*^{ge} and

Mp*erf13-2*^{ge}) exhibited no detectable abnormalities in thallus development and

reproductive growth; however, these mutants completely lacked oil bodies in the

Fig. 4 | MpERF13 transcription factor regulates oil body formation. (a - d) Maximum intensity projection images of BODIPY-stained gemmae of Tak-1 (a), Mp*erf13^GOF* (b), Mp*erf13-1^ge* (c), and Mp*erf13-2^ge* (d). (e) The number of oil bodies visualized by BODIPY-staining in gemmae. Bars indicate means ± SD. Statistical analyses between Tak-1 and each genotype were conducted using a two-tailed Welch's *t*-test. Sample numbers were 57 gemmae for Tak-1, 78 for Mp*erf13^GOF*, 61 for Mp*erf13-1^ge*, and 75 for Mp*erf13-2^ge*; and *p*-values are 1.19×10^{-43} for Mp*erf13^GOF*, 1.44×10^{-45} for Mp*erf13-1^ge*, and 1.44×10^{-45} for Mp*erf13-2^ge*. (f) A maximum intensity projection image of oil body and non-oil body cells in a gemma expressing tandemTomato (tdT)-NLS driven by the Mp*ERF13* promoter. (g and h) Single confocal images of thallus cells including oil body cells expressing mCitrine-MpABCG1 under its own regulatory elements (g) and the Mp*SYP13B* promoter (h). (i and j) Single confocal images of thallus cells expressing sec-mRFP under the Mp*ABCG1* (i) and the Mp*MYB02* (j) promoters. Green, magenta, and blue pseudo-colours indicate fluorescence from BODIPY or mCitrine, tdT or mRFP, and chlorophyll, respectively. (k) Ten-day-old thalli in indicated genotypes (Before) they were fed to starved pill bugs or 24 h after feeding (After). (l) Quantification of fluorescence signals images in panel (k). Ratios of thallus areas of "After" to "Before" pill bug feeding were calculated to indicate thallus area change ratios (n = 36 thalli for each genotype). Bars indicate means ± SD. Statistical analyses between Tak-1 and each genotype were conducted by two-tailed Welch's *t*-test. The *p*-values were 5.21×10^{-4} for Mp*erf13^GOF* and 1.05×10^{-16} for Mp*erf13-1^ge*. Bars = 100 µm in (a - d), 10 µm in (f - j), and 1 cm in (k).

gemma and thallus tissues (Figs. 4c-e and S4f, g). These phenotypes of gain- and

loss-of-function mutations suggested that MpERF13 is a major transcription factor

regulating oil body formation. Consistently, the Mp*ERF13* promoter was highly active

in oil body cells (Fig. 4f).

As MpERF13 is homologous to ERF/AP2 transcription factors, this protein

was expected to be involved in the transcriptional regulation of oil body formation

and/or the oil body cycle. To identify genes downstream of MpERF13, we analyzed

transcriptomes in Tak-1 (wild type) and Mp*erf13^GOF* and Mp*erf13-1^ge* mutant lines by

RNA-Seq. Through the comparison between the three groups using an ANOVA-like

test, we identified 136 differentially-expressed genes (DEGs) other than Mp*ERF13*

whose expression was higher than two $\log_2$-fold change (FC) in Mp*erf13*^{GOF} compared

with Tak-1, and lower than -2 $\log_2$FC in Mp*erf13-1*^{ge} than Tak-1 (FDR < 0.01)

(Supplementary Table S1). To verify the RNA-Seq result, we also performed the

quantitative RT-PCR analysis for selected 11 DEGs including Mp*ERF13* and

Mp*SYP12B*, which confirmed that all of these DEGs were highly expressed in

Mp*erf13*^{GOF}, but their expression was significantly lower or not detectable in

Mp*erf13-1*^{ge} and Mp*erf13-2*^{ge} compared with Tak-1 (Fig S5). These data strongly

suggested that these 136 genes function downstream of Mp*ERF13* in oil body cells,

which was further verified for the two genes Mapoly0083s0014.1 and MpMYB02

(Mapoly0006s0226). Mapoly0083s0014.1 was positively regulated by MpERF13 and

encodes a member of the G subfamily of ABC transporters[17], which we named

MpABCG1. Although ABCG members are generally targeted to the PM and mediate

efflux of secondary metabolites from intracellular to extracellular spaces[24,25],

mCitrine-MpABCG1 was targeted to the oil body membrane when expressed under its

own regulatory elements (Fig. 4g); however, this protein was targeted to the PM in

non-oil body cells when expressed under the Mp*SYP13B* promoter (Fig. 4h).

MpMYB02, whose expression was also positively regulated by MpERF13, is reported

to be a transcription factor responsible for the production of marchantin A and its

derivatives that accumulate in oil bodies[9,26]. Mp*ABCG1* and Mp*MYB02* are specifically

expressed in oil body cells and when sec-mRFP was driven by the promoters of these

genes, it was specifically targeted to the lumen of oil bodies. These results indicated that

MpABCG1 and MpMYB02, presumably together with other DEGs we identified, act

during oil body formation downstream of MpERF13.

Lastly, we asked why liverworts possess oil bodies. Using fresh and

alcohol-washed liverworts, the oil body was proposed to be an effective chemical

protection from herbivores[27], although its true function remains unknown because

alcohol can wash out substances as well as oil body constituents. We fed starved

Armadillidium vulgare (pill bug) with wild-type liverwort and genetically established

mutants that contain extra or no oil bodies. Thalli of Tak-1 and Mp*erf13^{GOF}* remained

almost intact after 24 h of herbivory, while the areas of Mp*erf13-1^{ge}* and Mp*erf13-2^{ge}*

thalli were significantly reduced after herbivory (Figs. 4k, l and S6). This result

demonstrated that the oil body is effective in protecting liverworts from herbivores.

We demonstrate that plants have acquired at least two organelles, the cell plate

and oil body, through a common evolutionary pathway of redirecting secretory patterns

from an outward to an inward direction in cells. The functions of these organelles are

totally different but they share several important traits that include specific SYP11/12

members that reside at their membranes, clathrin-coated vesicles generated from the

plasma and internal membranes, and both the cell plate and oil body are formed by

reorientation of secretory pathways. In both cases, orientation of the secretory pathway

is switched in response to transitioning cellular states, during the cell cycle for cell plate

formation and the oil body cycle for oil body formation. Orientation of the plant

secretory pathway is also modulated during plant-microbe interactions; symbiotic

arbuscular mycorrhizal fungi redirects the plant secretory pathway to form the arbuscule

in host plant cells that acts as an interface for sugar and mineral exchange[28,29]. Thus,

redirection of the secretory pathway is a common strategy for plants to acquire novel

organelles/cellular structures that had resulted in maximizing fitness for land plants

during evolution. The liverwort oil body is a nice system to evaluate how novel

organelles have been acquired during evolution, and the knowledge obtained from this

study is useful for engineering new organelles/cellular structures in plants to maximize

plant cellular functions and provides a powerful methodology for future organelle

technology.

Main References

All references:

1　Koumandou, V. L. *et al.* Molecular paleontology and complexity in the last

eukaryotic common ancestor. *Critical reviews in biochemistry and molecular*

biology **48**, 373-396, doi:10.3109/10409238.2013.821444 (2013).

2　Kanazawa, T. & Ueda, T. Exocytic trafficking pathways in plants: why and how

they are redirected. *The New phytologist* **215**, 952-957, doi:10.1111/nph.14613

(2017).

3　Klinger, C. M., Nisbet, R. E., Ouologuem, D. T., Roos, D. S. & Dacks, J. B.

Cryptic organelle homology in apicomplexan parasites: insights from

evolutionary cell biology. *Current opinion in microbiology* **16**, 424-431,

doi:10.1016/j.mib.2013.07.015 (2013).

4　L'Hernault, S. W. Spermatogenesis. *WormBook : the online review of C. elegans*

biology, 1-14, doi:10.1895/wormbook.1.85.1 (2006).

5　Hübener, J. W. P. Hepaticologia germanica, oder Beschreibung der Deutschen

lebermoose. *Schwan & Götz'sche Hofbuchlandlung, Mannheim* (1834).

6　Asakawa, Y., Ludwiczuk, A. & Nagashima, F. Phytochemical and biological

studies of bryophytes. *Phytochemistry* **91**, 52-80,

doi:10.1016/j.phytochem.2012.04.012 (2013).

7   He, X. L., Sun, Y. & Zhu, R. L. The Oil Bodies of Liverworts: Unique and

Important Organelles in Land Plants. *Critical Reviews in Plant Sciences* **32**,

293-302, doi:10.1080/07352689.2013.765765 (2013).

8   Suire, C. A comparative, transmission-electron microscopic study on the

formation of oil-bodies in liverworts. *J Hattori Bot Lab*, 209-232 (2000).

9   Tanaka, M. *et al.* Direct evidence of specific localization of sesquiterpenes and

marchantin A in oil body cells of *Marchantia polymorpha* L. *Phytochemistry*

130, 77-84, doi:10.1016/j.phytochem.2016.06.008 (2016).

10   Kanazawa, T. *et al.* SNARE Molecules in *Marchantia polymorpha*: Unique and

Conserved Features of the Membrane Fusion Machinery. *Plant & cell*

physiology **57**, 307-324, doi:10.1093/pcp/pcv076 (2016).

11   Assaad, F. F. *et al.* The PEN1 syntaxin defines a novel cellular compartment

upon fungal attack and is required for the timely assembly of papillae. *Mol Biol*

Cell **15**, 5118-5129, doi:10.1091/mbc.E04-02-0140 (2004).

12   Collins, N. C. *et al.* SNARE-protein-mediated disease resistance at the plant cell

wall. *Nature* **425**, 973-977, doi:10.1038/nature02076 (2003).

13   Enami, K. *et al.* Differential expression control and polarized distribution of

plasma membrane-resident SYP1 SNAREs in *Arabidopsis thaliana*. *Plant & cell*

physiology **50**, 280-289, doi:10.1093/pcp/pcn197 (2009).

14   Grefen, C. *et al.* A vesicle-trafficking protein commandeers Kv channel voltage

sensors for voltage-dependent secretion. *Nature plants* **1**, 15108,

doi:10.1038/nplants.2015.108 (2015).

15   Lauber, M. H. *et al.* The Arabidopsis KNOLLE protein is a cytokinesis-specific

syntaxin. *Journal of Cell Biology* **139**, 1485-1493, doi:DOI

10.1083/jcb.139.6.1485 (1997).

16   Lukowitz, W., Mayer, U. & Jurgens, G. Cytokinesis in the Arabidopsis embryo

involves the syntaxin-related KNOLLE gene product. *Cell* **84**, 61-71, doi:Doi

10.1016/S0092-8674(00)80993-9 (1996).

17   Bowman, J. L. *et al.* Insights into Land Plant Evolution Garnered from the

Marchantia polymorpha Genome. *Cell* **171**, 287-304 e215,

doi:10.1016/j.cell.2017.09.030 (2017).

18   Slane, D., Reichardt, I., El Kasmi, F., Bayer, M. & Jurgens, G. Evolutionarily

diverse SYP1 Qa-SNAREs jointly sustain pollen tube growth in Arabidopsis.

The Plant journal : for cell and molecular biology **92**, 375-385,

doi:10.1111/tpj.13659 (2017).

19   Reichardt, I. *et al.* Mechanisms of Functional Specificity Among

Plasma-Membrane Syntaxins in Arabidopsis. *Traffic* **12**, 1269-1280,

doi:10.1111/j.1600-0854.2011.01222.x (2011).

20   Ito, E. *et al.* Dynamic behavior of clathrin in *Arabidopsis thaliana* unveiled by

live imaging. *The Plant journal : for cell and molecular biology* **69**, 204-216,

doi:10.1111/j.1365-313X.2011.04782.x (2012).

21   Otegui, M. S., Mastronarde, D. N., Kang, B. H., Bednarek, S. Y. & Staehelin, L.

A. Three-dimensional analysis of syncytial-type cell plates during endosperm

cellularization visualized by high resolution electron tomography. *The Plant cell*

13, 2033-2051, doi:10.1105/tpc.010150 (2001).

22   Samuels, A. L., Giddings, T. H., Jr. & Staehelin, L. A. Cytokinesis in tobacco

BY-2 and root tip cells: a new model of cell plate formation in higher plants.

The Journal of cell biology **130**, 1345-1357, doi:10.1083/jcb.130.6.1345 (1995).

23   Park, M. *et al.* Concerted Action of Evolutionarily Ancient and Novel SNARE

Complexes in Flowering-Plant Cytokinesis. *Developmental cell* **44**, 500-511

e504, doi:10.1016/j.devcel.2017.12.027 (2018).

24   Adebesin, F. *et al.* Emission of volatile organic compounds from petunia flowers

is facilitated by an ABC transporter. *Science* **356**, 1386-1388,

doi:10.1126/science.aan0826 (2017).

25   Kang, J. *et al.* Plant ABC Transporters. *The arabidopsis book* **9**, e0153,

doi:10.1199/tab.0153 (2011).

26   Kubo, H. *et al.* Biosynthesis of riccionidins and marchantins is regulated by

R2R3-MYB transcription factors in *Marchantia polymorpha*. *Journal of plant*

research **131**, 849-864, doi:10.1007/s10265-018-1044-7 (2018).

27   Stahl, E. Pflanzen und Schnecken. *Jena. Z. Med. Naturwiss* **22**, 557 - 684

(1888).

28   Ivanov, S. & Harrison, M. J. A set of fluorescent protein-based markers

expressed from constitutive and arbuscular mycorrhiza-inducible promoters to

label organelles, membranes and cytoskeletal elements in *Medicago truncatula*.

The Plant journal : for cell and molecular biology **80**, 1151-1163,

doi:10.1111/tpj.12706 (2014).

29   Pumplin, N., Zhang, X., Noar, R. D. & Harrison, M. J. Polar localization of a

symbiosis-specific phosphate transporter is mediated by a transient reorientation

of secretion. *Proceedings of the National Academy of Sciences of the United*

States of America **109**, E665-672, doi:10.1073/pnas.1110215109 (2012).

30   Edgar, R. C. MUSCLE: a multiple sequence alignment method with reduced

time and space complexity. *BMC bioinformatics* **5**, 113,

doi:10.1186/1471-2105-5-113 (2004).

31   Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and

high throughput. *Nucleic acids research* **32**, 1792-1797, doi:10.1093/nar/gkh340

(2004).

32   Castresana, J. Selection of conserved blocks from multiple alignments for their

use in phylogenetic analysis. *Mol Biol Evol* **17**, 540-552, doi:DOI

10.1093/oxfordjournals.molbev.a026334 (2000).

33   Talavera, G. & Castresana, J. Improvement of phylogenies after removing

divergent and ambiguously aligned blocks from protein sequence alignments.

Syst Biol **56**, 564-577, doi:10.1080/10635150701472164 (2007).

34   Guindon, S. *et al.* New Algorithms and Methods to Estimate

Maximum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0.

Syst Biol **59**, 307-321, doi:10.1093/sysbio/syq010 (2010).

35   Ishizaki, K. *et al.* Development of Gateway Binary Vector Series with Four

Different Selection Markers for the Liverwort *Marchantia polymorpha*. *PloS*

one **10**, e0138876, doi:10.1371/journal.pone.0138876 (2015).

36   Naito, Y., Hino, K., Bono, H. & Ui-Tei, K. CRISPRdirect: software for

designing CRISPR/Cas guide RNA with reduced off-target sites. *Bioinformatics*

31, 1120-1123, doi:10.1093/bioinformatics/btu743 (2015).

37 Sugano, S. S. *et al.* Efficient CRISPR/Cas9-based genome editing and its

application to conditional genetic analysis in *Marchantia polymorpha. PloS one*

13, e0205117, doi:10.1371/journal.pone.0205117 (2018).

38 Ishizaki, K., Johzuka-Hisatomi, Y., Ishida, S., Iida, S. & Kohchi, T.

Homologous recombination-mediated gene targeting in the liverwort

Marchantia polymorpha L. *Scientific reports* **3**, 1532, doi:10.1038/srep01532

(2013).

39 Ishizaki, K., Chiyoda, S., Yamato, K. T. & Kohchi, T. Agrobacterium-mediated

transformation of the haploid liverwort *Marchantia polymorpha* L., an emerging

model for plant biology. *Plant & cell physiology* **49**, 1084-1091,

doi:10.1093/pcp/pcn085 (2008).

40 Kubota, A., Ishizaki, K., Hosaka, M. & Kohchi, T. Efficient

Agrobacterium-mediated transformation of the liverwort *Marchantia*

polymorpha using regenerating thalli. *Bioscience, biotechnology, and*

biochemistry **77**, 167-172, doi:10.1271/bbb.120700 (2013).

41 Minamino, N. *et al.* Dynamic reorganization of the endomembrane system

during spermatogenesis in *Marchantia polymorpha. Journal of plant research*

130, 433-441, doi:10.1007/s10265-017-0909-5 (2017).

42 Honkanen, S. *et al.* The Mechanism Forming the Cell Surface of Tip-Growing

Rooting Cells Is Conserved among Land Plants. *Curr Biol* **26**, 3238-3244,

doi:10.1016/j.cub.2016.09.062 (2016).

43 Liu, Y. G., Mitsukawa, N., Oosumi, T. & Whittier, R. F. Efficient Isolation and

Mapping of *Arabidopsis thaliana* T-DNA Insert Junctions by Thermal

Asymmetric Interlaced Pcr. *Plant Journal* **8**, 457-463, doi:DOI

10.1046/j.1365-313X.1995.08030457.x (1995).

44   Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic

RNA-seq quantification. *Nature biotechnology* **34**, 525-527,

doi:10.1038/nbt.3519 (2016).

45   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor

package for differential expression analysis of digital gene expression data.

Bioinformatics **26**, 139-140, doi:10.1093/bioinformatics/btp616 (2010).

46   Saint-Marcoux, D., Proust, H., Dolan, L. & Langdale, J. A. Identification of

reference genes for real-time quantitative PCR experiments in the liverwort

Marchantia polymorpha. *PloS one* **10**, e0118678,

doi:10.1371/journal.pone.0118678 (2015).

47   Nakazaki, A. *et al.* Leaf Endoplasmic Reticulum Bodies Identified in

Arabidopsis Rosette Leaves Are Involved in Defense against Herbivory. *Plant*

physiology **179**, 1515-1524, doi:10.1104/pp.18.00984 (2019).

48   Buschmann, H., Holtmannspotter, M., Borchers, A., O'Donoghue, M. T. &

Zachgo, S. Microtubule dynamics of the centrosome-like polar organizers from

the basal land plant *Marchantia polymorpha*. *The New phytologist* **209**,

999-1013, doi:10.1111/nph.13691 (2016).

49   Otani, K. *et al.* An evolutionarily conserved NIMA-related kinase directs rhizoid

tip growth in the basal land plant *Marchantia polymorpha*. *Development* **145**,

doi:10.1242/dev.154617 (2018).

50   Bowman, J. L. *et al.* The Naming of Names: Guidelines for Gene Nomenclature

in Marchantia. *Plant & cell physiology* **57**, 257-261, doi:10.1093/pcp/pcv193

(2016).

51   Montgomery, S. A. *et al*. Chromatin Organization in Early Land Plants Reveals

an Ancestral Association between H3K27me3, Transposons, and Constitutive

Heterochromatin. *Curr Biol* S0960-9822(19)31610-0.

doi:10.1016/j.cub.2019.12.015.

Fig. 1. MpSYP12A is important for cell plate formation in *M. polymorpha* thallus

cells

(a, c, and e) Single confocal images of Marchantia thallus cells expressing

mCitrine-MpSYP12A (a), mCitrine-MpSYP13A (c), and mCitrine-MpSYP13B (e)

stained with FM4-64. (b, d, and f) Line graphs showing relative fluorescence intensities

from mCitrine and FM4-64 along the white arrows in panels a (b), c (d), and e (f). (g)

Single confocal images of thallus cells of Tak-1 (left) and a putative knock-out mutant

(right) stained with FM4-64. (h and i) Accumulation of PM-resident MpSYP13 proteins

driven by the Mp*SYP12A* promoter at forming cell plates. Single confocal images of

FM4-64 staining in dividing thallus cells expressing mCitrine-MpSYP13A (h) or

mCitrine-MpSYP13B (i). Expression of the Mp*CYCB1* promoter results in

accumulation of MpSYP13A at the forming cell plate (j). A single confocal image of

FM4-64-stained dividing thallus cells expressing mCitrine-MpSYP13A are shown with

asterisks indicate forming cell plates. Green, magenta, and blue pseudo-colours indicate

fluorescence from mCitrine, FM4-64, and chlorophyll, respectively. Bars = 10 μm.

Fig. 2. The oil body membrane shares common properties with the plasma

membrane

(a) A five-day-old thallus expressing 2×Citrine (YFP) under the regulation of the

Mp*SYP12B* promoter observed with a light sheet microscope. (b) A single confocal

image of an oil body cell expressing mCitrine-MpSYP12B under its own regulatory

elements. The bright field image was merged with the fluorescent image. (c and d)

Immunoelectron micrographs of oil body cells in thalli of Tak-1 with (c) or without (d)

mCitrine-MpSYP12B expression. Cyt, cytosol, and OB, oil body. (e) A single confocal

image of an oil body cell expressing GFP-HDEL (an ER marker, magenta) and

mCitrine-MpSYP12B (green). (f) An electron micrograph of the oil body cell. The

arrowhead and arrow indicate a clathrin-coated vesicle and a clathrin-coated pit,

respectively. The magnified image of the boxed region is shown in the inset. (g) A

single confocal image of an oil body cell expressing MpCLC1-Citrine. Arrowheads

indicate clathrin-coated vesicles/pits forming at the oil body membrane. The bright field

image is also shown in the right panel. (h) A single confocal image of oil body cells

expressing mCitrine-MpPIP2 under its own regulatory elements. (i) A single confocal

image of an oil body cell expressing sec-mRFP driven by the Mp*EF1α* promoter. The

bright field image was merged with the fluorescent image. The blue pseudo-colour

indicates chlorophyll autofluorescence. Bars = 500 μm in (a), 10 μm in (b), (e), and (g -

i), and 200 nm in (c), (d), and (f).

Fig. 3. The oil body is formed by directional switching of the secretory pathway

(a and b) Single confocal images of thallus cells including oil body cells expressing

mCitrine-MpSYP13A (a) or mCitrine-MpSYP13B (b) under their own regulatory

elements. (c - e) Single confocal images of thallus cells including oil body cells

expressing mCitrine-MpSYP13A (c), mCitrine-MpSYP13B (d), or mCitrine-MpPIP2

(e) under the Mp*SYP12B* promoter. Bright field images are also shown in the right panel

for each fluorescent image. (f and g) Single confocal images of thallus cells including

oil body cells expressing sec-mRFP under the Mp*SYP12B* promoter (f) or the

Mp*SYP13B* promoter (g). Bright field images are merged with fluorescent images. (h -

j) Single confocal images of oil body cells at distinct developmental stages expressing

mCitrine-MpSYP12B and mRFP-MpSYP13B. Green, magenta, and blue

pseudo-colours indicate from mCitrine, mRFP, and chlorophyll signals, respectively.

Bars = 10 μm. (k) A scatter diagram of the areas of oil body cells on the X axis and the

areas of oil bodies on the Y axis. $R^2 = 0.9547$. (l and m) Schemes of the oil body cycle

hypothesis. Newly synthesized PM proteins and secreted cargos are targeted to the oil

body membrane in the oil body phase, and then to the PM and extracellular space

during the PM phase (l). Expression levels of MpSYP12B and MpSYP13B periodically

oscillate depending on the cell phases; whereas, MpSYP13A is constitutively expressed,

resulting in dual localization at the oil body membrane and the PM (m).

Fig. 4. MpERF13 transcription factor regulates oil body formation

(a - d) Maximum intensity projection images of BODIPY-stained gemmae of Tak-1 (a),

Mp*erf13^{GOF}* (b), Mp*erf13-1^{ge}* (c), and Mp*erf13-2^{ge}* (d). (e) The number of oil bodies

visualized by BODIPY-staining in gemmae. Bars indicate means ± SD. Statistical

analyses between Tak-1 and each genotype were conducted using a two-tailed Welch's

t-test. Sample numbers were 57 gemmae for Tak-1, 78 for Mp*erf13^{GOF}*, 61 for

Mp*erf13-1^{ge}*, and 75 for Mp*erf13-2^{ge}*; and *p*-values are 1.19×10^{-43} for Mp*erf13^{GOF}*,

1.44×10^{-45} for Mp*erf13-1^{ge}*, and 1.44×10^{-45} for Mp*erf13-2^{ge}*. (f) A maximum intensity

projection image of oil body and non-oil body cells in a gemma expressing

tandemTomato (tdT)-NLS driven by the Mp*ERF13* promoter. (g and h) Single confocal

images of thallus cells including oil body cells expressing mCitrine-MpABCG1 under

its own regulatory elements (g) and the Mp*SYP13B* promoter (h). (i and j) Single

confocal images of thallus cells expressing sec-mRFP under the Mp*ABCG1* (i) and the

Mp*MYB02* (j) promoters. Green, magenta, and blue pseudo-colours indicate

fluorescence from BODIPY or mCitrine, tdT or mRFP, and chlorophyll, respectively.

(k) Ten-day-old thalli in indicated genotypes (Before) they were fed to starved pill bugs

or 24 h after feeding (After). (l) Quantification of fluorescence signals images in panel

(k). Ratios of thallus areas of "After" to "Before" pill bug feeding were calculated to

indicate thallus area change ratios (n = 36 thalli for each genotype). Bars indicate means

± SD. Statistical analyses between Tak-1 and each genotype were conducted by

two-tailed Welch's *t*-test. The *p*-values were 5.21×10^{-4} for Mp*erf13*^{GOF} and 1.05×10^{-16}

for Mp*erf13-1*^{ge}. Bars = 100 μm in (a - d), 10 μm in (f - j), and 1 cm in (k).

Methods

Phylogenetic Analyses

Phylogenetic analyses were performed as previously described in Bowman *et al*, 2017 [17]

with minor modification. Previously published datasets in Kanazawa *et al*., 2016 and

Bowman *et al*., 2017, for phylogenetic analyses of SYP1 and ERF/AP2 members,

respectively[10,17]. Multiple sequence alignments were performed using the MUSCLE

program version 3.8.31[30,31] with the default parameter. The alignment gaps were

removed by Gblocks program version 0.91b[32,33] or manually. The maximum likelihood

phylogenetic analyses were performed using PhyML 3.0[34] under the LG model.

Bootstrap analyses were performed by resampling 1000 sets.

Vector Construction

Open reading frames (ORFs) and genomic sequences of Marchantia genes were

amplified by polymerase chain reaction (PCR) from cDNA and genomic DNA prepared

from the accession Tak-1, and the amplified products were subcloned into

pENTR/D-TOPO (ThermoFisher, Waltham, USA) according to the manufacturer's

instructions. The primer sequences and sizes of amplified products are listed in

Supplementary Table S2. For construction of pENTR *pro*Mp*ERF13*, the 5′ sequence

(promoter + 5′ UTR) was amplified and subcloned into pENTR/D-TOPO. The resultant

sequence was introduced into pMpGWB316[35] using the Gateway LR Clonase™ II

Enzyme Mix (ThermoFisher) according to the manufacturer's instructions. For

construction of pENTR genomic mGFP-MpSYP12A and mRFP-MpSYP13B constructs,

the cDNA for mGFP and mRFP were inserted into flanking *Sma*I and *Bam*HI sites in

the pENTR genomic XFP(*Sma*I)-MpSYP12A and pENTR genomic

XFP(*Bam*HI)-MpSYP13B, which were previously prepared[10] using the In-Fusion HD

Cloning System (Clontech, Shiga, Japan). For construction of pENTR genomic

mCitrine-MpPIP2, pENTR genomic mCitrine-MpSYP4, pENTR genomic

mCitrine-MpSEC22, and pENTR genomic mCitrine-MpABCG1, genomic sequences of

the protein-coding regions and 3′ flanking sequences were amplified with a *Sma*I site at

the 5′ end and subcloned into the pENTR vector. The 5′ sequences (promoter + 5′ UTR)

and cDNA of mCitrine were amplified and inserted into *Not*I and *Sma*I sites,

respectively, of the resultant pENTR vectors using the In-Fusion HD Cloning System.

The resultant chimeric genes were then introduced into pMpGWB301[35] using the

Gateway LR Clonase™ II Enzyme Mix. For construction of pENTR

*pro*Mp*SYP12B:mCitrine*, the *pro*Mp*SYP12B:mCitrine* sequence was amplified from the

pENTR genomic mCitrine-MpSYP12B vector, and subcloned into pENTR/D-TOPO.

The resultant sequence was introduced into pMpGWB307[35] using the Gateway LR

Clonase™ II Enzyme Mix to create the pMpGWB307 *pro*Mp*SYP12B:2×YFP* construct.

For construction of pENTR genomic MpCLC1-mCitrine, genomic sequence comprising

the 5′ flanking (promoter + 5′ UTR) and protein-coding sequences were amplified with

a *Sma*I site at the 3′ end and subcloned into the pENTR vector. The 3′ flanking sequence

including 3′ UTR and cDNA for mCitrine were amplified and inserted into *Asc*I and

*Sma*I sites, respectively, of the resultant pENTR vectors using the In-Fusion HD

Cloning System. The resultant chimeric genes were then introduced into

pMpGWB301[35] using the Gateway LR Clonase™ II Enzyme Mix. For construction of

pENTR sec-mRFP, sequences for the signal peptide and mRFP were independently

amplified by PCR, and the sequence of sec-mRFP was then amplified by PCR using the

mixture of the amplified products as templates. The amplified products were subcloned

into pENTR/D-TOPO. For the construction of pENTR mRFP-HDEL, the sequence of

SP-mRFP-HDEL was amplified by PCR using pENTR sec-mRFP as template, and

subcloned into pENTR/D-TOPO. For construction of pENTR *pro*Mp*SYP12B:*sec-mRFP,

pENTR *pro*Mp*SYP13B:*sec-mRFP, pENTR *pro*Mp*ABCG1:*sec-mRFP, and pENTR

*pro*Mp*MYB02:*sec-mRFP, the 5′ sequences (promoter + 5′ UTR) were amplified and

were inserted into *Not*I sites of the pENTR sec-mRFP vector using the In-Fusion HD

Cloning System. The resultant sequences were then introduced into pMpGWB301[35]

using the Gateway LR Clonase™ II Enzyme Mix. To construct pMpGWB301-derived

Gateway vectors (Fig. S2a), the promoter:*mCitrine* sequences were amplified from

genomic constructs described above, and inserted at a *Hin*dIII site of pMpGWB301

using the In-Fusion HD Cloning System. pENTR MpTUB2 was kindly provided from

Dr. R. Nishihama and introduced into pMpGWB301*pro*Mp*SYP2:mCitrine*-GW (Fig.

S2a) using the Gateway LR Clonase™ II Enzyme Mix. For construction of the

genome-editing vectors, the target sequences were selected using CRISPR direct

(https://crispr.dbcls.jp/)[36], and double-stranded oligonucleotides of the target sequences

were inserted into the pMpGE_En03 vector[37]. The resultant gRNA cassettes were

introduced into the pMpGE010 or pMpGE011 vectors[37] using the Gateway LR Clonase

II Enzyme Mix. For construction of the homologous recombination-mediated gene

targeting vector, the 3.5-kb homologous genomic sequences were amplified from the

Tak-1 genome and inserted at *Pac*I and *Asc*I sites of the pJHY-TMp1 vector[38] using the

In-Fusion HD Cloning System.

Plant materials and Transformation

Marchantia accession Takaragaike-1 (Tak-1, male) and Takaragaike-2 (Tak-2, female)

[39] were used throughout the study. The growth conditions and transformation methods

were previously described[10,39,40]. *M. polymorpha* expressing mCitrine-MpSYP12A,

mCitrine-MpSYP12B, mCitrine-MpSYP13A, mCitrine-MpSYP13B,

mCitrine-MpSYP2 or mCitrine-MpVAMP71 under the regulation of their own

regulatory elements (e.g. promoter, UTRs, and introns), and *M. polymorpha* expressing

mRFP-MpSYP6A or GFP-HDEL driven by the Mp*EF1α* promoter were previously

reported[10,35,41].

Genotyping

Genomic DNA was extracted from the thalli in an extraction buffer containing 1 M KCl,

100 mM Tris-HCl (pH 9.5), and 10 mM EDTA. This DNA was used as templates for

PCR. Genome fragments with putative mutation sites were amplified by PCR using

KOD FX Neo (TOYOBO, Osaka, Japan) according to the manufacturer's instructions.

The primers used in PCR-based genotyping are listed in Supplementary Table S2.

Confocal laser scanning microscopy

Two- and five-day-old thalli were used for imaging the cell plate the oil body,

respectively. Samples were mounted in distilled water and observed using an LSM780

confocal microscope (Carl Zeiss) equipped with an oil immersion lens (63×, numerical

aperture = 1.4) on an electrically driven stage. Gemmae were observed with a 10×

objective lens (numerical aperture = 0.45). Thalli were incubated in 10 μM

N-(3-triethylammoniumpropyl)-4-(6-(4-(diethylamino)phenyl)hexatrienyl) pyridinium

dibromide (FM4-64, Thermo Fisher) solution for 2 min. Samples were then washed

twice in water before observation. For

4,4-difluoro-1,3,5,7,8-pentamethyl-4-bora-3a,4a-diaza-s-indacene (BODIPY 493/503,

Thermo Fisher) staining, thalli or gemmae were incubated in 200 nM BODIPY 493/503

dissolved in water for 10 min and then washed twice in water before observation. The

samples were excited at 488 nm (Argon 488) and 561 nm (DPSS 561-10) lasers,

respectively, and the fluorescent emission was collected between 482-659 nm using

twenty GaAsP detectors. Spectral unmixing and constructing maximum intensity

projection images were conducted using the ZEN2012 software (Carl Zeiss). Image

processing was performed with ZEN2012 software, Photoshop software (Adobe

Systems), and ImageJ (National Institute of Health, https://imagej.nih.gov/ij/).

Electron microscopy

Five-day-old *M*. *polymorpha* thalli from Tak-1 and thalli expressing

mCitrine-MpSYP12B were used for the immunoelectron microscopic observation.

Sample preparation and observation were performed as previously reported[10].

Five-day-old Tak-1 thalli were used for morphological observation with a transmission

electron microscope. Sample preparation and observation were performed as previously

reported[41].

Light sheet microscopy

Five-day-old thalli expressing 2×Citrine driven by the Mp*SYP12B* promoter

were embedded in low melt agarose gel and observed using a Light sheet Z.1

microscope (Carl Zeiss) equipped with a water immersion lens (5×, numerical aperture

= 0.16). The samples were excited at 488 nm (Argon 488 laser). Acquisition and

construction of three-dimension images from multi-angle images were conducted using

the ZEN2014 software (Carl Zeiss). The images were processed digitally with Imaris

(Bitplane) and Photoshop software.

Fluorescent stereoscopic microscopy

Three-week-old wild-type and Mp*erf13^{ge}* thalli were stained by BODIPY493/503

(ThermoFisher) and observed using a fluorescent stereoscopic microscope (M165 FC,

Leica) equipped with a MC170 HD digital camera (Leica).

Forward genetic screening for the oil body formation mutants

The workflow of the mutant screening is described in Fig. S4a. Liverworts with

transfer DNA (T-DNA) insertions were generated by co-culture of *M. polymorpha*

sporelings with the agrobacteria strain GV2260 harbouring the binary vector

pCAMBIA1300

(https://cambia.org/welcome-to-cambialabs/cambialabs-projects/cambialabs-projects-le

gacy-pcambia-vectors-pcambia-legacy-vectors-1/cambialabs-projects-legacy-pcambia-v

ectors-list-of-legacy-pcambia-vectors-3/). T_1 plants were first selected on 1/2×

Gamborg's B5 plates containing 10 mgL^{-1} hygromycin and 100 mgL^{-1} cefotaxime.

Hygromycin-resistant T_1 plants were then transferred and incubated on hygromycin-free

medium for about one week, followed by BODIPY 493/503 staining to visualize oil

bodies. Visible screening for mutants defective in oil body formation was performed

using a fluorescent stereoscopic microscope (M165 FC, Leica). To identify flanking

sequences of T-DNA insertions, thermal asymmetric interlaced-PCR (TAIL-PCR) was

performed as described previously[39,42,43], with minor modifications using

crude-extracted DNA as templates. Flanking sequences were amplified using KOD FX

neo DNA polymerase and T-DNA-specific (TR1–3 and TL1–3 for right and left borders

of T-DNA, respectively) and universal adaptor (AD1–6) primers. The reaction cycles

are shown below. After agarose gel electrophoresis of the final TAIL-PCR products,

DNA bands were excised and purified using the Wizard SV Gel and PCR Clean-Up

System (Promega). Purified products were directly sequenced using TR3 or TL3

primers. The T-DNA insertion sites were identified using the genome sequence

registered in MarpolBase, genome version 3.1.

Cycling conditions for TAIL-PCR

Reaction	Cycle No.	Thermal settings
Primary	1	94°C 1 min, 95°C 1 min
	5	94°C 1 min, 65°C 1 min, 68°C 3 min
	1	94°C 1 min, 30°C 1min
		Ramping to 68°C 3 min, 68°C 3 min
	13	94°C 30 sec, 68°C 4 min
		94°C 30 sec, 68°C 4 min
		94°C 30 sec, 44°C 1 min, 68°C 3min

	1	68°C 5 min
Secondary	1	94°C 1 min, 95°C 1 min
	13	94°C 30 sec, 68°C 4 min
		94°C 30 sec, 68°C 4 min
		94°C 30 sec, 44°C 1 min, 68°C 3 min
	1	68°C 5 min
Tertiary	1	94°C 1 min, 95°C 1 min
	13	94°C 30 sec, 68°C 4 min
		94°C 30 sec, 68°C 4 min
		94°C 30 sec, 44°C 1 min, 68°C 3 min
	1	68°C 5 min

RNA extraction and RNA sequencing analysis

Total RNA from Tak-1, Mp*erf13*^{GOF} and Mp*erf13-1*^{ge} thalli was extracted by RNeasy

(QIAGEN) according to the manufacturer's instructions. The eluted total RNA samples

were treated with DNase I (Takara) to remove DNA contamination. The quality and

quantity of the total RNA were evaluated with a NanoDrop 1000 spectrophotometer

(Thermo Fisher Scientific), Qubit 2.0 Fluorometer (Thermo Fisher Scientific) and a

Bioanalyzer RNA6000 Nano Kit (Agilent Technologies). The sequence libraries were

prepared with the TruSeq RNA Sample Prep Kit v2 (Illumina) according to the

manufacturer's low sample protocol. The quality and quantity of each library were

determined using a Bioanalyzer with the High Sensitivity DNA kit (Agilent

Technologies) and the KAPA Library Quantification Kit for Illumina (Illumina). Equal

amounts of each library were mixed to make the 2 nM pooled library. Illumina

sequencing was performed using a HiSeq 1500 platform (Illumina) to produce 126-bp

single-end reads. Three biological replicates were prepared for the library construction

and RNA-Seq analysis. All reads are available through the Sequence Read Archive

(SRA) under the accession number DRA009193. The abundance of transcripts from

Illumina RNA-Seq data were quantified with the kallisto program (version 0.43.1)[44]

with the default parameters using the primary transcript dataset from the Marchantia

genome version 3.1. To estimate differentially expressed genes (DEGs), an

ANOVA-like test was performed by edgeR (version 3.18.1)[45]. Genes with FDR values

lower than 0.01 and an absolute $\log_2$ fold change (FC) > 2 were considered to be

differentially expressed. Among all combinations of DEGs ($\log_2$FC(Mp$erf13^{GOF}$/Tak-1)

> 2 and $\log_2$FC(Mp$erf13$-1^{ge}/Tak-1) > 2, $\log_2$FC(Mp$erf13^{GOF}$/Tak-1) > 2 and

$\log_2$FC(Mp$erf13$-1^{ge}/Tak-1) < -2, $\log_2$FC(Mp$erf13^{GOF}$/Tak-1) < -2 and

$\log_2$FC(Mp$erf13$-1^{ge}/Tak-1) > 2, $\log_2$FC(Mp$erf13^{GOF}$/Tak-1) < -2 and

$\log_2$FC(Mp$erf13$-1^{ge}/Tak-1) < -2), we selected the set $\log_2$FC(Mp$erf13^{GOF}$/Tak-1) > 2

and $\log_2$FC(Mp$erf13$-1^{ge}/Tak-1) < -2 as candidate genes regulated downstream of

MpERF13, which included both Mp*ERF13* and Mp*SYP12B*.

Quantitative reverse transcription(qRT)-PCR

For qRT-PCR, cDNA was synthesized from total RNA using SuperScript III Reverse

Transcriptase (Invitrogen) and an oligo dT(18) primer according to the manufacturer's

instructions. qRT-PCR was performed with LightCycler 480 (Roche) using FastStart

SYBR Green Master (Roche) according to the manufacturer's protocol. Sequences of

primers used are listed in Table S2. Mp*APT* (Mapoly0100s0027.1 / Mp3g25140.1) was

used as a housekeeping reference for normalization[46]. Three biological replicates were

prepared and three technical replicates were performed for each reaction.

Pill bug feeding assay

Feeding assay was performed according to Nakazaki *et al*. with modification[47].

Gemmae were cultured on 1/2× Gamborg's B5 medium containing 1% (w/v) agar and

1% (w/v) sucrose for five days at 22°C under continuous white light (50 μmol m^{-2}s^{-1}).

Five-day-old thalli were transferred onto 1/2× Gamborg's B5 medium containing 1%

(w/v) agar without sucrose and cultured for additional five days under the same

conditions. Pill bugs (*Armadillidium vulgare*) were collected in the Myodaiji area of

NIBB (Nishigonaka 38, Myodaiji, Okazaki 444-8585 Aichi, Japan). Pill bugs were

maintained on Prowipe (Daio Paper, Tokyo, Japan) moistened with sterilized water for

48 h at 22°C in the dark without feeding before the assay. Six pill bugs were introduced

into each medium plate containing 10-day-old thalli and kept for 24 h in the dark at

22°C. The thallus areas were calculated using ImageJ (National Institute of Health,

https://imagej.nih.gov/ij/).

Supplementary Materials

Gene accession numbers

Symbol	Gene ID (ver3.1)	Gene ID (ver5.1)
Mp*SYP12A*	Mapoly0163s0015	Mp6g00050
Mp*SYP12B*	Mapoly0101s0013	Mp4g20670
Mp*SYP13A*	Mapoly0009s0005	Mp7g13190
Mp*SYP13B*	Mapoly0055s0091	Mp2g19600
Mp*CLC1*	Mapoly0075s0069	Mp2g03080
Mp*PIP2*	Mapoly0041s0003	Mp4g17210
Mp*ERF13*	Mapoly0060s0052	Mp6g08690
Mp*ABCG1*	Mapoly0083s0014	Mp8g13070
Mp*MYB02*	Mapoly0006s0226	Mp3g07510
Mp*TUB2*	Mapoly0158s0010[48,49]	Mp2g09390
Mp*SYP6A*	Mapoly0140s0004	Mp3g18380
Mp*SYP2*	Mapoly0187s0013	Mp8g15260

Mp*SFT1*	Mapoly0006s0109	Mp3g06390
Mp*GOS11*	Mapoly0001s0245	Mp1g19070
Mp*SYP4*	Mapoly0042s0041	Mp2g14140
Mp*VAMP71*	Mapoly0064s0109	Mp8g00880
Mp*SEC20*	Mapoly0019s0137	Mp1g13670
Mp*USE1A*	Mapoly0001s0213	Mp1g18750
Mp*SEC22*	Mapoly0023s0071	Mp2g11050
Mp*CYCB1*	Mapoly0048s0068	Mp5g10030

We followed the nomenclature of genes, proteins, and mutants of Marchantia reported

in Bowman *et al*. (2016)[50]. Gene IDs were taken from MarpolBase

(http://marchantia.info/), genome version 3.1[17] and version 5.1[51].

Acknowledgments:

We thank Dr. Ikuko Hara-Nishimura (Konan University), Dr. Kimitsune Ishizaki (Kobe

University), Dr. Ryuichi Nishihama (Kyoto University), and Dr. Shigeo S. Sugano for

sharing vectors and plant materials, Ms. Miwako Matsumoto, Ms. Asaka Akita, and

Functional Genomics Facility (NIBB) for technical assistance in RNA-Seq experiments,

and Dr. Ken Naito (National Agriculture and Food Research Organization) for

instruction for RNA-Seq data analysis. We thank Dr. John L. Bowman (Monash

University) and Mr. Facundo Romani (Universidad Nacional del Litoral) for helpful

discussion. We also thank Dr. Teruyuki Niimi (NIBB) and Dr. Takahisa Miyatake

(Okayama University) for identification of the species of pill bug used in this study. The

support of plant cultivation rooms was provided by the Model Plant Research Facility of

NIBB.

Contributions:

T.Ka. performed the majority of the experiments, H.M. performed the experiment

presented in Fig. 1a-g, K.E. and T.L.S. provided essential ideas for experiments using

sec-mRFP targeting and oil body detection, respectively, S.I. generated the Mp*syp12b-1*

mutant, K.Y. and S.S. designed the RNA-seq experiment, N.M. examined the effect of

Mpe*rf13* mutations on reproductive growth, T.Ka. and T.U. wrote the manuscript, and

T.Ko., A.N., and T.U. supervised the study.

Funding:

This work was financially supported by Grants-in-Aid for Scientific Research from the

Ministry of Education, Culture, Sports, Science, and Technology of Japan (to T.U.,

19H05670, 19H05675, and 18H02470, and T.Ka., 18K14738).

Disclosures:

The authors have no conflicts of interest to declare.

 Extended Data Figures

Supplementary Fig. S1. Characterization of SYP1 member and clathrin light chain

localisation during cytokinesis in Marchantia thalli

(a) A maximum likelihood phylogenetic tree of SYP1 members in green plants.

Although the tree is unrooted, the proteins from Chlorophytes (*Ostreococcus* spp.,

Chlamydomonas reinhardtii, and *Volvox carteri*) are the outgroup to those of

streptophytes. The branch lengths are proportional to the estimated number of

substitutions per site. Bootstrap probability is indicated as a percentage on each branch

with at least 50% support. (b - e) Maximum intensity projection images of thalli

expressing mCitrine-MpSYP12A (b), mCitrine-MpSYP13A (c), mCitrine-MpSYP13B

(d), or mCitrine-MpSYP12B (e). White arrowheads indicate

mCitrine-MpSYP12B-expressing cells. (f) Single confocal images of thallus cells

expressing mCitrine-MpTUB2 and mGFP-MpSYP12A under the regulation of the

Mp*SYP2* and Mp*SYP12A* promoters, respectively, stained with FM-4-64. In mitotic

cells, spindles and phragmoplasts (solid and ring-shaped phases) were labelled by

mCitrine-MpTUB2, and endocytosed FM-4-64 accumulated at forming cell plates. (g)

Single confocal images of thallus cells expressing mCitrine-MpSYP12B stained with

FM4-64. The line graph indicates the relative fluorescence intensity along the white

arrow. Relative fluorescent intensity of mCitrine to fluorescence on the oil body

membrane is shown for mCitrine fluorescence. (h) The gene model and sequences of

MpSYP12A. The syntaxin domain, SNARE motif, and transmembrane domain (TMD)

are shown above the gene model. Black and magenta letters indicate genome and

translated amino acid sequences. The protospacer adjacent motif (PAM) sequence for

gRNA is underlined. (i) Expression by the Mp*CYCB1* promoter results in accumulation

of MpSYP13B at the forming cell plate. A single confocal image of FM4-64-stained

dividing thallus cells expressing mCitrine-MpSYP13B are shown. Asterisks indicate

forming cell plates. (j) A maximum intensity projection image of a thallus cell

expressing MpCLC1-mCitrine and mRFP-MpSYP6A under the regulation of their own

regulatory elements and the Mp*EF1α* promoter, respectively. (k) Time-lapse single

confocal images of thallus cells expressing MpCLC1-Citrine under the regulation of the

Mp*EF1α* promoter (upper panels) and merged images with FM4-64 signals.

Autofluorescence from chlorophyll is pseudo-coloured in blue. Bars = 50 μm in (b - e)

and 10 μm in (f), (g), (i), (j), and (k).

Supplementary Fig. S2. Subcellular localization of organelle markers in oil body

cells

(a) Gateway constructs used in this study. pMpGWB301 _pro_Mp*SYP2:*mCitrine-GW was

previously prepared[41]. (b) Maximum intensity projection images of thallus cells

including the oil body cell expressing mCitrine-fused markers for the Golgi apparatus

(MpSFT1 and MpGOS11), *trans*-Golgi network (MpSYP4 and MpSYP6A), and

tonoplast (MpSYP2 and MpVAMP71). The organelle markers were expressed under the

Mp*SYP12B*, Mp*SYP2*, and/or own promoters. Bright field images are also shown. (c)

Single confocal images of thallus cells including the oil body cell expressing

endoplasmic reticulum markers. mCitrine-MpSEC20 and mCitrine-MpUSE1A were

expressed under the Mp*SYP12B* promoter. mCitrine-MpSEC22 and mRFP-HDEL were

expressed using their own regulatory elements and the Mp*EF1α* promoter, respectively.

(d) A single confocal image of thallus cells expressing sec-mRFP expressed under the

Mp*EF1α* promoter. Green, magenta, and blue pseudo-colours indicate fluorescence

from mCitrine, mRFP, and chlorophyll, respectively. Bars = 10 μm.

Supplementary Fig. S3. Phenotype of Mp*syp12b* mutants

(a) Schematic representation of the Mp*SYP12B* gene structure. Gray and black boxes

indicate untranslated and coding regions, respectively, and arrowheads indicate

mutation sites in two independently generated mutants. The Mp*syp12b-1* mutant was

generated by a homologous recombination-mediated gene targeting method[38]. (b) The

genome (black letters) and translated amino acid (magenta letters) sequences of

Mp*SYP12B* and the Mp*syp12b-2*[ge] mutant, with the red letters indicating the mutated

region. The PAM sequence for gRNA is underlined. The 0 layer at the centre of the

SNARE motif is shown. (c) Single confocal images of thallus cells including the oil

body cell expressing mCitrine-MpSYP13A under its own regulatory elements, or

sec-mRFP under the Mp*SYP12B* promoter in Tak-1, Mp*syp12b-1*, and Mp*syp12b-2*[ge]

lines. Bright field images are also shown. (d) Single confocal images of thallus cells

802   including the oil body cell expressing mCitrine-MpSYP12B under its own regulatory

803   elements in Tak-1 and Mp*syp12b* mutants. Green, magenta, and blue pseudo-colours

804   indicate fluorescence from mCitrine, mRFP, and chlorophyll, respectively. Bars = 10

805   μm.

Supplementary Fig. S4. Screening and generation of Mp*erf13* mutants

(a) The flow chart of the screen for mutants defective in oil body formation from

T-DNA-insertion lines. (b) A maximum likelihood phylogenetic tree of proteins with

one ERF/AP2 domain. The colour code is shown in Fig S1a. The branch lengths are

proportional to the estimated number of substitutions per site. Bootstrap probability is

indicated as a percentage on each branch with at least 50% support. A more detailed tree

was presented previously[17]. (c) Schematic representation of the Mp*ERF13* gene

structure and mutations generated in this study. Gray and black boxes indicate the UTR

and coding sequences, respectively. Asterisks with numbers indicate sites of designed

gRNA to generate Mp*erf13-1^{ge}* (*1 and *4) and Mp*erf13-2^{ge}* (*2 and *3). (d)

PCR-based genotyping of Tak-1, Mp*erf13^{GOF}*, Mp*erf13-1^{ge}*, and Mp*erf13-2^{ge}*. The

combinations and annealing sites of primers (a to d) are shown in (c). (e) The genomic

and predicted amino acid sequences of the Mp*ERF13* locus in Tak-1 and Mp*erf13*

mutants. Light blue, dark blue, and magenta letters indicate UTR, coding, and predicted

amino acid sequences, respectively. The PAM sequences for gRNAs are underlined.

The caret indicates the indel site. (f) Fluorescent and bright-field images of

BODIPY-stained three-week-old thalli of Tak-1, Mp*erf13-1^{ge}*, and Mp*erf13-2^{ge}*. Bars =

0.5 mm. (g) The number of oil bodies visualized with BODIPY in a unit area (2.0 mm $\times$

2.0 mm). Bars indicate means $\pm$ SD. Statistical analyses between Tak-1 and each

genotype were conducted using a two-tailed Welch's *t*-test. Sample numbers were 23

thalli for Tak-1, 24 for Mp*erf13-1^{ge}*, and 26 for Mp*erf13-2^{ge}*. *p*-values are 5.14×10^{-11} for

Mp*erf13-1^{ge}* and 5.14×10^{-11} for Mp*erf13-2^{ge}*.

Supplementary Fig. S5. Gene expression in Mp*erf13* mutants

(a) List of selected differentially expressed genes (DEGs) for qRT-PCR analysis. The

full list of DEGs are shown as Supplementary Table S1. (b) The relative mRNA

expression of 11 DEGs in Tak-1, Mp*erf13^{GOF}*, Mp*erf13-1^{ge}*, and Mp*erf13-2^{ge}* measured

by qRT-PCR. Mp*APT* was used as an internal reference. Error bars represent ± SD.

Means are indicated above the bars. Statistical comparison between Tak-1 and each

genotype was conducted with a two-tailed Welch's *t*-test. **, $p < 0.01$. (c) Correlation

of the expression levels between the RNA-Seq analysis and qRT-PCR experiment. Both

the x- and y- axes are shown in a log$_2$FC scale. R^2 indicates the correlation coefficient.

Supplementary Fig. S6. Pill bug feeding assay

(a and b) Ten-day-old thalli of Tak-1, Mp$erf13^{GOF}$ and Mp$erf13\text{-}2^{ge}$ (a) were

co-cultivated with starved pill bugs for 24 hr (b). Bars = 1 cm. (c) Ratios of thallus area

before and after co-cultivation with pill bugs were calculated and indicated the change

in thallus area change (n = 36 thalli for each genotype). Bars indicate means ± SD.

Statistical analyses between Tak-1 and each genotype were conducted with a two-tailed

Welch's *t*-test. The *p*-values were 1.63×10^{-5} for Mp$erf13^{GOF}$ and 4.91×10^{-6} for

Mp$erf13\text{-}2^{ge}$.

Supplementary Table S1

The list of DEGs in Mp$erf13$ mutants identified in RNA-Seq analysis.

Supplementary Table S2

List of primers used in this study.

Supplementary movie S1. Formation of clathrin-coated vesicles at the oil body

membrane

A time-lapse movie of an oil body cell expressing MpCLC1-Citrine. Green and blue

pseudo-colours represent fluorescence from Citrine and chlorophyll, respectively. 32×

real time. Bar = 5 μm.

Supplementary movie S2. Expression of MpSYP12B in a Marchantia thallus

A five-day-old thallus expressing 2×YFP driven by the Mp*SYP12B* promoter observed

by light sheet microscopy. Green and blue pseudo-colours represent fluorescence from

Citrine and chlorophyll, respectively. Bar = 400 μm.