Title

Streptococcus pyogenes M1T1 variants induce an inflammatory phenotype in neutrophils

Authors/Affiliations

Jonathan G. Williams[1,2], Diane Ly[1,2], Nicholas J. Geraghty[1,2], Jason D. McArthur[1,2], Heema K. N.

Vyas[1,2], Jody Gorman[1,2], James A. Tsatsaronis[3], Ronald Sluyter[1,2], Martina L. Sanderson-Smith[1,2,4,*]

[1]Illawarra Health and Medical Research Institute, Wollongong, New South Wales, Australia.

[2]Molecular Horizons and School of Chemistry and Molecular Bioscience, University of

Wollongong, Wollongong, New South Wales, Australia.

[3]College of Science, Health and Engineering, La Trobe University, Bundoora, Victoria, Australia.

[4]Lead contact

Contact Info

*Correspondence: Illawarra Health and Medical Research Institute (IHMRI), School of Chemistry

and Molecular Bioscience, University of Wollongong, Wollongong, NSW, 2522, Australia. E-mail:

martina@uow.edu.au

Summary

Invasive infections due to Group A *Streptococcus* (GAS) advance rapidly causing tissue

degradation and unregulated inflammation. Neutrophils are the primary immune cells that

respond to GAS. M1T1 GAS isolate 5448 and animal passaged variant 5448AP, containing a *covS*

mutation, were assessed for their influence on neutrophils. 5448AP proliferated during

neutrophil co-incubation and induced lower neutrophil reactive oxygen species production when

compared with 5448. Infection with both strains invoked neutrophil death, however apoptosis

was reduced in response to 5448AP. Both strains induced neutrophil caspase-1 activation *in vitro*,

and elicited IL-1β and TNF-α release from neutrophils. GAS infection altered the cell-surface

expression of CD16 and CD31 on neutrophils. *In vivo* mouse studies confirmed activation of

caspase-1 in neutrophils, while 5448AP infection increased circulating IL-1β. GAS infections that

promote an inflammatory neutrophil phenotype may contribute to increased inflammation yet

ineffective bacterial eradication, contributing to the severity of invasive GAS infections.

Keywords

Polymorphonuclear leukocyte, PMN, apoptosis, pyroptosis, caspase-1, caspase-3, caspase-4, IL-

1β, TNF-α, IL-8, IL-18, cell-death, *covS*, 5448, 5448AP, LEGENDplex, FLICA, inflammation,

inflammasome

Introduction

Invasive infections due to the obligate human pathogen *Streptococcus pyogenes* (Group A

Streptococcus; GAS) are characterised by unregulated inflammation and high mortality (Davies et

al., 1996, Lamagni et al., 2008, Nelson et al., 2016, O'Grady et al., 2007, O'Loughlin et al., 2007,

Svensson et al., 2000). The highly virulent GAS M1T1 clone 5448 is well studied (Aziz and Kotb,

2008, Tart et al., 2007, Walker et al., 2007). The propensity of M1T1 GAS to cause invasive

infections is due, in part, to the increased frequency of spontaneous mutations in the two-

component control of virulence regulatory system (covRS) (Walker et al., 2007), resulting in

differential expression of multiple virulence factors, many of which are implicated in immune

defence (Kilsgård et al., 2016, Sumby et al., 2005, Sumby et al., 2006, Walker et al., 2007).

Compelling evidence describes the resistance of *covS* mutant GAS (mutation to the sensor protein

of covRS) to killing by human neutrophils (polymorphonuclear leukocytes, PMNs) (Ato et al.,

2008, Maamary et al., 2010, Walker et al., 2007). 5448AP (animal passaged) is one M1T1 *covS*

mutant strain that shows resistance to killing by human neutrophils, has increased bacterial

dissemination and results in decreased survival in mouse models of infection (Fiebig et al., 2015,

Maamary et al., 2010, Walker et al., 2007). Due to these characteristics, 5448AP can be used as a

model strain to explore *covS* induced hypervirulence and the host response to invasive GAS

infection.

Neutrophils are the primary innate immune cell-type that defend against bacterial pathogens,

with their presence contributing to the overarching inflammatory response (Kobayashi et al.,

2018). Proteins of neutrophil origin are highly abundant at sites of invasive GAS infection

(Edwards et al., 2018). The neutrophil life cycle is tightly regulated to avoid the release of

cytotoxic contents that can damage surrounding tissue and contribute further to inflammation

(Epstein and Weiss, 1989, Faurschou and Borregaard, 2003). Under normal conditions cytotoxic

reactive oxygen species (ROS) produced by neutrophils kill phagocytosed bacteria (Nguyen et al.,

2017). An increased production of ROS signals for the induction of the anti-inflammatory cell-

death pathway apoptosis (Lundqvist-Gustafsson and Bengtsson, 1999). During apoptosis,

neutrophils are carefully decommissioned, ensuring no unregulated release of material, then

cleared by a secondary phagocyte, macrophages, through a process known as efferocytosis

(Bratton and Henson, 2011). The cleavage of caspase-3 to the p17 form denotes the execution of

apoptosis (Nicholson et al., 1995). Increased neutrophil apoptosis is described during GAS

infection (Kobayashi et al., 2003). Pro-inflammatory neutrophil death has also been reported

during GAS infection (Tsatsaronis et al., 2015). One form of pro-inflammatory death is pyroptosis,

involving the formation of an inflammasome and subsequent caspase-1 or caspase-4 activation,

culminating in cell-lysis (Man and Kanneganti, 2016, Sollberger et al., 2012). Deregulation of (or

excessive) pro-inflammatory neutrophil-death may therefore contribute further to inflammatory

pathology, as demonstrated during infection with other pathogens, including *Pseudomonas*

aeruginosa (Ryu et al., 2017) and *Staphylococcus aureus* (Ghimire et al., 2018, Greenlee-Wacker

et al., 2017, Ryu et al., 2017). Inflammation is also heavily influenced by cell cytokine production,

where increased release can invoke neutrophil dysfunction, affecting neutrophil survival,

antibacterial function or even migration (Brown et al., 2006, Kipnis, 2013, Kovach and Standiford,

2012, Norrby-Teglund et al., 2000). Previous work outlines how GAS take advantage of

inadequate or exacerbated immune responses and focus upon the bacterial mechanisms involved

(Cole et al., 2011). Although models to investigate leukocyte inflammasome activation have been

proposed (Tran et al., 2019), a limited number of studies have investigated the neutrophil

response to GAS specifically (Kobayashi et al., 2003). Here, using an *in vitro* model, we temporally

describe the effect two M1T1 GAS strains have upon human neutrophil death, signalling and

inflammatory profile. We further characterise this interaction in murine neutrophils, using an *in*

vivo model of intradermal invasive GAS infection. Additionally, we identify differences in the

neutrophil response to GAS due to *covS* mutation. We hypothesise that changes to neutrophil

function by GAS can promote invasive GAS infection by exacerbating the inflammatory response,

decreasing antibacterial capacity and inducing inflammatory neutrophil death.

Results

GAS persistence and proliferation occurs during a dampened neutrophil ROS response

Neutrophil phagocytic ability is a predominant and well-established mechanism of immune

defence. Both GAS strains 5448 and 5448AP rapidly associated with human neutrophils, with

near maximal association observed within 30 min (Figures 1A and S1A-C). Higher percentages of

neutrophils were associated with 5448 compared to 5448AP (Figure 1A). 5448 and 5448AP also

induced ROS production in neutrophils, with less ROS produced by neutrophils incubated in the

presence and absence of 5448AP compared to those incubated with 5448 (Figure 1B). As such,

the rate of neutrophil ROS production was significantly increased compared to the neutrophil

only control during 5448, but not 5448AP, infection (p<0.01, Figure S1D), suggesting dampened

antibacterial function in neutrophils in response to 5448AP. This difference in GAS-induced ROS

production inversely corresponded to GAS proliferation in the presence of neutrophils, with high

numbers of 5448AP observed compared to 5448 over 180 min (Figure 1C). Differential GAS

proliferation required viable neutrophils, as 5448 and 5448AP proliferation in the presence of

neutrophil lysates was reduced and similar for both strains over 180 min (Figure 1D). Finally,

incubation of neutrophils with the phagocytosis inhibitor cytochalasin D significantly increased

5448, but not 5448AP, survival (p<0.001, Figure S1E), indicating that 5448 killing is

predominantly mediated via a phagocytic-related pathway. Collectively, these data indicate that

compared to 5448, 5448AP associates less readily with human neutrophils leading to reduced

ROS production and increased survival of this strain.

Figure 1. GAS persistence and proliferation occurs during a dampened neutrophil ROS

response. (A) Association of fluorescent GAS strains with human neutrophils (PMNs, n=6

donors), analysed via flow cytometry (Figure S1A-B). (B) Infection of human neutrophils with

GAS invokes ROS production. Representative of triplicate measurements from three separate

experiments (n=3 donors), shown as mean change in fluorescence units over time ($\Delta F.min^{-1}$). GAS

strains were incubated in the presence of (C) human neutrophils (n=8 donors) or (D) lysed

human neutrophils (n=4 donors) over 180 min with surviving bacterial concentration

determined as percentage over inoculum. Results are the pooled means±SD (of triplicate

measurements for panels C and D). **p<0.01 and ****p<0.0001.

Neutrophil death is delayed during infection with *covS* GAS

GAS can induce death of human neutrophils (Kobayashi et al., 2003, Tsatsaronis et al., 2015),

however the mechanism remains poorly described. Phosphatidylserine (PS) exposure is a

hallmark of the induction of multiple death pathways including apoptosis and pyroptosis (Man and Kanneganti, 2016, Wang et al., 2013). Human neutrophils were incubated with 5448, 5448AP or in the absence of GAS then the binding of annexin-V (AV), as a measure of PS exposure, and the binding of Zombie Fixable Viability Kit (Z), as a measure of membrane integrity, were analysed by flow cytometry (Figure 2). Cells were defined as viable (AV$^-$Z$^-$), PS exposed (AV$^+$Z$^-$) or dead (AV$^+$Z$^+$) (Figure 2C). Neutrophils incubated without GAS remained viable over 180 min (Figure 2D), with 13% exposing PS (Figure 2E) and 3% dead after this time (Figure 2F). In contrast, incubation of neutrophils with either 5448 or 5448AP resulted in increased PS exposure and cell-death over 180 min, with significantly more cell-death during 5448 infection compared to 5448AP infection at 30 and 60 min (p<0.05, Figure 2F). Delay in the initial induction of death in response to 5448AP suggests extended neutrophil viability during the early stages of infection (30-60 min).

GAS infection activates caspase-1 in neutrophils but reduced apoptosis occurs in response to *covS* GAS

The above data (Figure 2) suggests that both GAS strains induce rapid neutrophil death. To explore the mechanism of GAS-induced neutrophil death and whether this differs between GAS strains, human neutrophil lysates were separated by SDS-PAGE and screened for the abundance of active caspase-3, caspase-1 and caspase-4 via immunoblotting and quantified using densitometry (Figures 3A and S2A-D). Cleavage of executioner caspase-3 to p17 is a definitive hallmark of apoptosis induction (Nicholson et al., 1995). Conversely, caspase-1 and caspase-4 activity can indicate inflammasome activation, which in turn can cleave downstream effectors of pyroptosis (Man and Kanneganti, 2016, Sollberger et al., 2012). Moreover, a recent report has identified the caspase-1 p46 molecule as the principal active species during inflammasome activation in the cell (Boucher et al., 2018). Neutrophils incubated in the absence of GAS showed limited evidence of caspase-3 cleavage to p17 (Figures 3A and S2A), while levels of caspase-1 reduced over 180 min (Figures 3A and S2B-C). Infection of neutrophils with 5448 significantly

increased the presence of caspase-3 p17 at 180 min (p>0.0001) compared to both 5448AP-

infected and uninfected neutrophils (Figures 3A and S2A). In contrast, incubation of neutrophils

with either GAS strain caused an upregulation of caspase-1 (Figures 3A and S2B-C). Compared to

uninfected neutrophils, 5448 infection significantly increased pro-caspase-1 (50kDa) at 30 min,

60 min (p<0.05) and 180 min (p<0.01), whilst 5448AP infection only significantly increased pro-

caspase-1 at 180 min (p<0.001, Figures 3A and S2B). Further, 5448 and 5448AP infected

neutrophils sustain p46 expression over 180 min, though this difference was not statistically

significant (Figure 3A and S2C). Significant differences were not seen in the expression of

caspase-1 between 5448 and 5448AP infected neutrophils. Similar to caspase-1, the abundance

of full-length caspase-4 in neutrophils incubated in the absence of GAS declined over time

(Figures 3A and S2D). In contrast, 5448 infection increased caspase-4 expression at 30 (p<0.01),

60 (p<0.001) and 180 min compared to uninfected neutrophils (p<0.0001, Figures 2A and S2D).

5448AP infection however, only increased caspase-4 expression at 180 min (p<0.01, Figures 3A

and S2D).

These above data suggest both GAS strains induce pyroptosis in neutrophils, but the response to

5448AP is delayed. Therefore, to explore this possibility further, human neutrophils were

incubated with 5448, 5448AP or in the absence of GAS and caspase-1 activity assessed using a

fluorogenic flow cytometric caspase-1 assay (Figures 3B and S3A-C). 5448 infected neutrophils

showed increased active caspase-1 at 60 min and 180 min (p<0.05), whilst 5448AP infection

increased caspase-1 at 180 min (p<0.05) compared to uninfected neutrophils (Figure 3B). No

significant differences were observed between strains although caspase-1 activity was slightly

reduced during 5448AP infection compared to 5448 (Figure 3B), similar to the immunoblot data

for caspase-1 p46 (Figures 3A and S2C). Thus, GAS infection of human neutrophils corresponds

to increased caspase-1 activity and caspase-4 expression, consistent with inflammasome activation and pyroptosis.

Figure 2. GAS induced neutrophil death is delayed during infection with *covS* mutant. Human neutrophils (PMNs) were stained with Annexin V-FITC and Zombie Aqua Fixable Viability Kit via flow cytometry and initially gated upon (A) 'Singlets' (FSC-A vs. FSC-H, 10,000 events collected). 'All PMNs' (live and dead) were selected using (B) a gate excluding GAS. Cells were then compared for fluorescence of (C) annexin V-FITC/phosphatidylserine (PS) binding against Zombie live/dead viability dye. Purified PMNs were sampled over 180 min to measure (D) 'Viable' (AV$^-$Z$^-$), (E) 'PS exposed' (AV$^+$Z$^-$), and (F) 'Dead' (AV$^+$Z$^+$), neutrophils during GAS infection (n=3 donors). Results are pooled means±SD . *p<0.05, **p<0.01, ***p<0.001 and ****p<0.0001, with black asterisks denoting significance from control and grey asterisks between 5448 and 5448AP.

Proinflammatory cytokines IL-1β and TNF-α are released by neutrophils in response to GAS infection

The release of the inflammatory cytokine IL-1β occurs due to caspase-1 activation and is

associated with the induction of pyroptosis (Broz et al., 2010). Uncoordinated release of various

inflammatory cytokines can exacerbate infection (Tecchio et al., 2014). During GAS infection,

disease severity is negatively correlated to cytokine concentration (Norrby-Teglund et al., 2000).

Therefore, human neutrophils were incubated with 5448, 5448AP or in the absence of GAS and

supernatants assessed for cytokines using a flow cytometric bead assay. Neutrophils incubated

in the absence of GAS failed to release IL-1β (Figure 3C) and TNF-α (Figure 3D) despite releasing

increasing amounts of IL-8 over 360 min (Figure 3E). Incubation with GAS induced IL-1β release

from neutrophils at 180 min and 360 min, with significantly greater release during 5448AP

infection compared to 5448 at 180 and 360 min ($p<0.05$, $p<0.0001$, Figure 3C). Incubation with

GAS also induced significant release of TNF-α from neutrophils at 180 min ($p<0.001$) and 360 min

($p<0.0001$), with significantly greater release during 5448 infection compared to 5448AP at 360

min ($p<0.0001$, Figure 3D). 5448 induced significant release of IL-8 compared to uninfected from

neutrophils at 60 min and 180 min ($p<0.05$), with significantly greater release than 5448AP at 60

min ($p<0.05$, Figure 3E). Neutrophils released minimal amounts of IL-18 over 360 min, and

infection with both 5448 or 5448AP did not change this response (Figure 3F), suggesting that

neutrophils have no major role in IL-18 release over 360 min *in vitro*. These data indicate that

both 5448 and 5448AP induce IL-1β release from human neutrophils, consistent with the

hypothesis that both GAS strains induce pyroptosis in these cells. Moreover, the results indicate

differential cytokine release from human neutrophils infected with GAS, with 5448AP inducing

greater IL-1β release and 5448 inducing greater TNF-α and IL-8 release.

Figure 3. GAS infection activates caspase-1 in human neutrophils and induces proinflammatory cytokine IL-1β and TNF-α release. (A) Human neutrophil (PMN) lysates were prepared at 30, 60 and 180 min during GAS infection and compared to uninfected neutrophils (0, 30, 60, 180 min) via immunoblotting of caspase-3, caspase-1 and caspase-4. Images shown are from a single donor and are representative of triplicate experiments using different donors. Immunoblot bands were quantified (ImageJ) and normalised over total protein (Figure S2). SE=short exposure, LE=long exposure. (B) Caspase-1 activation in human neutrophils due to GAS infection was confirmed via flow cytometry using FLICA (FAM-YVAD-FMK) (n=3 donors). Flow cytometry gating strategy used singlets/PMNs (figure S3A) and FAM-FLICA fluorescence (Figure S3B). The release of cytokines from neutrophils during GAS infection was measured using the LEGENDplex™ human inflammation cytometric bead assay over 360 min. Neutrophils differentially released (C) IL-1β, (D) TNF-α, (E) IL-8 and (F) IL-18 in response to GAS

infection (duplicate measurements, n=6 donors). (B-F) Results are the pooled means±SD. *p<0.05, **p<0.01, ***p<0.001 and ****p<0.0001.

GAS infection may cause neutrophil dysfunction

Neutrophil adhesion and activation are central to the innate immune response, with CD11b (Mac-1) and CD66b (CEACAM8) playing key respective roles (Arnaout, 1990, Power et al., 2001, Skubitz et al., 1996). CD16 (FcγRIII) facilitates opsonisation of pathogens and is down-regulated during apoptosis (Dransfield et al., 1994, Fossati et al., 2002), whilst down-regulation of CD31 (PECAM-1) facilitates clearance of apoptotic neutrophils and the resolution of inflammation (Brown et al., 2002, Kurosaka et al., 2003). Therefore, to further explore the impact of GAS on neutrophil function, human neutrophils were incubated with 5448, 5448AP or in the absence of GAS and the expression of cell-surface CD11b, CD66b, CD16 and CD31 was assessed using flow cytometry (Figure 4A-F). In the absence of GAS, neutrophils displayed minor increases in CD11b expression over time, while incubation with GAS resulted in minor decreases over time, with 5448 inducing a greater loss of CD11b than 5448AP, though these differences were not statistically significant (p=0.289, Figure 4C). Neutrophils incubated in the absence of GAS revealed minor increases in cell-surface CD66b expression over time (Figure 4D). Incubation with GAS further increased expression of CD66b, with 5448 inducing a slightly greater increase than 5448AP, though these differences were not statistically significant (p=0.162, Figure 4D). Incubation of neutrophils in the absence of GAS resulted in a steady decline in CD16 expression over time, a loss significantly increased by co-incubation with either GAS strain 180 min post-infection (p<0.0001, Figure 4E). Notably, incubation of neutrophils with 5448 induced a significantly greater loss of CD16 expression than 5448AP incubation at all earlier time points (Figure 4E). Similar to CD16, incubation of neutrophils in the absence of GAS resulted in a steady decline in CD31 expression over time, a loss significantly increased by co-incubation with either GAS strain 180 min post-infection (p<0.0001, Figure 4F). Moreover, incubation of neutrophils with 5448 induced a significantly greater loss of CD31 expression than 5448AP incubation at earlier time points.

Collectively, these data indicate that GAS has minimal impact upon human neutrophil CD11b and

CD66b, but increases the loss of CD16 and CD31 expression, which may impact opsonisation of

GAS and subsequent efferocytosis of neutrophils by macrophages, during infection.

Figure 4. GAS infection changes neutrophil functionality. Human neutrophils (PMNs) were

analysed using flow cytometry, sequentially gated upon (A) 'Singlets' (FSC-A vs. FSC-H, 10,000

events collected) then (B) viable 'PMNs' (FSC-A vs. FSC-H). Purified neutrophils infected with GAS

were sampled over 180 min for the cell-surface expression of (C) CD11b-FITC (n=4 donors), (D)

CD66b-PerCP/Cy5.5 (n=5 donors), (E) CD16-FITC (n=4 donors) and (F) CD31-PE/Cy7 (n=4

donors). Results are the pooled means±SD. *p<0.05, **p<0.01, ***p<0.001 and ****p<0.0001, with

black asterisks denoting significance from control and grey asterisks between 5448 and 5448AP.

GAS infection activates caspase-1 in neutrophils *in vivo*

248 The above data indicate that GAS influences human neutrophils *in vitro.* To explore if GAS may

249 alter neutrophils *in vivo*, we assessed the neutrophil response in a mouse model of GAS infection

250 (Ly et al., 2014, Maamary et al., 2010, Tsatsaronis et al., 2014). C57BL/6J mice were injected

251 intradermally with 5448, 5448AP or saline and euthanised at 6 or 24 hours post infection, with

252 neutrophil populations (CD45[+]/CD11b[+]/Ly-6G[+]) and caspase-1 activation characterised by flow

cytometry (Figures 5A-B and S4A-F). Infection of mice with either GAS strain increased neutrophil populations at sites of injection, however only 5448AP infection resulted in significantly greater neutrophil numbers than the saline control at 24 h (p<0.01, Figure 5A). The percentage of circulating neutrophils in the blood was also significantly increased due to 5448 (p<0.05) and 5448AP (p<0.01) infection at 6 h, however returned to baseline after 24 h (Figure S4G), indicating the importance of neutrophils during the early stages of infection. Murine neutrophils lavaged from sites of saline injection had low amounts of active caspase-1 (Figure 5B). Infection with 5448AP, but not 5448, significantly increased caspase-1 activation at 6 h (p<0.05), while both strains increased caspase-1 activation at 24 h (p<0.01, Figure 5B). Cytokines were assessed in murine serum using a flow cytometric bead assay. IL-1β concentrations remained low for saline control and 5448 infected mice but were significantly increased at 24 h during 5448AP infection (p<0.0001, Figure 5C).

Figure 5. GAS infection *in vivo* activates caspase-1 in murine neutrophils. (A) Percentage of CD45+ cells lavaged from site of injection characterised as neutrophils (PMNs, CD11b+/Ly-6G+) by flow cytometry (Figure S4A-E). (B) Caspase-1 activation in neutrophils at site of GAS infection was confirmed via flow cytometry using FLICA 660 (660-YVAD-FMK, Figure S4F). Each data point represents the results from a single mouse where n=4. (C) The release of IL-1β in mouse serum during GAS infection was measured using the LEGENDplex™ mouse inflammation cytometric bead assay at 6 and 24 h post infection where duplicate measurements are shown and n=4. Results are the mean±SD. **p<0.01, ***p<0.001 and ****p<0.0001 where significance is denoted from control or as indicated by a line.

Discussion

Here we have mapped the human neutrophil response to GAS, addressing cell-death, signalling and the inflammatory profile during the early stages of M1T1 (5448 and 5448AP) infection. For the first time, we describe an inflammatory neutrophil phenotype, as evidenced by caspase-1 activation, that may promote inflammation and exacerbate GAS disease. Most notably, changes to the expression of cell-surface proteins CD16 and CD31 may impede the removal of bacteria and clearance of neutrophils at sites of infection, hindering the resolution of inflammation. Further, we elucidate differences between 5448 and 5448AP that aid in understanding the survival and proliferation of GAS harbouring *covS* mutations in the presence of human neutrophils.

The temporal expression of neutrophil cell surface markers has not previously been investigated in the context of GAS infection. Our data show that significant reductions to both CD16 and CD31 occur in neutrophils due to GAS infection *in vitro*. CD16 functions as a receptor for opsonised bacteria and immune complexes (Fossati et al., 2002), whilst the loss of CD16 has also been associated with the induction of apoptosis (Dransfield et al., 1994). The induction of apoptosis following bacterial phagocytosis is advantageous to the host as it promotes controlled removal of bacteria and the resolution of inflammation (Kobayashi et al., 2018). We hypothesise that the retention of CD16 during infection promotes a reduction in apoptosis and prolonged degranulation at the site of infection. Further, CD31 expression is reduced during GAS infection, facilitating the eventual removal of neutrophils from sites of infection (Kobayashi et al., 2003). Parallel to CD16, CD31 is retained during 5448AP infection of neutrophils to a greater extent than 5448 infection. This may result in neutrophils remaining at sites of infection longer, further increasing inflammation. Additionally, we report a trend towards increased cell-surface expression of CD11b and CD66b during GAS infection. Migration of neutrophils to sites of infection, facilitated by CD11b-mediated neutrophil adhesion (Arnaout, 1990), may therefore be affected during GAS infection. Down-regulation of CD11b can prevent accumulation of neutrophils at inflammatory sites (Huston et al., 2009), however retention, such as that seen

during 5448AP infection may contribute to prolonged inflammation. Neutrophil activation can be determined through the expression of cell-surface CD66b following stimulation (Skubitz et al., 1996). Reduced cell-surface CD66b expression in response to 5448AP infection may therefore indicate a reduction to neutrophil function and further explain the resistance to neutrophil-mediated killing. Collectively, the neutrophil response is altered and inflammation prolonged at sites of *covS* mutant GAS infection.

Disruption of the role neutrophils play during the resolution of GAS infection and induction of inflammatory death pathways has been reported previously (Kobayashi et al., 2003, Tsatsaronis et al., 2015). Proinflammatory activation of the NLRP3 inflammasome in macrophages has been demonstrated for M1 GAS (Valderrama et al., 2017), here we provide the first evidence that GAS activates caspase-1 in human and murine neutrophils, consistent with inflammasome activation and pyroptosis. Caspase-1 activation was consistent during infection with both M1T1 GAS isolates, however the release of inflammatory cytokine IL-1β, a hallmark of clinical invasive infection (LaRock and Nizet, 2015), was significantly increased in response to 5448AP infection *in vitro* and *in vivo*. In murine bone marrow-derived dendritic cells caspase-1 activation is essential for IL-1β release (Schneider et al., 2017). The activation of caspase-1 in human and murine neutrophils does not correlate directly to the amount of IL-1β released, and an alternative mechanism may facilitate IL-1β maturation. Type I interferons are known regulators of IL-1β in GAS infected mice (Castiglia et al., 2016), where increased IL-1β release in murine serum seen in this study may also be attributed to disruption of this homeostatic relationship. Increased IL-1β, being a neutrophil chemoattractant, therefore may further promote inflammation at sites of infection following release (Chen et al., 2007), supporting the hypothesis that *covS* mutant GAS promote greater levels of inflammation during GAS infection.

Previous studies have characterised differences between 5448 and 5448AP, demonstrating resistance to neutrophil-mediated killing *in vitro* and increased bacterial dissemination and decreased survival *in vivo* following the acquisition of *covS* mutations (Fiebig et al., 2015,

Maamary et al., 2010, Walker et al., 2007). Here, we demonstrate that 5448AP is not only resistant

to neutrophil-mediated killing but proliferates in the presence of functional neutrophils. GAS

survival has been demonstrated intracellularly in murine neutrophils and human macrophages

(Medina et al., 2003a, Medina et al., 2003b, O'Neill et al., 2016) and can even be facilitated by host

red blood cells (Wierzbicki et al., 2019). Increased 5448AP proliferation may result from the

reduction in neutrophil ROS response during infection. GAS has previously been shown to display

resistance to ROS through numerous virulence mechanisms (Henningham et al., 2015). The

absence of a hostile environment may further permit bacterial replication. Additionally, the

increased production of ROS can induce neutrophil death via apoptosis (Geering and Simon,

2011). Reduced neutrophil ROS production during 5448AP infection may therefore explain the

reduction seen in caspase-3 activation (apoptosis). Previously it has been demonstrated that GAS

M1 strain MGAS5005 modulates neutrophil apoptosis (Kobayashi et al., 2003), a finding which is

supported by the current study.

Changes to the function of neutrophils during the initial innate immune response have the

potential to hinder mechanisms essential for the resolution of infection. Disruption to these

processes during GAS infection may induce this neutrophil phenotype we have described, with

increased inflammatory properties and potential to exacerbate infection. Caspase-1 activation

during GAS infection of neutrophils indicates possible inflammasome activation and death via

pyroptosis. Further studies should explore the precise cell death mechanism. A reduction in

apoptosis is evident during *covS* GAS infection, as has been reported for non-M1T1 GAS

(Tsatsaronis et al., 2015), whilst modulation of apoptosis has previously been implicated as a

factor contributing to GAS survival (Kobayashi et al., 2003). The current study aids in

understanding the complex host-pathogen interaction identifying host factors that may

contribute to severe pathology of invasive GAS infection. Future studies may be able to isolate

host-specific targets that could be exploited to control inflammation and tissue damage during

invasive GAS infection.

Acknowledgements

We thank Professor Marijka Batterham (University of Wollongong) for her statistical

consultation. We thank Associate Professor Thomas Proft (University of Auckland, New Zealand)

for pLZ12Km2-P23R:TA. We also thank our generous blood donors for their participation and

time.

Author Contributions

Conceptualisation, M.S.S, R.S., J.A.T., J.G.W.; Methodology, M.S.S., R.S., J.G.W., J.G., J.D.M., D.L.;

Investigation, J.G.W., D.L., N.J.G., H.K.N.V.; Data and figure curation, J.G.W., D.L., R.S., M.S.S.; Writing-

Original draft, J.G.W.; Writing-Review and editing, M.S.S., R.S., J.G.W., D.L., N.J.G., J.D.M., H.K.N.V.,

J.G., J.A.T.; Funding acquisition, M.S.S and D.L.; Resources, M.S.S., J.D.M.; Supervision, M.S.S., R.S.,

J.G.

Declaration of Interests

The authors declare no competing interests.

STAR Methods

Ethics statement

All experiments involving the use of human blood were conducted with informed consent of

healthy volunteers, approved and authorised by the University of Wollongong Human Research

Ethics Committee (Protocol HE08/250). All experiments involving the use of animals were

approved and authorised by the University of Wollongong Animal Ethics Committee (Protocol

AE18/10).

Bacterial strains and culture

Escherichia coli MC106 were grown in Luria–Bertani Broth (LB) at 37 °C with constant shaking.

M1T1 *S. pyogenes* invasive clinical isolate 5448 (*emm1*) and hypervirulent animal passaged

5448AP (containing a non-functional control of virulence regulator mutation) have been

described previously (Aziz et al., 2004, Chatellier et al., 2000). GAS strains were routinely cultured

at 37°C on horse-blood agar (Thermo Fisher) and enumerated on yeast supplemented (1% w/v)

Todd-Hewitt (THY, Benton Dickson) agar. Static overnight cultures were grown at 37°C in THY

then sub-inoculated (1:10) into fresh THY and grown to mid-logarithmic phase, antibiotics were

added when required (100 μg/ml ampicillin and 200 μg/ml kanamycin). Bacterial pellets were

washed twice with phosphate buffered saline (PBS) then resuspended at the specified

multiplicity of infection (MOI) and assay specific media conditions.

Construction of GFP expression vector pLZ12Km2-P23R-TA:GFP for GAS

In brief, stable GFP expression by GAS was created by synthesizing the ribosomal binding site

(RBS) and *gfp* gene from pDC*erm*-GFP (Ly et al., 2014) into the pUC57 plasmid (GenScript),

resulting in pUC57-RBSGFP plasmid. RBS and *gfp* from pUC57-RBSGFP were then sub-cloned

using NotI into the toxin–antitoxin (TA) stabilized expression plasmid pLZ12Km2-P23R:TA,

kindly provided by Associate Professor Thomas Proft (University of Auckland, New Zealand) (Loh

and Proft, 2013), to produce pLZ12Km2-P23R-TA:GFP. The resultant plasmid system utilises the

Streptococcal ω–ε–ζ TA cassette to achieve segregational plasmid stability under non-selective

conditions (Lioy et al., 2010).

Specifically, plasmids pLZ12Km2-P23R:TA and pUC57-RBSGFP (engineered with RBS and *gfp*

from pDC*erm*-GFP) were first transformed into chemically competent *E. coli* MC106 via

electroporation. The plasmids were then retrieved and purified from *E. coli* cultures using a

Wizard® Plus SV Minipreps DNA Purification System (Promega). The pLZ12Km2-P23R:TA

plasmid was digested with 20 U NotI enzyme and further treated with 5 U shrimp alkaline

phosphatase. The pUC57-RBSGFP plasmid was also incubated with 20 U NotI to excise the RBS

and *gfp* gene from pUC57-RBSGFP, and these were then ligated into digested pLZ12km2-P23R:TA

plasmid. The resulting plasmid, pLZ12Km2-P23R-TA:GFP, was then transformed and cloned in

MC106 *E. coli*, and later extracted and purified. pLZ12Km2-P23R-TA:GFP was transformed into

GAS using standard GAS electroporation techniques (McLaughlin and Ferretti, 1995).

Transformed GAS was confirmed for GFP expression via flow cytometry.

Isolation of human neutrophils

Venous human blood was drawn into 10 mL lithium heparin-coated Vacutainer (Benton Dickson)

tubes and layered over equal volumes of Polymorphprep (Axis Shield) and centrifuged as per

manufacturer's instructions. The resulting layer of neutrophils was isolated and erythrocytes

hypotonically lysed. Isotonic concentration was restored with Hank's Balanced Salt solution

(without Ca^{2+} or Mg^{2+}, Corning Inc.). Prior to experimentation neutrophils were resuspended at

specified concentrations in complete medium, Roswell Park Memorial Institute (RPMI)-1640

medium (Life Technologies) containing 2% heat-inactivated autologous plasma and 2 mM L-

glutamine (Life Technologies), unless otherwise stated. Neutrophil viability was assessed via

Trypan Blue (Sigma-Aldrich) staining and neutrophil purity was assessed using a Benton Dickson

LSR Fortessa X-20 flow cytometer via distinct forward and side scatter profiles or CD66b-

peridinin chlorophyll protein (PerCP)/Cy5.5 (clone G10F5, BioLegend) expression. Neutrophils

were maintained at room temperature throughout processing.

***In vitro* infection of neutrophils with GAS**

417   Purified human neutrophils were seeded in either 96-well plates at MOI (GAS:neutrophil) 1:10

418   (survival and proliferation) or 10:1 (ROS production and cytokine release), 24-well plates at 10:1

419   (phagocytosis, Annexin-V/Viability staining, CD expression and FAM-FLICA caspase-1 activation)

420   or 6-well plates at 10:1 (immunoblotting) and incubated at 37°C in 5% CO_2.

GAS survival and proliferation

422   Human neutrophils were co-cultured with GAS or lysed via three freeze-thaw cycles prior to

423   incubation. To block phagocytosis neutrophils were preincubated with 10 μm cytochalasin D

424   (Cayman Chemicals) in complete medium at 37°C for 30 min prior to infection. At indicated time

425 points neutrophils were hypotonically lysed and surviving bacteria serially diluted before plating

426 on THY agar.

GAS phagocytosis

428 Human neutrophils were co-cultured with GAS expressing GFP for the times indicated in

429 complete medium. Cells were removed and washed twice with 10% (v/v) heat-inactivated foetal

430 bovine serum (FBS, Bovogen Biologicals) diluted in PBS, followed by data acquisition via flow

431 cytometry.

ROS production

433 ROS production was assessed as previously described (Kobayashi et al., 2003). Briefly, prior to

434 infection, human neutrophils were incubated in complete medium containing 25 μM

435 dichlorofluorescein (DCF, Molecular Probes) for 45 min at room temperature (RT). ROS

436 production was measured fluorometrically ($_{ex}$485 nm $_{em}$520 nm) using a POLARstar Omega plate

437 reader (BMG Labtech).

Flow cytometry

For human neutrophil in vitro assays, flow cytometry data was acquired using a BD LSR Fortessa

X-20 with excitation lasers; violet (405 nm), blue (488 nm), yellow/green (561 nm) and red (640

nm). Bandpass filters for Zombie Aqua Fixable Viability Kit (525/50), fluorescein isothiocynate

(FITC)/GFP/FAM FLICA (525/50), PerCP-Cy5.5 (695/40) R-phycoerythrin (PE, 586/15),

propidium iodide (PI, 610/20), PE-Cy7 (780/60) and allophycocyanin (APC, 670/30) were used.

For murine infection studies, flow cytometry data was acquired using an Invitrogen Attune NxT

with excitation lasers; violet (405 nm), blue (488 nm), yellow (561 nm) and red (638 nm).

Bandpass filters VL1 (440/50), VL2 (512/25), BL1 (530/30), YL1 (585/16), YL3 (695/40) and

RL2 (720/30) were used. Data was analysed using FlowJo software V10.6.1 (TreeStar Inc.).

Annexin-V/viability staining

To assess cell viability human neutrophils at indicated times were washed with PBS and incubated with Zombie Aqua Fixable Viability Kit (BioLegend) for 15 min at RT. Cells were washed again once with PBS, then 10% (v/v) FBS in PBS and stained with Annexin-V-FTIC (BioLegend) for 15 min at RT. Cells were analysed immediately via flow cytometry.

Immunoblotting and antibodies

Following co-culture in the presence or absence of GAS, human neutrophil lysates were prepared in RIPA buffer (150 mM NaCl, 5 mM EDTA, 50 mM Tris, 1.0% (v/v) Triton X-100, 0.1% (w/v) SDS, 0.5% (w/v) sodium deoxycholate, 2x cOmplete protease inhibitor cocktail (Roche), 1 mM phenylmethylsulphonyl fluoride, 5 mM sodium pyrophosphate, 5 mM sodium molybdate and 5 mM β-glycerophosphate) following co-culture in the presence or absence of GAS and incubated for 30 min on ice with intermittent vortexing. Soluble fractions were separated via centrifugation at 4°C, snap frozen in liquid N_2 and stored at -80°C. Protein concentrations were determined by comparison to bovine serum albumin standards using the DC Protein Assay (Bio-Rad) and absorbance (A_{750nm}) measured using a POLARstar Omega plate reader. Neutrophil lysates (20 µg) were separated on 4-20% TGX Stain-Free protein gels (Bio-Rad) as per the manufacturer's running conditions then activated for 5 min under UV to determine total protein (Bio-Rad ChemiDoc XR, Image Lab Software). Proteins were transferred to PVDF membranes (Bio-Rad), blocked, then incubated overnight at 4°C with caspase-1 polyclonal (1:1000, #2225), caspase-3 polyclonal (1:1000, #9662), caspase-4 polyclonal (1:1000, #4450) or caspase-8 monoclonal (1:1000, #4790) antibodies (Cell Signalling Technology). PVDF membranes were washed thrice between incubations for 5 min with tris-buffered saline containing 0.1% (v/v) Tween 20. PVDF membranes were incubated with horseradish peroxidase-conjugated goatαrabbit IgG (1:5000, Invitrogen) for 1 h at RT and detected using Clarity or Clarity Max Western ECL Blotting Substrate (Bio-Rad) and imaged (Amersham AI600). Bands were quantified using ImageJ software (National Institutes of Health) as area under the peak and normalised over UV determined total protein.

FAM-FLICA caspase-1 activation

Caspase-1 activation was measured using FAM-FLICA Caspase-1 assay kit (ImmunoChemistry Technologies) as per the manufacturer's instructions. In brief, following co-culture of human neutrophils with GAS, cells were removed and incubated in serum free medium (RPMI-1640) containing 1 x FAM-YVAD-FMK for 60 min at 37°C. Cells were then analysed for caspase-1 activation via flow cytometry.

Cytometric cytokine bead assay

Human neutrophil supernatants were at indicated times during GAS infection, snap frozen with liquid N_2 and stored at -80°C for a period no longer than 5 days. Neutrophils were assessed for the release of cytokines IL-1β, IL-8, IL-18 and TNF-α using the LEGENDplex™ human inflammation panel bead-based immunoassay (BioLegend) and flow cytometry as per manufacturer's instructions. In brief, samples and standards were incubated with beads and detection antibodies for 120 min at room temperature, with shaking (600 rpm) protected from light. PE-conjugated streptavidin was then added and incubated for a further 30 min at room temperature, with shaking (600 rpm) protected from light. Beads were washed twice with LEGENDplex™ wash buffer and data collected via flow cytometry. Beads were initially gated upon FSC-A and SSC-A (A and B populations). Cytokines have signature APC fluorescence with quantitative expression determined using PE fluorescence in comparison to cytokine standards. Data was analysed using the LEGENDplex™ software V8.0 (VigeneTech Inc.).

Cluster of differentiation expression

Human neutrophil cell surface CD expression was assessed during GAS infection. Neutrophils at indicated times were washed with PBS then routinely incubated with Zombie Aqua Fixable Viability Kit for 15 min at RT. Cells were washed once again in PBS, followed by 10% (v/v) FBS in PBS. Neutrophils were stained with fluorochrome-conjugated antibodies CD11b-FITC (Clone ICRF44), CD31-PE/Cy7 (Clone WM59), or CD66b-PerCP/Cy5.5 (Clone G10F5) (BioLegend) and

500   CD16-FITC (Clone 3G8, Benton Dickson) for 15 min at RT. Neutrophils were washed with 10%

501   (v/v) FBS in PBS then analysed via flow cytometry.

Murine intradermal GAS infection model

Intradermal GAS challenge of C57BL/6J mice has been described previously (Ly et al., 2014,

Maamary et al., 2010, Tsatsaronis et al., 2014). In brief, equal numbers of 6-8 week old male and

female C57BL/6J mice (Australian BioResources) were anesthetised via isoflurane inhalation,

followed by two intradermal injections into shaved left and right flanks with either 1×10^8 CFU

of mid-logarithmic GAS or 100 μL sterile 0.7% saline. Mice were euthanised via slow-fill CO_2

asphyxiation at 6 or 24 h post-infection. Blood collected via cardiac puncture was separated into

two aliquots for serum and flow cytometric analysis. Serum was collected by clotting for 1 h at RT

then placed on ice until centrifugation at 1200 x g for 10 min at 4°C, then immediately stored at -

80°C until use. Remaining blood samples were mixed with 20 μL 0.5% sodium citrate (w/v) per

mL of blood and placed on ice. Equal volume of PBS was added to the blood sample and

centrifuged at 350 x g for 5 min. For erythrocyte lysis, samples were incubated with 1 mL of

ammonium-chloride-potassium lysing buffer (150 mM NH_4Cl, 1 mM $KHCO_3$, 0.1 mM Na_2CO_3, pH

7.3) for 5 min with gentle agitation and centrifuged at 350 x g for 5 min. This process was repeated

and cells were then washed with PBS and resuspended in 500 μL PBS and placed on ice.

Additionally, sites of infection were lavaged with 1 mL of sterile 0.7% saline, and fluid collected

from both right and left flanks was pooled to increase cell numbers. Pooled samples were kept

cold until centrifugation at 350 x G for 5 min, resuspended in 500 uL of PBS and placed on ice.

Both blood and lavage fluid samples were analysed via flow cytometry.

Flow cytometric analysis of murine cells

Cell viability was assessed as described above. To assess caspase-1 activation and CD expression,

cells were initially washed with 1 mL of PBS, followed by 1 mL of 10% (v/v) FBS in PBS. Cells

were then centrifuged and simultaneously stained with 1 x 660-YVAD-FMK (FLICA 660 Caspase-

1 assay kit, ImmunoChemistry Technologies) for caspase-1 activation and with fluorochrome-

conjugated antibodies CD45-BV421 (clone 30-F11), CD11b-PE/Cy5 (clone M1-70) and Ly-6G-PE

(clone 1A8) (Biolegend) in RPMI 1640 media for 30 min at RT . Cells were washed with 10% (v/v)

FBS in PBSand analysed via flow cytometry.

Murine serum IL-1β concentration

530   IL-1β release was measured in murine serum using the LEGENDplex™ mouse inflammation panel

531   bead-based immunoassay (BioLegend) and flow cytometry as described above except shaking

532   was performed at 800 rpm.

Statistical analyses

Graphs were created using Prism 6 (GraphPad Software Inc.) and statistical analysis was

performed using Prism 6 and IBM SPSS Statistics 25 (IBM® corporation). Data was analysed

using one-way and two-way ANOVA to determine significant differences and adjusted using

Tukey HSD corrections or as stated otherwise. To account for the nested nature of the human

neutrophil Annexin-V/Zombie and CD data with repeated measures over time in the same donor,

a linear mixed model was used to determine significant interaction between treatment and time.

Post-hoc tests were performed by multiple one-way ANOVA at each time point where multiple

comparisons were adjusted using Tukey HSD corrections. *In vivo* data was analysed by two-way

ANOVA where differences are shown between treatment and time, also using Tukey HSD

corrections. Values presented for 'p' represent interaction (treatment*time) unless otherwise

stated. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.

 Supplemental Information

Supplementary figure 1. The association of fluorescent GAS (GFP) to human neutrophils

determined via flow cytometry. (A) Primary neutrophil (PMN) 'Singlets' cell selection (FSC-A

vs. FSC-H) and (B) viable 'PMNs' discrimination (FSC-A vs. SSC-A). (C) Representative histogram

showing relative fluorescence (GFP) of GAS infected neutrophils and gating for the selection of

GAS-neutrophil association. (D) Rate of ROS production by neutrophils between 30-60 min (n=6

donors). (E) GAS killing by human neutrophils is inhibited at 30 min following pre-incubation

with actin polymerising molecule cytochalasin D (n=4 donors, Sidak's multiple comparison).

Results are the pooled means±SD (of triplicate measurements for panels D and E). **$p < 0.01$,

***$p < 0.001$.

Supplementary figure 2. Quantification of protein bands in human neutrophil lysates

identified through immunoblotting. (A) Caspase-3 p17, (B) caspase-1 full-length, (D) caspase-

1 p46 and (D) caspase-4 immunoblots were imaged then analysed using ImageJ where area under

the curve was normalised over total protein. Results are the means±SD where 3 donors were

used. *p<0.05, **p<0.01, ***p<0.001 and ****p<0.0001, with black asterisks denoting significance

to control and grey asterisks between 5448 and 5448AP.

Supplementary figure 3. Human neutrophils were assessed for the activation of caspase-1

and release of cytokines via and flow cytometry. (A) Primary neutrophil (PMN) 'Singlet' cell

selection (FSC-A vs. FSC-H) and (B) viable 'PMNs' discrimination (FSC-A vs. SSC-A). (C)

Representative plot showing unstained and GAS infected neutrophils (Singlets/PMNs) with

FLICA (FAM-YVAD-FMK) fluorescence. Gating strategy for LEGENDplex cytometric bead assay

where (D) beads A and B were gated (FSC-A vs. SSC-A). Gates for beads A (shown) and B were

assessed for (E) PE vs. APC fluorescence where cytokines had unique APC fluorescence and

expression was measured via PE fluorescence when compared to a standard curve.

Supplementary figure 4. Flow cytometric sequential gating strategy for identification of

murine neutrophils determining caspase-1 activation. (A) Primary 'Singlets' cell selection

(FSC-A vs. FSC-H) and (B) 'Cells' discrimination (FSC-A vs. SSC-A) of murine blood and lavage

fluid. 'Cells' were further gated upon (C) CD45-BV421 expression and (D) CD11b-PE/Cy5

expression. Neutrophils (PMNS) were defined as (E) Ly-6G-PE+ cells from the CD45+/CD11b+

population. Neutrophils were assessed for (F) caspase-1 activation using FLICA 660 (660-YVAD-

FMK). (G) Percentage of neutrophils (CD11b+/Ly6G+) of CD45+ leukocytes in the blood where

each data point represents the result from a single mouse where n=4. Results are the means±SD.

*p<0.05 and **p<0.01 where significance is depicted from control or as indicated by a line.

References

Arnaout, M. A. 1990. Structure and function of the leukocyte adhesion molecule CD11/CD18. *Blood,*
75**,** 1037-1050.
Ato, M., Ikebe, T., Kawabata, H., Takemori, T. & Watanabe, H. 2008. Incompetence of neutrophils to
invasive group A streptococcus is attributed to induction of plural virulence factors by
dysfunction of a regulator. *PLoS ONE,* 3.
Aziz, R. K. & Kotb, M. 2008. Rise and persistence of global M1T1 clone of Streptococcus pyogenes.
Emerging Infectious Diseases, 14**,** 1511-1517.
Aziz, R. K., Pabst, M. J., Jeng, A., Kansal, R., Low, D. E., Nizet, V. & Kotb, M. 2004. Invasive M1T1
group A Streptococcus undergoes a phase-shift in vivo to prevent proteolytic degradation of
multiple virulence factors by SpeB. *Molecular Microbiology,* 51**,** 123-134.
Boucher, D., Monteleone, M., Coll, R. C., Chen, K. W., Ross, C. M., Teo, J. L., Gomez, G. A., Holley, C.
L., Bierschenk, D., Stacey, K. J., Yap, A. S., Bezbradica, J. S. & Schroder, K. 2018. Caspase-1
self-cleavage is an intrinsic mechanism to terminate inflammasome activity. *Journal of*
Experimental Medicine, 215**,** 827-840.
Bratton, D. L. & Henson, P. M. 2011. Neutrophil clearance: When the party is over, clean-up begins.
Trends in Immunology, 32**,** 350-357.
Brown, K., Brain, S., Pearson, J., Edgeworth, J., Lewis, S. & Treacher, D. 2006. Neutrophils in
development of multiple organ failure in sepsis. *Lancet,* 368**,** 157-169.
Brown, S., Heinisch, I., Ross, E., Shaw, K., Buckley, C. O. & Savill, J. 2002. Apoptosis disables CD31-
mediated cell detachment from phagocytes promoting binding and engulfment. *Nature,* 418**,**
200-203.
Broz, P., Von Moltke, J., Jones, J. W., Vance, R. E. & Monack, D. M. 2010. Differential requirement for
caspase-1 autoproteolysis in pathogen-induced cell death and cytokine processing. *Cell Host*
and Microbe, 8**,** 471-483.
Castiglia, V., Piersigilli, A., Ebner, F., Janos, M., Goldmann, O., Damböck, U., Kröger, A., Weiss, S.,
Knapp, S., Jamieson, A. M., Kirschning, C., Kalinke, U., Strobl, B., Müller, M., Stoiber, D.,
Lienenklaus, S. & Kovarik, P. 2016. Type i Interferon Signaling Prevents IL-1β-Driven Lethal
Systemic Hyperinflammation during Invasive Bacterial Infection of Soft Tissue. *Cell Host and*
Microbe, 19**,** 375-387.
Chatellier, S., Ihendyane, N., Kansal, R. G., Khambaty, F., Basma, H., Norrby-Teglund, A., Low, D. E.,
Mcgeer, A. & Kotb, M. 2000. Genetic relatedness and superantigen expression in group A
streptococcus serotype M1 isolates from patients with severe and nonsevere invasive
diseases. *Infection and Immunity,* 68**,** 3523-3534.
Chen, C. J., Kono, H., Golenbock, D., Reed, G., Akira, S. & Rock, K. L. 2007. Identification of a key
pathway required for the sterile inflammatory response triggered by dying cells. *Nature*
Medicine, 13**,** 851-856.
Cole, J. N., Barnett, T. C., Nizet, V. & Walker, M. J. 2011. Molecular insight into invasive group A
streptococcal disease. *Nature Reviews Microbiology,* 9**,** 724-736.
Davies, H. D., Mcgeer, A., Schwartz, B., Green, K., Cann, D., Simor, A. E. & Low, D. E. 1996. Invasive
group A streptococcal infections in Ontario, Canada. *New England Journal of Medicine,* 335**,**
547-554.
Dransfield, I., Buckle, A. M., Savill, J. S., Mcdowall, A., Haslett, C. & Hogg, N. 1994. Neutrophil
apoptosis is associated with a reduction in CD16 (FcγRIII) expression. *Journal of Immunology,*
153**,** 1254-1263.
Edwards, R. J., Pyzio, M., Gierula, M., Turner, C. E., Abdul-Salam, V. B. & Sriskandan, S. 2018.
Proteomic analysis at the sites of clinical infection with invasive Streptococcus pyogenes.
Scientific Reports, 8.
Epstein, F. H. & Weiss, S. J. 1989. Tissue Destruction by Neutrophils. *New England Journal of*
Medicine, 320**,** 365-376.

Faurschou, M. & Borregaard, N. 2003. Neutrophil granules and secretory vesicles in inflammation.
Microbes and Infection, 5**,** 1317-1327.
Fiebig, A., Loof, T. G., Babbar, A., Itzek, A., Koehorst, J. J., Schaap, P. J. & Nitsche-Schmitz, D. P. 2015.
Comparative Genomics of Streptococcus pyogenes M1 isolates differing in virulence and
propensity to cause systemic infection in mice. *International Journal of Medical*
Microbiology, 305**,** 532-543.
Fossati, G., Moots, R. J., Bucknall, R. C. & Edwards, S. W. 2002. Differential role of neutrophil Fcγ
receptor IIIb (CD16) in phagocytosis, bacterial killing, and responses to immune complexes.
Arthritis and Rheumatism, 46**,** 1351-1361.
Geering, B. & Simon, H. U. 2011. Peculiarities of cell death mechanisms in neutrophils. *Cell Death*
and Differentiation, 18**,** 1457-1469.
Ghimire, L., Paudel, S., Jin, L., Baral, P., Cai, S. & Jeyaseelan, S. 2018. NLRP6 negatively regulates
pulmonary host defense in Gram-positive bacterial infection through modulating neutrophil
recruitment and function. *PLoS Pathogens,* 14.
Greenlee-Wacker, M. C., Kremserová, S. & Nauseef, W. M. 2017. Lysis of human neutrophils by
community-associated methicillin-resistant Staphylococcus aureus. *Blood,* 129**,** 3237-3244.
Henningham, A., Döhrmann, S., Nizet, V. & Cole, J. N. 2015. Mechanisms of group A Streptococcus
resistance to reactive oxygen species. *FEMS Microbiology Reviews,* 39**,** 488-508.
Huston, J. M., Rosas-Ballina, M., Xue, X., Dowling, O., Ochani, K., Ochani, M., Yeboah, M. M.,
Chatterjee, P. K., Tracey, K. J. & Metz, C. N. 2009. Cholinergic neural signals to the spleen
down-regulate leukocyte trafficking via CD11b. *Journal of Immunology,* 183**,** 552-559.
Kilsgård, O., Karlsson, C., Malmström, E. & Malmström, J. 2016. Differential compartmentalization of
Streptococcus pyogenes virulence factors and host protein binding properties as a
mechanism for host adaptation. *International Journal of Medical Microbiology*.
Kipnis, E. 2013. Neutrophils in sepsis: Battle of the bands. *Critical Care Medicine,* 41**,** 925-926.
Kobayashi, S. D., Braughton, K. R., Whitney, A. R., Voyich, J. M., Schwan, T. G., Musser, J. M. & Deleo,
F. R. 2003. Bacterial pathogens modulate an apoptosis differentiation program in human
neutrophils. *Proceedings of the National Academy of Sciences of the United States of*
America, 100**,** 10948-10953.
Kobayashi, S. D., Malachowa, N. & Deleo, F. R. 2018. Neutrophils and Bacterial Immune Evasion.
Journal of Innate Immunity, 10**,** 432-441.
Kovach, M. A. & Standiford, T. J. 2012. The function of neutrophils in sepsis. *Current Opinion in*
Infectious Diseases, 25**,** 321-327.
Kurosaka, K., Takahashi, M., Watanabe, N. & Kobayashi, Y. 2003. Silent Cleanup of Very Early
Apoptotic Cells by Macrophages. *Journal of Immunology,* 171**,** 4672-4679.
Lamagni, T. L., Darenberg, J., Luca-Harari, B., Siljander, T., Efstratiou, A., Henriques-Normark, B.,
Vuopio-Varkila, J., Bouvet, A., Creti, R., Ekelund, K., Koliou, M., Reinert, R. R., Stathi, A.,
Strakova, L., Ungureanu, V., Schalén, C., Jasir, A., Ioannou, Y., Kriz, P., Motlova, J.,
Hammerum, A., Kaltoft, M. S., Iivonen, J., Loubinoux, J., Mihaila, L., Van Der Linden, M.,
Lütticken, R., Papaparaskevas, J., Zachariadou, L., Baldassarri, L., Orefici, G., Straut, M.,
Norrby-Teglund, A., Keshishian, C. & Neal, S. 2008. Epidemiology of severe Streptococcus
pyogenes disease in Europe. *Journal of Clinical Microbiology,* 46**,** 2359-2367.
Larock, C. N. & Nizet, V. 2015. Inflammasome/IL-1β responses to streptococcal pathogens. *Frontiers*
in Immunology, 6.
Lioy, V. S., Pratto, F., De La Hoz, A. B., Ayora, S. & Alonso, J. C. 2010. Plasmid pSM19035, a model to
study stable maintenance in Firmicutes. *Plasmid,* 64**,** 1-17.
Loh, J. M. S. & Proft, T. 2013. Toxin-antitoxin-stabilized reporter plasmids for biophotonic imaging of
Group A streptococcus. *Applied Microbiology and Biotechnology,* 97**,** 9737-9745.
Lundqvist-Gustafsson, H. & Bengtsson, T. 1999. Activation of the granule pool of the NADPH oxidase
accelerates apoptosis in human neutrophils. *Journal of Leukocyte Biology,* 65**,** 196-204.

Ly, D., Taylor, J. M., Tsatsaronis, J. A., Monteleone, M. M., Skora, A. S., Donald, C. A., Maddocks, T., Nizet, V., West, N. P., Ranson, M., Walker, M. J., Mcarthur, J. D. & Sanderson-Smith, M. L. 2014. Plasmin(ogen) Acquisition by Group A Streptococcus Protects against C3b-Mediated Neutrophil Killing. *Journal of Innate Immunity,* 6**,** 240-250.

Maamary, P. G., Sanderson-Smith, M. L., Aziz, R. K., Hollands, A., Cole, J. N., Mckay, F. C., Mcarthur, J. D., Kirk, J. K., Cork, A. J., Keefe, R. J., Kansal, R. G., Sun, H., Taylor, W. L., Chhatwal, G. S., Ginsburg, D., Nizet, V., Kotb, M. & Walker, M. J. 2010. Parameters governing invasive disease propensity of non-M1 serotype group A Streptococci. *Journal of Innate Immunity,* 2**,** 596-606.

Man, S. M. & Kanneganti, T. D. 2016. Converging roles of caspases in inflammasome activation, cell death and innate immunity. *Nature Reviews Immunology,* 16**,** 7-21.

Mclaughlin, R. E. & Ferretti, J. J. 1995. Electrotransformation of Streptococci. *Methods in Molecular Biology (Clifton, N.J.),* 47**,** 185-193.

Medina, E., Goldmann, O., Toppel, A. W. & Chhatwal, G. S. 2003a. Survival of Streptococcus pyogenes within host phagocytic cells: A pathogenic mechanism for persistence and systemic invasion. *Journal of Infectious Diseases,* 187**,** 597-603.

Medina, E., Rohde, M. & Chhatwal, G. S. 2003b. Intracellular survival of Streptococcus pyogenes in polymorphonuclear cells results in increased bacterial virulence. *Infection and Immunity,* 71**,** 5376-5380.

Nelson, G. E., Pondo, T., Toews, K. A., Farley, M. M., Lindegren, M. L., Lynfield, R., Aragon, D., Zansky, S. M., Watt, J. P., Cieslak, P. R., Angeles, K., Harrison, L. H., Petit, S., Beall, B. & Van Beneden, C. A. 2016. Epidemiology of invasive group a streptococcal infections in the United States, 2005-2012. *Clinical Infectious Diseases,* 63**,** 478-486.

Nguyen, G. T., Green, E. R. & Mecsas, J. 2017. Neutrophils to the ROScue: Mechanisms of NADPH oxidase activation and bacterial resistance. *Frontiers in Cellular and Infection Microbiology,* 7.

Nicholson, D. W., Ali, A., Thornberry, N. A., Vaillancourt, J. P., Ding, C. K., Gallant, M., Gareau, Y., Griffin, P. R., Labelle, M., Lazebnik, Y. A., Munday, N. A., Raju, S. M., Smulson, M. E., Yamin, T. T., Yu, V. L. & Miller, D. K. 1995. Identification and inhibition of the ICE/CED-3 protease necessary for mammalian apoptosis. *Nature,* 376**,** 37-43.

Norrby-Teglund, A., Chatellier, S., Low, D. E., Mcgeer, A., Green, K. & Kotb, M. 2000. Host variation in cytokine responses to superantigens determine the severity of invasive group A streptococcal infection. *European Journal of Immunology,* 30**,** 3247-3255.

O'grady, K. a. F., Kelpie, L., Andrews, R. M., Curtis, N., Nolan, T. M., Selvaraj, G., Passmore, J. W., Oppedisano, F., Carnie, J. A. & Carapetis, J. R. 2007. The epidemiology of invasive group A streptococcal disease in Victoria, Australia. *Medical Journal of Australia,* 186**,** 565-569.

O'loughlin, R. E., Roberson, A., Cieslak, P. R., Lynfield, R., Gershman, K., Craig, A., Albanese, B. A., Farley, M. M., Barrett, N. L., Spina, N. L., Beall, B., Harrison, L. H., Reingold, A. & Van Beneden, C. 2007. The epidemiology of invasive group A streptococcal infection and potential vaccine implications: United States, 2000-2004. *Clinical Infectious Diseases,* 45**,** 853-862.

O'neill, A. M., Thurston, T. L. M. & Holden, D. W. 2016. Cytosolic replication of group a Streptococcus in human macrophages. *mBio,* 7.

Power, C., Wang, J. H., Sookhai, S., Wu, Q. D. & Redmond, H. P. 2001. Proinflammatory effects of bacterial lipoprotein on human neutrophil activation status, function and cytotoxic potential in vitro. *Shock,* 15**,** 461-466.

Ryu, J. C., Kim, M. J., Kwon, Y., Oh, J. H., Yoon, S. S., Shin, S. J., Yoon, J. H. & Ryu, J. H. 2017. Neutrophil pyroptosis mediates pathology of P. aeruginosa lung infection in the absence of the NADPH oxidase NOX2. *Mucosal Immunology,* 10**,** 757-774.

Schneider, K. S., Groß, C. J., Dreier, R. F., Saller, B. S., Mishra, R., Gorka, O., Heilig, R., Meunier, E., Dick, M. S., Ćiković, T., Sodenkamp, J., Médard, G., Naumann, R., Ruland, J., Kuster, B., Broz,

P. & Groß, O. 2017. The Inflammasome Drives GSDMD-Independent Secondary Pyroptosis and IL-1 Release in the Absence of Caspase-1 Protease Activity. *Cell Reports,* 21**,** 3846-3859.

Skubitz, K. M., Campbell, K. D. & Skubitz, A. P. N. 1996. CD66a, CD66b, CD66c, and CD66d each independently stimulate neutrophils. *Journal of Leukocyte Biology,* 60**,** 106-117.

Sollberger, G., Strittmatter, G. E., Kistowska, M., French, L. E. & Beer, H. D. 2012. Caspase-4 is required for activation of inflammasomes. *Journal of Immunology,* 188**,** 1992-2000.

Sumby, P., Porcella, S. F., Madrigal, A. G., Barbian, K. D., Virtaneva, K., Ricklefs, S. M., Sturdevant, D. E., Graham, M. R., Vuopio-Varkila, J., Hoe, N. P. & Musser, J. M. 2005. Evolutionary origin and emergence of a highly successful clone of serotype M1 group A Streptococcus involved multiple horizontal gene transfer events. *Journal of Infectious Diseases,* 192**,** 771-782.

Sumby, P., Whitney, A. R., Graviss, E. A., Deleo, F. R. & Musser, J. M. 2006. Genome-wide analysis of group A streptococci reveals a mutation that modulates global phenotype and disease specificity. *PLoS Pathogens,* 2**,** 0041-0049.

Svensson, N., Öberg, S., Henriques, B., Holm, S., Källenius, G., Romanus, V. & Giesecke, J. 2000. Invasive group A streptococcal infections in Sweden in 1994 and 1995: Epidemiology and clinical spectrum. *Scandinavian Journal of Infectious Diseases,* 32**,** 609-614.

Tart, A. H., Walker, M. J. & Musser, J. M. 2007. New understanding of the group A Streptococcus pathogenesis cycle. *Trends in Microbiology,* 15**,** 318-325.

Tecchio, C., Micheletti, A. & Cassatella, M. A. 2014. Neutrophil-derived cytokines: Facts beyond expression. *Frontiers in Immunology,* 5.

Tran, T. a. T., Grievink, H. W., Lipinska, K., Kluft, C., Burggraaf, J., Moerland, M., Tasev, D. & Malone, K. E. 2019. Whole blood assay as a model for in vitro evaluation of inflammasome activation and subsequent caspase-mediated interleukin-1 beta release. *PLoS ONE,* 14.

Tsatsaronis, J. A., Ly, D., Pupovac, A., Goldmann, O., Rohde, M., Taylor, J. M., Walker, M. J., Medina, E. & Sanderson-Smith, M. L. 2015. Group A Streptococcus Modulates Host Inflammation by Manipulating Polymorphonuclear Leukocyte Cell Death Responses. *Journal of Innate Immunity,* 7**,** 612-622.

Tsatsaronis, J. A., Walker, M. J. & Sanderson-Smith, M. L. 2014. Host Responses to Group A Streptococcus: Cell Death and Inflammation. *PLoS Pathogens,* 10.

Valderrama, J. A., Riestra, A. M., Gao, N. J., Larock, C. N., Gupta, N., Ali, S. R., Hoffman, H. M., Ghosh, P. & Nizet, V. 2017. Group A streptococcal M protein activates the NLRP3 inflammasome. *Nature Microbiology,* 2**,** 1425-1434.

Walker, M. J., Hollands, A., Sanderson-Smith, M. L., Cole, J. N., Kirk, J. K., Henningham, A., Mcarthur, J. D., Dinkla, K., Aziz, R. K., Kansal, R. G., Simpson, A. J., Buchanan, J. T., Chhatwal, G. S., Kotb, M. & Nizet, V. 2007. DNase Sda1 provides selection pressure for a switch to invasive group A streptococcal infection. *Nature Medicine,* 13**,** 981-985.

Wang, Q., Imamura, R., Motani, K., Kushiyama, H., Nagata, S. & Suda, T. 2013. Pyroptotic cells externalize eat-me and release find-me signals and are efficiently engulfed by macrophages. *International Immunology,* 25**,** 363-372.

Wierzbicki, I. H., Campeau, A., Dehaini, D., Holay, M., Wei, X., Greene, T., Ying, M., Sands, J. S., Lamsa, A., Zuniga, E., Pogliano, K., Fang, R. H., Larock, C. N., Zhang, L. & Gonzalez, D. J. 2019. Group A Streptococcal S Protein Utilizes Red Blood Cells as Immune Camouflage and Is a Critical Determinant for Immune Evasion. *Cell Reports*.