Adipose-tissue derived signals control bone remodelling.

He Fu[1], Maria-Bernadette Madel[2,3], Dominique D. Pierroz[4], Mariano Schiffrin[1], Carine Winkler[1], Anne Wilson[5], Cécile Pochon[6], Barbara Toffoli[1], Jean-Yves Jouzeau[6], Federica Gilardi[1], Serge Ferrari[7], Nicolas Bonnet[7], Claudine Blin-Wakkach[2,3], Béatrice Desvergne[1*] and David Moulin[6*]

[1]Center for Integrative Genomics, Genopode, Lausanne Faculty of Biology and Medicine CH-1015 Lausanne, Switzerland; [2] CNRS UMR7370, Laboratoire de PhysioMédecine Moléculaire, Nice, France; [3]Université Côte d'Azur, Nice, France; [4]International Osteoporosis Foundation, 1260 Nyon, Switzerland; [5]Ludwig Centre for Cancer Research, Department of Oncology, University of Lausanne, 1066 Epalinges, Switzerland, [6]Université de Lorraine, CNRS, IMoPA, F-54000 Nancy, France ; and [7]Division of Bone Diseases, Department of Internal Medicine Specialties, Geneva University Hospital & Faculty of Medicine, Geneva 14, Switzerland

*These authors contributed equally and are corresponding authors.

Running title
Lack of adiponectin drives hyperosteoclastogenesis in lipoatrophic mice

Summary

Long bones from mammals host blood cell formation and contain multiple cell types, including adipocytes. Overabundance of adipocytes in bone marrow is associated with pathological conditions such as age-related osteoporosis or marrow aplasia induced by irradiation or chemotherapy. However, physiological functions of bone marrow adipocytes are poorly documented. Herein, we investigated the consequence of total adipocyte deficiency on bone homeostasis in mice. By generating adipocyte-deficient mice using PPARγ deletion, we found that lipoatrophy leads to dramatic alterations of trabecular and cortical femoral bone. More specifically, cortical bone of these mice is extremely porous and poorly defined due to the excessive presence of active bone-resorbing osteoclasts. Thus, despite the supposed role of PPARγ in osteoclast formation, osteoclastogenesis is highly active in this mouse model. Two independent models of lipoatrophy recapitulated this phenotype, demonstrating that hyperosteoclastogenesis is not intrinsically linked to PPARγ deficiency. We further showed that adiponectin, a cytokine produced by adipocytes and mesenchymal stromal cells, is a potent inhibitor of osteoclastogenesis *in vitro* and *in vivo*. Consistently, pharmacological activation of adiponectin receptors by the synthetic agonist AdipoRon inhibits mature osteoclast activity both in mouse and human by blocking podosome formation. This adiponectin-inhibiting action on mature osteoclasts occurs though AMPK activation, thereby demonstrating that even fully differentiated and active osteoclasts are sensitive to adipose-derived signals. Finally, we showed that AdipoRon inhibits bone erosion in vivo in a murine model of inflammatory bone loss. Collectively, these data reveal that adiponectin-producing cells are key regulators of bone homeostasis, and that preserving functional bone marrow adiponectin pathway can improve bone integrity in the context of metabolic and inflammatory disorders.

Keywords: Adiponectin, Osteoclast, Bone Marrow Adiposity, Cortical Bone porosity

Introduction

Bone homeostasis is a result of constant remodelling activities, with a balance of bone resorption and bone formation. The most prevalent disorder of bone homeostasis is osteoporosis, affecting men and women, with a particular high prevalence in post-menopausal women. The low efficacy of actual anti-osteoporotic therapeutic options highlights the need of better understanding the pathophysiological processes at work.

The role of adipocyte in bone homeostasis is one of the factors raising great interest, but it faces quite some complexity. An illustration of this complexity, is the fact that obesity is traditionally considered to be protective against osteoporosis, consequently to an increased proliferation and differentiation of osteoblasts and osteocytes stimulated by mechanical load (Felson et al., 1993). However, obesity in postmenopausal women and in men correlates with increased marrow adiposity, alteration of the bone microarchitecture, and increased risk of fracture (Sheu and Cauley, 2011) .Thus, the respective roles of ageing, increased BMI, and visceral *vs* bone marrow fat on bone homeostasis remain difficult to disentangle (Cornish et al., 2018; Horowitz et al., 2017).

Peroxisome proliferator-activated receptors (PPARs) are ligand-activated transcription factors that belong to the nuclear hormone receptor superfamily. PPARγ is a key factor and master regulator in adipocytes development and functions (Tontonoz and Spiegelman, 2008). Indeed, loss-of-function studies both *in vitro* and *in vivo* have clearly demonstrated that PPARγ is essential for the formation of adipose tissue (Barak et al., 1999; E. D. Rosen et al., 1999; F. Wang et al., 2013) and mature adipocyte homeostasis (Imai et al., 2004). Recently, we demonstrated that mice carrying a full body deletion of PPARγ (*Pparg*$^{\Delta/\Delta}$) are totally devoid of adipocytes (Sardella et al., 2018). The activity of PPARγ is also central in bone homeostasis by modulating both bone formation by osteoblasts and bone resorption due to osteoclast activity. First, adipocytes and osteoblasts share same mesenchymal progenitors, and the cell fate decision between adipocyte and osteoblast lineages depends on various signals (Calo et al., 2010; Idris et al., 2009; Kawai and C. J. Rosen, 2010; Wan, 2010). Activation of PPARγ inhibits osteoblast formation, and thus bone formation, by favouring adipocyte differentiation (Ali et al., 2005; Bianco et al., 2001; Rzonca et al., 2004). Along this line, *in vivo* evidence indicates that mice heterozygous for a null allele of the gene encoding PPARγ (*Pparg*) exhibit increased osteoblastogenesis, which results in increased bone mass with a doubling of the rate of new bone formation, when compared to control mice. Moreover, PPARγ also directly inhibits osteoblast activity, independently of adipogenesis, as demonstrated by the consequence of

osteoblast-specific *Pparg* deletion in osteoblast differentiation and trabecular bone formation (Sun et al., 2013). Second, in bone resorption, PPARγ promotes osteoclastogenesis, as illustrated by increased osteoclast numbers in aged mice treated with thiazolidinediones (Lazarenko et al., 2007). Conversely, mice in which *Pparg* is specifically deleted in endothelial and hematopoietic cells, (i.e., affecting osteoclast progenitors but not osteoblasts), exhibit decreased osteoclastogenesis compared to wild-type mice, provoking osteopetrosis with impaired bone resorption and reduced medullary cavities (Wan et al., 2007). However, a recent controversy on the role of PPARγ in osteoclast was raised by Zou et al. who found that only the pharmacological, and not the physiological activation of PPARγ affects osteoclastogenesis (Evans and Wan, 2016; Zou et al., 2016). Finally, hyperactivation of PPARγ via agonist treatment causes accumulation of adipocytes in the bone marrow in thiazolidinedione-treated rodents, whereas increased bone loss and bone fractures have been correlated with thiazolidinedione therapy in humans (Grey, 2008; Kahn et al., 2008; Lazarenko et al., 2007).

Herein, we used mice carrying a PPARγ full body deletion to explore the consequences of the lack of adipocyte on bone homeostasis. This model, together with two other complementary models of genetically-induced lipoatrophic mice, i.e. mice carrying an adipose-tissue specific deletion of PPARγ and AZIP mice, allowed to explore the cross-talk between adipocytes and osteoclasts, whose alteration leads to hyperosteoclastogenesis and osteoporosis.

Results

In order to evaluate the impact of adipose tissue on bone homeostasis, we studied lipoatrophic mice that we recently generated. These mice, hereby called *Pparg^{Δ/Δ}* carry a whole body deletion of PPARγ and were obtained through epiblastic (Sox2 promoter) *cre* deletion of the Pparγ allele (Sardella et al., 2018). Bone analyses first show that these mice displayed dramatic alterations of both trabecular and cortical bones. Three-dimensional images generated by trabecular bone microcomputed tomography (3D micro-CT) reconstruction of the femurs of one-year old male and female mice highlighted a remarkably high trabecularization of the bone architecture in *Pparg^{Δ/Δ}* mice, which extended along the shaft and almost reached the mid-diaphysis (VIDEO 1, Fig. 1a). In contrast, the cortical bone was found to be extremely porous at the epiphysis and the diaphysis. Owing to hypertrabecularization and high cortical porosity and as a result, the cortical bone is poorly delimited in these mice.

Video of 3D micro-CT reconstruction of genetically induced lipoatrophic mice showing alterations of both trabecular and cortical femoral bone.

Figure 1: Epiblastic PPARγ-deletion induces trabecular and cortical bone alteration

Figure 1: Epiblastic PPARγ-deletion induces trabecular and cortical bone alteration
a. Micro-CT analysis of femur from *Pparg*$^{Δ/Δ}$ (γ$^{Δ/Δ}$) mice or control littermates (CTL). Histograms on the right show for the different genotypes (black bars for CTL and grey bars for γ$^{Δ/Δ}$) the percentage bone volume/trabecular volume (BV/TV) in % and connectivity density (ConnDens) of distal femoral trabecular bone (upper panels), and the cortical bone volume (BV) in mm^3 and the percentage of cortical porosity of femur (lower panels); n = 5 WT and 5 γ$^{Δ/Δ}$. All mice were 1-year-old. **b**. RT-qPCR analysis of Alkaline phosphatase (*AlpI*) gene expression in femur from CTL and γ$^{Δ/Δ}$; n = 6 WT and 7 γ$^{Δ/Δ}$. **c.** ELISA measurements of serum Alkaline phosphatase (ALP) and osteocalcin (OCN); n = 8 CTL and 8 γ$^{Δ/Δ}$. **d.** RT-qPCR analysis of cathepsin K (*Ctsk*) expression in femur from CTL and γ$^{Δ/Δ}$; n = 7 CTL and 7 γ$^{Δ/Δ}$. **e.** ELISA measurements of serum Carboxy-terminal Collagen I telopeptide (CTX); n= 8 CTL and 8 γ$^{Δ/Δ}$. **f.** Representative images of TRAcP staining of decalcified sections of femurs, at mid-shaft levels, from control (CTL) or *Pparg*$^{Δ/Δ}$ (γ$^{Δ/Δ}$) mice. ⊙ = cortical bone; ◆ = bone marrow. Data are presented as mean ± S.E.M. Statistical significance was determined by two-tailed unpaired *t*-test.

Molecular analyses confirmed an increased trabecular bone formation, as exemplified by increased levels of alkaline phosphatase mRNA (*Alp1*) in the bone, evaluated by RT-qPCR and increased levels of alkaline phosphatase (ALP) and osteocalcin (OCN) in the serum measured by ELISA (Fig.1b and c). Three-D micro-CT of the fourth lumbar vertebrae clearly demonstrated increased trabecular bone volume fraction and trabecular connectivity density, indicating increased osteoblastogenesis in *Pparg*$^{\Delta/\Delta}$ and in agreement with previous report (Akune et al., 2004) (see also Supplemental Data Fig. 1). These results are consistent with the pivotal role of PPARγ on the fate of bone marrow mesenchymal stromal cells, with previous reports showing increased osteoblastogenesis when PPARγ signalling is pharmacologically or genetically blocked (Takada et al., 2009).

Remarkably, increased expression of bone resorbing markers such as cathepsin K (*Ctsk*) (Fig. 1d) and carboxy-terminal collagen I telopeptide (CTX) (Fig. 1e), measured by RT-qPCR in bone extracts and ELISA in serum, respectively, suggest a parallel increase in osteoclastogenesis. Indeed, TRAcP staining on femoral sections (Fig. 1f) confirmed the increased presence of bone-resorbing osteoclasts in cortical bone, explaining the observed cortical porosity.

PPARγ is known to play a role in activating osteoclastogenesis. More specifically, Tie2^{Cre}-directed deletion of PPARγ severely impaired osteoclastogenesis, whereas PPARγ synthetic agonists exacerbated it (Wan et al., 2007). This role of PPARγ in osteoclast has been recently questioned by the study of Zou *et al.*, showing that the *in vivo* effect of PPARγ on osteoclast formation is only seen upon pharmacological activation of PPARγ but neither on physiological nor on pathological osteoclast formation (Zou et al., 2016). This raised the question whether the elevated osteoclast activity observed in cortical bone is attributable to lipoatrophy, or to unforeseen intrinsic effects of PPARγ deficiency in the osteoclast lineage. To disentangle these two possibilities, we used another PPARγ-independent lipodystrophic mouse model. AZIP$^{tg/+}$ mouse is a hemizygote transgenic mouse strain in which a dominant negative protein of key B-ZIP transcription factors (A-ZIP) blocks adipocyte development. AZIP$^{tg/+}$ mice are born with no white adipose tissue and markedly decreased brown adipose tissue (Moitra et al., 1998). In support of the lipoatrophy hypothesis, micro-CT analyses of femur of AZIP$^{tg/+}$ mice revealed an increased cortical porosity (Fig 2a). Finally, we generated lipoatrophic mice with wild-type hematopoietic lineage by invalidating PPARγ in adipose tissue (Adipoq^{cre} PPARγ fl/fl, γ^{FatKO} (F. Wang et al., 2013), called herein PPARγ^{adipoKO}). TRAcP staining of cortical bone and micro-

CT analysis in these mice revealed increased osteoclast numbers and activity with an enhanced cortical porosity, as observed in $Pparg^{\Delta/\Delta}$ mice (Fig. 2a-c and Supplemental Data Fig. 2a-c). These two independent mouse models confirmed that adipocyte deficiency is pivotal to the cortical bone phenotype observed here.

To investigate how adipocytes regulate osteoclastogenesis, and to evaluate whether factors produced by adipocytes within the bone marrow can influence osteoclastogenesis, we performed an osteoclastogenesis assay using bone marrow cells from wild-type mice cultured in the presence of either control medium or adipocyte-conditioned medium (ACM in figure 2). Adipocyte-conditioned medium was isolated from culture of adipocyte-differentiated 3T3-L1 cells whereas control medium (CTL in figure 2) was obtained from cultures of undifferentiated pre-adipocytes 3T3-L1 cells. As seen in Figure 2d, the presence in bone marrow cells culture of adipocyte-conditioned medium, compared to pre-adipocyte-conditioned medium, provoked a substantial decrease in the number of mature osteoclasts. Consistently, adipocyte-conditioned medium significantly inhibited bone resorption as assessed by the decrease number of resorption pits in the calvaria resorption assay (Fig. 2ef). These data demonstrate that adipocyte-secreted mediators are responsible for the inhibition of RANKL-induced osteoclast differentiation *in vitro*.

Next, we searched for component(s) of ACM responsible for this anti-osteoclastogenic activity. Interestingly, the ACM inhibitory activity was maintained after filtering out all components larger than 35 kDa from the conditioned medium. The molecular weights of the major adipokines, adiponectin and leptin, are below 35 kDa and their circulating levels are not or barely detectable in $Pparg^{\Delta/\Delta}$ mice ((Gilardi et al., 2019) and Supplemental Data Fig. 2d). Unlike *Leptin*, *Adiponectin* is preferentially expressed by adipocyte and by bone-marrow MSC, and is a well-characterized target gene of PPARγ (Iwaki et al., 2003). Western-blot analyses confirmed that adiponectin was present in large amounts in ACM but not in CTL medium (Supplemental Data Fig. 2d). We therefore tested whether blocking adiponectin would affect ACM-mediated inhibition of osteoclastogenesis *in vitro*. As shown in Figure 2f, the addition of a blocking adiponectin antibody reversed the reduction in osteoclast number in a dose-dependent manner. These results support that the soluble factors secreted by adipocytes, in particular adiponectin, are inhibitors of osteoclastogenesis. Consistently, the total lack of adipocytes, including in the bone marrow, results in a complete absence of this adipokine, leading to the loss of its repressive activity in osteoclast differentiation.

Figure 2: Lipoatrophy is responsible for unregulated osteoclastogenesis

Figure 2: Lipoatrophy is responsible for unregulated osteoclastogenesis
a. . Representative 3D sections of midshaft femur from A-ZIP[tg/+] (n=6) and CTL (n=6) littermates. **b.** Femoral cortical porosity measured by micro-CT analysis of femur from A-ZIP[tg/+] (n=6) and control (CTL; n=6) littermates. **c.** Representative images of TRAcP staining of longitudinal sections of decalcified femurs, at mid-shaft levels, from control (CTL) or A-ZIP[tg/+] mice. ⊙ = cortical bone; ◆ = bone marrow. **d.** Experimental design of

the *in vitro* assay used to test anti-osteoclastogenic activity from adipocyte secretome. **e.** Calvaria resorption pit assay. Representative sections of skulls from 14-day-old wild-type pups cultured for 2 weeks in complete αMEM in the presence of conditioned medium from either undifferentiated 3T3-L1 preadipocytes (CTL, n=3) or mature 3T3-L1 adipocytes (ACM, n=3). Data are presented as mean ± s.e.m. **f.** One day after plating, wild type BM cells were cultured for 6 more days in the presence of M-CSF and RANKL (osteoclast differentiation media) in addition to conditioned medium prepared from differentiated (ACM: adipocyte-conditioned medium) or undifferentiated (CTL: control-conditioned medium) 3T3-L1 preadipocytes. Different doses of a specific blocking antibody against adiponectin (1.5–12.0 µg/ml) were added to the conditioned medium (ACM50) during osteoclast differentiation (ACM 100 means pure ACM; ACM50 means 1:1 diluted ACM with osteoclast differentiation medium).

AdipoRon is a non-peptidic agonist of adiponectin receptors (Okada-Iwabu et al., 2013). We thus perform the reverse experiment of the antibody blocking assay, adding AdipoRon to culture of bone marrow cells. Consistently, AdipoRon recapitulated the blocking effect of ACM on osteoclast differentiation irrespective of the origin of the osteoclast progenitors (bone marrow, spleen, or macrophage-enriched fraction) with IC50 in the micromolar range (Fig. 3a, b Supplementary Fig. 3). Furthermore, AdipoRon blocked osteoclast differentiation in *Pparg* deficient cells, confirming that intrinsic alterations do not play a role in the observed phenotype (Fig. 3c).

Previous studies demonstrated *in vivo* that mice treated with adiponectin expressing adenovirus have reduced osteoclast numbers and bone-resorption markers (Oshima et al., 2005). Using a macrophage cell line prone to differentiate in osteoclasts, the authors further showed that adiponectin blocked *in vitro* osteoclastogenesis of RAW264 by inhibiting NFATc1. Blockade of NFATc1 was suggested to occur through activation of the AMPK signalling pathway, a key integrator of metabolic signals and target of adiponectin signalling (Yamaguchi et al., 2008).

We thus tested whether this mechanism would apply *in vivo* and to primary cells. Flow cytometry analysis of osteoclasts generated from murine bone marrow cells confirmed that the adiponectin receptor AdipoR1 is expressed also by mature osteoclasts generated from murine bone marrow cells (Fig. 3d). In agreement, western-blot analysis of AMPK phosphorylation confirmed that the Adiponectin signalling pathway is active in both precursors and mature osteoclasts (Fig. 3e-g).

Figure 3: Adiponectin signalling activation blocks OCL differentiation

Figure 3: Adiponectin signalling activation blocks osteoclast differentiation
a. Representative images of TRAcP stained BM-derived osteoclasts differentiated in the presence or absence of AdipoRon (5μM). Scale bars=100μm. **b.** Dose response effect of AdipoRon on osteoclast differentiation using RAW264.7 cells (n=3). **c**. Effect of AdipoRon on osteoclast differentiation assay using bone marrow cells from *Pparg*$^{Δ/Δ}$ (γ$^{Δ/Δ}$) mice and control littermates (CTL). **d**. Expression of Type 1 Adiponectin Receptor (AdipoR1) at the surface of bone marrow derived osteoclasts demonstrated by flow cytometry. Grey curve represents unstained negative control; blue curve represents BM-derived osteoclasts. **e.** Western-blot analysis of bone marrow-derived osteoclasts treated with AdipoRon (5μM). AdipoRon treatment activates AMPK phosphorylation (p-AMPK) in murine BM-derived osteoclasts (n=2 CTL; n=2 AdipoRon stimulated) **f**. Ratio of P-AMPK over total AMPK measured by densitometry **g**. Kinetic of AMPK phosphorylation (P-AMPK) after AdipoRon challenge in RAW264.7-derived osteoclasts. Data are presented as mean ± S.E.M. Statistical significance was determined by two-tailed unpaired *t*-test.

AdipoRon was thus used to investigate the consequences of adiponectin receptor activation on mature osteoclasts. AICAR was used as a synthetic activator of AMPK. As seen in Fig 4a,b each of these molecules inhibited the formation of podosomes that are essential for bone resorption. Moreover, dorsomorphin, a reversible AMP-kinase inhibitor, could reverse the AdipoRon inhibitory effect (Fig. 4b). This is in line with the recently published effect of recombinant adiponectin on RAW264.7 cells, confirming the impact of adiponectin on both osteoclasts differentiation and podosome formation (Chen et al., 2018) and endorsing the use of AdipoRon as a surrogate activator for Adiponectin receptors. Finally, adiponectin signalling regulates numerous cellular metabolic pathways including mitochondrial functions (Iwabu et al., 2010). AdipoRon treatment provoked disruption of the perinuclear mitochondrial network when added to culture of mature osteoclasts (Fig. 4c). Altogether, these results indicate that, beside its inhibiting action in osteoclast differentiation, adiponectin signalling exerts a pivotal role in mature osteoclast energy homeostasis regulation.

The inhibitory effect of AdipoRon was further tested *in vivo* on the well-described model of osteoclast bone resorption induced by lipopolysaccharide (LPS) injection into the mouse calvaria (Fig. 4d). Both micro-CT (Fig. 4d) and TRAcP staining (Fig. 4e) showed that AdipoRon administration resulted in a strong decrease in osteoclast numbers and bone resorption pits on calvaria.

Figure 4: Adiponectin signalling regulates OCL podosome formation and activity

Figure 4: Adiponectin regulates OCL activity and podosome formation
a. Confocal microscopy of phalloidin-marked podosomes in RAW264.7-OCL in the presence of increasing concentrations of AdipoRon (representative images of n=3 per concentration) Scale bars=100 μm. **b.** Confocal microscopy of phalloidin-marked podosomes in the presence of AMPK modulating drugs (AdipoRon, 10μM; AICAR, 50nM; Dorsomorphin, 1.2μM)

(representative image of n=3 per conditions white asterisks mark podosomes). Scale bars=50 μm. Blue arrows indicate podosome rings. **c**. Confocal microscopy of MitoTracker Orange in mature RAW264.7-OCL after AdipoRon (10μM) challenge at different time points. Scale bars=20 μm. **d**. Micro-CT of mouse skulls collected 3 days after mice received daily *s.c.* injections on the calvaria. Mice were injected with either 100 μg lipopolysaccharide from *Salmonella abortus equi* (LPS), or 100 μg LPS + 500 ng AdipoRon (LPS + AdipoRon) or vehicle (PBS) injection (control, CTL). (n=3 animals per group) **e**.TRAcP staining of mouse skull after LPS +/- AdipoRon challenge, (representative sections, n=3 animals per group). Scale bars=100 μm.

We thus aimed to validate our findings on human cells and therefore investigated the impact of AdipoRon on hPBMC-derived osteoclasts. Consistent with the results obtained with mouse cells, low dose of AdipoRon significantly inhibited RANKL-induced human osteoclast differentiation (Fig.S4 a and b). AdipoRon also activated AMPK phosphorylation in human PBMC-derived osteoclasts whithin minutes (SFig.S4 c). At later time-point, i.e. within hours after the challenge with AdipoRon, the number of actin-containing podosome was drastically reduced in human osteoclasts (SFig.4 d). These original findings offer new therapeutic perspectives in inflammatory bone loss management by targeting adiponectin/AdipoR.

Discussion

In the present report, using lipodystrophic mouse models, we evaluate how the total lack of adipose tissue may affect bone homeostasis. Taken together our results demonstrate that mature osteoclasts are sensitive to extrinsic adipose-derived metabolic signals such as adipokine and that bone marrow adiposity must be considered as a physiological regulator of bone homeostasis. We show that AdipoRon, a pharmacological agonist of adiponectin receptors, inhibits osteoclast differentiation of precursors from different origins (bone-marrow, spleen and monocytic fraction). Besides its effect on osteoclast differentiation, AdipoRon blocks mature osteoclast activity via AMPK signaling. Activation of AMPK induced mitochondrial network alteration and podosome formation blockade, whereas AMPK blockade reverted these effects. We also demonstrate that adiponectin signaling inhibits human osteoclast differentiation as well as activity of human mature osteoclast. These original findings reveal a conserved adiponectin-mediated metabolic control of osteoclast activity and are clinically promising, in particular with regard to the treatment and prevention of excessive pathological bone-resorption conditions, such as in osteoporotic patients.

The first model used in the present study, the PPARγ null mice, is bringing up some decisive arguments with respect to the controversy on the role of PPARγ in osteoclastogenesis. Whereas initial studies reported that PPARγ is a key factor for osteoclastogenesis (Wan et al., 2007) *Zou et al.* could not confirm such activity and proposed that only pharmacological activation of PPARγ had impact on osteoclastogenesis (Zou et al., 2016)(Evans and Wan, 2016). Herein, we definitely demonstrate that osteoclastogenesis can occur in the absence of PPARγ signal, and that other adipose tissue-derived signals, such as adipokines, affect osteoclast differentiation and activity. In contrast, the increase trabecular bone formation seen is vertebrae, in femoral epiphyses and distal diaphysis of *Pparg$^{Δ/Δ}$* mouse is consistent with the known role of PPARγ in inhibiting osteogenesis.

Using three complementary genetic models (2 distinct conditional null mice of PPARγ and the AZIP$^{tg/+}$ mice, a PPARγ-independent lipoatrophy model), we show that the dramatic cortical bone alteration due to an excessive osteoclastogenesis in mice is linked to the lack of adipocyte and adipocyte-derived signals. The systemic and local activities of adipokines such as leptin and adiponectin have been recently reviewed (Cornish et al., 2018). Although leptin can act as a positive regulator of osteogenesis in *in vitro* systems, other studies in mice show that systemic leptin acting via the central nervous system is a potent inhibitor of bone formation (Ducy et al., 2000). Systemic adiponectin was reported to promote bone formation through its depressive

action on the sympathetic tone in mice in age-dependent manner (Kajimura et al., 2013). However, *in vitro* differentiation assay showed that adiponectin inhibits osteoclast formation (Oshima et al., 2005; Williams et al., 2009; Yamaguchi et al., 2008). The observations *in vivo* are more controversial. Adiponectin-null mice exhibit an increased trabecular volume and fewer osteoclasts (Q.-P. Wang et al., 2014), although Shinoda *et al.* (Shinoda et al., 2006) found normal bone development in these mice. More recently, adiponectin-null female mice were shown to exhibit a decreased cortical thickness (Naot et al., 2016). Thus, adiponectin may have different effects on bone remodelling, depending on the activated autocrine, paracrine or systemic pathways. Our results from the osteoclastogenesis assay using bone marrow-derived cells and adiponectin-specific blocking antibodies support the important paracrine role of adiponectin in the inhibition of osteoclastogenesis in the normal BM environment. Direct paracrine activity is also consistent with the presence of the adiponectin receptors, AdipoR1 and AdipoR2, on the surface of differentiating osteoclasts (Pacheco-Pantoja et al., 2013). Nonetheless, we cannot exclude that other adipokines may also contribute to the paracrine effects of adipocytes on osteoclasts.

Identifying the source of adiponectin in the vicinity of osteoclast precursors and/or mature osteoclasts in bone marrow and bone compartment is challenging. Indeed, levels of adiponectin production by BM adipocytes remain controversial. Cawthorn et al. (Cawthorn et al., 2014)reported that BM adipocytes serve as a major source of circulating adiponectin exerting systemic effects during caloric restriction whereas others groups reported that bone marrow adipocytes express lower levels of adiponectin than adipocytes in adipose tissue (Li et al., 2019; Liu et al., 2011; Poloni et al., 2013). Others sources of bone marrow adiponectin have been recently described. Bone marrow PDGFRβ[+]VCAM-1[+] stromal cells have been identified as major producers of adiponectin in the bone marrow (Mukohira et al., 2019). Single cell transcriptomic profiling of the mouse bone marrow stromal compartment indicated that LepR[+] bone marrow mesenchymal stromal cells, which form a functional continuum with osteolineage cells, are expressing adiponectin (Baryawno et al., 2019). Beside their proliferative action on HSC and antiproliferative effects on myelomonocytic and early B lymphoid cells, these adiponectin-producing cells appears to be essential to the hematopoietic niche. In this context, our finding that adiponectin regulates osteoclast activity and differentiation in bone further strengthens the argument that a metabolic pathway driven by adiponectin controls many osteoimmunology processes, including remodeling of bone marrow hematopoietic stem cell niche, with potential implications for multiple human disease states including cancer.

Finally, the inhibitory effects of adiponectin signaling on osteoclast activity and differentiation appear to be promising from a therapeutic outlook, in particular with regard to the treatment and prevention of excessive pathological bone-resorption conditions, such as in osteoporotic patients. The present study implies that preserving functional adiponectin sensitivity in patients suffering from metabolic diseases (e.g. diabetes, obesity, metabolic syndrome) might be also beneficial for preventing bone frailty.

Supplemental data

Figure S1

Supplemental Data Figure 1: Osteoblastic activity is increased in mice with epiblastic deletion of PPARγ. a. Representative 3-D micro-CT of the fourth lumbar vertebrae of CTL and *Pparg*$^{Δ/Δ}$ (γ$^{Δ/Δ}$) mice. n = 5 WT and 5 γ$^{Δ/Δ}$. **b.** Trabecular bone volume fraction measured by micro-CT analysis of vertebra from γ$^{Δ/Δ}$ (n=5) and control (CTL; n=5) littermates. **c.** Trabecular connectivity density measured by micro-CT analysis of vertebra from γ$^{Δ/Δ}$ (n=5) and control (CTL; n=5) littermates. **d.** Primary osteoblast differentiation and proliferation in culture of calvaria pre-osteoblast from of newborn CTL and γ$^{Δ/Δ}$ pups. Alizarin red S staining of calcium deposition was conducted in osteoblasts cultured 3 weeks after changing to osteogenic medium (containing 50 µg/ml ascorbic acid and 100 nM β-glycerolphosphate). **e.** Primary adipocyte differentiation from culture of bone marrow mesenchymal stem cells obtained from newborn CTL and γ$^{Δ/Δ}$ pups, 3 weeks after changing the culture medium to adipogenic medium (containing dexamethasone (2.5 µM), 3-isobutyl-1-methylxanthine (0.5 mM) and insulin (10 µg/mL)). Blue arrows indicate lipid droplets.

Figure S2

Supplemental Data Figure 2: Increased cortical porosity in two lipodystrophic mouse models: A-ZIP^{tg/+} mice and mice with fat-specific deletion of PPARγ. a. Representative 3D sections of midshaft femur from γ^{adipoKO} (n=3) and CTL (n=3) littermates. **b.** Femoral cortical porosity measured by micro-CT analysis of femur from γ^{adipoKO} (n=3) and control (CTL; n=3) littermates. **c.** Representative images of TRAcP staining of longitudinal sections of decalcified femurs, at mid-shaft levels, from control (CTL) or γ^{adipoKO} mice. ⊙ = cortical bone; ◆ = bone marrow. **d.** RT-qPCR analysis of *Leptin* and *AdipoQ* mRNA levels in long bone from A-ZIP^{tg/+}(n=8) and their CTL littermates (n=8) and from γ^{Δ/Δ} (n=7) and their CTL littermates (n=6). Data are presented as mean ± S.E.M. Statistical significance was determined by two-tailed unpaired *t*-test. **e**. Western-blot analysis of conditioned medium from undifferentiated (CTL) and differentiated 3T3-L1 adipocyte (A-CM) against adiponectin. Recombinant adiponectin (200ng rmAdiponectin) was added to demonstrate specificity of the signal.

Figure S3

Supplemental Data Figure 3: AdipoRon inhibits osteoclast (OCL) differentiation of multiple origins.
Left panels: representative images of TRAcP stained (**a.**) BM-derived, (**b.**) spleen derived and (**c.**) macrophages enriched fraction osteoclasts differentiated in the presence or absence of AdipoRon (5µM). *Right panels*: the mean IC50 of AdipoRon inhibition of OCL differentiation was determined from the curve with the error of the fit (S.E.M.) Data are mean ± S.E.M. (n = 3 biological replicates).

Figure S4: AdipoRon inhibits human OCL differentiation and activity

Figure S4: AdipoRon inhibits human osteoclast (OCL) differentiation and activity through AMPK activation. a. Dose response effect of AdipoRon on osteoclast differentiation from hPBMC (n=5). **b.** Representative images of TRAcP staining of hPBMC derived osteoclast differentiation in the presence or absence of AdipoRon (5μM). **c.** Western-blot analysis of hPBMC-derived osteoclasts treated with AdipoRon (5μM). AdipoRon treatment (5', 15', 30') activates AMPK phosphorylation (p-AMPK) in hPBMC. **d.** Fluorescence microscopy of phalloidin-marked podosomes in hPBMC derived OCL in the presence of AdipoRon (5μM, 12h) (representative images of n=5).

Material & Methods

Key ressources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Neutralizing adiponectin receptor 1 polyclonal antibody	Enzo Life Science	ALX-210-645-C200
Adiponectin rabbit monoclonal antibody	Cell Signaling Technology	#2789
AMPKα Rabbit mAb (D63G4)	Cell Signaling Technology	#5832
Phospho-AMPKα (Thr172) (40H9) Rabbit mAb	Cell Signaling Technology	#2535
Chemicals, Peptides, and Recombinant Proteins		
Fetal Bovine Serum	GIBCO	10082139
Alpha MEM with NTP, dNTP	GIBCO	22571
AdipoRon	SIGMA	SML0998
mRANKL	R&D systems	462-TEC
hRANKL	R&D systems	6449-TEC
Critical Commercial Assays		
Acid Phosphatase, Leukocyte (TRAP) assay	SIGMA	387A
murine serum CTX ELISA	LSBio	F21349
murine serum OCN ELISA	LSBio	F12227
murine serum PTH ELISA	LSBio	F23085
Experimental Models: Cell Lines		
3T3-L1 pre adipose murine cell line	ATCC	CL-173
RAW 264.7 murine macrophage	ATCC	TIB-71
Experimental Models: Organisms/Strains		
Sox2-Cre mouse	The Jackson Laboratory	004783
Adipoq-Cre mouse	The Jackson Laboratory	010803
PPARg L2/L2 mouse	B Desvergne	PMID:15070754
AZIP/F1 tg mouse	C Vinson	PMID:9784492
Oligonucleotides		
Gapdh		Sup Table1
ActB		Sup Table1
AdipoQ		Sup Table1
Leptin		Sup Table1
Software and Algorithms		
CTan CTVol	Skyscan	
ImageJ	Fiji	

Mouse models

Genotype denomination follows the rules recommended by the Mouse Genome Database (MGD) Nomenclature Committee. Animal procedures were authorized by the Canton of Vaud veterinary service. Sox2CRE transgenic mice (SOX2CRE$^{tg/+}$; Tg(Sox2-cre)1Amc/J) and AdipoQCRE (*Adipoq-Cre*$^{tg/+}$) from The Jackson Laboratory, were maintained in the University of Lausanne Animal Facility. The series of matings allowing the generation of the

SOX2CRE$^{tg/+}$*Pparg$^{em\Delta/\Delta}$* mouse (hereafter called *Pparg$^{\Delta/\Delta}$*), which expressed the SOX2CRE transgene and have no *Pparg* functional alleles, as well as their control littermates (CTL) that have no SOX2CRE transgene and two functional *Pparg* alleles (*Pparg$^{fl/+}$*) have been described in (Gilardi et al., 2019). PPARγ^{adipoKO} and their control littermates were generated as described in (F. Wang et al., 2013). AZIP/F1 mice on FVB background (Tg(AZIP/F)1Vsn in MGD, herein called AZIP$^{tg/+}$) and wild type FVB controls were a kind gift from Dr. Charles Vinson and were generated as previously described(Moitra et al., 1998).

Primary cell culture

Bone marrow cells were isolated by crushing entire long bones into DMEM/3% FCS (GIBCO). Bone fragments were removed by filtration through a 70 μm filter mesh. Splenocyte suspensions were obtained by mashing the spleen through sieves into DMEM/3% FCS, washing by centrifugation then filtering through 70 μm filter caps. Primary osteoblasts from neonatal calvaria of *Pparg$^{fl/+}$* and *Pparg$^{\Delta/\Delta}$* were harvested by sequential collagenase type II (3 mg/mL) digestions of individual calvarium from 2 to 3-day-old mice. After genotyping, calvaria cells of the same genotype were pooled and plated accordingly. The cells were incubated at 37° C with 5% CO2 and media was changed every two days until they reached 80% confluency. To assess proliferation, primary osteoblasts were plated in 24-well plates at a concentration of 10,000 cells/well in EMEM media (GIBCO) containing 10% FBS, 100 U/ml penicillin and 100 ug/ml streptomycin (GIBCO).

Osteoclastogenesis Assays and TRAcP staining of murine and human primary cells

Erythrocytes were eliminated from murine splenocyte and BM preparation or human PBMC using ACK lysis buffer. Cells were cultured in complete αMEM with 40 ng/ml M-CSF (ProSpec-Tany TechnoGene, Rehovat, Israel) and l0 ng/ml RANKL (R&D Systems, MN). Tartrate-resistant acid phosphatase (TRAcP) staining was performed after 4-6 days, using the Acid Phosphatase, Leukocyte (TRAP) Kit according to the manufacturer's instructions (Sigma). Mature osteoclasts were identified as multi-nucleate TRAcP^{+} cells using light microscopy. 3T3-L1 cells (ATCC) were grown to sub-confluence in DMEM/10% FBS. Cells were then cultured in adipogenic conditions (DMEM/10%FBS, 0.5 mM 3-isobutyl-1-methylxanthine, 1 μM dexamethasone (Sigma), and 10 μg/ml bovine insulin (Sigma)) for 2 weeks, after which the medium was changed back to DMEM/10% FBS. Adipocyte Conditioned Medium (A-CM) was collected after 24 hours exposure to differentiated cells. Similarly, conditioned control medium (CTL) was obtained from undifferentiated 3T3-L1 cells. A-CM

and CTL were concentrated 5X using Centriprep^{TM} (Millipore). Conditioned medium from either differentiated or undifferentiated 3T3-L1 cells was added during osteoclastogenesis from multipotent progenitor cells. For neutralisation experiments, 50% CM was incubated with different amounts of anti-adiponectin antibody (Enzo Life Sciences) for 30 min at room temperature then added during osteoclastogenesis. TRAcP staining to evaluate osteoclast numbers was performed as mentioned above.

Osteoclast differentiation and TRAcP staining of RAW 264.7 cells

For osteoclastogenesis, a total of $1x10^4$ RAW 264.7 (ATCC) cells were seeded per well on 24-well plates in 500 µl αMEM containing 5% FCS (Hyclone, GE Healthcare), 1% penicillin-streptomycin (Gibco) as well as 50 µM β-mercaptoethanol (Gibco) and 30 ng/ml murine RANKL (R&D Systems). To study the impact of adiporon on osteoclast differentiation, 10 µM AdipoRon (Sigma-Aldrich) were added to the culture medium. Tartrate-resistant acid phosphatase (TRAcP) activity was examined using the leucocyte acid phosphatase kit (Sigma-Aldrich) according to manufacturer's specifications. TRAcP$^+$ cells with $\geq$3 nuclei were considered as osteoclasts. In addition, $5x10^3$ RAW 264.7 cells were seeded per well in a Nunc Lab-Tek II 8-well Chamber Slide system (ThermoFisher Scientific) in a total of 300 µl osteoclast differentiation medium containing 30 ng/ml RANKL. Fully differentiated osteoclasts were treated with 10 µM AdipoRon for the indicated time points. FACS analysis as well as sorting of mature OCLs for subsequent Western-blot analyses was performed as described previously (Madel et al., 2018).

Measurement of bone morphology and microarchitecture

Bone mineral density (BMD mg/cm^2) and bone mineral content (BMC mg) of femurs were measured *ex vivo* using dual energy X-ray absorptiometry (pDXA; PIXImus, GE- Lunar Corp.). High-resolution micro-computed tomography (UCT40, ScancoMedical AG, Bassdorf, Switzerland) was used to scan the femur and the 5th lumber vertebral body. Cortical and trabecular bones were evaluated using isotropic 12µm voxels. For trabecular bones, we assessed bone volume fraction (BV/TV, %), trabecular thickness (Tb.Th, µm), trabecular number (Tb.N, mm^{-1}), and trabecular separation (Tb.Sp, µm). For cortical midshaft femur, we measured the average total volume (TV, mm^3), bone volume (BV, mm^3) and average cortical thickness (µm). SkyScan software (CTAn and CTVol) was employed to calculate cortical bone porosity of femur using a dedicate method after acquisition using a Skyscan 1272 (SkyScan 1272, Bruker, Brussels, Belgium) at a resolution of 10 µm/pix and visualized in 3D.

Serum protein assays

Blood was obtained by cardiac puncture immediately after euthanasia. After clotting and centrifugation, serum was collected and stored at -80°C until use. Serum CTX (carboxy-terminal collagen crosslinks), osteocalcin (OCN), alkaline phosphatase (ALP) and parathyroid hormone (PTH) were measured by ELISA assays according to the manufacturer's protocol (LifeSpan Biosciences, Inc., Seattle, WA).

Calvaria injection

Nine 6-week old C57/BL6 male mice were purchased from Envigo and housed in the local animal facility under a 12h light/12h dark cycle. The animals were randomly divided into three experimental groups, each group comprising three mice. Mice received daily *s.c.* injections on the calvaria before analysis on the third day. Group 1 was injected with 100 µg lipopolysaccharide from *Salmonella abortus equi* (LPS, Sigma-Aldrich), group 2 with 100 µg LPS + 500 ng AdipoRon (Sigma-Aldrich) and experimental group 3 underwent vehicle (PBS) injection. Calvariae were collected and fixed overnight in 4% formaldehyde at 4°C and then stored in 70% EtOH. Subsequent micro-CT analysis were used to evaluate the resorbed area. Calvariae were decalcified in 10% EDTA, embedded in paraffin and 5-micron sections were cut and stained for tartrate-resistant acid phosphatase (TRAcP, Sigma-Aldrich).

Quantitative RT-PCR

Total RNA was isolated from total long bones, including the crude BM, or primary cell cultures, using TRIzol LS (Invitrogen) and purified with the RNeasy kit (Qiagen). RNA quality was verified by microfluidic (Agilent 2100 Bioanalyzer) and concentration determined with a NanoDrop spectrophotometer (Wilmington). Total RNA (500 ng/µg) was reverse transcribed using Script™ cDNA synthesis kit (Bio-Rad) according to the manufacturer's instructions. Real time qPCR was performed with SYBR® Green PCR mastermix using an ABI PRISM® 7900 PCR machine (Applied Biosystems). Results were normalized to glyceraldehyde-3-phosphate dehydrogenase (*Gapdh*) or beta-actin beta (*Actb*).

Western blot anlysis

Sorted mature OCLs were washed twice with ice-cold PBS and scraped in 1X Laemmli buffer. Cells were disrupted by sonication and centrifuged at 3000 rpm for 10 min. Protein samples were analyzed by SDS/PAGE (gradient 4-15% acrylamide, Criterion, BioRad) and electroblotted onto 0.2% PVDF membrane using Transblot Turbo (BioRad). After 1 h in

blocking buffer (Tris-buffered saline (TBS)–Tween 20 with 5% fat free dry milk), membranes were blotted overnight at 4°C with Rabbit mAb against AMPKα (D63G4) (dilution 1:1 000, Cell Signaling Technology) or Phospho-AMPKα (Thr172) (40H9) (dilution 1 :1000, Cell Signaling Technology), diluted in TBS–Tween with 5% BSA. After three washing steps with TBS–Tween 20, blot was incubated for 1 h at room temperature with anti-rabbit IgG conjugated with horseradish peroxidase diluted 1:2000 in blocking buffer. After four additional washing steps with TBS–Tween 20, protein bands were detected by chemiluminescence using Chemi Doc XRS+ (BioRad).

Imaging

Podosomes were visualized by phalloidin staining (Sigma-Aldrich) following manufacturer's instructions. After removing unbound phalloidin conjugates, cells were labelled with 1 µg/ml DAPI (Sigma-Aldrich). Microscopic imaging was performed using a fluorescence microscope (Zeiss Axio Observer D1). For mitochondrial analysis, live cells were labeled with 200 nM MitoTracker Orange (ThermoFisher) according to manufacturer's recommendations. After staining, cells were fixed in 4% formaldehyde, labelled with 1 µg/ml DAPI and subjected to z-stack confocal microscopy analyses (Zeiss LSM 710 on an inverted Axio Observer.Z1 stand). Images were processed using Fiji/Image J software(Schindelin et al., 2012).

Statistical analysis

The data were analyzed using one- or two-factor ANOVA for multiple comparisons as appropriate. Two-group comparisons were performed using the Student's *t*-test. ns: non-significant, * $p<0.05$, ** $p<0.01$, *** $p< 0.001$, **** $p<0.0001$.

Acknowledgements

This work was supported by the Etat de Vaud (BD and AW) the FNRS (BD), the Région Grand Est (J-YJ), the Fondation Arthritis (DM) and by the french PIA project Lorraine Université d'Excellence, reference ANR-15-IDEX-04-LUE (J-YJ and DM).

Authors contribution

HF designed the experimental plan, performed the experiments, analyzed the data, and contribute to the original draft preparation. MBM, DDP, MS, CW, AW, CP, and BT performed experiments, analyzed the data, and prepared the vizualisation. JYJ, FG, SF, NB and CBW contributed to the conceptualization, analyzed the data, and reviewed/edited the manuscript. BD and DM conceived the study, designed the experimental plan, supervised the study, analyzed the data, wrote the manuscript, and acquired financial support for the project. All authors read and edited the manuscript. BD and DM are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analyses.

References

Akune, T., Ohba, S., Kamekura, S., Yamaguchi, M., Chung, U.-I., Kubota, N., Terauchi, Y., Harada, Y., Azuma, Y., Nakamura, K., Kadowaki, T., Kawaguchi, H., 2004. PPARgamma insufficiency enhances osteogenesis through osteoblast formation from bone marrow progenitors. J. Clin. Invest. 113, 846–855. doi:10.1172/JCI19900

Ali, A.A., Weinstein, R.S., Stewart, S.A., Parfitt, A.M., Manolagas, S.C., Jilka, R.L., 2005. Rosiglitazone Causes Bone Loss in Mice by Suppressing Osteoblast Differentiation and Bone Formation. Endocrinology 146, 1226–1235. doi:10.1210/en.2004-0735

Barak, Y., Nelson, M.C., Ong, E.S., Jones, Y.Z., Ruiz-Lozano, P., Chien, K.R., Koder, A., Evans, R.M., 1999. PPAR gamma is required for placental, cardiac, and adipose tissue development. 4, 585–595.

Baryawno, N., Przybylski, D., Kowalczyk, M.S., Kfoury, Y., Severe, N., Gustafsson, K., Kokkaliaris, K.D., Mercier, F., Tabaka, M., Hofree, M., Dionne, D., Papazian, A., Lee, D., Ashenberg, O., Subramanian, A., Vaishnav, E.D., Rozenblatt-Rosen, O., Regev, A., Scadden, D.T., 2019. A Cellular Taxonomy of the Bone Marrow Stroma in Homeostasis and Leukemia. Cell 177, 1915–1932.e16. doi:10.1016/j.cell.2019.04.040

Bianco, P., Riminucci, M., Gronthos, S., Robey, P.G., 2001. Bone marrow stromal stem cells: nature, biology, and potential applications. Stem Cells 19, 180–192. doi:10.1634/stemcells.19-3-180

Calo, E., Quintero-Estades, J.A., Danielian, P.S., Nedelcu, S., Berman, S.D., Lees, J.A., 2010. Rb regulates fate choice and lineage commitment in vivo. Nature 466, 1110–1114. doi:10.1038/nature09264

Cawthorn, W.P., Scheller, E.L., Learman, B.S., Parlee, S.D., Simon, B.R., Mori, H., Ning, X., Bree, A.J., Schell, B., Broome, D.T., Soliman, S.S., DelProposto, J.L., Lumeng, C.N., Mitra, A., Pandit, S.V., Gallagher, K.A., Miller, J.D., Krishnan, V., Hui, S.K., Bredella, M.A., Fazeli, P.K., Klibanski, A., Horowitz, M.C., Rosen, C.J., MacDougald, O.A., 2014. Bone Marrow Adipose Tissue Is an Endocrine Organ that Contributes to Increased Circulating Adiponectin during Caloric Restriction. Cell Metab. 20, 368–375. doi:10.1016/j.cmet.2014.06.003

Chen, G., Huang, L., Wu, X., Liu, X., Xu, Q., Li, F., Dai, M., Zhang, B., 2018. Adiponectin inhibits osteoclastogenesis by suppressing NF-κB and p38 signaling pathways. Biochemical and Biophysical Research Communications 503, 2075–2082. doi:10.1016/j.bbrc.2018.07.162

Cornish, J., Wang, T., Lin, J.-M., 2018. Role of Marrow Adipocytes in Regulation of Energy Metabolism and Bone Homeostasis. Curr Osteoporos Rep 16, 116–122. doi:10.1007/s11914-018-0425-0

Ducy, P., Amling, M., Takeda, S., Priemel, M., Schilling, A.F., Beil, F.T., Shen, J., Vinson, C., Rueger, J.M., Karsenty, G., 2000. Leptin inhibits bone formation through a hypothalamic relay: a central control of bone mass. Cell 100, 197–207. doi:10.1016/s0092-8674(00)81558-5

Evans, R.M., Wan, Y., 2016. Reply to "PPAR-γ regulates pharmacological but not physiological or pathological osteoclast formation". Nat. Med. 22, 1205–1205. doi:10.1038/nm.4207

Felson, D.T., Zhang, Y., Hannan, M.T., Anderson, J.J., 1993. Effects of weight and body mass index on bone mineral density in men and women: the Framingham study. J. Bone Miner. Res. 8, 567–573. doi:10.1002/jbmr.5650080507

Gilardi, F., Winkler, C., Quignodon, L., Diserens, J.-G., Toffoli, B., Schiffrin, M., Sardella, C., Preitner, F., Desvergne, B., 2019. Systemic PPARγ deletion in mice provokes lipoatrophy, organomegaly, severe type 2 diabetes and metabolic inflexibility. Metab. Clin. Exp. 95, 8–20. doi:10.1016/j.metabol.2019.03.003

Grey, A., 2008. Skeletal consequences of thiazolidinedione therapy. Osteoporos Int 19, 129–137. doi:10.1007/s00198-007-0477-y

Horowitz, M.C., Berry, R., Holtrup, B., Sebo, Z., Nelson, T., Fretz, J.A., Lindskog, D., Kaplan, J.L., Ables, G., Rodeheffer, M.S., Rosen, C.J., 2017. Bone marrow adipocytes. Adipocyte 6, 193–204. doi:10.1080/21623945.2017.1367881

Idris, A.I., Sophocleous, A., Landao-Bassonga, E., Canals, M., Milligan, G., Baker, D., van't Hof, R.J., Ralston, S.H., 2009. Cannabinoid receptor type 1 protects against age-related osteoporosis by regulating osteoblast and adipocyte differentiation in marrow stromal cells. Cell Metab. 10, 139–147. doi:10.1016/j.cmet.2009.07.006

Iwabu, M., Yamauchi, T., Okada-Iwabu, M., Sato, K., Nakagawa, T., Funata, M., Yamaguchi, M., Namiki, S., Nakayama, R., Tabata, M., Ogata, H., Kubota, N., Takamoto, I., Hayashi, Y.K., Yamauchi, N., Waki, H., Fukayama, M., Nishino, I., Tokuyama, K., Ueki, K., Oike, Y., Ishii, S., Hirose, K., Shimizu, T., Touhara, K., Kadowaki, T., 2010. Adiponectin and AdipoR1 regulate PGC-1alpha and mitochondria by Ca(2+) and AMPK/SIRT1. Nature 464, 1313–1319. doi:10.1038/nature08991

Kahn, S.E., Zinman, B., Lachin, J.M., Haffner, S.M., Herman, W.H., Holman, R.R., Kravitz, B.G., Yu, D., Heise, M.A., Aftring, R.P., Viberti, G., for the A Diabetes Outcome Progression Trial (ADOPT) Study Group, 2008. Rosiglitazone-Associated Fractures in Type 2 Diabetes: An analysis from A Diabetes Outcome Progression Trial (ADOPT). Diabetes Care 31, 845–851. doi:10.2337/dc07-2270

Kajimura, D., Lee, H.W., Riley, K.J., Arteaga-Solis, E., Ferron, M., Zhou, B., Clarke, C.J., Hannun, Y.A., DePinho, R.A., Guo, X.E., Guo, E.X., Mann, J.J., Karsenty, G., 2013. Adiponectin regulates bone mass via opposite central and peripheral mechanisms through FoxO1. Cell Metab. 17, 901–915. doi:10.1016/j.cmet.2013.04.009

Kawai, M., Rosen, C.J., 2010. PPARγ: a circadian transcription factor in adipogenesis and osteogenesis. Nat Rev Endocrinol 6, 629–636. doi:10.1038/nrendo.2010.155

Lazarenko, O.P., Rzonca, S.O., Hogue, W.R., Swain, F.L., Suva, L.J., Lecka-Czernik, B., 2007. Rosiglitazone Induces Decreases in Bone Mass and Strength that Are Reminiscent of Aged Bone. Endocrinology 148, 2669–2680. doi:10.1210/en.2006-1587

Li, Y., Meng, Y., Yu, X., 2019. The Unique Metabolic Characteristics of Bone Marrow Adipose Tissue. Front Endocrinol (Lausanne) 10, 69. doi:10.3389/fendo.2019.00069

Liu, L.-F., Shen, W.-J., Ueno, M., Patel, S., Kraemer, F.B., 2011. Characterization of age-related gene expression profiling in bone marrow and epididymal adipocytes. BMC Genomics 12, 212. doi:10.1186/1471-2164-12-212

Madel, M.-B., Ibáñez, L., Rouleau, M., Wakkach, A., Blin-Wakkach, C., 2018. A Novel Reliable and Efficient Procedure for Purification of Mature Osteoclasts Allowing Functional Assays in Mouse Cells. Frontiers in immunology 9, 2567. doi:10.3389/fimmu.2018.02567

Moitra, J., Mason, M.M., Olive, M., Krylov, D., Gavrilova, O., Marcus-Samuels, B., Feigenbaum, L., Lee, E., Aoyama, T., Eckhaus, M., Reitman, M.L., Vinson, C., 1998. Life without white fat: a transgenic mouse. Genes & Development 12, 3168–3181. doi:10.1101/gad.12.20.3168

Mukohira, H., Hara, T., Abe, S., Tani-ichi, S., Sehara-Fujisawa, A., Nagasawa, T., Tobe, K., Ikuta, K., 2019. Mesenchymal stromal cells in bone marrow express adiponectin and are efficiently targeted by an adiponectin promoter-driven Cre transgene. International immunology 45, 1219–13. doi:10.1093/intimm/dxz042

Naot, D., Watson, M., Callon, K.E., Tuari, D., Musson, D.S., Choi, A.J., Sreenivasan, D., Fernandez, J., Tu, P.T., Dickinson, M., Gamble, G.D., Grey, A., Cornish, J., 2016. Reduced Bone Density and Cortical Bone Indices in Female Adiponectin-Knockout Mice. Endocrinology 157, 3550–3561. doi:10.1210/en.2016-1059

Okada-Iwabu, M., Yamauchi, T., Iwabu, M., Honma, T., Hamagami, K.-I., Matsuda, K., Yamaguchi, M., Tanabe, H., Kimura-Someya, T., Shirouzu, M., Ogata, H., Tokuyama, K., Ueki, K., Nagano, T., Tanaka, A., Yokoyama, S., Kadowaki, T., 2013. A small-molecule AdipoR agonist for type 2 diabetes and short life in obesity. Nature 503, 493–499. doi:10.1038/nature12656

Oshima, K., Nampei, A., Matsuda, M., Iwaki, M., Fukuhara, A., Hashimoto, J., Yoshikawa, H., Shimomura, I., 2005. Adiponectin increases bone mass by suppressing osteoclast and activating osteoblast. Biochemical and Biophysical Research Communications 331, 520–526. doi:10.1016/j.bbrc.2005.03.210

Pacheco-Pantoja, E.L., Waring, V.J., Wilson, P.J.M., Fraser, W.D., Gallagher, J.A., 2013. Adiponectin receptors are present in RANK-L-induced multinucleated osteoclast-like cells. J. Recept. Signal Transduct. Res. 33, 291–297. doi:10.3109/10799893.2013.828070

Poloni, A., Maurizi, G., Serrani, F., Mancini, S., Zingaretti, M.C., Frontini, A., Cinti, S., Olivieri, A., Leoni, P., 2013. Molecular and functional characterization of human bone marrow adipocytes. Experimental hematology 41, 558–566.e2. doi:10.1016/j.exphem.2013.02.005

Rosen, E.D., Sarraf, P., Troy, A.E., Bradwin, G., Moore, K., Milstone, D.S., Spiegelman, B.M., Mortensen, R.M., 1999. PPAR gamma is required for the differentiation of adipose tissue in vivo and in vitro. Mol. Cell 4, 611–617. doi:10.1016/s1097-2765(00)80211-7

Rzonca, S.O., Suva, L.J., D, G., DC, M., Lecka-Czernik, B., 2004. Bone is a target for the antidiabetic compound rosiglitazone. Endocrinology 145, 401–406.

Sardella, C., Winkler, C., Quignodon, L., Hardman, J.A., Toffoli, B., Giordano Attianese, G.M.P., Hundt, J.E., Michalik, L., Vinson, C.R., Paus, R., Desvergne, B., Gilardi, F., 2018. Delayed Hair Follicle Morphogenesis and Hair Follicle Dystrophy in a Lipoatrophy Mouse Model of Pparg Total Deletion. J. Invest. Dermatol. 138, 500–510. doi:10.1016/j.jid.2017.09.024

Sheu, Y., Cauley, J.A., 2011. The role of bone marrow and visceral fat on bone metabolism. Curr Osteoporos Rep 9, 67–75. doi:10.1007/s11914-011-0051-6

Shinoda, Y., Yamaguchi, M., Ogata, N., Akune, T., Kubota, N., Yamauchi, T., Terauchi, Y., Kadowaki, T., Takeuchi, Y., Fukumoto, S., Ikeda, T., Hoshi, K., Chung, U.-I., Nakamura, K., Kawaguchi, H., 2006. Regulation of bone formation by adiponectin through autocrine/paracrine and endocrine pathways. J. Cell. Biochem. 99, 196–208. doi:10.1002/jcb.20890

Sun, H., Kim, J.K., Mortensen, R., Mutyaba, L.P., Hankenson, K.D., Krebsbach, P.H., 2013. Osteoblast-Targeted suppression of PPARγ increases osteogenesis through activation of mTOR signaling 31, 2183–2192. doi:10.1002/stem.1455

Takada, I., Kouzmenko, A.P., Kato, S., 2009. Wnt and PPARγ signaling in osteoblastogenesis and adipogenesis 5, 442–447. doi:10.1038/nrrheum.2009.137

Tontonoz, P., Spiegelman, B.M., 2008. Fat and beyond: the diverse biology of PPARgamma. Annual review of biochemistry 77, 289–312. doi:10.1146/annurev.biochem.77.061307.091829

Wan, Y., 2010. PPARγ in bone homeostasis. Trends Endocrinol. Metab. 21, 722–728. doi:10.1016/j.tem.2010.08.006

Wan, Y., Chong, L.-W., Evans, R.M., 2007. PPAR-gamma regulates osteoclastogenesis in mice. Nat. Med. 13, 1496–1503. doi:10.1038/nm1672

Wang, F., Mullican, S.E., DiSpirito, J.R., Peed, L.C., 2013. Lipoatrophy and severe metabolic disturbance in mice with fat-specific deletion of PPAR. Proceedings of the National Academy of Sciences 110, 18656–18661. doi:10.1073/pnas.1314863110

Wang, Q.-P., Li, X.-P., Wang, M., Zhao, L.-L., Li, H., Xie, H., Lu, Z.-Y., 2014. Adiponectin exerts its negative effect on bone metabolism via OPG/RANKL pathway: an in vivo study. Endocrine 47, 845–853. doi:10.1007/s12020-014-0216-z

Williams, G.A., Wang, Y., Callon, K.E., Watson, M., Lin, J.-M., Lam, J.B.B., Costa, J.L., Orpe, A., Broom, N., Naot, D., Reid, I.R., Cornish, J., 2009. In vitro and in vivo effects of adiponectin on bone. Endocrinology 150, 3603–3610. doi:10.1210/en.2008-1639

Yamaguchi, N., Kukita, T., Li, Y.-J., Kamio, N., Fukumoto, S., Nonaka, K., Ninomiya, Y., Hanazawa, S., Yamashita, Y., 2008. Adiponectin inhibits induction of TNF-alpha/RANKL-stimulated NFATc1 via the AMPK signaling. FEBS Lett 582, 451–456. doi:10.1016/j.febslet.2007.12.037

Zou, W., Rohatgi, N., Chen, T.H.-P., Schilling, J., Abu-amer, Y., Teitelbaum, S.L., 2016. PPAR-γ regulates pharmacological but not physiological or pathological osteoclast formation. Nat. Med. 22, 1203–1205. doi:10.1038/nm.4208