Development and evaluation of a loop-mediated isothermal amplification (LAMP) assay for

the detection of *Tomato brown rugose fruit virus* (ToBRFV)

Alian Sarkes[1], Heting Fu[1], David Feindel[1], Michael W. Harding[2], and Jie Feng[1]

[1] Alberta Plant Health Lab, Crop Diversification Centre North, Alberta Agriculture and Forestry,

Edmonton, AB, T5Y 6H3, Canada;

[2] Crop Diversification Centre South, Alberta Agriculture and Forestry, Brooks, AB, T1R 1E6,

Canada

Correspondence to:

Michael Harding, email: michael.harding@gov.ab.ca

Jie Feng, email: jie.feng@gov.ab.ca

Abstract

Tomato brown rugose fruit virus (ToBRFV) is a member of *Tobamovirus* infecting tomato and pepper. Within North America, both the United States and Mexico consider ToBRFV to be a regulated pest. In Canada, the presence of ToBRFV has been reported, but an efficient diagnostic system has not yet been established. Here, we describe the development and assessment of a loop-mediated isothermal amplification (LAMP)-based assay to detect ToBRFV. The LAMP test was efficient and robust, and results could be obtained within 35 min with an available RNA sample. Amplification was possible when either water bath or oven were used to maintain the temperature at isothermal conditions (65°C), and results could be read by visual observation of colour change. Detection limit of the LAMP was eight target RNA molecules. Under the experimental conditions tested, LAMP was as sensitive as qPCR and 100 times more sensitive than the currently used rt-PCR. We recommend this sensitive, efficient LAMP protocol to be used for routine lab testing of ToBRFV.

Introduction

Tomato (*Solanum lycopersicum* L.) is one of the most important vegetable crops in the world [1].

Canada is the main producer of greenhouse tomatoes in North America, with an annual yield of

more than 250,000 t and a value of around $500 million (Agriculture & Agri-Food Canada,

www.agr.gc.ca/pmc-cropprofiles). Diseases caused by viruses are one of the most critical factors

affecting tomato production worldwide [2]. *Tomato brown rugose fruit virus* (ToBRFV) is a

member of the *Tobamovirus* genus with a host range including tomato, pepper and some weeds

such as nightshades [3]. It was first detected in Israel in 2014 [4] and later in countries in North

America [5,6], Asia [7-9] and Europe [10,11]. Within North America, both the United States [12]

and Mexico [13] consider ToBRFV to be a regulated pest. In Canada, ToBRFV was first

identified in Ontario in 2019 (http://thegrower.org/news/tomato-brown-rugose-fruit-virus-

identified-ontario). Since then the Alberta Plant Health Lab (APHL) began testing of tomato

samples submitted from Alberta greenhouses, but no positive samples had been found at the time

this report was submitted.

ToBRFV has a single-stranded positive-sense RNA of ~6,400 nucleotides (nt), with a typical

tobamovirus genome organization that consists in four open reading frames (ORFs) encoding

two replication-related proteins [7]. On tomato, ToBRFV causes symptoms including leaf

interveinal yellowing and deformation, mosaic staining, young leaves deformation and necrosis,

sepal necrosis and deformation, young fruits discoloration, deformation and necrosis [3]. Besides

the significant damage to yield and fruit quality, ToBRFV is of special concern compared to

other *tobamoviruses* because of its ability to overcome the *R* genes *Tm-2* and *L*, which are

deployed in tomato and pepper varieties, respectively, for *tobamoviruses* resistance [4].

54 An effective disease management program is dependent on timely and proper identification to

55 the causal agent of the disease. Enzyme-linked immunosorbent assay (ELISA), reverse

56 transcriptase polymerase chain reaction (rt-PCR), next-generation sequencing (NGS) and

57 transmission electron microscopy (TEM) have been used for ToBRFV detection [3]. In Canada,

58 the Alberta Plant Health Lab, Alberta Agriculture and Forestry, had utilized the rt-PCR methods

59 developed by Ling et al. [6] for diagnosis on suspected tomato samples. Recently, protocols

60 employing real time rt-PCR for ToBRFV detection were developed and demonstrated to be

61 highly specific and sensitive [14].

63 As an alternative of the above mentioned diagnostic methods, loop-mediated isothermal

64 amplification (LAMP) was not yet available for ToBRFV. This technique was originally

65 designed by Notomi et al. [15], in which six specific primers named internal primers (F2-

66 F1c/B2-B1c), external primers (F3/B3) and loop-specific primers (LoopF/LoopB) were used.

67 Compared to other methods, LAMP has advantages due to its rapidity, specificity and simplicity.

68 For example, a typical LAMP reaction can be completed within 30 min, and it is highly specific

69 as it can detect six gene regions of the target sequence by the six primers. Furthermore, LAMP

70 does not need specialized equipment, such as a thermocycler, but can be conducted in warm

71 water bath, thermal blocks, or oven, and does not require electrophoresis for detection and

72 identification of the reaction product(s).

74 LAMP diagnostic protocols have been developed for many plant diseases [16-20]. However,

75 there was no such protocol available for ToBRFV. The purpose of the present study was to

develop a LAMP test for detection of ToBRFV and compare it with PCR and qPCR tests. Our

results indicated that the LAMP method was sensitive and specific with potential to be developed

into a field-friendly diagnostic test.

Materials and Methods

Chemicals and standard techniques

All chemicals and equipment were purchased from Fisher Scientific Canada (Ottawa, ON) unless

otherwise specified. Extraction of genomic DNA and total RNA from plant samples was

conducted using a DNeasy plant mini kit (Qiagen Canada, Toronto, ON) and a RNeasy plant

mini kit (Qiagen Canada), respectively. Synthesis of the first-strand cDNA from the extracted

total RNA was conducted using a QuantaBio qScript cDNA synthesis kit (VWR Canada,

Edmonton, AB). Primers and gBlocks were synthesized by Integrated DNA Technologies

(Coralville, IA). LAMP, polymerase chain reaction (PCR) and reverse transcriptase (rt)-PCR

were conducted in a Proflex 96-well PCR cycler. Quantitative (q) PCR was performed in a

CFX96 touch real-time PCR detection system (Bio-Rad Canada, Mississauga, ON).

LAMP primer design and pre-selection

A query of the national center for biotechnology information (NCBI) whole genome sequence

(WGS) database (https://www.ncbi.nlm.nih.gov/genome) with the term "tomato brown rugose

fruit virus" resulted in one entry (tested on Feb 25, 2020) with an International Nucleotide

Sequence Database Collaboration (INSDC) number KT383474. Based on this WGS, ten sets of

LAMP primers were designed using LAMP Designer

(http://www.premierbiosoft.com/isothermal/lamp.html), an online tool designing LAMP primers,

and screening them for specificity. Using the WGS of KT383474 as query, a BLASTn was

conducted against the NCBI database. From the result of the BLAST, all available ToBRFV

WGS (seven in total as confirmed by Feb 25, 2020), three selected TMV WGS and two selected

ToMV WGS were retrieved. Alignment of these 12 sequences was conducted using Clustal

Omega (https://www.ebi.ac.uk/Tools/msa/clustalo). The output of the alignment was transferred

into a Microsoft Word file and the file was named Align.doc. Based on Align.doc, the specificity

of all primers in the ten primer sets was manually checked. If the last five nucleotides at the 3'

end of a primer matched with any of the non-targets (the three TMV and the two ToMV) or did

not match with one of the targets (the seven ToBRFV), the entire primer set that this primer

belongs to was discarded. After this pre-selection, three LAMP primer sets remained and one set

was selected for use in the subsequent studies. The selected primer set was hereafter referred as

LAMP primers and their locations and sequences are given in Fig. 1 and Table 1, respectively.

Design of gBlocks and qPCR primers

In the file Align.doc, one genomic region of 999 nt containing all binding sites of the LAMP

primers was selected. Three 999-nt double-stranded DNA fragments, corresponding to the RNA

sequence of this genomic region in one selected ToBRFV strain (INSDC no: KT383474; nt

1657-2655), one selected TMV strain (INSDC no: FR878069; nt 1649-2647) and one selected

ToMV strain (INSDC no: MH507166; nt 1653-2651) were synthesized as gBlocks. These

gBlocks were named gBlock^{ToBRFV}, gBlock^{TMV} and gBlock^{ToMV}, respectively. In addition, two

gBlocks, in which the sequences of the binding site of primer F1c or B2 on gBlock^{ToBRFV} were

replaced with the corresponding sequences in one of the ToMV strains (DQ873692), were

synthesized and named gBlock$^{\Delta F1c}$ and gBlock$^{\Delta B2}$, respectively (Table 1 and Fig. 1). Based on

the sequence of gBlock^ToBRFV, a pair of qPCR primers was designed using the Primer3 software

(http://primer3.ut.ee). This primer pair was name qF1/qR1 and their sequences are given in Table

1.

LAMP assays

All LAMP reactions were conducted in the WarmStart colorimetric LAMP master mix (NEB

Canada, Whitby, ON). Each reaction was 25 µL in total and contained 1 µL template. Quantities

of other components in each reaction followed the NEB's instructions for the master mix. All

reactions were conducted in individual 200-µL PCR tubes. The reaction program consisted of

only one step in which the tubes were incubated at 65°C for 30 minutes. After the incubation, the

reactions were checked visually and the results were recorded by photography.

PCR, rt-PCR and qPCR

Each reaction of PCR, rt-PCR and qPCR was 25 µL containing 1 µL template and 0.25 µM of

each primer. PCR and qPCR was conducted in Promega PCR master mix and SsoAdvanced

universal SYBR green supermix (Bio-Rad Canada), respectively. The PCR program consisted of

an initial denaturation at 94°C for 3 min, followed by 40 cycles of denaturation at 94°C for 30 s,

annealing at 55°C for 45 s and extension at 72°C for 1 min, and a final extension at 72°C for 5

min. The qPCR program consisted of an initial denaturation step of 95°C for 2 min, followed by

40 cycles of 5 s at 95°C and 1 min at 60°C. After completion of the qPCR amplification, a

melting curve analysis was run to evaluate the amplification specificity. rt-PCR was conducted

with a OneStep RT-PCR kit (Qiagen Canada) by following the manufacturer's instructions.

Assays on gBlocks

Each gBlock was dissolved in water and the concentration was adjusted to 100 fM, which

roughly equals to 60000 molecules/µL. From this concentration a set of 10× serial dilutions were

prepared for each gBlock down to a concentration of 0.06 molecules/µL (seven solutions for

each gBlock). Using the serial dilutions of all gBlocks as templates, LAMP was conducted.

Using the serial dilutions of gBlock-ToBRFV as templates, PCR and qPCR were conducted with

primers ToBRFV-F/ToBRFV-R [6](Ling et al. 2019) and qF1/qR1 (Table 1), respectively. In all

assays, sterile, DNA-free water was used as the negative control. All LAMP, PCR and qPCR

were repeated using the same preparations of serial dilutions.

Assays on genomic DNA from health plants

Genomic DNA was extracted from young leaves of Tomato (var. Cherry Nebula), pepper (var.

Tabasco), potato (*Solanum tuberosum* var. Norland) and tobacco (*Nicotiana tabacum* unknown

variety). Using 40 ng of genomic DNAs as templates, LAMP was conducted individually for

each of the four host plants. The LAMP assay was repeated on alternative DNA preparations.

Assays on RNA of infected plants

Total RNA was extracted from tomato plants with symptoms of ToBRFV infection. Five RNA

samples, confirmed to be ToBRFV positive, were provided to APHL by the Plant Disease Clinic

of University of Guelph and named 'original RNA samples'. The APHL tested the five original

RNA samples again with rt-PCR using the primer pair ToBRFV-F/ToBRFV-R, and with LAMP

using the designed LAMP primers. In addition, a 2-µL subsample was taken from each of the

five original RNA samples; the five subsamples were pooled and diluted 30× to form a 300-µL

RNA mixture. From this mixture, a set of eight 10-fold dilution series was prepared. rt-PCR and

LAMP were conducted on all eight dilutions using the above indicated primers. In all assays,

sterile, RNA-free water was used as the negative control. All assays were repeated using the

same RNA preparations.

Assays on cDNA derived from the infected plants

A 2-µL subsample was taken from each of the five original RNA samples; the five subsamples

were pooled, from which the first-strand cDNA was synthesized and the final cDNA solution

was adjusted to 300 µL. From this cDNA solution, a set of eight 10-fold dilution series was

prepared. These eight dilutions were used as template for PCR, LAMP and qPCR reactions. The

primer pair ToBRFV-F/ToBRFV-R was used in PCR. Primer pairs ToB5520F/ToB5598R and

qF1/qF2 were used in qPCR. In all assays, sterile DNA-free water was used as the negative

control. All assays were repeated using the same cDNA preparations.

Statistics

In all qPCR assays, each sample was tested with three technical repeats. The average and

standard deviation of the quantification cycle (Cq) values from the three repeats were calculated

using MS Excel. Calculation of Poisson distribution was conducted with an online Poisson

distribution calculator (https://stattrek.com/online-calcµLator/poisson.aspx).

Results

LAMP on the gBlock

In all LAMP reactions, changing color of the reaction mixture from pink to yellow indicated a

positive result, which was hereafter referred to as positive signal. With the designed LAMP

primers, no positive signal could be generated from gBlock^{TMV} and gBlock^{ToMV} (Fig. 2A).

Positive signals were generated from gBlock^{ToBRFV} when its molecule number was more than or

equal to 6 per 25-µL reaction (Fig. 2A). When gBlock$^{\Delta F1c}$ and gBlock$^{\Delta B2}$ were used as the

templates, positive signals could be generated when their molecule numbers were more than or

equal to 600 per 25-µL reaction; there was no positive signal when the molecule numbers were

less than or equal to 60 per 25-µL reaction (Fig. 2B). These results confirmed that the LAMP

primers were ToBRFV-specific. When the 6 molecules/µL gBlock^{ToBRFV} sample was repeatedly

tested by LAMP, 35 out of 40 reactions were found to be positive (Fig. 2C). This result indicated

that the LAMP assay could detect as little as 6 molecules in a 25-µL reaction more than 87% of

the time. In contrast, PCR could detect the target when the molecule number of gBlock^{ToBRFV}

was more than or equal to 600 per 25-µL reaction (Fig. 2D). In qPCR assay, Cq values were

generated from all of the three repeated reactions when the molecule number of gBlock ToBRFV

was more than or equal to 6 per 25 µL; when the molecule number was 0.6 per 25 µL, only one

out of the three reactions generated a Cq value (Fig. 2E). These results indicated that the

sensitivity of the LAMP assay on the gBlock was similar to that of qPCR but 100 times higher

than that of PCR. Repeated assays of LAMP, PCR and qPCR produced results similar to Figs.

2A and 2C, Fig. 2D and Fig. 2E, respectively.

LAMP on plant genomic DNA

When genomic DNA from healthy plants of four *Solanaceae* species was used as the template in

LAMP, no positive signal could be generated (data not shown). This result confirmed the

specificity of the LAMP primers.

LAMP on RNA samples

When RNA was used as the template, LAMP generated positive signals from all five original

RNA samples (Fig. 3A). This result was confirmed by rt-PCR (Fig. 3B). When the serial

dilutions were used as the templates, LAMP could generate positive signals from the dilutions 1-

7 (Fig. 3C); in contrast, rt-PCR could only produce a band from dilutions 1-5 (Fig. 3D). This

result indicated that LAMP was 100 times more sensitive than rt-PCR on RNA samples.

Repeated assays of LAMP and PCR produced results similar to Figs. 3A and 3C and Figs. 3B

and 3D, respectively.

Efficiency of LAMP on cDNA

When the serial dilutions of cDNA were used as the templates, LAMP generated positive signals

from samples 1-6 dilutions (Fig. 4A). In contrast, PCR generated positive bands from samples 1-

4 (Fig. 4B). qPCR assays using alternative primer sets showed similar sensitivity as LAMP (Fig.

4C). Repeated assays of LAMP, PCR and qPCR produced results similar to Fig. 4A, Fig. 4B and

Fig. 4C, respectively. These results indicated that the sensitivity of the LAMP assay on the

cDNA samples was similar to that of qPCR but 100 times higher than that of PCR.

Discussion

Specificity of the LAMP

LAMP is more specific than other PCR-based techniques because LAMP has six specific

primers. In this study, all LAMP primers were designed to be specific to all ToBRFV strains

with an available WGS, and non-specific to TMV and ToMV. TMV and ToMV are closely

related to ToBRFV [3], which is supported by the fact that WGS of TMV and ToMV were

always the top BLAST hits when the NCBI database was queried with the ToBRFV WGS. When

the LAMP primers were tested on the gBlock^{TMV} and gBlock^{ToMV}, as well as genomic DNA

from various *Solanaceae* plants, no signal could be generated, further confirming the specificity

of the LAMP primers to ToBRFV. Moreover, when the LAMP primers were tested on

gBlock$^{\Delta F1c}$ and gBlock$^{\Delta B2}$, which differed from gBlock^{ToBRFV} by only on a few nt on the binding

sites of primer F1c and B2, respectively, no positive signal could be generated at concentrations

of ≤ 60 molecules/25µL. This result supported the conclusion that the LAMP primers were

highly specific to ToBRFV.

Sensitivity of the LAMP

On the serial dilutions of the gBlocks or cDNA, LAMP was 100 times more sensitive than rt-

PCR and similar in sensitivity as qPCR. As well, on the serial dilutions of the RNA samples,

LAMP was 100 times more sensitive than rt-PCR. These results were comparable with those of

LAMP testing on other RNA viruses. For examples, LAMP was 100 times more sensitive than

rt-PCR on the detection of *Citrus leaf blotch virus* [21] and *Little cherry virus 1* [22]; on *Potato*

virus X [23] and *Sugarcane streak mosaic virus* [24], LAMP was 10 times more sensitive than rt-

PCR. Compared to rt-qPCR, LAMP was similarly sensitive for detection of *Onion yellow dwarf*

virus [25] and *Potato leafroll virus* [26].

Testing primers on gBlocks allows more accurate calculation of primer efficiency. In this study,

we obtained data that 35 out of 40 LAMP reactions generated positive signals when the template

gBlock concentration was 6 molecules/µL. At a very low concentration, the distribution of DNA

particles in aliquots follows a Poisson distribution [27]. Using the Poisson distribution calculator,

we could calculate that the possibility of obtaining ≥ 4 gBlock molecules in a 1-µL aliquot from

a 6 molecules/µL solution was 85%, which is comparable to the data from the LAMP test (35/40

= 88%). Since gBlock is double-stranded and the virus RNA is single-stranded, we concluded

that the LAMP could detect the virus RNA in a concentration of as low as 8 viruses per reaction.

This is comparable to qPCR results obtained in this study and to the theoretical limit of qPCR

detection (3 molecules) proposed by Forootan et al. [28. The estimated efficiency of the RNA

extraction kit is approximately 40 to 80% (the ratio between the molecule numbers of extracted

genomic RNA and the number of virus particles in the sample) when the sample was 100-mg

tomato leaf tissue and the final RNA volume was 50 µL (Feng, unpublished data). Since 1 µL

template was used in the LAMP test, we concluded that the LAMP test could detect ToBRFV

when a 100-mg plant sample contains as little as 1,000 virus particles (8÷40%×50). While this

level of detection is encouraging, it is important to note that the minimal number of virus

particles necessary to cause plant infection could be very low. For example, one Tobacco etch

virus (TEV) particle is sufficient to initiate a systemic infection [29]. We want to emphasize here

that more improvement is required on the commonly used diagnostic protocols, and that the

efficiency of a molecular-based diagnostic protocol is determined not only by the efficiency of

the detection technique only, but also by other factors such as the efficiency of DNA/RNA

extraction.

Time-saving and simplicity of LAMP

Compared to PCR, rt-PCR or rt-qPCR, LAMP is time saving. For example, in the present study, testing one DNA, RNA or cDNA sample by LAMP could be completed within 35 min. Testing eight samples could be completed in less than one hour. In contrast, testing one or eight samples by PCR or rt-PCR needed more than three hours (for PCR and electrophoresis) and testing by rt-qPCR would need more than 2 hours. All LAMP results reported in the present study were generated in a PCR thermocycler. However, identical results were obtained when the LAMP tests were conducted in an oven or a heat block. Thus, for labs where PCR or qPCR thermocycler is not available, LAMP protocol developed in this study is highly recommended for ToBRFV testing.

Acknowledgements

We thank Dr. Shannon Xuechan Shan from the Plant Disease Clinic of University of Guelph for providing all RNA samples used in this study. This study was supported by a grant from the Canadian Agricultural Partnership (no. 601322) to JF.

References

1. Foolad MR, Panthee DR. Marker-assisted selection in tomato breeding. Crit Rev Plant Sci. 2012;31: 93-123.

2. Hanssen IM, Lapidot M, Thomma BP. Emerging viral diseases of tomato crops. Mol Plant-Microbe Interact. 2010;23: 539-548.

3. Oladokun J, Halabi M, Barua, P, Nath PJPP. Tomato brown rugose fruit disease: current distribution, knowledge and future prospects. Plant Pathol. 2019;68: 1579-1586.

4.   Luria N, Smith E, Reingold V, Bekelman I, Lapidot M, Levin I, et al. A new Israeli

Tobamovirus isolate infects tomato plants harboring *Tm-22* resistance genes. PLoS One.

2017;12: e0170429.

5.   Cambrón-Crisantos JM, Rodríguez-Mendoza J, Valencia-Luna JB, Rangel SA, De Jesús

García-Ávila C, López-Buenfil JA. First report of *Tomato brown rugose fruit virus*

(ToBRFV) in Michoacan, Mexico. Mex J Phytopathol. 2018;37: 185-192.

6.   Ling KS, Tian T, Gurung S, Salati R, Gilliard A. First report of *Tomato brown rugose fruit*

virus infecting greenhouse tomato in the United States. Plant Dis. 2019;103: 1439.

7.   Salem N, Mansour A, Ciuffo M, Falk BW, Turina M. A new *tobamovirus* infecting tomato

crops in Jordan. Arch Virol. 2016;161: 503-506.

8.   Alkowni R, Alabdallah O, Fadda Z. 2019. Molecular identification of *Tomato brown rugose*

fruit virus in tomato in Palestine. J Plant Pathol. 2019;101: 719-723.

9.   Fidan H, Sarikaya P, Calis O. First report of *Tomato brown rugose fruit virus* on tomato in

Turkey. New Dis Rep. 2019;39: 18.

10. Menzel W, Knierim D, Winter S, Hamacher J, Heupel M. First report of *Tomato brown*

rugose fruit virus infecting tomato in Germany. New Dis Rep. 2019;39: 1.

11. Panno S, Caruso AG, Davino S. First report of *Tomato brown rugose fruit virus* on tomato

crops in Italy. Plant Dis. 2019;103: 1443.

12. USDA. Federal order for U.S. imports of tomato (*Solanum lycopersicum*) and pepper

(*Capsicum* spp.) hosts of *Tomato brown rugose fruit virus* (ToBRFV). 2019.

https://www.aphis.usda.gov/import_export/plants/plant_imports/federal_order/downloads/20

19/DA-2019-28.pdf.

13. Puente M. AMSAC - Mexican Seed Trade Association. Presentation delivered at NAPPO

AGM. Interactions between government and industry facing the impacts of an emerging

pest-the case of ToBRFV. Wednesday, Oct. 30, 2019.

14. Panno S, Ruiz-Ruiz S, Caruso AG, Alfaro-Fernandez A, San Ambrosio MI, Davino S. Real-

time reverse transcription polymerase chain reaction development for rapid detection of

Tomato brown rugose fruit virus and comparison with other techniques. PeerJ. 2019;7:

e7928.

15. Notomi T, Okayama H, Masubuchi H, Yonekawa T, Watanabe K, Amino N, et al. Loop-

mediated isothermal amplification of DNA. Nuc Acids Res. 2000;28: 63-70.

16. Kogovšek P, Mehle N, Pugelj A, Jakomin T, Schroers HJ, Ravnikar M. et al. Rapid loop-

mediated isothermal amplification assays for grapevine yellows phytoplasmas on crude leaf-

vein homogenate has the same performance as qPCR. Eur J Plant Pathol. 2017;148: 75-84.

17. Okiro LA, Tancos MA, Nyanjom SG, Smart CD, Parker ML. Comparative evaluation of

LAMP, qPCR, conventional PCR, and ELISA to detect *Ralstonia solanacearum* in Kenyan

potato fields. Plant Dis. 2019;103: 959-965.

18. Romero JL, Carver GD, Johnson PA, Perry KL, Thompson JR. A rapid, sensitive and

inexpensive method for detection of grapevine red blotch virus without tissue extraction

using loop-mediated isothermal amplification. Arch Virol. 2019;164: 1453-1457.

19. Yan H, Zhang J, Ma D, Yin J. qPCR and loop mediated isothermal amplification for rapid

detection of *Ustilago tritici*. PeerJ. 2019;7: e7766.

20. Zhang L, Gleason C. Loop-mediated isothermal amplification for the diagnostic detection of

Meloidogyne chitwoodi and *M. fallax*. Plant Dis. 2019;103: 12-18.

21. Liu H, Wu W, Tan J, Li Y, Mi W, Jiang L. et al. Development and evaluation of a one-step

reverse transcription loop-mediated isothermal amplification for detection of *Citrus leaf*

blotch virus. J Virol Methods. 2019;270: 150-152.

22. Tahzima R, Foucart Y, Peusens G, Belien T, Massart S, De Jonghe K. New sensitive and

fast detection of *Little cherry virus 1* using loop-mediated isothermal amplification (RT-

LAMP). J Virol Methods. 2019;265: 91-98.

23. Raigond B, Verma A, Roach S, Kochhar T, Arjunan J, Kumar R, et al. One-step reverse

transcription loop-mediated isothermal amplification: a simple, sensitive and rapid assay for

detection of *Potato virus X* in potato leaves and tubers. Indian Phytopathol. 2019;72: 321-

328.

24. Wang KL, Deng QQ, Chen JW, Shen WK. Development of a reverse transcription loop-

mediated isothermal amplification assay for rapid and visual detection of *Sugarcane streak*

mosaic virus in sugarcane. Crop Protect. 2019;119: 38-45.

25. Tiberini A, Tomlinson J, Micali G, Fontana A, Albanese G, Tomassoli L. Development of a

reverse transcription-loop-mediated isothermal amplification (LAMP) assay for the rapid

detection of *Onion yellow dwarf virus*. J Virol Methods. 2019;271: 113680.

26. Raigond B, Verma A, Jandrajupalli S, Kochhar T, Sharma S, Chakrabarti SK. Squash print

reverse transcription loop-mediated isothermal amplification assay for detection of *Potato*

leafroll virus in single aphid and in potato. Potato Res. 2019: 1-4.

27. Wang Z, Spadoro J. Determination of target copy number of quantitative standards used in

PCR-based diagnostic assays. In: Ferr F, editor. Gene Quantification. 1998. Boston:

Birkhäuser; pp. 31-43.

28. Forootan A, Sjöback R, Björkman J, Sjögreen B, Linz L, Kubista M. Methods to determine

limit of detection and limit of quantification in quantitative real-time PCR (qPCR). Biomol

Detect Quantif. 2017;12: 1-6.

29. Zwart MP, Daròs JA, Elena SF. One is enough: in vivo effective population size is dose-

dependent for a plant RNA virus. PLoS Pathog. 2011;7: e1002122.

Author Contributions

Conceptualization: Jie Feng.

Data Curation: Jie Feng.

Formal Analysis: Jie Feng.

Funding Acquisition: Jie Feng, Michael W. Harding, David Feindel.

Investigation: Alian Sarkes, Heting Fu, Jie Feng.

Methodology: Jie Feng.

Project Administration: Jie Feng.

Resources: Jie Feng, David Feindel.

Supervision: Jie Feng.

Validation: Jie Feng.

Visualization: Jie Feng.

Writing – Original Draft Preparation: Jie Feng.

Writing – Review & Editing: Jie Feng, Michael W. Harding.

Table 1. Primers used in this study

Name	Sequence	Size (bp)	Reaction	Reference
F3	TTGGAGTCTTAGATGTTGCG	279		
B3	GGACACCGTCAACTAGGA			
FIP(F1c+F2)	CCTTCTCCAACTGTCGCAAGTCAC	n/a	LAMP	This study
	ATGCTAGGAAGTACCAC			
BIP(B1c+B2)	CCGTGAGTTCTGAGTCAATGGTTG			
	AGGCTCACCATCTCTTAA			
LoopF	CTCCATGCTCATCATACTCCAA	n/a		
LoopB	GCTCAGAACACTGAGGAGATT			
ToBRFV-F	GAAGTCCCGATGTCTGTAAGG	842	PCR & rt-PCR	[6]
ToBRFV-R	GTGCCTACGGATGTGTATGA			
ToB5520F	GTAAGGCTTGCAAAATTTCGTTCG	79	qPCR	[14]
ToB5598R	CTTTGGTTTTTGTCTGGTTTCGG			
qF1	CGGTGGATATGCCAGTGCTTG	99	qPCR	This study
qR1	TGTCCGACTCCTTGAGCACAG			

 Figure legends

 Fig. 1. Alignment of partial genomic sequences of selected ToBRFV, TMV and ToMV strains.

 Locations of LAMP primers were highlighted and the primer name indicated above. Degenerated

 nucleotides within the primer binding sites were bolded. Italic letters indicate the sequences

 present in gBlock$^{\Delta F1c}$ and gBlock$^{\Delta B2}$ that differentiate the two gBlocks from gBlock^{ToBRFV}. Line

 1, represents five ToRBFV strains. Lines 2 and 3, are sequences from two ToBRFV strains.

 Lines 4-6 are sequences from three TMV strains. Lines 7 and 8 are sequences from two ToMV

 strains. (Detailed description of the virus strains is list in Table S1).

 Fig. 2. LAMP (A-C), PCR (D) and qPCR (E) tests on gBlocks. A, LAMP test on serial dilutions

 of gBlock^{TMV}, gBlock^{ToMV} and gBlock^{ToBRFV}. B, LAMP test on two dilutions of gBlock$^{\Delta F1c}$ and

 gBlock$^{\Delta B2}$. C, Repeated LAMP tests on gBlock^{ToBRFV} at the concentration of 6 molecules per

 reaction. 40 repeats were conducted but only eight were shown; 35 out of the 40 were positive. D,

 PCR test on serial dilutions of gBlock^{ToBRFV} using the primer pair ToBRFV-F/ToBRFV-R. M,

 GeneRuler Express DNA Ladder. E, qPCR test on serial dilutions of gBlock^{ToBRFV} using the

 primer pair qF1/qR1.

 Fig. 3. LAMP (A and C) and rt-PCR (B and D) tests on total RNA extracted from five tomato

 samples infected by ToBRFV. A and B, Tests on five original RNA samples. a, from tomato fruit;

 b, from tomato leaf; c-e, from tomato flesh. M, GeneRuler express DNA ladder. C and D, Tests

 on serial dilutions (1-8) of a RNA sample pooled from same volumes of each of the five original

 RNA samples. M', TrackIt 100 bp DNA ladder.

419   **Fig. 4.** LAMP (A), PCR (B) and qPCR (C) tests on serial dilutions (1-7) of cDNA synthesized

420   from a RNA sample pooled from same volumes of each of the five original RNA samples. M,

421   GeneRuler express DNA ladder. Primer pair used in PCR was ToBRFV-F/ToBRFV-R and those

422   used in qPCR were qF1/qR1 for Test 1 and ToB5520F/ToB5598R for Test 2.

```
                                        F3
1 TAGATCTCGAAACCCGTCAGAAGTTTGGAGTCTTAGATGTTGCGACCAAAAGATGGTTAAT
2 TAGATCTCGAAACCCGTCAGAAGTTTGGAGTCTTAGATGTTGCGACCAAAAGATGGTTAAT
3 TAGATCTCGAAACCCGTCAGAAGTTTGGAGTCTTAGATGTTGCGACCAAAAGATGGTTAAT
4 TAGATCTCGAAACCCGTCAAAAGTTCGGAGTTCTGGATGTTGGCATCTAGGAAGTGGTTGAT
5 TTGACCTTGAAACCCGTCAAAAGTTTGGAGTCTTGGATGTTGCGATCTAGGAAGTGGTTGAT
6 TTGACCTTGAAACCCGTCAAAAGTTTGGAGTCTTGGATGTTCGCATCTAGGAAGTGGTTAAT
7 TTGACCTTGAAACTCGTCAAAAGTTCGGAGTCTTGGATGTTGCTTCGAAAAGGTGGCTAGT
8 TTGACCTTGAAACTCGTCAAAAGTTCGGAGTTCTGGATGTTGCTTCGAAAAGGTGGCTAGT

                                                     F2
1 TAAACCTTTAGCCAAGAATCACGCATGGGGCGTTATTGAAACACATGCTAGGAAGTACCAC
2 TAAACCTTTAGCCAAGAATCACGCATGGGGCGTTATTGAAACACATGCTAGAAAGTACCAC
3 TAAACCTTTAGCCAAGAATCACGCATGGGGCGTTATTGAAACACATGCTAGAAAGTACCAC
4 TAAACCGACTGCTAAGAGTCACGCATGGGGCGTTGTTGAGACTCATGCGAGGAAATATCAC
5 CAAACCAACGGCCAAGAGTCATGCATGGGGTGTTGTCGAAACCCACGCGAGGAAGTACAT
6 TAAACCGGCCAAGAGTCATGCATGGGGTGTTGTTGAAACCCACGCGAGGAAGTTCAT
7 TAAACCATCCGCAAAGAACCATGCATGGGGGGTTGTTGAGACTCATGCGAGGAAATATCAC
8 TAAACCATCCGCAAAGAACCATGCATGGGGGGTTGTTGAGACTCATGCGAGGAAATATCAC

              LoopF                              F1c
1 GTTGCACTTTTGGAGTATGATGAGCATGGAGTGGTAACTTGCGACAGTTGGAGAAGGGTGG
2 GTTGCACTTTTGGAGTATGATGAGCATGGAGTGGTAACTTGCGACAGTTGGAGAAGGGTGG
3 GTTGCACTTTTGGAGTATGATGAGCATGGAGTGGTAACTTGCGACAGTTGGAGAAGGGTGG
4 GTTGCACTTCTGGAGTACGATGAGCAGGGTATAGTAACTTGCGACAACTGGAGAAGAGTTG
5 GTGGCGCTTTTGGAATATGATGAGCAGGGTGTGGTAACATGCGATGATTGGAGAAGAGTAG
6 GTGGCGCTTCTGGAATATGATGAGCAGGGTGTGGTGACATGCGATAATTGGAGAAGAGTAG
7 GTCGCATTACTGGAGCACGATGAATTGGCATTATTACGTGCGATAACTGGCGACGGGTGG
8 GTCGCATTACTGGAGCACGATGAATTGGCATTATTACGTGCGATAACTGGCGACGGGTGG

         B1c                                LoopB
1 CCGTGAGTTCTGAGTCAATGGTTTATTCTGATATGGCAAAGCTCAGAACACTGAGGAGATT
2 CCGTGAGTTCTGAGTCAATGGTTTATTCTGATATGGCAAAGCTCAGAACACTGAGGAGATT
3 CCGTGAGTTCTGAGTCAATGGTTTATTCTGATATGGCAAAGCTCAGAACACTGAGGAGATT
4 CTGTTAGTTCTGAGTCAGGTTGTATATTCTGATATGGCGAAAACTCAGAACTCTGAGGAAATT
5 CTGTCAGCTCTGAGTCTGTTGTTTATTCCGACATGGCGAAAACTCAGAACTCTGCGCAGACT
6 CTGTCAGCTCTGAGTCTGTTGTTTATTCCGACATGGCGAAAACTCAGAACTCTGCGCAGACT
7 CTGTGAGTTCTGAGTCGGTAGTATATTCTGATATGGCTAAAACTTAGGACTCTGAGGAAGATT
8 CTGTGAGTTCTGAGTCGGTAGTATATTCTGATATGGCTAAAACTCAGGACTCTGAGGAAGATT

         B2                              B3
1 ATTAAGAGATGGTGAGCCTCATGTCAGCAGTGCTAAAGTCGTCCTAGTTGACGGTGTCCCG
2 ATTAAGAGATGGTGAGCCTCATGTCAGCAGTGCTAAAGTCGTCCTAGTTGACGGTGTCCCG
3 ATTAAGAGATGGAGAGCCTCATGTCAGCAGTGCTAAAGTCGTCCTAGTTGACGGTGTCCCG
4 GCTTCAAAATGGTGAACCTCATGTTAGTAACGCAAAGGTTGTTCTCGTGGACGGTGTCCCA
5 GCTTCGAAACGGTGAACCGCATGTCAGTAGCCGCAAAGGTTGTTCTTGTGGACGGAGTTCCG
6 GCTTCGAAACGGAGAACCGCATGTCAGTAGCCGCAAAGGTTGTTCTTGTGGACGGAGTTCCG
7 GCTCAAAGGTGAGAACCACACGTTAGTTCAGCAAAGGTTGTTTGGTGGATGGCATTCCA
8 GCTTAAAGATGGAGAACCACACGTTAGTTCAGCAAAGGTGGTTTGGTGGATGGCGTTCCA
```

A

Copy number of the target gBlock in the reaction

	60000	6000	600	60	6	0.6	0.06	water

TMV

ToMV

ToBRFV

B

gBlock ΔF1c   gBlock ΔB2

600	60	600	60

C

8 repeats of the 6-copies/reaction treatment

D

Copy number of the target gBlock in the reaction

M 60000 6000 600 60 6 0.6 0.06 water

1000
750
(bp)

E

	Copy number of the target gBlock in the reaction							
	60000	6000	600	60	6	0.6	0.06	water
Cq (mean±SD)	21.86 ±0.02	25.26 ±0.03	28.65 ±0.12	32.10 ±0.19	35.09 ±0.36	38.98	n/a	n/a
#repeats with a Cq	3	3	3	3	3	1	0	0