Ongoing exposure to peritoneal dialysis fluid alters resident peritoneal macrophage phenotype and activation propensity

Tara E. Sutherland[1,2], Tovah N. Shaw[1,2], Rachel Lennon[3,4], Sarah E. Herrick[1,4], Dominik Rückerl[1]

1 Lydia Becker Institute for Immunology and Infection, Faculty of Biology, Medicine and Health, Division of Infection, Immunity and Respiratory Medicine, University of Manchester, Manchester, UK
2 Manchester Collaborative Centre for Inflammation Research (MCCIR), University of Manchester, Manchester, UK
3 Wellcome Centre for Cell-Matrix Research, University of Manchester, Manchester, UK
4 Division of Cell Matrix Biology and Regenerative Medicine, Faculty of Biology Medicine and Health, School of Biological Sciences, University of Manchester, Manchester, UK

Running title: Tissue resident macrophages in PD

Corresponding author: Dr. Dominik Rückerl, Dominik.Ruckerl@manchester.ac.uk

Abstract:

Peritoneal dialysis (PD) is a more continuous alternative to haemodialysis, for patients with chronic kidney disease, with considerable initial benefits for survival, patient independence and healthcare cost. However, longterm PD is associated with significant pathology, negating the positive effects over haemodialysis. Importantly, peritonitis and activation of macrophages is closely associated with disease progression and treatment failure. However, recent advances in macrophage biology suggest opposite functions for macrophages of different cellular origins. While monocyte-derived macrophages promote disease progression in some models of fibrosis, tissue resident macrophages have rather been associated with protective roles. Thus, we aimed to identify the relative contribution of tissue resident macrophages to PD induced inflammation in mice. Unexpectedly, we found an incremental loss of homeostatic characteristics, anti-inflammatory and efferocytic functionality in peritoneal resident macrophages, accompanied by enhanced inflammatory responses to external stimuli. Thus, alterations in tissue resident macrophages may render longterm PD patients sensitive to developing peritonitis and consequently fibrosis/sclerosis.

Introduction:

An estimated 5-10 million people worldwide die every year due to chronic kidney disease (Luyckx, Tonelli, & Stanifer, 2018). In Europe, an average of 850 people per million

population (pmp) require renal replacement therapy (RRT) and 120 new patients pmp commence treatment annually (Kramer et al., 2019). The average 5-year survival rate of patients receiving RRT is only 50.5%. This can be improved to over 90% if patients receive a kidney transplant, but rates of transplantation remain low (32 pmp), primarily due to organ availability and the majority of patients rely on dialysis as a therapy to substitute excretory kidney function (Kramer et al., 2019).

Peritoneal dialysis (PD), utilizing the body's own peritoneal membrane as a filter during dialysis, is a cost-effective alternative to haemodialysis (HD). Although PD has been associated with better initial survival rates (van de Luijtgaarden et al., 2016), lower costs for the health system (Chui et al., 2013) and increased patient autonomy (François & Bargman, 2014) as compared to HD, the incidence rate of PD over several decades has dropped in Europe (van de Luijtgaarden et al., 2016). There are a variety of reasons for this reduction in PD uptake, but the significant risk of adverse effects and in some cases fatal outcomes, has limited the general adoption of PD in adult patients across Europe (Bartosova & Schmitt, 2019). Treatment failure is commonly associated with repeated episodes of peritonitis (i.e. inflammation) and a progressive thickening and vascularisation of the peritoneum, leading to impaired filtration and thus reduced efficacy of PD (Bartosova & Schmitt, 2019; Morelle et al., 2015). In rare cases, the fibrotic changes to the peritoneum become so extreme that they form a fibrous cocoon encapsulating the internal organs leading to persistent or recurring adhesive bowel obstruction (Johnson et al., 2010; Moinuddin, Summers, Van Dellen, Augustine, & Herrick, 2014; Morelle et al., 2015). The recognition of this Encapsulating Peritoneal Sclerosis (EPS) is an indication to discontinue PD with the mortality approaching 50% one year after diagnosis (Danford, Lin, Smith, & Wolf, 2018). Management of EPS includes surgery but there is a relatively high frequency of symptom recurrence (Danford et al., 2018). In contrast, immunosuppressive treatments and the use of anti-fibrotic agents, like tamoxifen, have shown noticeable benefits to patient survival (Danford et al., 2018). Therefore, aberrant activation of the immune system appears to be linked to both alteration of peritoneal structure and PD treatment failure, as well as the progression of the fibrotic sequelae. Indeed, experimental rodent models of the disease, have suggested inappropriate or excessive activation of macrophages as a major cause of the pathology (Q. Li et al., 2018; Liao et al., 2017; Minutti et al., 2017; Jie Wang et al., 2013).

Recent advances in macrophage biology have highlighted the intrinsic heterogeneity of

macrophage populations (Gordon & Plüddemann, 2017). Grossly simplified, macrophages

can be split into tissue resident macrophages (MΦres) which are present in tissues during

homeostasis, and monocyte-derived macrophages (MΦmono), which are recruited to the

tissue during inflammatory conditions (Yang Zhao, Zou, Du, & Zhao, 2018). Both

macrophage populations can and do respond to external stimuli, like infection, but they

possess distinct response profiles and adopt distinct functional properties (Gundra et al.,

2014; Ruckerl et al., 2017).

In health, peritoneal MΦres are essential for maintaining tissue homeostasis by silently

removing apoptotic cells through efferocytosis (Allen & Ruckerl, 2017; Uderhardt et al.,

2012) and by providing a source of tissue-reparative cells that infiltrate surrounding

tissues (e.g. the liver) during injury (Jing Wang & Kubes, 2016). Importantly, upon

encountering inflammatory signals, MΦres undergo the Macrophage Disappearance

Reaction (MDR) (Barth, Hendrzak, Melnicoff, & Morahan, 1995). Following injection of

inflammatory agents (e.g. bacterial antigens) or infection the number of peritoneal MΦres

detectable in peritoneal lavage will drop significantly within a few hours (Accarias et al.,

2016). The exact mechanism underlying MDR is still poorly understood, but it has been

proposed that MΦres undergo activation-induced cell death (Ruckerl et al., 2017) or

adhere to the mesothelial lining of the peritoneal cavity reducing their recovery (Jonjic et

al., 1992; Jing Wang & Kubes, 2016; N. Zhang et al., 2019). Of note, the degree of

detectable MΦres -loss directly correlates to the amount of inflammatory stimulus (e.g.

cfu of bacteria) and the recruitment of inflammatory MΦmono (Ruckerl et al., 2017).

Following resolution of the inflammation, MΦres can re-populate the peritoneal cavity and

return to homeostatic numbers through proliferative expansion of remaining MΦres

(Davies et al., 2013).

In models of fibrotic disorders of the lung or kidney, influx of monocytes and monocyte-

derived macrophages has been linked to disease progression and induction of pathology

(Braga et al., 2018; Misharin et al., 2017; Puranik et al., 2018). Similarly, in rodent models

of peritoneal fibrosis, preventing the influx of monocytes or depleting all macrophages

limits the degree of peritoneal thickening and improves glomerular filtration (Y.-T. Chen,

Hsu, Lin, Pan, Liu, Wu, Tsai, Liao, Cheng, Chiang, Chen, Chu, & Lin, 2019a; Lee et al.,

2012; Jie Wang et al., 2013). Moreover, injection of MΦmono, activated ex vivo using

bacterial antigens (i.e. Lipopolysaccharide), often referred to as M1 macrophages, exacerbates disease progression (Q. Li et al., 2018). Together these data highlight the role of inflammation and infiltration of MΦmono in the progression of peritoneal fibrosis and seem to provide a cohesive picture explaining the enhanced risk of PD failure associated with repeated episodes of peritonitis (Lambie et al., 2016). In this context, it is interesting to note that continuing peritoneal irrigation and thus continual removal of peritoneal cells, including any inflammatory infiltrate, in patients discontinuing PD, has been suggested to prevent subsequent EPS formation (Moriishi et al., 2002).

Other studies have found a role for anti-inflammatory, IL-4 activated macrophages (also called M2), characterised by the expression of CD206, Arg1 and Ym1, in promoting peritoneal dialysis fluid (PD fluid) induced fibrosis (Bellón et al., 2011; Chandrasekaran et al., 2019; Jie Wang et al., 2013). Additionally, chronic fibrotic kidney disease is linked to a switch from inflammatory M1 to predominantly M2 activated macrophages (Han, Ma, Tesch, Manthey, & Nikolic-Paterson, 2013). Indeed, both types of macrophage activation seem to have the capacity to promote kidney fibrosis, possibly acting during different phases of the pathology (Nikolic-Paterson, Wang, & Lan, 2015). Importantly, some markers used to define M2 are differentially expressed on MΦres and MΦmono. For example CD206 is expressed constitutively by MΦmono but not MΦres (Gundra et al., 2014). Thus, the described role of M2 macrophages may merely reflect an enhanced influx of MΦmono rather than IL-4 mediated activation. In fact, transfer of exogenously activated M2 macrophages showed no impact on disease progression in a model of peritoneal fibrosis (Q. Li et al., 2018) and even reduced pathology in a model of renal fibrosis (Y Wang et al., 2007). Moreover, using different strategies to deplete macrophages in kidney fibrosis yielded opposing results with regard to disease progression indicating that depletion of different subsets of macrophage may lead to different outcomes (Ferenbach et al., 2012). Indeed, renal MΦres as compared to MΦmono have been shown to be protective in a model of kidney pathology (Puranik et al., 2018). This suggests that the cellular origin of macrophages may play a prominent part in determining their role during fibrotic disorders and influence the overall disease outcome.

Here we analysed the effects of repeated PD fluid injection on macrophage population dynamics and responses to activating signals. Our data indicate a significant change in MΦres phenotype over time during PD fluid administration. MΦres lost expression of anti-

inflammatory and efferocytic markers correlating with the duration of PD fluid treatment. Moreover, MΦres from PD fluid treated animals showed enhanced inflammatory responses to external stimulation. Importantly, the enhanced inflammatory phenotype of MΦres persisted even when PD fluid administration was stopped. Thus, repeated exposure to PD fluid may render patients more susceptible to peritonitis, and by extension to peritoneal fibrosis, due to exaggerated inflammatory responses of MΦres.

Results:

Dialysis fluid induces the disappearance of tissue resident macrophages

To determine the impact of peritoneal dialysis fluid (PD fluid) injection on peritoneal macrophage populations (see suppl. fig 1 for gating strategy) we first characterised the cellular dynamics induced after a single application of PD fluid. Six hours after PD fluid injection, tissue resident macrophages (MΦres) underwent a pronounced disappearance reaction with the number of F4/80 high MΦres reduced to approximately 30% of the levels found in naive control animals (Fig 1A). Simultaneously, a significant influx of neutrophils and Ly6C high monocytes were detected, indicative of an inflammatory response (Fig 1B&C). Of note, similar to previous reports (Accarias et al., 2016) tissue dwelling, monocyte-derived MΦmono (F4/80 low MHC-II high) also underwent a disappearance reaction and were reduced in numbers by approximately 40 % (Fig 1D). F4/80 high MΦres displayed limited signs of activation at this early time point post PD fluid injection. No increase in expression of MHC-II could be detected (Fig 1E). In contrast intracellular Ym1 and CD206 expression were significantly enhanced following PD fluid injection, but expression was restricted to less than 10 % of cells (Fig 1F&G). The disappearance of MΦres in this context seemed in part due to enhanced cell death as indicated by increased levels of Annexin V staining (Fig 1H). By 24 hours post PD fluid injection, the numbers of MΦres had returned to baseline levels and expression of CD206 was no longer significantly different, whereas Ym1 expression remained elevated compared to naïve controls (Fig S2).

Overall, these data are consistent with the induction of low grade inflammation caused by PD fluid injection accompanied by a significant change in the prevalence of various myeloid cell populations within the peritoneal cavity.

Figure 1: **Injection of dialysis fluid alters peritoneal myeloid cell composition**

C57BL/6 mice were injected with Physioneal (orange circles) or left untreated (purple squares) and whole PEC isolated 6 h later. Cells were counted and analysed by flow cytometry to determine the number of A) MΦres (lineage-CD11b+, F4/80high,MHC-II low), B) neutrophils (lineage +, SSC mid, MHC-II-,F4/80-,CD11b+), C) monocytes (lineage-CD11b+, F4/80-,MHC-II low, Ly6C high) and D) monocyte-derived macrophages (lineage-CD11b+, F4/80low,MHC-II high). E) Histogram and quantitative data of fluorescence mean intensity of MHC-II expression by MΦmono (orange, dashed line) or MΦres from Physioneal injected (orange, solid line) or naive animals (purple, solid line). F-H) Expression of CD206, Ym1 and binding of Annexin V by MΦres identified in A) assessed by flow cytometry. Datapoints depict individual animals and bars indicate mean and SD. Data pooled from 4 independent experiments using 3-5 animals per group. Data analysed using a Mann-Whitney-U test after transformation. n.s.: not significant; *: p<0.05; **: p<0.01; ****: p<0.0001

lineage: TCRβ,CD19,Siglec-F, Ly6G

Repeated PD fluid treatment induces a gradual change in MΦres phenotype

We next sought to determine the long term effect of PD fluid administration on peritoneal macrophage populations. For this animals were injected once per day, 5 times a week with PD fluid i.p. and the peritoneal exudate cells (PEC) analysed 24 hours after 1, 4, 9 and 14 injections, respectively (Fig 2A). Over the course of the experiment the number of MΦres remained consistently lower in Physioneal treated animals as compared to naive mice (Fig 2B). Simultaneously, enhanced influx of Ly6C monocytes and successive accumulation of F4/80 low MΦmono could be detected in the peritoneal cavity following multiple rounds of PD fluid injection (Fig 2C& D).

Figure 2: Repeated injection of PD fluid leads to a progressive change in MPres phenotype

C57BL/6 mice were injected with PD fluid (orange circle) or left untreated (purple square) 5 times a week for the indicated number of injections. At each timepoint whole PEC were isolated 24 h after the last injection and analysed by flow cytometry. A) Schematic depiction of experimental timeline. B-D) Number of MΦres, MΦmono and Ly6C high monocytes. E-H) Expression of MΦres associated cell-surface markers on F4/80 high macrophages. I) Intracellular expression of Gata6 in MΦres identified in A).

Datapoints depict individual animals and lines indicate (B-H) median or (I) mean with SD. Data from a single experiment using 3 animals per group per timepoint. Data analysed using 2-way ANOVA followed by Tukey's post-hoc test after transformation. n.s.: not significant; *: $p < 0.05$; **: $p < 0.01$; ****: $p < 0.0001$

Despite the overall reduced numbers of MΦres, repeated injection of PD fluid led to marked induction of Ki67 expression in MΦres, indicative of proliferative expansion (Fig. S3). This is in line with a previous report demonstrating repopulation of the peritoneal cavity by MΦres following an inflammatory insult through proliferation (Davies et al., 2011). Thus, it is likely that PD fluid injection leads to a repeated cycle of MΦres disappearance followed by re-population. Similarly, it is likely that the gradual influx of monocytes and MΦmono underlies cyclic fluctuations after each round of PD fluid injection. Thus, this data indicated that macrophage populations within the peritoneal cavity undergo dynamic but limited changes following PD fluid instillation, with a slight accumulation of MΦmono over time.

Closer investigation of the MΦres phenotype, however, revealed a progressive loss of characteristics that define the tissue resident phenotype. In particular markers associated with the efferocytic and anti-inflammatory function of MΦres were expressed at increasingly lower levels following repeated injection of PD fluid (Fig 2E - G). T cell

immunoglobulin and mucin domain containing 4 (Tim4), a molecule associated with the

efficient removal of apoptotic cells (Miyanishi, Segawa, & Nagata, 2012; Miyanishi et al.,

2007) as well as V-set and immunoglobulin domain–containing 4 (Vsig4), associated with

limiting inflammatory responses (J. Li et al., 2017; Yuan, Yang, Dong, Yamamura, & Fu,

2017)], are specifically expressed in MΦres during steady state conditions (Heng, Painter,

Immunological Genome Project Consortium, 2008). MΦres gradually lost expression of

these markers with increasing number of PD fluid injections (Fig 2 E & G). Similarly, CD73,

an anti-inflammatory effector molecule specifically expressed by MΦres (Haskó et al.,

2011; Rosas et al., 2014) was found to be significantly reduced after 14 injections of PD

fluid (Fig 2 F).

In contrast, expression of CD102 (Icam2) was not altered by PD fluid injection (Fig 2 H). A

similar phenotype (loss of MΦres marker expression with sustained CD102 expression)

has previously been described in mice lacking the transcription factor Gata6, indicating

that repeated PD fluid injection may exert its effects via affecting Gata6 expression

(Okabe & Medzhitov, 2014). However, no significant difference in Gata6 expression by

MΦres was found after 9 injections of PD fluid measured in subsequent experiments (Fig

2 I). Notably, Gata6 expression levels varied following PD fluid injection with diminished

expression detected in approximately 50 % of animals (Fig 2 I). Thus, any effect of PD

fluid injection on Gata6 expression may be transient or reflect the impact of other factors,

like the degree of inflammation.

Taken together, following repeated PD fluid injection, MΦres, while retaining their tissue

identity (F4/80 high CD102+ Gata6+), lose some of their functional characteristics and in

particular anti-inflammatory and efferocytosis associated functions.

Repeated PBS injection but not chronic bacterial infection leads to a loss of MΦres

characteristics

As shown in Figure 1 and previously reported (Q. Li et al., 2018), repeated injection of PD

fluid is associated with the induction of low grade inflammation as evidenced by influx of

neutrophils and Ly6C+ monocytes. Thus, to assess whether the loss of MΦres functional

Figure 3: Chronic inflammation induced by S.Typhimurium infection does not phenocopy repeated PD fluid injection

C57BL/6 mice were infected orally with 1x10^8 cfu S.Typhimurium (orange squares) or left untreated (blue circles) and peritoneal cells analysed 34 (d34) and 55 days (d55) post infection. Dotplot (d55) and quantification of MHC-II (A) and Sca-1 (B) expression by MΦres assessed by flow cytometry C) Number of peritoneal neutrophils. D) Enumeration of bacterial colony forming units present in the peritoneal cavity. E-G) Expression of MΦres associated cell surface markers assessed by flow cytometry.

Datapoints depict individual animals and bars / lines indicate mean with SD. Data from a single experiment using 3-4 animals per group per timepoint. Data analysed using 2-way ANOVA followed by Tukey's post-hoc test after transformation. n.s.: not significant; *: p<0.05; **: p<0.01; ***: p<0.001

markers (i.e. Tim4, CD73, Vsig4) was due to the elicited inflammatory response, we analysed peritoneal exudate cells from animals subjected to prolonged bacterial infection. Animals were infected orally with attenuated *Salmonella enterica* ser. Typhimurium (SL3261, ΔaroA) and peritoneal cells analysed during the chronic phase of infection, 34 and 55 days post infection. In line with previous data (Ruckerl et al., 2017) and confirming the inflammatory environment, MΦres from SL3261 infected animals showed clear upregulation of MHC-II as well as enhanced expression of Sca-1 on day 55 (Fig 3A & B). Moreover, persistent influx of neutrophils was detected in the peritoneal cavity of SL3261 infected animals (Fig 3C) as well as live, cfu-forming bacteria (Fig 3D) confirming an ongoing pro-inflammatory activation. However, unlike following repeated PD fluid injection, MΦres isolated from animals harbouring *S. Typhimurium* did not show any noticeable loss of Tim4, CD73 or Vsig4 expression in the chronic phase of the infection (Fig 3 E-G). Rather to the contrary MΦres

from SL3261 infected animals expressed elevated levels of Tim4 (Fig 3 E). Thus, chronic

inflammatory conditions alone did not cause the progressive loss of MΦres phenotype as

observed following repeated Physioneal injection.

Importantly, injection of sterile PBS instead of PD fluid induced similar, albeit less

pronounced changes in peritoneal macrophage phenotype (Fig 4A). This would indicate

that repeated disturbance of the peritoneal immune system alone, rather than specific

constituents of the dialysis fluid, was sufficient to drive the observed alterations in

peritoneal MΦres. However, PD fluid enhanced these effects (Fig 4A).

Figure 4: Repeated injection of sterile saline induces similar alterations to dialysis fluid

A) Female C57BL/6 mice were injected 5 times a week for a total of nine injections with Physioneal (orange

circles), sterile PBS (green triangles) or left untreated (purple squares). 24 h after the last injection whole

PEC were isolated and analysed by flow cytometry for expression of MΦres associated cell-surface

markers. Data pooled from two independent experiments.

B) Male (♂) and female (♀) C57BL/6 mice were injected 5 times a week for a total of nine injections with PD

fluid (orange symbols) or left untreated (purple symbols). 24 h after the last injection whole PEC were

isolated and analysed as described for A).

Datapoints depict individual animals and lines indicate mean and SD. Data from a single experiment.

Data analysed using 2-way ANOVA followed by Tukey's multiple comparison test after transformation. n.s.:

not significant; *: $p<0.05$; **: $p<0.01$; ****: $p<0.0001$

Furthermore, it has been suggested that male peritoneal dialysis patients have a

significantly reduced survival rate on PD than female patients (Kitterer, Segerer, Braun,

Alscher, & Latus, 2017). Although the reason for this discrepancy is unknown and likely

due to multiple factors, differences in the inflammatory response may contribute to the observed effects. Moreover, male and female mice differ considerably in the maintenance and cellular dynamics of peritoneal macrophages (Bain et al., 2016). Thus, we assessed the effect of repeated PD fluid injection on myeloid cell populations in male and female mice. MΦres from naive male or female mice showed comparable expression levels of CD73 and Vsig4. In contrast naive male mice showed considerably lower expression of Tim4 (Figure 4B-D). However, independent of these differences in the steady state, Physioneal induced the loss of Tim4, CD73 and Vsig4 in both sexes to a similar degree. Taken together this data shows that repeated disturbance of the peritoneal environment triggers low grade inflammation which alters the MΦres phenotype.

Prolonged PD fluid injection alters the response of MΦres to external stimuli

Our previous data had highlighted a significant loss of Tim4 as well as Vsig4 and CD73, markers which have been associated with MΦres core functions (i.e. efferocytosis and anti-inflammatory activity) (Lopez-Castejón, Baroja-Mazo, & Pelegrin, 2011; Miyanishi et al., 2007; Yuan et al., 2017). Thus, we next aimed to assess whether repeated injection of dialysis fluid altered macrophage functional responses.

To test the capacity of MΦres to take up and ingest apoptotic cells, whole PEC from animals injected for various times with Physioneal or from naive controls were incubated in the presence of apoptotic thymocytes labelled with pHrodo. pHrodo labelled cells emit a very low, nearly undetectable fluorescent signal after staining, but will become brightly fluorescent and clearly detectable by flow cytometry after encountering an acidic environment, as found inside a phagolysosome (Miksa, Komura, Wu, Shah, & Wang, 2009). Thus, use of pHrodo allows the reliable detection of ingested apoptotic cells as compared to labeled cells bound to the surface of a phagocyte.

Repeated injection of Physioneal gradually reduced the proportion of myeloid cells (total CD11b+lin-) capable of ingesting apoptotic cells (Fig 5A). This was in part due to the increased proportion of Ly6C high monocytes and F4/80 low, monocyte-derived macrophages within the myeloid cell pool (Fig 2C&D), cells which possess reduced efferocytic activity (Gautier, Ivanov, Lesnik, & Randolph, 2013). However, when the analysis was restricted to MΦres a similar progressively reduced capacity to ingest apoptotic cells was detected (Fig 5B). Thus, MΦres from Physioneal injected animals

Figure 5: MΦres from PD fluid-injected animals show progressively enhanced responses to stimulation

C57BL/6 mice were injected with PD fluid (orange circle) or left untreated (purple square) 5 times a week for the indicated number of injections. At each timepoint whole PEC were isolated 24 h after the last injection and incubated in vitro in the presence of pHrodo labelled apoptotic thymocytes for 90 minutes (A & B) or stimulated with LPS/IFNγ (C & D; 6 h) or recombinant IL-4 (E & F; 24 h).

Uptake of apoptotic cells by total CD11b+ myeloid cells (A) or MΦres (CD102+I-A/I-E low) (B) as well as expression of MΦres associated cell-surface markers (C-F) assessed by flow cytometry.

Datapoints depict individual animals and lines indicate mean and SD. Data from a single experiment. Data analysed using 2-way ANOVA followed by Tukey's post-hoc test after transformation. n.s.: not significant; *: $p < 0.05$; **: $p < 0.01$; ****: $p < 0.0001$

seemed to lose the functionality to carry out efferocytosis efficiently. Of note, this loss of efferocytic capacity was not restricted to the use of dialysis fluid, as repeated injection of sterile PBS induced a similar reduction in the capacity of MΦres to ingest apoptotic cells (Fig S4A).

To further analyse whether MΦres from Physioneal injected animals were in general less responsive to external stimuli, we subjected whole PEC to in vitro stimulation with LPS and rIFNγ. MΦres from PD fluid treated animals showed a gradually increasing inflammatory response towards bacterial stimulation (Fig 5C&D). The proportion of cells

expressing NOS2 or Sca-1, markers of pro-inflammatory, M1 activation (Menzies, Henriquez, Alexander, & Roberts, 2010; S. Zhang, Kim, Batra, McKerrow, & Loke, 2010), was consistently higher in cells from PD fluid treated animals as compared to cells from naive animals (Fig 5C&D). Indeed, Sca-1 was not found to be upregulated on naive MΦres after 6h stimulation with IFNγ/LPS in vitro, indicating repeated Physioneal injections resulted in a stronger and more rapid response of MΦres to pro-inflammatory stimuli.

Importantly, an enhanced activation profile was not only observed in response to pro-inflammatory stimuli, but also in response to rIL-4, a driver of M2, anti-inflammatory macrophage activation (Ruckerl & Cook, 2019). MΦres stimulated with rIL-4 for 24h showed increased expression of Ym1 and at a later timepoint also Relmα (Fig 5E&F).

Thus, MΦres altered their functional repertoire following repeated exposure to dialysis fluid with a reduced efferocytic capacity and enhanced responses to both M1 and M2 polarising stimuli.

Altered PD fluid-induced responsiveness of MΦres is maintained after treatment is discontinued

Next we examined whether the phenotypical and functional changes observed in MΦres following repeated PD fluid injection were temporary or persisted even after treatment ceased. For this mice were injected 5 times a week with Physioneal or PBS for a total of 9 injections, a timepoint when the altered MΦres phenotype is evident (Fig 2), and then rested for seven days. Subsequently, the PEC were collected and analysed for the expression of activation markers as well as re-stimulated in vitro to analyse their response to external stimuli as described above. MΦres from discontinued Physioneal or PBS injected animals maintained a reduced expression of CD73, Tim4 and Vsig4 compared to naive controls (Fig 6A). Similarly, MΦres from Physioneal and PBS injected animals showed an enhanced response to in vitro stimulation with LPS/IFNγ (NOS2, Sca1; stimulated for 6 h) (Fig 6B). These data highlight that repeated PD fluid injection sensitise peritoneal macrophages and, thus, potentially exacerbates the severity and the detrimental sequelae of peritonitis events.

In order to verify that repeated exposure to PD fluid alters the response of peritoneal macrophages to inflammatory stimuli in vivo we utilised a model of sterile inflammation.

Figure 6: PD fluid conditioned MPres maintain altered activation phenotype after treatment is discontinued and show enhanced inflammatory responses in vivo

(A) C57BL/6 mice were injected 5 times a week for a total of nine injections with Physioneal (orange circles), sterile PBS (green triangles) or left untreated (purple squares). 7 days after the last injection whole PEC were isolated and analysed by flow cytometry for expression of MΦres associated cell-surface markers.

B&C) The cells isolated in A) were subjected to in vitro stimulation with LPS/IFNγ (6 h, NOS2 & Sca1) and analysed by flow cytometry.

D) Schematic depiction of the experimental timeline for experiments with intermittent induction of inflammation.

E-G) Animals were injected 3 times a week for a total of nine injections with Physioneal (orange circles) or left untreated (purple squares). Some Physioneal injected animals received after the third dose an additional single i.p. injection of sterile 4 % thioglycollate (black & orange circles, shaded bars) to induce inflammation. 24 h after the last injection whole PEC were isolated and analysed by flow cytometry.

Datapoints depict individual animals and lines indicate mean & SD. (A - C) Data pooled from 2 independent experiments (except C from a single experiment). Data analysed using 2-way ANOVA followed by Tukey's multiple comparison test after transformation. (E-G) Data from a single experiment analysed by 1-way ANOVA.

n.s.: not significant; *: p<0.05; **: p<0.01; ***: p<0.001; ****: p<0.0001

Animals were subjected to repeated PD fluid injection (3 times per week for a total of 9 injections) and an inflammatory response was induced with a single dose of sterile 4 % Brewer's modified thioglycollate medium at day 6, mimicking an intermittent event of

peritonitis (see Fig 6C). Twenty-four hours after the last PD fluid injection MΦres were collected and analysed for the expression of Tim4, CD206 and Ym1. As expected, repeated injection of dialysis fluid led to a significant loss of Tim4 expression on MΦres with limited induction of CD206 or Ym1 expression (Fig 6D-F). Additionally, in line with our in vitro findings (Fig 5), intermittent induction of inflammation did not further alter MΦres phenotype as determined by Tim4 expression, but led to enhanced activation of MΦres adopting a pro-fibrosis associated phenotype (Fig 6D-F). Importantly, expression of CD206, which has been closely linked to induction of fibrosis (Bellón et al., 2011; Chandrasekaran et al., 2019; Jie Wang et al., 2013), was thought to not be expressed by MΦres (Gundra et al., 2014).

Overall these findings indicate that prolonged and repeated exposure to PD fluid renders peritoneal macrophages more responsive to stimulation potentially exacerbating detrimental inflammatory responses.

Discussion:

Macrophages are versatile cells implicated in many pathologies (Sica, Erreni, Allavena, & Porta, 2015). They play both essential protective as well as detrimental / pathological roles, often within the same disease setting (Guillot & Tacke, 2019; Ribatti, 2013; Wynn & Vannella, 2016). Thus, the regulatory mechanisms governing macrophage responses have become the focus of current research aiming to dissect these contradictory behaviours. Moreover, scientists have proposed macrophages as an excellent target for therapeutic interventions, as altering their phenotype may not only improve disease outcome, but actively revert associated pathologies.

Recent discoveries have highlighted the inherent diversity of the macrophage pool, indicating the presence of different types of macrophages with different ontogeny (Bain & Jenkins, 2018), response profiles (Gundra et al., 2014) and functional roles (Ruckerl et al., 2017). In particular, MΦres and acutely recruited MΦmono have been identified as distinct mediators of pathology. During lung fibrosis recruited MΦmono have been shown to be essential drivers of the fibrotic pathology and disease progression (Misharin et al., 2017). Similarly, in murine models of peritoneal dialysis associated fibrosis, infiltration of MΦmono has been shown to be detrimental to disease outcome (Y.-T. Chen, Hsu, Lin, Pan, Liu, Wu, Tsai, Liao, Cheng, Chiang, Chen, Chu, & Lin, 2019b; Lee et al., 2012; Q. Li

et al., 2018; Jie Wang et al., 2013). Furthermore, significant changes in the prevalence of

specific macrophage/monocyte populations have been described in dialysis patients,

dependent on the history of peritonitis episodes (Liao et al., 2017). Thus, the pathology

inducing effect of peritonitis associated inflammation in peritoneal dialysis seem to be due

to the recruitment and accumulation of fibrosis promoting cells like MΦmono. Due to

legal restrictions on the maximum number of i.p. injections allowed, we did not observe

any clinical signs of fibrosis in our experimental setup. However, similar to the findings

above, we observed low-grade inflammation and infiltration of MΦmono in our system. In

addition, we demonstrated a drastic change in phenotype of MΦres. Importantly this

change in phenotype translated into an enhanced response profile boosting both pro-

inflammatory as well as M2 associated effector molecule production. Thus, the simple

dichotomy of *good* and *bad* macrophages cannot be upheld in these settings. The

increased risk of developing pathological sequelae following repeated episodes of

peritonitis in PD patients is likely due to the damaging effect of the inflammatory response

(Williams et al., 2002). Our data implies, that patients on longterm PD treatment are at an

increased risk to develop clinical symptoms of peritonitis, not only because of an

increased risk of exposure to bacterial antigens, but also due to an enhanced

inflammatory response of peritoneal resident cells. Whereas in healthy people small

numbers of infiltrating pathogens are silently removed by MΦres without triggering an

inflammatory response, PD fluid conditioned MΦres may have lost some of this regulatory

capacity leading to exacerbated pathology-promoting inflammation. Thus, targeting

MΦres to restore some of their homeostatic, anti-inflammatory capabilities may provide a

feasible approach to extend the therapeutic window of PD. Indeed, adding soluble factors

(e.g. soluble TLR2) to the dialysis fluid targeting the inflammatory response has previously

been suggested (Raby & Labéta, 2018) and could be adapted to specifically target

MΦres.

Intriguingly in PD patients receiving oral supplementation with vitamin D, a factor involved

in the expansion of peritoneal MΦres in mice (Song et al., 2016), antibacterial responses

were enhanced (Bacchetta et al., 2014), potentially through improved phagocytosis of

invading bacteria (Lishko, Moreno, Podolnikova, & Ugarova, 2016). MΦres in virtually all

tissues of the body are seeded during embryonic development and maintained locally

through proliferative expansion. However, in many tissues, including the peritoneal cavity,

these embryonic derived macrophages are, over time, replaced by monocyte-derived

cells adopting features of tissue resident cells (Bain et al., 2016). Thus, it may be feasible to develop therapeutic approaches fostering the non-inflammatory removal of bacteria by promoting MΦres differentiation. However, the conversion of MΦmono in the peritoneal cavity under steady state seems inefficient and leads to a gradual accumulation of Gata6+, F4/80 high MΦres with low expression of Tim4 (Bain et al., 2016). Our data from *Salmonella enterica* ser. Typhimurium infected mice indicated that the late stages of infection were linked to an increase in Tim4 expressing MΦres as compared to naive animals. Similarly, previously published results from helminth infected animals indicate enhanced expression of Tim4 on MΦres during the chronic phase of these infection (Campbell et al., 2018). Hence, factors associated with the resolution of inflammation or chronic Th2 immune responses may allow to 'rejuvenate' monocyte-derived MΦres or promote the conversion of MΦmono. Further research is required to determine the specific factors driving the MΦres phenotype as well as their potential use as therapeutic adjunct during PD.

In this context it is important to consider that peritonitis or PD induced inflammation will drive the disappearance of MΦres and, thus, may undo any previous accumulation / conversion attempts. It has previously been suggested that the loss of MΦres is at least in part due to enhanced cell death of resident cells during inflammation (Ruckerl et al., 2017). Importantly, the degree of loss of MΦres is directly correlated to the degree of inflammation as well as the number of MΦres present within the peritoneal cavity (Ruckerl et al., 2017). Thus, targeted expansion of MΦres prior to or during PD fluid treatment may provide an option to retain enhanced numbers of MΦres during inflammatory events. In contrast, recent work suggests that the apparent reduction in the number of MΦres detectable in the peritoneal wash is due to the adherence of these cells to the peritoneum rather than an actual loss in numbers (N. Zhang et al., 2019). This is especially noteworthy in the context of peritoneal dialysis as it would bring these cells in close contact to the main site of pathology. Moreover, targeted disruption of the coagulation cascade reduced the apparent loss of MΦres and prevented their adherence (N. Zhang et al., 2019) potentially opening a window for therapeutic intervention. Currently neither approach (increasing MΦres numbers or preventing the MDR) is likely suitable for use in human patients due to systemic and off-target effects of the applied methodology. However, both approaches indicate intriguing opportunities to develop novel techniques specifically targeting peritoneal MΦres.

Independent of the therapeutic potential discussed above, analysing the immune cells

contained in the effluent of PD patients may yield a useful biomarker strategy. Changes in

the composition of myeloid cells as described by Liao et al. (Liao et al., 2017) as well as

the assessment of cellular activation markers as described in mice here, may allow risk-

based patient stratification. Patients likely to develop pathological sequelae of PD

treatment could then be prioritised for kidney transplant or transferred to haemodialysis

prior to PD failure or before overt pathological changes occur.

Lastly, our data comparing PBS injected animals to Physioneal 40 injection indicated a

limited impact of the composition of the PD fluid on the observed physiological changes.

This seems in contrast to the commonly held believe that the constituents of PD fluid, in

particular lactate, glucose-degradation-products (GDP) and low pH, are instrumental in

driving PD related pathology (Krishnan, Tam, Wu, Breborowicz, & Oreopoulos, 2005;

Mortier, Faict, Lameire, & De Vriese, 2005). However, our data is in line with a recent

report investigating the use of so-called biocompatible PD fluid in patients. Despite

considerably lower levels of GDP and lactate in these solutions, the morphological

changes observed were very similar to those seen in patients utilising standard PD fluid

and even an early increase in microvascular density was detected (Schaefer et al., 2018).

Of note, it has been suggested that this adverse effect of biocompatible PD fluid may be

due to an altered inflammatory response (Blake, 2018). Thus, together with our data this

suggests it may be more effective to prevent pathological changes by targeting the

elicited immune response to PD fluid instillation, even in the absence of peritonitis.

Taken together we have shown here that repeated exposure of the peritoneal

microenvironment to PD fluid instillation in mice led to a gradual change in phenotype as

well as activation response of peritoneal resident cells. These changes were relatively

long lasting and resulted in a more vigorous response to inflammatory triggers. Thus,

targeting MΦres and preventing excessive inflammatory responses in PD patients may

pose an exciting novel approach to limit PD-related pathology.

Materials and Methods:

Ethics Statement

All animal experiments were performed in accordance with the UK Animals (Scientific Procedures) Act of 1986 under a Project License (70/8548) granted by the UK Home Office and approved by the University of Manchester Ethical Review Committee.

Mice and in vivo treatments

Eight to 13 week old male and female C57BL/6 mice were obtained from a commercial vendor (Envigo, Hillcrest, UK) and maintained in groups of 4-6 animals in specific pathogen–free facilities at the University of Manchester. Experimental mice were randomly allocated to treatment groups using a computer-based randomization technique and age and sex matched. Euthanasia was performed by asphyxiation in carbon dioxide in a rising concentration. Animal numbers were based on initial power calculations of key determinants derived from preliminary experiments.

Animals were injected i.p. with 500 µL peritoneal dialysis fluid (Physioneal 40, 3.86% glucose, Baxter HealthCare Ltd., Compton, UK) for the indicated number of injections. Injections were carried out daily or every other day for three or five days per week for up to 4 weeks (maximum 14 injections per animal). Control animals were left untreated or received an equal volume of PBS i.p. as indicated.

For the induction of inflammatory responses PD fluid treated mice received intra-peritoneal injections of 400 µL 4% Brewer modified thioglycollate medium (BD Biosciences, San Jose, CA) or sterile saline as control. The attenuated *Salmonella enterica* serovar Typhimurium strain SL3261 (ΔaroA) (Hoiseth & Stocker, 1981) was cultured overnight at 37^0C in a shaking incubator from frozen stock in Luria-Bertani broth with 50 µg/mL streptomycin. The following morning, culture was diluted in fresh Luria-Bertani broth with 50 µg/mL streptomycin and incubated at 37^0C in a shaking incubator to ensure the bacteria were in the growth phase. CFU/mL was estimated by the OD600 reading. Animals were pre-treated with 20 mg streptomycin 1 day prior to oral infection with ~1×10^8 CFU *Salmonella* Typhimurium diluted in PBS. Infectious doses and peritoneal bacterial burdens were enumerated by plating inocula or peritoneal exudate cells in 10-fold serial dilutions in PBS on LB-Agar plates.

Cell-isolation

Peritoneal cavity exudate cells (PEC) were obtained by washing the cavity with 10 mL lavage media comprised of RPMI 1640 (Sigma-Aldrich, Dorset, UK) containing 2 mM EDTA and 1% L-Glutamine (Thermo Fisher Scientific, Waltham, MA). Erythrocytes were removed by incubating with red blood cell lysis buffer (Sigma-Aldrich). Cellular content

was assessed by cell counting using Viastain AO/PI solution on a Cellometer ® Auto 2000 Cell Counter (Nexcelom Bioscience, Manchester, UK) in combination with multicolor flow cytometry.

Flow cytometry

Equal numbers of cells were stained with Zombie UV viability assay (Biolegend, London, UK). All samples were then blocked with 5 µg/mL anti CD16/32 (93; Biolegend) and heat-inactivated normal mouse serum (1:10, Sigma-Aldrich) in flow cytometry buffer (0.5% BSA and 2 mM EDTA in Dulbecco's PBS) before surface staining on ice with antibodies to F4/80 (BM8), SiglecF (E502440, BD Biosciences), Ly6C (HK1.4), Ly-6G (1A8), TCRβ (H57-597), CD11b (M1/70), CD11c (N418), I-A/I-E (M5/114.15.2), CD19 (6D5), CD115 (AFS98), CD73 (TY/11.8), CD102 (3C4 (MIC2/4)), Tim4 (RMT4-54), CD206 (MR6F3), Vsig4 (NLA14; Thermo Fisher Scientific), CD226 (10E5), Sca1/Ly6A (D7). All antibodies were purchased from Biolegend unless stated otherwise.

Detection of intracellular activation markers was performed directly ex vivo. Cells were stained for surface markers then fixed and permeabilized for at least 16 h using the eBioscience™ Foxp3 / Transcription Factor Staining Buffer Set (Thermo Fisher Scientific). Cells were then stained with directly labeled Abs to NOS2 (CXNFT; Thermo Fisher Scientific), Arg1 (polyclonal, R&D Systems, Abingdon, UK), Ki67 (B56, BD Biosciences), Annexin V (Biolegend), Gata-6 (D61E4; Cell Signaling Technology Europe, Leiden, Netherlands) or purified polyclonal rabbit antiRelmα (PeproTech EC Ltd., London, UK) and biotinylated anti-Ym1/2 (polyclonal, R&D Systems) followed by Zenon anti–rabbit reagent (Thermo Fisher Scientific) or streptavidin BUV 737 (BD Biosciences), respectively. Samples were acquired on a BD LSR II or BD FACSymphony using BD FACSDiva software (BD Biosciences) and post-acquisition analysis performed using FlowJo v10 software (BD Biosciences).

Cell-culture experiments

For in vitro stimulation of PD fluid-conditioned cells, whole PEC were counted as described above and seeded to 96-well U bottom plates at 3×10^5 cells per well in RPMI 1640 containing 5 % foetal bovine serum, 2 mM L-glutamine, 100 U/mL penicillin and 100 µg/mL streptomycin and stimulated with lipopolysaccharide (LPS, 100 ng/mL; *Salmonella enterica* ser. Typhimurium; Sigma-Aldrich) and recombinant murine Interferon γ (IFNγ, 20 ng/mL PeproTech EC Ltd.) or medium alone for 6 h or with murine recombinant Interleukin–4 (rIL-4, 20 ng/mL, PeproTech EC Ltd.) for 24 h and analysed for MΦ activation markers by flow cytometry.

Apoptotic cell uptake assay

Uptake of apoptotic cells by peritoneal MΦres was assessed as previously described (Miksa et al., 2009). Briefly, thymocytes were collected from naive animals by mincing thymi through 2 µm gauze until completely homogenized. Erythrocytes were removed by incubating with red blood cell lysis buffer (Sigma-Aldrich). Thymocytes were resuspended at 1×10^7 cells/mL in complete DMEM and incubated in the presence of 0.1 µM dexamethasone (Sigma-Aldrich) at 37 °C for 18 h. This produced >90 % apoptosis, as assessed by Viastain AO/PI staining measured on a Cellometer ® Auto 2000 Cell Counter (Nexcelom Bioscience). Subsequently, apoptotic thymocytes were washed twice with PBS and resuspended in PBS at 10^6 cells/mL containing 40 ng/mL pHrodo-SE (Thermo Fisher Scientific) and incubated at RT for 30 min. Thereafter the cells were washed twice with PBS and resuspended in RPMI containing 5 % foetal bovine serum, 2 mM L-glutamine, 100 U/mL penicillin and 100 µg/mL streptomycin. Unstained apoptotic cells served as staining control.

Statistical analysis

Statistical analysis was performed using Prism 8 for Mac OS X (v8.2.1, GraphPad Software, San Diego, CA). Differences between groups were determined by t-test or ANOVA followed by Tukey's or Dunn's multiple comparison-test. In some cases data was log-transformed to achieve normal distribution as determined by optical examination of residuals. Where this was not possible a Mann-Whitney or Kruskal-Wallis test was used. Percentages were subjected to arcsine transformation prior to analysis. Differences were assumed statistically significant for *P* values of less than 0.05.

Author contributions
- Tara E Sutherland (Resources, Writing – Review & Editing, Funding Acquisition)
- Tovah N Shaw (Investigation, Writing – Review & Editing)
- Rachel Lennon (Resources, Writing – Review and Editing, Clinical advice)
- Sarah E Herrick (Conceptualization, Writing – Review and Editing)
- Dominik Ruckerl (Conceptualization, Formal Analysis, Validation, Investigation, Writing - Original Draft Preparation, Writing – Review and Editing, Visualization, Project Administration, Funding Acquisition)

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

This work was supported by the Medical Research Council UK (MR/P02615X/1, DR) and

the Medical Research Foundation / Asthma UK (MRFAUK-2015-302, TES). TNS was

supported by a BBSRC Discovery Research Fellowship (BB/S01103X/1, TNS), RL was

supported by a Wellcome Trust Senior Fellowship award (202860/Z/16/Z, RL) and SEH

and DR were supported by an MRC Research Grant (MRC MR/S02560X/1, SEH).

We thank Dr. Gareth Howell from the Flow Cytometry Core Facility at the University of

Manchester for assistance with the flow cytometry analyses. We'd also like to thank

Sister Trish Smith from the Royal Manchester Children's Hospital for the provision of

Physioneal 40.

References:

Accarias, S., Genthon, C., Rengel, D., Boullier, S., Foucras, G., & Tabouret, G. (2016). Single-cell
analysis reveals new subset markers of murine peritoneal macrophages and highlights
macrophage dynamics upon Staphylococcus aureusperitonitis. *Innate Immunity*, 22(5), 382–
392. http://doi.org/10.1177/1753425916651330
Allen, J. E., & Ruckerl, D. (2017). The Silent Undertakers: Macrophages Programmed for
Efferocytosis. *Immunity*, 47(5), 810–812. http://doi.org/10.1016/j.immuni.2017.10.010
Bacchetta, J., Chun, R. F., Gales, B., Zaritsky, J. J., Leroy, S., Wesseling-Perry, K., et al. (2014).
Antibacterial responses by peritoneal macrophages are enhanced following vitamin D
supplementation. *PLoS ONE*, 9(12), e116530. http://doi.org/10.1371/journal.pone.0116530
Bain, C. C., & Jenkins, S. J. (2018). The biology of serous cavity macrophages. *Cellular
Immunology*. http://doi.org/10.1016/j.cellimm.2018.01.003
Bain, C. C., Hawley, C. A., Garner, H., Scott, C. L., Schridde, A., Steers, N. J., et al. (2016). Long-
lived self-renewing bone marrow-derived macrophages displace embryo-derived cells to inhabit
adult serous cavities. *Nature Communications*, 7, ncomms11852.
http://doi.org/10.1038/ncomms11852
Barth, M. W., Hendrzak, J. A., Melnicoff, M. J., & Morahan, P. S. (1995). Review of the
macrophage disappearance reaction. *Journal of Leukocyte Biology*, 57(3), 361–367.
Bartosova, M., & Schmitt, C. P. (2019). Biocompatible Peritoneal Dialysis: The Target Is Still Way
Off. *Frontiers in Physiology*, 9, 2047–13. http://doi.org/10.3389/fphys.2018.01853
Bellón, T., Martínez, V., Lucendo, B., del Peso, G., Castro, M. J., Aroeira, L. S., et al. (2011).
Alternative activation of macrophages in human peritoneum: implications for peritoneal
fibrosis. *Nephrology, Dialysis, Transplantation : Official Publication of the European Dialysis
and Transplant Association - European Renal Association*, 26(9), 2995–3005.
http://doi.org/10.1093/ndt/gfq771
Blake, P. G. (2018). Is the peritoneal dialysis biocompatibility hypothesis dead? *Kidney
International*, 94(2), 246–248. http://doi.org/10.1016/j.kint.2018.04.014
Braga, T. T., Correa-Costa, M., Silva, R. C., Cruz, M. C., Hiyane, M. I., da Silva, J. S., et al. (2018).
CCR2 contributes to the recruitment of monocytes and leads to kidney inflammation and
fibrosis development. *Inflammopharmacology*, 26(2), 403–411. http://doi.org/10.1007/s10787-
017-0317-4
Campbell, S. M., Knipper, J. A., Ruckerl, D., Finlay, C. M., Logan, N., Minutti, C. M., et al.
(2018). Myeloid cell recruitment versus local proliferation differentiates susceptibility from
resistance to filarial infection. *eLife*, 7, 955. http://doi.org/10.7554/eLife.30947
Chandrasekaran, P., Izadjoo, S., Stimely, J., Palaniyandi, S., Zhu, X., Tafuri, W., & Mosser, D. M.
(2019). Regulatory Macrophages Inhibit Alternative Macrophage Activation and Attenuate

Pathology Associated with Fibrosis. *The Journal of Immunology*, *203*(8), 2130–2140. http://doi.org/10.4049/jimmunol.1900270

Chen, Y.-T., Hsu, H., Lin, C.-C., Pan, S.-Y., Liu, S.-Y., Wu, C.-F., Tsai, P.-Z., Liao, C.-T., Cheng, H.-T., Chiang, W.-C., Chen, Y.-M., Chu, T.-S., & Lin, S.-L. (2019a). Inflammatory macrophages switch to CCL17-expressing phenotype and promote peritoneal fibrosis. *The Journal of Pathology*, path.5350. http://doi.org/10.1002/path.5350

Chen, Y.-T., Hsu, H., Lin, C.-C., Pan, S.-Y., Liu, S.-Y., Wu, C.-F., Tsai, P.-Z., Liao, C.-T., Cheng, H.-T., Chiang, W.-C., Chen, Y.-M., Chu, T.-S., & Lin, S.-L. (2019b). Inflammatory macrophages switch to CCL17-expressing phenotype and promote peritoneal fibrosis. *The Journal of Pathology*, path.5350. http://doi.org/10.1002/path.5350

Chui, B. K., Manns, B., Pannu, N., Dong, J., Wiebe, N., Jindal, K., & Klarenbach, S. W. (2013). Health care costs of peritoneal dialysis technique failure and dialysis modality switching. *American Journal of Kidney Diseases : the Official Journal of the National Kidney Foundation*, *61*(1), 104–111. http://doi.org/10.1053/j.ajkd.2012.07.010

Danford, C. J., Lin, S. C., Smith, M. P., & Wolf, J. L. (2018). Encapsulating peritoneal sclerosis. *World Journal of Gastroenterology*, *24*(28), 3101–3111. http://doi.org/10.3748/wjg.v24.i28.3101

Davies, L. C., Rosas, M., Jenkins, S. J., Liao, C.-T., Scurr, M. J., Brombacher, F., et al. (2013). Distinct bone marrow-derived and tissue-resident macrophage lineages proliferate at key stages during inflammation. *Nature Communications*, *4*, 1886. http://doi.org/10.1038/ncomms2877

Davies, L. C., Rosas, M., Smith, P. J., Fraser, D. J., Jones, S. A., & Taylor, P. R. (2011). A quantifiable proliferative burst of tissue macrophages restores homeostatic macrophage populations after acute inflammation, *41*(8), 2155–2164. http://doi.org/10.1002/eji.201141817

Ferenbach, D. A., Sheldrake, T. A., Dhaliwal, K., Kipari, T. M. J., Marson, L. P., Kluth, D. C., & Hughes, J. (2012). Macrophage/monocyte depletion by clodronate, but not diphtheria toxin, improves renal ischemia/ reperfusion injury in mice. *Kidney International*, *82*(8), 928–933. http://doi.org/10.1038/ki.2012.207

François, K., & Bargman, J. M. (2014). Evaluating the benefits of home-based peritoneal dialysis. *International Journal of Nephrology and Renovascular Disease*, *7*, 447–455. http://doi.org/10.2147/IJNRD.S50527

Gautier, E. L., Ivanov, S., Lesnik, P., & Randolph, G. J. (2013). Local apoptosis mediates clearance of macrophages from resolving inflammation in mice. *Blood*, *122*(15), 2714–2722. http://doi.org/10.1182/blood-2013-01-478206

Gordon, S., & Plüddemann, A. (2017). Tissue macrophages: heterogeneity and functions. *BMC Biology*, *15*(1), 53. http://doi.org/10.1186/s12915-017-0392-4

Guillot, A., & Tacke, F. (2019). Liver Macrophages: Old Dogmas and New Insights. *Hepatology Communications*, *3*(6), 730–743. http://doi.org/10.1002/hep4.1356

Gundra, U. M., Girgis, N. M., Ruckerl, D., Jenkins, S., Ward, L. N., Kurtz, Z. D., et al. (2014). Alternatively activated macrophages derived from monocytes and tissue macrophages are phenotypically and functionally distinct. *Blood*, *123*(20), e110–22. http://doi.org/10.1182/blood-2013-08-520619

Han, Y., Ma, F. Y., Tesch, G. H., Manthey, C. L., & Nikolic-Paterson, D. J. (2013). Role of macrophages in the fibrotic phase of rat crescentic glomerulonephritis. *AJP: Renal Physiology*, *304*(8), F1043–53. http://doi.org/10.1152/ajprenal.00389.2012

Haskó, G., Csóka, B., Koscsó, B., Chandra, R., Pacher, P., Thompson, L. F., et al. (2011). Ecto-5'-nucleotidase (CD73) decreases mortality and organ injury in sepsis. *The Journal of Immunology*, *187*(8), 4256–4267. http://doi.org/10.4049/jimmunol.1003379

Heng, T. S. P., Painter, M. W., Immunological Genome Project Consortium. (2008). The Immunological Genome Project: networks of gene expression in immune cells. *Nature Immunology*, *9*(10), 1091–1094. http://doi.org/10.1038/ni1008-1091

Hoiseth, S. K., & Stocker, B. A. (1981). Aromatic-dependent Salmonella typhimurium are non-virulent and effective as live vaccines. *Nature*, *291*(5812), 238–239.

Johnson, D. W., Cho, Y., Livingston, B. E. R., Hawley, C. M., McDonald, S. P., Brown, F. G., et al.
(2010). Encapsulating peritoneal sclerosis: incidence, predictors, and outcomes. *Kidney*
International, *77*(10), 904–912. http://doi.org/10.1038/ki.2010.16
Jonjic, N., Peri, G., Bernasconi, S., Sciacca, F. L., Colotta, F., Pelicci, P., et al. (1992). Expression
of adhesion molecules and chemotactic cytokines in cultured human mesothelial cells. *The*
Journal of Experimental Medicine, *176*(4), 1165–1174. http://doi.org/10.1084/jem.176.4.1165
Kitterer, D., Segerer, S., Braun, N., Alscher, M. D., & Latus, J. (2017). Gender-Specific Differences
in Peritoneal Dialysis. *Kidney and Blood Pressure Research*, *42*(2), 276–283.
http://doi.org/10.1159/000477449
Kramer, A., Pippias, M., Noordzij, M., Stel, V. S., Andrusev, A. M., Aparicio-Madre, M. I., et al.
(2019). The European Renal Association - European Dialysis and Transplant Association
(ERA-EDTA) Registry Annual Report 2016: a summary. *Clinical Kidney Journal*, *12*(5), 702–
720. http://doi.org/10.1093/ckj/sfz011
Krishnan, M., Tam, P., Wu, G., Breborowicz, A., & Oreopoulos, D. G. (2005). Glucose degradation
products (GDP's) and peritoneal changes in patients on chronic peritoneal dialysis: will new
dialysis solutions prevent these changes? *International Urology and Nephrology*, *37*(2), 409–
418. http://doi.org/10.1007/s11255-004-1392-1
Lambie, M. R., Chess, J., Summers, A. M., Williams, P. F., Topley, N., & Davies, S. J. (2016).
Peritoneal inflammation precedes encapsulating peritoneal sclerosis: results from the GLOBAL
Fluid Study. *Nephrology, Dialysis, Transplantation : Official Publication of the European*
Dialysis and Transplant Association - European Renal Association, *31*(3), 480–486.
http://doi.org/10.1093/ndt/gfv440
Lee, S. H., Kang, H.-Y., Kim, K. S., Nam, B. Y., Paeng, J., Kim, S., et al. (2012). The monocyte
chemoattractant protein-1 (MCP-1)/CCR2 system is involved in peritoneal dialysis-related
epithelial-mesenchymal transition of peritoneal mesothelial cells. *Laboratory Investigation; a*
Journal of Technical Methods and Pathology, *92*(12), 1698–1711.
http://doi.org/10.1038/labinvest.2012.132
Li, J., Diao, B., Guo, S., Huang, X., Yang, C., Feng, Z., et al. (2017). VSIG4 inhibits
proinflammatory macrophage activation by reprogramming mitochondrial pyruvate
metabolism. *Nature Communications*, *8*(1), 1322. http://doi.org/10.1038/s41467-017-01327-4
Li, Q., Zheng, M., Liu, Y., Sun, W., Shi, J., Ni, J., & Wang, Q. (2018). A pathogenetic role for M1
macrophages in peritoneal dialysis-associated fibrosis. *Molecular Immunology*, *94*, 131–139.
http://doi.org/10.1016/j.molimm.2017.12.023
Liao, C.-T., Andrews, R., Wallace, L. E., Khan, M. W. A., Kift-Morgan, A., Topley, N., et al.
(2017). Peritoneal macrophage heterogeneity is associated with different peritoneal dialysis
outcomes. *Kidney International*, *91*(5), 1088–1103. http://doi.org/10.1016/j.kint.2016.10.030
Lishko, V. K., Moreno, B., Podolnikova, N. P., & Ugarova, T. P. (2016). Identification of Human
Cathelicidin Peptide LL-37 as a Ligand for Macrophage Integrin αMβ2 (Mac-1, CD11b/CD18)
that Promotes Phagocytosis by Opsonizing Bacteria. *Research and Reports in Biochemistry*,
2016(6), 39–55. http://doi.org/10.2147/rrbc.s107070
Lopez-Castejón, G., Baroja-Mazo, A., & Pelegrin, P. (2011). Novel macrophage polarization
model: from gene expression to identification of new anti-inflammatory molecules. *Cellular*
and Molecular Life Sciences : CMLS, *68*(18), 3095–3107. http://doi.org/10.1007/s00018-010-
0609-y
Luyckx, V. A., Tonelli, M., & Stanifer, J. W. (2018). The global burden of kidney disease and the
sustainable development goals. *Bulletin of the World Health Organization*, *96*(6), 414–422D.
http://doi.org/10.2471/BLT.17.206441
Menzies, F. M., Henriquez, F. L., Alexander, J., & Roberts, C. W. (2010). Sequential expression of
macrophage anti-microbial/inflammatory and wound healing markers following innate,
alternative and classical activation. *Clinical and Experimental Immunology*.
http://doi.org/10.1111/j.1365-2249.2009.04086.x
Miksa, M., Komura, H., Wu, R., Shah, K. G., & Wang, P. (2009). A novel method to determine the

engulfment of apoptotic cells by macrophages using pHrodo succinimidyl ester. *Journal of*
Immunological Methods, *342*(1-2), 71–77. http://doi.org/10.1016/j.jim.2008.11.019
Minutti, C. M., Jackson-Jones, L. H., García-Fojeda, B., Knipper, J. A., Sutherland, T. E., Logan,
N., et al. (2017). Local amplifiers of IL-4Rα–mediated macrophage activation promote repair in
lung and liver. *Science*, *356*(6342), 1076–1080. http://doi.org/10.1126/science.aaj2067
Misharin, A. V., Morales-Nebreda, L., Reyfman, P. A., Cuda, C. M., Walter, J. M., McQuattie-
Pimentel, A. C., et al. (2017). Monocyte-derived alveolar macrophages drive lung fibrosis and
persist in the lung over the life span. *Journal of Experimental Medicine*, *214*(8), 2387–2404.
http://doi.org/10.1084/jem.20162152
Miyanishi, M., Segawa, K., & Nagata, S. (2012). Synergistic effect of Tim4 and MFG-E8 null
mutations on the development of autoimmunity. *International Immunology*, *24*(9), 551–559.
http://doi.org/10.1093/intimm/dxs064
Miyanishi, M., Tada, K., Koike, M., Uchiyama, Y., Kitamura, T., & Nagata, S. (2007).
Identification of Tim4 as a phosphatidylserine receptor. *Nature*, *450*(7168), 435–439.
http://doi.org/10.1038/nature06307
Moinuddin, Z., Summers, A., Van Dellen, D., Augustine, T., & Herrick, S. E. (2014).
Encapsulating peritoneal sclerosis-a rare but devastating peritoneal disease. *Frontiers in*
Physiology, *5*, 470. http://doi.org/10.3389/fphys.2014.00470
Morelle, J., Sow, A., Hautem, N., Bouzin, C., Crott, R., Devuyst, O., & Goffin, E. (2015).
Interstitial Fibrosis Restricts Osmotic Water Transport in Encapsulating Peritoneal Sclerosis.
Journal of the American Society of Nephrology : JASN, *26*(10), 2521–2533.
http://doi.org/10.1681/ASN.2014090939
Moriishi, M., Kawanishi, H., Kawai, T., Takahashi, S., Hirai, T., Shishida, M., et al. (2002).
Preservation of peritoneal catheter for prevention of encapsulating peritoneal sclerosis.
Advances in Peritoneal Dialysis. Conference on Peritoneal Dialysis, *18*, 149–153.
Mortier, S., Faict, D., Lameire, N. H., & De Vriese, A. S. (2005). Benefits of switching from a
conventional to a low-GDP bicarbonate/lactate-buffered dialysis solution in a rat model. *Kidney*
International, *67*(4), 1559–1565. http://doi.org/10.1111/j.1523-1755.2005.00237.x
Nikolic-Paterson, D. J., Wang, S., & Lan, H. Y. (2015). Macrophages promote renal fibrosis
through direct and indirect mechanisms. *Kidney International Supplements*, *4*(1), 1–5.
http://doi.org/10.1038/kisup.2014.7
Okabe, Y., & Medzhitov, R. (2014). Tissue-specific signals control reversible program of
localization and functional polarization of macrophages. *Cell*, *157*(4), 832–844.
http://doi.org/10.1016/j.cell.2014.04.016
Puranik, A. S., Leaf, I. A., Jensen, M. A., Hedayat, A. F., Saad, A., Kim, K.-W., et al. (2018).
Kidney-resident macrophages promote a proangiogenic environment in the normal and
chronically ischemic mouse kidney. *Scientific Reports*, *8*(1), 13948.
http://doi.org/10.1038/s41598-018-31887-4
Raby, A.-C., & Labéta, M. O. (2018). Preventing Peritoneal Dialysis-Associated Fibrosis by
Therapeutic Blunting of Peritoneal Toll-Like Receptor Activity. *Frontiers in Physiology*, *9*,
1692. http://doi.org/10.3389/fphys.2018.01692
Ribatti, D. (2013). Mast cells and macrophages exert beneficial and detrimental effects on tumor
progression and angiogenesis. *Immunology Letters*, *152*(2), 83–88.
http://doi.org/10.1016/j.imlet.2013.05.003
Rosas, M., Davies, L. C., Giles, P. J., Liao, C.-T., Kharfan, B., Stone, T. C., et al. (2014). The
transcription factor Gata6 links tissue macrophage phenotype and proliferative renewal.
Science, *344*(6184), 645–648. http://doi.org/10.1126/science.1251414
Ruckerl, D., & Cook, P. C. (2019). Macrophages assemble! But do they need IL-4R during
schistosomiasis? *European Journal of Immunology*, *49*(7), 996–1000.
http://doi.org/10.1002/eji.201948158
Ruckerl, D., Campbell, S. M., Duncan, S., Sutherland, T. E., Jenkins, S. J., Hewitson, J. P., et al.
(2017). Macrophage origin limits functional plasticity in helminth-bacterial co-infection. *PLoS*

Pathogens, *13*(3), e1006233. http://doi.org/10.1371/journal.ppat.1006233

Schaefer, B., Bartosova, M., Macher-Goeppinger, S., Sallay, P., Vörös, P., Ranchin, B., et al.
(2018). Neutral pH and low-glucose degradation product dialysis fluids induce major early
alterations of the peritoneal membrane in children on peritoneal dialysis. *Kidney International*,
94(2), 419–429. http://doi.org/10.1016/j.kint.2018.02.022

Sica, A., Erreni, M., Allavena, P., & Porta, C. (2015). Macrophage polarization in pathology.
Cellular and Molecular Life Sciences : CMLS, *72*(21), 4111–4126.
http://doi.org/10.1007/s00018-015-1995-y

Song, L., Papaioannou, G., Zhao, H., Luderer, H. F., Miller, C., Dall'Osso, C., et al. (2016). The
Vitamin D Receptor Regulates Tissue Resident Macrophage Response to Injury.
Endocrinology, *157*(10), 4066–4075. http://doi.org/10.1210/en.2016-1474

Uderhardt, S., Herrmann, M., Oskolkova, O. V., Aschermann, S., Bicker, W., Ipseiz, N., et al.
(2012). 12/15-lipoxygenase orchestrates the clearance of apoptotic cells and maintains
immunologic tolerance. *Immunity*, *36*(5), 834–846.
http://doi.org/10.1016/j.immuni.2012.03.010

van de Luijtgaarden, M. W. M., Jager, K. J., Segelmark, M., Pascual, J., Collart, F., Hemke, A. C.,
et al. (2016). Trends in dialysis modality choice and related patient survival in the ERA-EDTA
Registry over a 20-year period. *Nephrology, Dialysis, Transplantation : Official Publication of*
the European Dialysis and Transplant Association - European Renal Association, *31*(1), 120–
128. http://doi.org/10.1093/ndt/gfv295

Wang, Jie, Jiang, Z.-P., Su, N., Fan, J.-J., Ruan, Y.-P., Peng, W.-X., et al. (2013). The role of
peritoneal alternatively activated macrophages in the process of peritoneal fibrosis related to
peritoneal dialysis. *International Journal of Molecular Sciences*, *14*(5), 10369–10382.
http://doi.org/10.3390/ijms140510369

Wang, Jing, & Kubes, P. (2016). A Reservoir of Mature Cavity Macrophages that Can Rapidly
Invade Visceral Organs to Affect Tissue Repair. *Cell*, *165*(3), 668–678.
http://doi.org/10.1016/j.cell.2016.03.009

Wang, Y, Wang, Y. P., Zheng, G., Lee, V. W. S., Ouyang, L., Chang, D. H. H., et al. (2007). Ex
vivo programmed macrophages ameliorate experimental chronic inflammatory renal disease.
Kidney International, *72*(3), 290–299. http://doi.org/10.1038/sj.ki.5002275

Williams, J. D., Craig, K. J., Topley, N., Ruhland, Von, C., Fallon, M., Newman, G. R., et al.
(2002). Morphologic changes in the peritoneal membrane of patients with renal disease.
Journal of the American Society of Nephrology, *13*(2), 470–479.

Wynn, T. A., & Vannella, K. M. (2016). Macrophages in Tissue Repair, Regeneration, and Fibrosis.
Immunity, *44*(3), 450–462. http://doi.org/10.1016/j.immuni.2016.02.015

Yuan, X., Yang, B.-H., Dong, Y., Yamamura, A., & Fu, W. (2017). CRIg, a tissue-resident
macrophage specific immune checkpoint molecule, promotes immunological tolerance in NOD
mice, via a dual role in effector and regulatory T cells. *eLife*, *6*, ra120.
http://doi.org/10.7554/eLife.29540

Zhang, N., Czepielewski, R. S., Jarjour, N. N., Erlich, E. C., Esaulova, E., Saunders, B. T., et al.
(2019). Expression of factor V by resident macrophages boosts host defense in the peritoneal
cavity. *Journal of Experimental Medicine*, *216*(6), 1291–1300.
http://doi.org/10.1084/jem.20182024

Zhang, S., Kim, C. C., Batra, S., McKerrow, J. H., & Loke, P. (2010). Delineation of diverse
macrophage activation programs in response to intracellular parasites and cytokines. *PLoS*
Neglected Tropical Diseases, *4*(3), e648. http://doi.org/10.1371/journal.pntd.0000648

Zhao, Yang, Zou, W., Du, J., & Zhao, Y. (2018). The origins and homeostasis of monocytes and
tissue-resident macrophages in physiological situation. *Journal of Cellular Physiology*,
233(10), 6425–6439. http://doi.org/10.1002/jcp.26461