Synonymous Dinucleotide Usage: A Codon-Aware Metric for Quantifying Dinucleotide

Representation in Viruses

Spyros Lytras [1,*] and Joseph Hughes [1]

[1]MRC – University of Glasgow Centre for Virus Research, Glasgow, United Kingdom

*Correspondence: s.lytras.1@research.gla.ac.uk

Abstract

Distinct patterns of dinucleotide representation, such as CpG and UpA suppression, are characteristic

of certain viral genomes. Recent research has uncovered vertebrate immune mechanisms that select

against specific dinucleotides in targeted viruses. This evidence highlights the importance of

systematically examining the dinucleotide composition of viral genomes. We have developed a novel

metric, called Synonymous Dinucleotide Usage (SDU), for quantifying dinucleotide representation in

coding sequences. Our method compares the abundance of a given dinucleotide to the null

hypothesis of equal synonymous codon usage in the sequence. We present a Python3 package,

DinuQ, for calculating SDU and other relevant metrics. We have applied this method on two sets of

invertebrate- and vertebrate-specific flaviviruses and rhabdoviruses. The SDU shows that the

vertebrate viruses exhibit consistently greater under-representation of CpG dinucleotides in all three

codon positions in both datasets. In comparison to existing metrics for dinucleotide quantification, the

SDU allows for a biologically informed interpretation of its values by comparing representation to an

expectation based on the codon table. Here we apply the method to viruses, but coding sequences

of other living organisms can be analysed in the same way.

Introduction

Certain dinucleotides, two nucleotides adjacent in a sequence, are known to be over- or under-

represented in the genomes of living organisms, creating distinct compositional patterns [1].

Organisms with methylated genomes such as vertebrates and plants have low levels of CpG

dinucleotides. This is not the case for methylase-absent organisms like invertebrates, bacteria and

fungi [2,3]. The cause of CpG suppression lies in the DNA methylation mechanisms of vertebrates

and plants, where cytosine is frequently converted to thymine by DNA methyltransferases [4,5]. UpA

deprivation is consistently present in most living organisms, including prokaryotes. This bias is

suspected to be due to UpA-rich mRNA being unstable and more prone to degradation by cytoplasmic

RNAses [6,7].

Similar patterns of dinucleotide composition have been observed in RNA and DNA viruses, and

appear to have a functional role in the infection and propagation of the viruses [8]. Studies have

shown that experimentally increasing UpA and CpG levels in RNA viruses leads to a decrease in

replication and subsequent viral attenuation, while decreasing their abundance has the opposite

effect [9–11]. Influenza CpG-/UpA-rich sequences have also been shown to have reduced replication

in vivo, causing a more powerful immune response, and in the case of CpG increase, showing

reduced clinical severity [12]. A similar decrease in replication and virulence has been observed in

CpG-/UpA-increased yellow fever virus [13].

Recently, a host immune mechanism of vertebrates which acts on viral genomes on the dinucleotide

level, has been uncovered. The Zinc-finger Antiviral Protein (ZAP) appears to be integral to this

system by targeting CpG-rich infecting viral RNA [14]. This host-driven force selects against viruses

rich in CpG dinucleotides and drives the observed under-representation of CpG in HIV-1 [14], the

model RNA virus Echovirus 7 [15] and even DNA viruses [16]. Additionally, UpA is mildly under-

represented in most viral genomes [17]. Even though our understanding of this dinucleotide's

depletion is limited, there is some evidence suggesting that a similar antiviral mechanism might also

be responsible for this pattern [15,18].

The genetic code is degenerate, meaning that pairs of codons encode the same amino acid. These

codons are called synonymous, because changes between them do not alter the resulting peptide

sequence. The frequency in which synonymous codons are found in a coding sequence can be

biased, as a result of various mutational or selective pressures acting on the genome [19]. Metrics for

measuring codon usage bias exist, such as the Relative Synonymous Codon Usage (RSCU), which

quantify the bias in synonymous codon frequencies [20]. However, these measures do not inform on

the underlying dinucleotide bias in the sequence. If such forces are acting on the dinucleotide and

not the peptide level in viral genomes, adaptation towards, for example, a less CpG-rich genome,

would preferentially take place through synonymous changes in the sequence. This would lead to a

genome with a reduced abundance of dinucleotide patterns recognized by host immunity, whilst still

maintaining the protein sequence. Thus, the dinucleotide-acting compositional bias would produce a

skew in the frequencies of synonymous codon usage in the genome.

So far, quantifying dinucleotide representation in research has been performed by calculating the

relative dinucleotide abundance. This is simply the odds ratio between the frequency of a dinucleotide

XY and the product of the frequencies of each single nucleotide X and Y across a whole sequence

[21]. The recent evidence of a host mechanism selecting against dinucleotides in viruses highlights

the importance of developing a method for quantifying dinucleotide representation that has a direct

biological interpretation. We propose the Synonymous Dinucleotide Usage (SDU), a metric that

compares the occurrences of a given dinucleotide to the null hypothesis that there is equal usage of

synonymous codons. This method provides a way of measuring the extent to which a host-driven

force acting on the dinucleotide level of viral genomes has skewed the synonymous codon usage of

the protein sequence and can be applied in all dinucleotide combinations and sequence frame

positions.

Materials and Methods

Synonymous Dinucleotide Usage (SDU)

A coding sequence can have three distinct dinucleotide frame positions. We define the dinucleotide

frame position 1 as the first and second nucleotide position of a codon, dinucleotide frame position 2

as the second and third nucleotide position of a codon, and dinucleotide bridge position as the third

nucleotide position of a codon and the first position of the downstream codon. Each one of these

positions can take a set of different dinucleotides without changing the amino acid (positions 1 and

2) or amino acid pair (bridge position) in the protein sequence, we define these as a set of

synonymous dinucleotides. For example, threonine has four synonymous codons ACU, ACC, ACA,

ACG. In dinucleotide position 2 of a codon encoding for threonine there are four synonymous

dinucleotides, CU, CC, CA, and CG. As such, the expected proportion of CpU occurring in position 2

coding for threonine under the null hypothesis of equal synonymous codon usage is: $e_i = 0.25$. The

SDU compares the observed proportion of a synonymous dinucleotide of interest (o_i) to that expected

under equal synonymous codon usage (e_i) for a given dinucleotide frame position. The ratio between

o_i and e_i is calculated for each different amino acid or amino acid pair (for the bridge position) and the

SDU is defined as the weighted arithmetic mean of the ratios, weighted by the abundance of each

amino acid in the sequence (Equation 1).

Table 1. Notation used to define SDU and RSDU.

Symbol	Description
i	Amino acid or amino acid pair
j	Dinucleotide
h	Dinucleotide frame position
n_i	Number of occurrences of amino acid or amino acid pair i in the sequence
k	Set of different amino acids or amino acid pairs present in the sequence
$o_{i,j,h}$	Synonymous proportion of dinucleotide j in frame position h for amino acid or amino acid pair i observed in the sequence
$e_{i,j,h}$	Synonymous proportion of dinucleotide j in frame position h for amino acid or amino acid pair i expected under equal synonymous codon usage
N	Total number of amino acids or amino acid pairs present in the sequence

$$SDU_{j,h} = \frac{\sum_{i=1}^{k} n_i \times \frac{o_{i,j,h}}{e_{i,j,h}}}{N} \tag{1}$$

The set *j* includes 16 possible dinucleotide combinations. With three frame positions *h*, the matrix of

$SDU_{j,h}$ has 48 possible combinations. The result of the SDU directly reflects the overall synonymous

dinucleotide representation in each frame position of a given sequence:

• an SDU of 1 indicates that the representation of the dinucleotide of interest in the given frame

position is equal to that expected under the null hypothesis of equal synonymous codon

usage;

• an SDU of 0 indicates that the dinucleotide of interest is completely absent in the given frame

position across the sequence;

• an SDU greater than 1 indicates that the dinucleotide of interest is over-represented in the

given frame position, compared to the representation expected under the null hypothesis;

• an SDU between 0 and 1 indicates that the dinucleotide of interest is under-represented in

the given frame position, compared to the representation expected under the null hypothesis.

Relative Synonymous Dinucleotide Usage (RSDU)

The number of amino acids or amino acid pairs that can be synonymously produced by a certain

dinucleotide varies between dinucleotides and frame positions. This means that SDU measurements

for different positions and dinucleotides can reach different maximum values. Under- (SDU < 1) and

over-representation (SDU > 1) can still be consistently interpreted between positions and

dinucleotides, since an SDU of 1 always reflects complete agreement with the null hypothesis.

However, the magnitude of over-representation cannot be compared between different positions and

dinucleotides. In order to make this comparison possible we extended the SDU to the Relative

Synonymous Dinucleotide Usage (RSDU). This is essentially the calculated SDU value, normalised

by the maximum SDU for this position and dinucleotide (Equation 2). This extension to the metric

does not have a consistent scale for comparing to the null hypothesis, instead allows for comparing

the extent of the difference from the null hypothesis between the different parameters.

$$RSDU_{j,h} = \frac{\sum_{i=1}^{k} n_i \times \frac{o_{i,j,h}}{e_{i,j,h}}}{\sum_{i=1}^{k} n_i \times \frac{1}{e_{i,j,h}}} \qquad (2)$$

126   The results of the RSDU are as follows:

127   •   an RSDU of 1 indicates that only the dinucleotide of interest is being used in the sequence,

128   all the other synonymous dinucleotides being absent for the given position;

129   •   an RSDU of 0, similar to the SDU, indicates that the dinucleotide of interest is completely

130   absent in the given frame position.

Measuring the null distribution

The variance of the metric is expected to vary depending on the sequence length and amino acid

composition. Thus, we have developed a method of measuring the null distribution by randomly

populating the given sequence with synonymous codons without changing the amino acid sequence.

This process takes place for a number of iterations, calculating the SDU (or RSDU) for each model

sequence. This produces a distribution that represents the random error of the metric for the given

sequence abiding to the null hypothesis.

DinuQ Python package

We have developed a Python3 package called *DinuQ* (Dinucleotide Quantification) that includes

modules for calculating the SDU and RSDU with the respective error distribution for any set of coding

sequences provided in a FASTA file. The package further includes modules for calculating the RSCU

[20] and relative dinucleotide abundance (RDA, sometimes called DRA) [22], so that the user can

make comparisons between metrics. The calculated values can be provided in a Python dictionary to

manipulate within the Python framework, or simply exported in a table format. The Biopython package

[23] is being used by the tool to facilitate sequence input. The package can be installed through the

Python Package Index (PyPI). Full documentation and code are available online at

https://github.com/spyros-lytras/dinuq.

Applying the SDU on viral sequences

To test the uses of the SDU we have collected polyprotein coding sequences of vertebrate- and

invertebrate-specific viruses of the *Flaviviridae* family, analysed in previous research [24]. Our

analysis focuses on two representatives of the dataset: the insect-specific *Aedes flavivirus* (AEFV,

Genbank accession: AB488408.1) that usually affects *Aedes spp.* mosquitoes [25] and the *Apoi virus*

(APOIV, Genbank accession: AF160193.1) that has no known insect vector and affects rodents of

the *Apodemus* genus [26]. To extend the analysis, we assembled a second dataset by collecting the

coding sequences of all members of the *Rhabdoviridae* family included in the ICTV Virus Metadata

Resource (version November 27, 2019; MSL34) [27], being labelled as having a vertebrate or

invertebrate host. The SDU and RSCU values where calculated using the DinuQ Python package.

General Linear Models (GLM) for statistical comparisons were performed using the R coding

language.

Results

Effect of Sequence Length on SDU Error

Because the SDU splits up the sequence information into many categories (i.e. independent

proportions of the dinucleotide of interest for each amino acid or amino acid pair), if the given

sequence is short there will be less information in each category and subsequently more

variability/error in the calculated SDU value. We have randomly simulated 10 amino acid sequences

of different lengths sequentially increasing by 10% of the longest sequence's length (i.e. from 700 to

7000 amino acids long). We then implemented our method of measuring error and calculated the

SDU for each random sample. Figure 1a and b illustrate how the error distributions and their standard

deviations vary between sequences of different lengths. By principle, the standard deviation of the

error distribution should approach 0 as the sequence length increases. Our simulation experiment,

however, shows that the magnitude of error is consistently very low for sequences longer than about

17,000bp (standard deviation < 0.05).

Differences in SDU between an insect- and a vertebrate-specific flavivirus

Previous analysis [24] showed that the host environment differently affects the nucleotide, codon and

dinucleotide composition in viruses of the *Flaviviridae* family. In particular, APOIV and other

flaviviruses with no known insect vector show an under-representation of CpG, possibly a result of

the ZAP-related vertebrate immune response targeting CpG-rich viruses. In contrast, AEFV and other

insect-specific flaviviruses do not exhibit this bias. In Figure 2, we present the SDU patterns for all

dinucleotides and frame positions of the two viruses. The SDU of all three frame positions of CpG in

AEFV fall within the error distribution ($SDU_{CpGbridge} = 0.96$, $error_{MIN} = 0.85$, $error_{MAX} = 1.22$; $SDU_{CpGpos1}$

$= 1.08$, $error_{MIN} = 0.84$, $error_{MAX} = 1.15$; $SDU_{CpGpos2} = 0.87$, $error_{MIN} = 0.80$, $error_{MAX} = 1.22$), whereas

the respective SDU values in APOIV, the rodent infecting virus, are well below 1 ($SDU_{CpGbridge} = 0.41$,

$error_{MIN} = 0.82$, $error_{MAX} = 1.19$; $SDU_{CpGpos1} = 0.56$, $error_{MIN} = 0.82$, $error_{MAX} = 1.19$; $SDU_{CpGpos2} = 0.30$,

$error_{MIN} = 0.79$, $error_{MAX} = 1.21$), indicating significant CpG under-representation. These observations

are in agreement with the previous research and the hypothesis of a CpG-targeting antiviral

mechanism in vertebrates.

The SDU of a given dinucleotide should directly reflect the RSCU of the codons that contain it (or in

case of the bridge position: the first nucleotide in the third codon position and second nucleotide in

the first codon position of the downstream amino acid). To illustrate this relation between the two

metrics, we have calculated the RSCU for all codons of AEFV and APOIV (Table 2). ApG in position

1 seems to be over-represented in APOIV ($SDU_{ApGpos1} = 1.51$), which is not the case in AEFV

($SDU_{ApGpos1} = 0.95$). This is clearly depicted in the RSCU values of all AG-starting codons being higher

in APOIV (Table 2). AGA is highly over-represented in the APOIV genome exclusively (APOIV: RSCU

$= 2.19$; AEFV: RSCU $= 1.00$), which also explains the high SDU value for GpA in frame position 2

(APOIV: $SDU_{GpApos2}$ = 1.63; AEFV: $SDU_{GpApos2}$ = 0.96). All CG-starting and -ending codons are under-

represented in APOIV (Table 1), consistent with CpG depletion and low $SDU_{CpGpos1}$ and $SDU_{CpGpos2}$

in the vertebrate-infecting virus. While it is seemingly easy to trace such dinucleotide bias patterns in

the 1st and 2nd frame position only by using the RSCU metric, this becomes much harder for the bridge

position. For example, the SDU values for the bridge positions of ApG, GpA and CpG also fall outside

the null hypothesis distribution in APOIV, but not in AEFV (Figure 2). This highlights the importance

of having a metric for detecting compositional biases on the dinucleotide.

Table 2. RSCU values for each codon, calculated for the APOIV and AEFV coding sequences.

Highlighted in bold are the values mentioned in the text. No values have been calculated for stop

codons (UAA, UAG, UGA), since only one coding sequence was used for each virus.

	APOIV	AEFV		APOIV	AEFV		APOIV	AEFV		APOIV	AEFV
UUU	1.11	1.08	**UCU**	1.01	0.75	**UAU**	0.98	1.00	**UGU**	0.95	1.04
UUC	0.89	0.92	**UCC**	0.73	1.13	**UAC**	1.02	1.00	**UGC**	1.05	0.96
UUA	0.38	0.56	**UCA**	1.57	0.96	**UAA**	STOP	STOP	**UGA**	STOP	STOP
UUG	1.39	1.23	**UCG**	**0.34**	**1.06**	**UAG**	STOP	STOP	**UGG**	1.00	1.00
CUU	1.08	0.83	**CCU**	1.19	0.89	**CAU**	1.25	1.05	**CGU**	**0.48**	**1.19**
CUC	1.00	1.42	**CCC**	0.86	1.00	**CAC**	0.75	0.95	**CGC**	**0.39**	**1.19**
CUA	0.67	0.79	**CCA**	1.69	1.32	**CAA**	0.91	1.23	**CGA**	**0.68**	**1.05**
CUG	1.48	1.17	**CCG**	**0.25**	**0.79**	**CAG**	1.09	0.77	**CGG**	**0.68**	**0.88**
AUU	1.10	1.22	**ACU**	1.13	0.98	**AAU**	0.91	0.84	**AGU**	**1.04**	**0.96**
AUC	1.15	1.05	**ACC**	1.27	1.07	**AAC**	1.09	1.16	**AGC**	**1.32**	**1.13**
AUA	0.75	0.73	**ACA**	1.29	1.00	**AAA**	1.01	1.23	**AGA**	**2.19**	**1.00**
AUG	1.00	1.00	**ACG**	**0.32**	**0.95**	**AAG**	0.99	0.77	**AGG**	**1.58**	**0.69**
GUU	1.08	1.29	**GCU**	1.65	1.07	**GAU**	0.97	0.67	**GGU**	0.73	0.74
GUC	1.00	0.98	**GCC**	1.03	1.62	**GAC**	1.03	1.33	**GGC**	0.73	0.73
GUA	0.29	0.62	**GCA**	1.05	0.68	**GAA**	1.09	1.07	**GGA**	1.75	1.57
GUG	1.63	1.11	**GCG**	**0.27**	**0.63**	**GAG**	0.91	0.93	**GGG**	0.80	0.96

SDU shows consistent CpG differences between insect- and vertebrate-specific viruses

Since there is evidence for a vertebrate immune response selecting against CpG dinucleotides in

viral genomes, we decided to further explore this trend between viruses specific to and absent in

vertebrate hosts using the SDU. First, we calculated the SDU of CpG for all frame positions for the

two sets of insect-specific and vertebrate-specific (no known insect vector) viruses used by Simón *et*

al. [24]. Figure 3a shows the SDU values for the two groups plotted against the total GC content of

the viral coding sequences. As indicated by the graph, the CpG SDU correlates with the overall GC

content with slopes that are not significantly different between hosts (GLM: $F_{1,57} = 1.492$, P = 0.227).

However, the intercepts of the fitted lines are significantly different between the host groups with the

host and the GC content predictors explaining 89% of the variance. The vertebrate-infecting viruses

have lower SDU values, thus more CpG depletion than invertebrate-specific flaviviruses (GLM: R^2 =

0.89, $F_{2,57} = 238.3$, P < 0.0001).

This analysis was replicated on a set of viruses of the *Rhabdoviridae* family (Figure 3b). The trend

observed is similar to that seen with the *Flaviviridae*. Both GC content and host-specificity explain

51% of the variance (GLM: $R^2 = 0.51$, $F_{2,201} = 105.4$, P < 0.0001) around the CpG representation as

quantified by our metric. The difference in rhabdovirus CpG depletion is not as large as in flaviviruses.

CpG representation in invertebrate-infecting *Flaviviridae* is consistent with the null hypothesis of equal

synonymous dinucleotides (mean $SDU_{CpG} = 0.94$), while vertebrate-infecting members of the family

show a higher CpG suppression (mean $SDU_{CpG} = 0.31$). On the contrary, both sets of *Rhabdoviridae*

have depleted CpG levels, vertebrate-specific ones having slightly more suppression (invertebrate-

specific: mean $SDU_{CpG} = 0.42$, vertebrate-specific: mean $SDU_{CpG} = 0.41$).

Discussion

In this paper we propose SDU as a novel method for quantifying dinucleotide representation in a coding sequence by comparing the observed frequency of a synonymous dinucleotide to that expected under the null hypothesis of equal synonymous codon usage. We further extend this metric for comparisons between different dinucleotides with the RSDU formula and provide a means of measuring statistical error for the calculated values.

Using this novel metric, we have explored the representation of different dinucleotides in a set of previously examined members of the *Flaviviridae* family and produced results in agreement with previous research [24]. Based on suggestions from current research that a host-mediated antiviral response in vertebrates selects against CpGs in viral RNA, we focused on examining CpG representation in the *Flaviviridae* and a set of *Rhabdoviridae* viruses. In both datasets, the SDU calculations show a greater depletion of CpG in the vertebrate-specific viruses. Even though there is no experimental evidence that a CpG-recognising immune response is acting on flaviviruses or rhabdoviruses in vertebrates, other RNA viruses seem to be affected by such a mechanism [10,12,28]. Thus, we hypothesise that this immune mechanism is responsible for the difference in CpG SDU values between the two virus groups.

It is important to consider that factors other than a dinucleotide-targeting immune response are likely to bias viral genomic composition and such biases might also be reflected in the SDU values. For example, most invertebrate flaviviruses show weak or no CpG under-representation, while vertebrate-infecting flaviviruses show high CpG depletion. This pattern is less clear in the rhabdoviruses, where CpG is under-represented in both groups. We suggest this might be a result of other rhabdovirus-

specific host mechanisms, or simply a lineage-specific mutational bias or the base composition of the

Rhabdoviridae.

Finally, the SDU allows distinct examination of the different frame positions. This feature should be

useful for exploring cases where selective or mutational processes are acting on specific positions in

the coding sequence, or on longer motifs including the dinucleotide of interest. In comparison to the

relative dinucleotide abundance, the SDU allows for a biologically relevant interpretation of its values,

by comparing dinucleotide abundance to a null hypothesis of synonymous codon usage.

Interpretations of the metric should still be made with caution, since it does not disentangle between

the forces affecting synonymous dinucleotide representation, whether that is selection or mutation on

the dinucleotide level, or forces targeting codons and longer motifs, reflected on the dinucleotide

representation. Whilst we have developed this metric for expanding our understanding of dinucleotide

biases in viruses, the SDU can have broader applications, for example in the host coding sequences.

Computer Code and Software: The *DinuQ* Python package is available to install through PyPI,

under the name 'dinuq' (https://pypi.org/project/dinuq/). All code is available on GitHub

(https://github.com/spyros-lytras/dinuq).

Supplementary Materials: Table S1: NCBI accessions and SDU values for Figure 3.

Author Contributions: Conceptualization, S.L. and J.H.; methodology, S.L.; software, S.L.;

validation, S.L.; formal analysis, S.L. and J.H.; writing—original draft preparation, S.L.; writing—

review and editing, J.H.; visualization, S.L.; supervision, J.H.; project administration, J.H. All authors

have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Medical Research Council (MC_UU_12014/12).

Acknowledgments: We thank Richard Reeve, Paul Johnson and Richard Orton for feedback on the

approach and Oscar MacLean for providing feedback on the manuscript.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the

design of the study; in the collection, analyses, or interpretation of data; in the writing of the

manuscript, or in the decision to publish the results.

References

1. Beutler, E.; Gelbart, T.; Han, J.; Koziol, J.A.; Beutler, B. Evolution of the genome and the genetic

code: Selection at the dinucleotide level by methylation and polyribonucleotide cleavage. *Proc. Natl.*

Acad. Sci. U. S. A. **1989**, *86*, 192–196.

2. Karlin, S.; Doerfler, W.; Cardon, L.R. Why is CpG suppressed in the genomes of virtually all small

eukaryotic viruses but not in those of large eukaryotic viruses? *J. Virol.* **1994**.

3. Cheng, X.; Virk, N.; Chen, W.; Ji, S.; Ji, S.; Sun, Y.; Wu, X. CpG Usage in RNA Viruses: Data and

Hypotheses. *PLoS One* **2013**, *8*.

4. Bird, A.P. DNA methylation and the frequency of CpG in animal DNA. *Nucleic Acids Res.* **1980**, *8*,

1499–1504.

5. Cooper, D.N.; Krawczak, M. Cytosine methylation and the fate of CpG dinucleotides in vertebrate

genomes. *Hum. Genet.* **1989**, *83*, 181–8.

6. Shaw, G.; Kamen, R. A conserved AU sequence from the 3′ untranslated region of GM-CSF mRNA

mediates selective mRNA degradation. *Cell* **1986**.

7. Duan, J.; Antezana, M.A. Mammalian Mutation Pressure, Synonymous Codon Choice, and mRNA

Degradation. *J. Mol. Evol.* **2003**, *57*, 694–701.

8. Simmonds, P.; Xia, W.; Baillie, J.; McKinnon, K. Modelling mutational and selection pressures on

dinucleotides in eukaryotic phyla –selection against CpG and UpA in cytoplasmically expressed RNA

and in RNA viruses. *BMC Genomics* **2013**, *14*, 610.

9. Atkinson, N.J.; Witteveldt, J.; Evans, D.J.; Simmonds, P. The influence of CpG and UpA

dinucleotide frequencies on RNA virus replication and characterization of the innate cellular pathways

underlying virus attenuation and enhanced replication. *Nucleic Acids Res.* **2014**, *42*, 4527–4545.

10. Tulloch, F.; Atkinson, N.J.; Evans, D.J.; Ryan, M.D.; Simmonds, P. RNA virus attenuation by

codon pair deoptimisation is an artefact of increases in CpG/UpA dinucleotide frequencies. *Elife* **2014**,

3, e04531.

11. Witteveldt, J.; Martin-Gans, M.; Simmonds, P. Enhancement of the replication of hepatitis C virus

replicons of genotypes 1 to 4 by manipulation of CpG and UpA dinucleotide frequencies and use of

cell lines expressing SECL14L2 for antiviral resistance testing. *Antimicrob. Agents Chemother.* **2016**,

60, 2981–2992.

12. Gaunt, E.; Wise, H.M.; Zhang, H.; Lee, L.N.; Atkinson, N.J.; Nicol, M.Q.; Highton, A.J.; Klenerman,

P.; Beard, P.M.; Dutia, B.M.; et al. Elevation of CpG frequencies in influenza A genome attenuates

pathogenicity but enhances host response to infection. *Elife* **2016**, *5*.

13. Klitting, R.; Riziki, T.; Moureau, G.; Piorkowski, G.; Gould, E.A.; de Lamballerie, X. Exploratory

re-encoding of yellow fever virus genome: new insights for the design of live-attenuated viruses. *Virus*

Evol. **2018**, *4*, vey021.

14. Takata, M.A.; Gonçalves-Carneiro, D.; Zang, T.M.; Soll, S.J.; York, A.; Blanco-Melo, D.; Bieniasz,

P.D. CG dinucleotide suppression enables antiviral defence targeting non-self RNA. *Nature* **2017**,

550, 124–127.

15. Odon, V.; Fros, J.J.; Goonawardane, N.; Dietrich, I.; Ibrahim, A.; Alshaikhahmed, K.; Nguyen, D.;

Simmonds, P. The role of ZAP and OAS3/RNAseL pathways in the attenuation of an RNA virus with

elevated frequencies of CpG and UpA dinucleotides. *Nucleic Acids Res.* **2019**, *47*, 8061–8083.

16. Lin, Y.-T.; Chiweshe, S.; McCormick, D.; Raper, A.; Wickenhagen, A.; DeFillipis, V.; Gaunt, E.;

Simmonds, P.; Wilson, S.J.; Grey, F. Human cytomegalovirus evades ZAP detection by suppressing

CpG dinucleotides in the major immediate early genes. *bioRxiv* **2020**, 2020.01.07.897132.

17. Shackelton, L.A.; Parrish, C.R.; Holmes, E.C. Evolutionary Basis of Codon Usage and Nucleotide

Composition Bias in Vertebrate DNA Viruses. *J. Mol. Evol.* **2006**, *62*, 551–563.

18. Fros, J.J.; Dietrich, I.; Alshaikhahmed, K.; Passchier, T.C.; Evans, D.J.; Simmonds, P. CpG and

UpA dinucleotides in both coding and non-coding regions of echovirus 7 inhibit replication initiation

post-entry. *Elife* **2017**, *6*.

19. Sharp, P.M.; Averof, M.; Lloyd, A.T.; Matassi, G.; Peden, J.F. DNA-sequence evolution - the

sounds of silence. *Philos. Trans. R. Soc. B-Biological Sci.* **1995**, *349*, 241–247.

20. Sharp, P.M.; Tuohy, T.M.F.; Mosurski, K.R. Codon usage in yeast - cluster analysis clearly

differentiates highly and lowly expressed genes. *Nucleic Acids Res.* **1986**, *14*, 5125–5143.

21. Karlin, S.; Burge, C. Dinucleotide relative abundance extremes: a genomic signature. *Trends

Genet.* 1995, *11*, 283–290.

22. Karlin, S.; Burge, C. Dinucleotide relative abundance extremes: a genomic signature. *Trends

Genet.* 1995, *11*, 283–290.

23. Cock, P.J.A.; Antao, T.; Chang, J.T.; Chapman, B.A.; Cox, C.J.; Dalke, A.; Friedberg, I.;

Hamelryck, T.; Kauff, F.; Wilczynski, B.; et al. Biopython: Freely available Python tools for

computational molecular biology and bioinformatics. *Bioinformatics* **2009**.

24. Simón, D.; Fajardo, A.; Sóñora, M.; Delfraro, A.; Musto, H. Host influence in the genomic

composition of flaviviruses: A multivariate approach. *Biochem. Biophys. Res. Commun.* **2017**, *492*,

572–578.

25. Blitvich, B.J.; Firth, A.E. Insect-specific flaviviruses: A systematic review of their discovery, host

range, mode of transmission, superinfection exclusion potential and genomic organization. *Viruses*

2015, *7*, 1927–1959.

26. Billoir, F.; De Chesse, R.; Tolou, H.; De Micco, P.; Gould, E.A.; De Lamballerie, X. Phylogeny of

the genus Flavivirus using complete coding sequences of arthropod-borne viruses and viruses with

no known vector. *J. Gen. Virol.* **2000**, *81*, 781–790.

27. International Committee on Taxonomy of Viruses (ICTV) Available online:

https://talk.ictvonline.org/taxonomy/vmr/ (accessed on Feb 27, 2020).

28. Ibrahim, A.; Fros, J.; Bertran, A.; Sechan, F.; Odon, V.; Torrance, L.; Kormelink, R.; Simmonds,

P. A functional investigation of the suppression of CpG and UpA dinucleotide frequencies in plant

RNA virus genomes. *Sci. Rep.* **2019**, *9*.

Figure 1. Comparison of error for the SDU_{CpGbridge} of 10 simulated amino acid sequences of different lengths with 1000 random samples of nucleotide sequences for each amino acid sequence: (**a**) standard deviation of the mean of the SDU error distributions; (**b**) Violin plots of the error distribution for each simulated sequence.

Figure 2. SDU values for all informative dinucleotides and frame positions plotted for the AEFV and APOIV coding sequences (labelled respectively). Dot points indicate observed SDU values and violin plots indicate SDU error distributions around the null hypothesis (1000 random samples for each SDU value). Position 1 of dinucleotides CpC, CpA, GpC, GpG, GpU, GpA, UpG, UpA, ApC, ApU, ApA are excluded because they can only produce one amino acid (non-informative).

Figure 3. Comparison of SDU$_{CpG}$ for all frame positions between invertebrate- and vertebrate-specific

(**a**) *Flaviviridae* and (**b**) *Rhabdoviridae*, plotted against the overall GC content of the coding

sequences.