Chromatin remodelling by INO80 at transcription pause sites promotes premature

termination of mRNA synthesis

Sara Luzzi[1,*], Ugo Szachnowski[2,*], Sarah Greener[1], Kenny Schumacher[3,4,5,6], Stuart Fulton[7],

Camille Gautier[2], Kang Hoo Han[8], Jack Darke[1], Rossana Piccinno[9], Anne Lafon[2,9], B.

Franklin Pugh[8,10], Didier Devys[3,4,5,6], Laszlo Tora[3,4,5,6], Antonin Morillon[2†], and Manolis

Papamichos-Chronakis[4,11†]

Affiliations:

[1]Biosciences Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon

Tyne, NE2 4HH, United Kingdom

[2]ncRNA, Epigenetic, and Genome Fluidity, Institut Curie, PSL Research University, CNRS

UMR 3244, Pierre et Marie Curie University, 26 rue d'Ulm, 75248 Paris Cedex 05, France

[3]Institut de Génétique et de Biologie Moléculaire et Cellulaire, 67404, Illkirch, France

[4]Centre National de la Recherche Scientifique, UMR7104, 67404, Illkirch, France

[5]Institut National de la Santé et de la Recherche Médicale, U964, 67404, Illkirch, France

[6]Université de Strasbourg, 67404, Illkirch, France

[7]Department of Molecular Physiology and Cell Signaling, Institute of Systems, Molecular,

and Integrative Biology, University of Liverpool, Liverpool L69 7ZB, UK

[8]Center for Eukaryotic Gene Regulation, Department of Biochemistry and Molecular

Biology, The Pennsylvania State University, University Park, PA 16802, USA

[9]Institut Curie, PSL Research University, CNRS, UMR3664, 26 rue d'Ulm, 75248 Paris,

France

[10]Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853,

USA

[11]Lead Contact

*These authors contributed equally to this work

†Correspondence to: antonin.morillon@curie.fr and manolis.papamichos@newcastle.ac.uk

Abstract

How co-transcriptional RNA quality control is regulated remains poorly understood. Here, we report that in *S. cerevisiae* premature transcription termination of mRNAs is regulated by the evolutionarily conserved ATP-dependent chromatin remodeling INO80 complex. Loss of INO80 leads to an increase in promoter-proximally paused RNA Polymerase II and defective progression into the gene body. We show that promoter-proximal transcriptional pausing correlates with loading of RNA surveillance and transcription termination factors to mRNA transcripts. Cells lacking INO80 are defective for the Nrd1-Nab3-Sen1 (NNS)-dependent pathway for transcription termination at snRNA genes and promoter-proximally sites of mRNA genes. We demonstrate that INO80 promotes the association of the RNA surveillance and termination factor Nab2 with short promoter-proximal mRNA transcripts. We provide evidence that co-transcriptional recruitment of Nab2 to chromatin is regulated by INO80, which enables the interaction of Nab2 with the histone variant H2A.Z. Our work suggests a chromatin mechanism for premature transcription termination at promoter-proximally pausing sites, linking RNA quality control to the transcriptional process.

INTRODUCTION

Elimination of aberrant mRNAs is essential for proper gene expression. In eukaryotic cells nuclear RNA surveillance and quality control mechanisms monitor mRNA biogenesis co-transcriptionally and terminate transcription prematurely for degradation of the nascent RNA transcript[1]. Elegant genome-wide studies in the budding yeast *Saccharomyces cerevisiae* have suggested that unproductive transcriptional elongation of protein coding genes is wide-spread and common[2,3]. Furthermore, promoter-proximal short mRNAs enriched in 3'-oligo(A) tails have been identified[4], indicating extensive premature termination across the yeast protein-coding genome[5]. Likewise, only 10% of promoter-proximally paused polymerases was shown to enter productive elongation in human cells, with the remaining being prematurely terminated during abortive elongation[6,7].

The major nuclear RNA quality control pathway in yeast is mediated by the RNA surveillance complex Nrd1-Nab3-Sen1 (NNS)[8] and the Rrp6-dependent nuclear exosome complex with 3' exonuclease activity[9]. The NNS-dependent nuclear RNA quality control pathway acts prominently to restrict the extensive and pervasive transcription of non-coding RNAs (ncRNAs)[10]. Nrd1 and Nab3 have also been found to be recruited at active protein-coding genes with a preference near their 5' end and bind most mRNA transcripts[11-15], suggesting a genome-wide role for NNS in mRNA transcription termination[15] and co-transcriptional control of mRNA metabolism. However, this model has been challenged by a separate study, which proposed that only a few mRNA genes are subject to transcriptional attenuation by the NNS complex[16]. Thus, although the RNA surveillance and termination pathways have been characterized in molecular detail, the mechanisms regulating premature transcription termination during mRNA synthesis by Pol II remain unclear.

In metazoans, short, promoter-associated polyadenylated mRNA transcripts[17] have been found to associate with Pol II pausing[18], with premature termination and exosome-dependent

degradation coinciding with promoter-proximal pausing sites[19]. Experiments with chemical

transcriptional inhibitors indicated high rates of turnover and premature termination of

promoter-proximal paused Pol II[6,20,21], suggesting regulation of gene expression by

transcriptional attenuation. Even though promoter-proximal pausing plays an important

regulatory role in eukaryotic transcription[22-25], how pausing is linked to premature

termination remains poorly understood.

Eukaryotic transcription is controlled by the underlying chromatin structure. ATP-dependent

chromatin remodelling enzymes shape the chromatin landscape and act at almost all stages of

RNA production controlling transcription activation and silencing, modulation of

transcriptional rates, transcription termination and RNA export from the nucleus[26-30]. In

mammals, premature termination of mRNA genes and promoter-proximal transcriptional

pausing have been associated with nucleosome organization[19]. However, the role of

chromatin remodelers in co-transcriptional RNA quality control remains elusive.

The multisubunit ATP-dependent chromatin remodelling complex INO80 has a well-

established and important role in transcription across eukaryotes[31]. INO80 is enriched at the

transcription start sites (TSS) of most active and poised mRNA genes in yeast and

mammals[32,33]. The nucleosome remodelling activity of INO80 is important for the exchange

of the histone variant H2A.Z for H2A in nucleosomes and for controlling nucleosome

positions[31]. While it is generally proposed that the function of INO80 during transcription is

to control transcription initiation, INO80 physically interacts with the elongating RNA

Polymerase II machinery[34,35] and binds to mRNAs[36,37]. This indicates a -yet- undefined role

for INO80 in the transcriptional process.

Here we investigate the role of INO80 in transcription regulation in *S. cerevisiae*. Using

Native Elongating Transcript sequencing (NET-seq) we analyse how the process of

transcription is regulated by INO80 at nucleotide resolution. We show that INO80 enables

the progression of Pol II from the promoter-proximal region into the gene body. Our analysis

indicates that promoter-proximal pausing of Pol II is linked to loading of transcription

termination and RNA processing factors to mRNA transcripts. Mechanistically our study

reveals that INO80 modulates the interaction between the RNA surveillance and termination

factor Nab2 and H2A.Z, enabling recruitment of Nab2 to chromatin. Overall, our results

suggest that the chromatin remodelling function of INO80 promotes premature transcription

termination.

RESULTS

Defective progression of Pol II from the promoter proximal region into the gene body in

the absence of INO80

To illuminate the role of INO80 in transcription, we performed Native Elongating Transcript

NET-seq analysis in Wild-Type (WT) and *ino80Δ* yeast cells expressing a FLAG-tag version

of the largest subunit of the RNA Polymerase II complex, Rpb1 (Extended Data Fig. 1a, b).

NET-seq captures nascent RNAs bound to transcriptionally engaged Pol II and sequences the

3'-end of the purified RNAs, thus providing a detailed map of elongating Pol II across the

genome at nucleotide resolution[38]. Therefore, and to avoid confusion with the term "nascent

RNAs" that is commonly used to describe newly synthesized RNAs that are not necessarily

associated with Pol II[39], we will henceforth refer to the Pol II-associated transcripts captured

by NET-seq as Pol II density.

Following removal of duplicated reads and DEseq normalization, NET-seq densities were

computed as normalized counts over gene length and their distribution across the gene was

compared between the two strains. Consistent with previous reports[38,40], promoter-proximal

region showed sense Pol II density, with a subsequent reduction further into the gene body in

WT cells (Fig. 1a). Interestingly, in *ino80Δ* cells, while sense Pol II density was reduced at

the promoter-proximal peak, it decreased further downstream the promoter-proximal region,

compared to WT (Fig 1a). Changes in the abundance of sense and antisense NET-seq reads at

mRNA genes in *ino80Δ* did not correlate (Extended Data Fig. 1c, r=0.04), indicating that

altered sense Pol II density in the absence of INO80 is independent of antisense transcription.

We explored whether the decrease in Pol II density downstream the promoter proximal

region in *ino80Δ* indicates defective progression of RNA Polymerase II into the gene body.

To test this possibility, we calculated Traveling Ratio (TR) values for all mRNA genes by

comparing the Pol II density in the gene body relative to the promoter-proximal region of the

first 200bp after the TSS (Fig. 1b). In WT cells, 68% of protein-coding genes exhibited

higher Pol II density at the promoter-proximal region compared to Pol II density in the gene

body for (TR<1), indicating accumulation of Pol II proximally to promoter for most of the

genes. Loss of INO80 led to an overall significant reduction in TR, with 809 genes exhibiting

a decrease in TR of at least 1.5 fold-change compared to WT (Fig. 1b). Overall, 73% of genes

demonstrated a TR<1 in *ino80Δ*. This indicates a genome-wide defect in progression of Pol II

from the promoter-proximal region into the gene body in the absence of INO80.

INO80 physically interacts with the elongating RNA Polymerase II complex[34,35] and is

enriched within gene bodies[33,41]. To evaluate its role during transcription elongation, we

asked whether the enrichment of INO80 inside gene bodies correlates with the intragenic

levels of Pol II. ChIP-exo analysis provides near single-nucleotide resolution of factor

binding to DNA[42] and can distinguish between recruitment inside and outside the gene body

with great accuracy. When the Ino80[41] and Rpb1 ChIP-exo densities within the gene bodies

were compared, a strong, positive correlation between their enrichment was observed (Fig.

1c, r=0.68). This indicates that the presence of INO80 inside the gene is associated with

transcription elongation activity.

Chromatin remodelling activity and functionality of the INO80 complex relies on the INO80-

specific subunits Arp5 and Arp8[31]. Cells lacking either *INO80* or the INO80-specific subunits

ARP5 or *ARP8* grew poorly in the presence of 6-azauracil (6-AU) (Fig. 1d and Extended

Data Fig. 1d, e), an inhibitor of GTP biosynthesis which sensitizes transcriptional elongation

mutants[43]. Therefore, these results suggest that INO80 promotes transcription elongation,

supporting a genome-wide, post-initiation role for INO80 in gene expression.

Global downregulation of Pol II transcription in the absence of INO80 is associated

with defective mRNA degradation

Increase in transcription initiation can lead to enhanced pausing of Pol II at predetermined

sites[44]. We therefore asked whether transcription rates are increased in the absence of INO80,

by comparing the levels of newly synthesized mRNAs in WT and *ino80Δ* cells. Cells grown

in rich medium were pulse-labelled with 4-thiouracil (4tU) and mixed in a fixed ratio with

labelled *S.pombe* cells (spike-in) for normalization[45]. In parallel, 4tU labelling was also

conducted in *ino80Δ* cells expressing wild-type *INO80* from a plasmid, serving as a control

for *ino80Δ*-mediated changes in RNA expression. Purified total (steady state) and labelled

(newly synthesized) RNA from all samples were subjected to sequencing.

After normalization to the *S. pombe* spiked-in signal and applying a cutoff of 1.5-fold change

and p-value < 0.05, only 3.6% of all protein coding genes showed significant increase in

newly synthesized mRNA levels in *ino80Δ* (n=178, Fig. 3a). Contrary, 34% of the protein-

coding genome demonstrated a significant reduction in mRNA synthesis in *ino80Δ* compared

to WT (n=1706, Fig. 3a). The downregulation of mRNA synthesis rates was INO80-

dependent, since in *ino80Δ* cells expressing *INO80* from an episomal plasmid, less than 4%

of genes had markedly decreased newly synthesized mRNA levels (n=193, Extended Data

Fig. 3a). Therefore, deletion of INO80 induces genome-wide downregulation of mRNA

synthesis rates, arguing against increased transcription initiation as the underlying cause of

enhanced Pol II pausing in *ino80Δ*.

Interestingly, despite the global decrease in newly synthesized mRNA levels in *ino80Δ*, the

cellular, total mRNA levels for 81% of the protein-coding genes did not significantly change,

while 13% were increased in the absence of INO80 (Fig. 3b). This disentanglement between

newly synthesized and total RNAs suggests buffering of mRNA transcript levels[46] in *ino80Δ*

by a compensatory mechanism. Cellular mRNA levels are determined by the rates of mRNA

synthesis and degradation. To understand why cellular mRNA levels remain high in *ino80Δ*,

we conducted comparative dynamic transcriptome analysis (cDTA)[47], which measures

mRNA synthesis and decay rates (SR and DR respectively) for all protein-coding genes.

Comparison of the SR and DR changes in *ino80Δ* relative to WT revealed that the decrease in

SR observed in *ino80Δ* positively correlated with a concomitant decrease in DR (r=0.47, p-

value=$2.9e^{-233}$, Fig 3c). This indicates that the defect in Pol II transcription in *ino80Δ* is

associated with compromised mRNA degradation, in a similar fashion to what has been

observed for the rpb1-N488D mutant[47], which reduces the processivity of Pol II[48].

To verify that mRNAs are stabilized in the absence of INO80, we analysed mRNA decay

rates following transcription inhibition with thiolutin in WT and ino80Δ cells. Thiolutin was

used at concentration below the threshold reported to affect mRNA half-life[49] and mRNA

values were normalized relative to the stable scR1 RNA, which is transcribed by Pol III. The

relative abundance of *NRD1*, *RNA1* and *HPT1* transcripts, which showed decreased DR in the

cDTA analysis (Extended data Fig. 3b), was markedly higher in *ino80Δ* compared to WT

after transcription inhibition (Fig. 3d and Extended data Fig. 3c). Therefore, the concomitant

change in mRNA synthesis and degradation in *ino80Δ* suggest that the transcriptional role of

INO80 is linked to RNA decay by buffering steady state mRNA levels.

Promoter-proximal Pol II pausing is linked to increased loading of RNA surveillance

and termination factors to mRNA transcripts

The results from our NET-seq and 4tU-seq analysis prompted us to evaluate whether

promoter-proximal pausing is linked to nuclear RNA quality control and premature

transcription termination.

The nuclear RNA cleavage and termination factor Nab2[50] and the Trf4/Trf5-Air1/Air2-Mtr4

polyadenylation (TRAMP) complex are required for processing and delivering nascent

transcripts to the nuclear exosome for degradation[51-55]. Nab2, Mtr4 and the Nab3 subunit of

the NNS complex bind most of mRNAs[4,56]. We retrieved published genomic data for RNA

binding (CRAC) for Nab3, Nab2 and Mtr4[4,56]. and corrected their binding to mRNA

transcripts for their respective gene transcription levels as measured by CRAC-Rpb1[56] (Fig.

4a). This normalisation generated an index of Transcript Instability (TI) values for all

protein-coding genes, allowing us to quantitatively assess the level of premature transcription

termination taking place within each mRNA gene.

We evaluated whether accumulation of Pol II at the promoter-proximal region is associated

with premature termination. TI values were calculated for the promoter-proximal region (TSS

to TSS+200) and the downstream gene body (TSS+200 to PAS). Genes were clustered in

four groups based on their TR values in WT conditions (Fig. 4b) and $TIs^{Mtr4/Nab3/Nab2}$ were

analysed for the genes of each group. Intriguingly, a reciprocal relationship between TR and

promoter-proximal TI was observed: genes with increased accumulation of Pol II at the

promoter-proximal region relative to the gene body (low TR) had high $TI^{Mtr4/Nab3/Nab2}$, while

genes with little or no promoter-proximal accumulation of Pol II (high TR) had low TI (Fig.

4c). Contrary, no decrease in the $TI^{Mtr4/Nab3/Nab2}$ values was observed within the gene body in

the four TR groups (Fig. 4c). Therefore, defective progression of Pol II from the promoter-

proximal region correlates with early transcription termination at the same region.

Functional and physical interactions link INO80 to RNA quality control and NNS-

dependent transcription termination pathway

Since in *ino80Δ* cells Pol II is stalled at genomic sites associated with increased binding of

RNA surveillance factors to RNA, we interrogated whether the role of INO80 in gene

expression is functionally linked to the RNA quality control process. Abd1 is an essential

m7G cap methyltransferase that prevents mRNA destabilization and degradation[57]. Deletion

of *ARP8* in the *abd1-DAmP* strain, which expresses a hypomorphic allele of *ABD1*[58], led to

synthetic lethality in non-permissive conditions (Fig. 5a), suggesting that INO80 is essential

for cellular viability when mRNAs are destabilized. Deletion of *ARP8* in the *nrd1-V368G*

(designated as *nrd1-5*) mutant strain, which is deficient in transcription termination[8] led to

synthetic lethality in 6-AU (Fig. 5b). In addition, the *arp8Δ* and *nrd1-5* mutants were

epistatic for *INO1* expression (Fig. 5c) suggesting that the INO80 and NNS complexes work

in the same pathway in gene regulation. These genetic interactions link INO80 to co-

transcriptional RNA quality control, prompting us to test whether the INO80 complex

physically associates with the transcription termination machinery.

Protein co-immunoprecipitation analysis after nucleic acid removal demonstrated that the

INO80 complex physically interacts with both Nrd1 and Nab2 *in vivo* (Fig. 5d, e). Contrary,

we could not detect a physical interaction between the INO80 complex and the component of

cleavage and polyadenylation factor IA (CF IA) complex Pcf11 (Extended Data Fig. 4a),

which promotes transcription termination at the PAS.

INO80 promotes NNS-dependent transcription termination

To explore the possibility that INO80 is involved in Nrd1-dependent termination, we

employed the well-characterized *ACT-SNR^{TTS}-CUP* reporter system[8,59] (Extended Data Fig.

5a). Briefly, cells were transformed with ACT-CUP plasmids without or with the

Transcription Termination Site (TTS) of the snoRNA gene *SNR13* (*SNR^{TTS}*) inserted prior to

the CUP1 gene, which confers resistance to copper. Termination of snoRNAs, including

SNR13, is regulated by the NNS complex[8,60]. INO80 is co-enriched with Nrd1 at the

endogenous *SNR13*^{TTS} region (Extended Data Fig. 4b), making the *ACT-SNR^{TTS}-CUP*

reporter a suitable system to test the role of INO80 in NNS-dependent transcription

termination. Contrary to WT, *ino80Δ* cells transformed with the *ACT-SNR^{TTS}-CUP* plasmid

were able to grow in the presence of copper, suggesting read-through transcription of *CUP1*

(Fig. 6a). RT-qPCR analysis using primer pairs that target the regions before and after the

SNR13^{TTS} insertion site in the reporter plasmids showed a significant increase in the relative

transcription of *CUP1* in the *ACT-SNR^{TTS}-CUP* plasmid in *ino80Δ* and *arp8Δ* mutants by 2.8-

fold and 3.5-fold respectively (Fig. 6b and Extended Data Fig. 5c). Similarly, a 3-fold

increase was observed in *nrd1-5* cells, in accordance to previous studies[8] (Fig. 6b and

Extended Data Fig. 5c). Inducible degradation of Ino80 also resulted in an acute increase in

relative *CUP1* levels from the *ACT-SNR^{TTS}-CUP* but not from the control *ACT-CUP* plasmid

lacking the transcription termination site (Extended Data Fig. 5d). Deletion of either *INO80*

or *ARP8* did not significantly alter gene expression of any of the known transcription

termination and RNA surveillance factors (Extended Data Table 2). These results support a

direct role for INO80 in transcription termination at the *SNR13^{TTS}* termination site.

To further substantiate a role for INO80 in NNS-dependent termination, we evaluated read-

through transcription at the termination sites of endogenous snoRNA genes. Using paired-

end, spiked-in RNA sequencing we mapped the sense RNA-seq reads encompassing the TTS

of the snoRNA genes in WT and *ino80Δ*. In the absence of INO80 read-through transcripts

are significantly increased for the majority of the snoRNA genes, but not for the ribosomal

protein coding gene *YNL096C* (Fig. 6c, d, and Extended Data Fig. 5e and Extended Data

Table 3). The increase in readthrough transcripts in *ino80Δ* is not due to defective

degradation, as termination of snoRNAs in cells deleted for the 5'-3' exoribonuclease *XRN1*

was normal, while deletion of the nuclear exosome subunit *RRP6* compromised termination

only at a small number of specific snoRNA genes, in accordance with previous studies[61,62]

(Fig. 6d and Extended Data Table 2). These results suggest that INO80 promotes NNS-

dependent transcription termination.

To investigate the effect of INO80 loss on premature termination of mRNA synthesis, we

focused our analysis on the *NRD1* gene, which is a known target for promoter-proximal

transcription termination by the NNS/Rrp6 pathway[8,61]. Ino80 is enriched at the promoter-

proximal region of *NRD1* (Extended Data Fig. 5f) and loss of INO80 resulted in increased

promoter-proximal Pol II density at *NRD1* compared to WT (Extended Data Fig. 5g). To

assess termination proximally to promoters and at the end of the gene we conducted

quantitative 3' rapid amplification of cDNA ends (3' RACE) using gene-specific primers for

either the promoter-proximal or the PAS region of the *NRD1* gene respectively. Deletion of

RRP6 increased the relative abundance of promoter-proximally terminated *NRD1* mRNAs

compared to WT, while also upregulated the expression of *NRD1* transcripts terminated at the

end of the gene, in line with RNA-seq data (Fig. 6e and Extended Data Fig. 5h). Notably,

deletion of either *INO80* or *APR8* attenuated the expression of early terminated transcripts

compared to the PAS-terminated *NRD1* transcripts (Fig. 6e). This suggests that the INO80

complex promotes premature transcription termination at mRNA genes.

INO80 promotes chromatin recruitment and association of Nab2 with H2A.Z

To understand the mechanism underlying the function of INO80 in premature termination,

we hypothesized that INO80 regulates the recruitment of the RNA quality control machinery

to chromatin. The RNA surveillance factor Nab2 promotes transcription termination[50]. Nab2

binds promoter-proximal mRNAs that are prematurely terminated[4], including *NRD1*

(Extended Data Fig. 6a), while it is co-transcriptionally recruited to the gene body of actively

transcribed mRNA genes[12]. We therefore tested by ChIP-qPCR whether association of Nab2

with chromatin depends on INO80. Co-transcriptional recruitment of Nab2 to the promoter-

proximal region of *NRD1* was reduced in the absence of INO80 (Fig. 7a and Extended Data

Fig. 6b). Likewise, Nab2 co-transcriptional recruitment at the *SNR13* gene was also

compromised in *ino80Δ* (Extended Data Fig. 6c). Co-immunoprecipitation for Nab2

confirmed that the physical interaction between Nab2 and Rpb1 was independent of INO80

(Extended Data Fig. 6d), excluding the possibility that the decreased recruitment of Nab2 to

chromatin is caused by compromised binding to the Pol II machinery. To evaluate whether

loss of INO80 disrupts the association of Nab2 with the promoter-proximal *NRD1* transcript,

we conducted quantitative RNA-IP for Nab2. In *ino80Δ* cells the binding of Nab2 to

promoter-proximal *NRD1* RNA was significantly reduced compared to its binding to the

PAS-proximal *NRD1* region (*3'-NRD1*) (Fig. 7b). These results suggest that INO80 regulates

recruitment of Nab2 to chromatin to promote its binding to nascent RNAs.

We sought to gain insight into how INO80 promotes recruitment of Nab2 to chromatin.

Promoter-proximal nucleosomes are enriched in the histone variant H2A.Z[63]. H2A.Z has

been suggested to have a role in RNA quality control and non-PAS transcription termination

in *D.melanogaster* and *S.pombe*[64-66]. INO80 preferentially remodels nucleosomes containing

H2A.Z and controls the dynamic association of H2A.Z with chromatin[33,67-69], raising the

possibility that H2A.Z might be involved in NNS-dependent termination. Cells lacking

HTZ1, which encodes for H2A.Z in budding yeast, grew in the presence of copper when

transformed with the *ACT-SNR^{TTS}-CUP* plasmid (Fig.7c and Extended Data Fig. 6f),

suggesting that H2A.Z suppresses read-through transcription from NNS-dependent

termination sites. Promoter-proximally terminated *NRD1* transcripts were decreased in *htz1Δ*

compared to WT (Fig. 7d), indicating that loss of H2A.Z compromises premature

transcription termination of mRNA genes. In line with defective early termination, co-

transcriptional recruitment of Nab2 at the promoter-proximal region of *NRD1* was reduced in

htz1Δ cells (Fig. 7e and Extended Data Fig. 6g). Co-immunoprecipitation assays after nucleic

acid removal demonstrated that H2A.Z interacts with Nab2, even after non-physiological

high-salt wash conditions, but not with Pcf11 (Fig. 7f and Extended Data Fig. 7h, i). Since

our evidence indicate that H2A.Z is involved in premature transcription termination and RNA

quality control at mRNA genes, we asked whether INO80 controls the interaction between

H2A.Z and Nab2. Remarkably, binding of Nab2 to H2A.Z was decreased in *ino80Δ* (Fig. 7f).

This suggests that INO80 enables the recruitment of the Nab2 RNA surveillance machinery

to chromatin by promoting its interaction with H2A.Z.

DISCUSSION

Here, we elucidate a role for INO80 in transcription. We report that INO80 enables recruitment of the RNA quality control machinery to transcribed mRNA genes for early transcription termination, promoting transcription elongation. Our data suggest that by removal of Pol II and termination of unproductive transcription, INO80 defines a chromatin-based pathway for co-transcriptional mRNA quality control. Based on our results we propose that control of premature transcriptional termination by INO80-dependent chromatin remodelling is a genome-wide regulatory mechanism of eukaryotic gene expression.

Linking mRNA quality control to chromatin regulation

What is the role of Pol II accumulation at the promoter-proximal region? Our analysis links promoter-proximal accumulation of Pol II with RNA surveillance and transcription termination. Binding of the RNA quality control factors Nab3, Mtr4 and Nab2 to mRNA transcripts is enhanced at promoter-proximal regions. Therefore, a model emerges that increased dwelling of Pol II at specific promoter-proximal sites provides the time for RNA surveillance and termination factors to be loaded to nascent transcripts efficiently. Our cumulative findings that in the absence of INO80 (i) Pol II accumulates at promoter proximal regions, (ii) Nab2 recruitment to chromatin and binding to RNA are defective and (iii) early termination is compromised, suggest that removal of Pol II molecules that are not engaged in productive transcription is mediated by INO80. Given that loss of INO80 results in reduced transcription rates, our results substantiate the hypothesis that chromatin promotes transcription by orchestrating the co-transcriptional RNA quality control pathway. It is therefore tempting to speculate that chromatin acts as a molecular switch that dictates whether Pol II will either progress into productive elongation or be removed to allow for the next round of transcription to take place.

Chromatin regulation of premature transcription termination by INO80

How does INO80 promote premature termination? The physical interactions of the INO80

complex with the transcription termination and RNA surveillance factors Nrd1 and Nab2, and

the enrichment of INO80 at NNS-dependent termination regions suggest a direct involvement

for INO80 in the co-transcriptional RNA quality control pathway. Nab2 associates with the

RNA cleavage factor Hrp1[70] and both factors are involved in mRNA quality control and

NNS-dependent termination[4,71,72]. The INO80-dependent recruitment of Nab2 to chromatin

could therefore promote co-transcriptional cleavage of the nascent RNA, leading to

destabilization of the transcription elongation complex and subsequent release of Pol II from

chromatin. As INO80 interacts with and facilitates release of poly-ubiquitinated Rpb1 from

chromatin[35], an intriguing possibility is that Rpb1 ubiquitination is involved in the process,

providing specificity for selective eviction of Pol II molecules by INO80.

In agreement with a role for H2A.Z in RNA quality control and non-poly(A) transcription

termination[64-66], our results suggest that INO80 promotes recruitment of Nab2 to chromatin

by regulating its interaction with H2A.Z (Figure 7). This implies that remodelling of H2A.Z-

containing nucleosomes by INO80[67-69] is important for premature termination. It is therefore

likely that intermediate steps in the H2A.Z exchange reaction by INO80, for example

disposing of DNA from the surface of H2A.Z-H2B dimers[69], might be involved in the

process. How H2A.Z promotes recruitment of the RNA quality control machinery to

chromatin is not clear. It would not be surprising if regulation of premature termination

entails posttranslational modifications on H2A.Z, which would command its role in the RNA

quality control pathway.

Promoter-proximal transcriptional pausing has recently been linked to premature termination

in metazoans[6,19-21]. Uncontrolled transcription elongation and dysregulated Pol II pausing

have been implicated in human disease[73,74]. Mutations in the RNA surveillance machinery

have been associated with increased transcriptional stress and genomic instability[75], both

hallmarks of cancer[76]. Dysregulated expression of INO80 leads to uncontrolled gene

expression and is associated with disease[31,77,78]. We therefore anticipated that our reported

chromatin mechanism of co-transcriptional RNA quality control has general relevance for

regulation of eukaryotic gene expression and in disease.

Competing interests

The authors declare that they have no competing interests. BFP has a financial interest in

Peconic, LLC, which utilizes the ChIP-exo technology implemented in this study and could

potentially benefit from the outcomes of this research.

Funding

MPC is supported by Newcastle and Liverpool University. AM is supported by the Agence

Nationale de la Recherche (ANR) (DNA-Life) and the European Research Council

(EPIncRNA starting grant, DARK consolidator grant). This work has received support from

the U.S. National Institutes of Health HG004160 to BFP, and from the ANR (AAPG2019

PICen to L.T., ANR-18-CE12-0026 and ANR-20-CE12-0014-02 to D.D.).

Acknowledgements

We thank D. Shapira (Newcastle University) for assisting with experiments and constructive

discussions. We thank S. Baulande and P. Legoix-Né (NGS platform, Institute Curie). We

also thank Maxime Wéry and Nicolas Vogt for outstanding help with the NET-seq and RNA-

seq assays and critical reading of the manuscript, David A. Brow for the generous gift of the

pGAC24 and pGAC24-SNR13TTS plasmids and the *nrd1-5* yeast strain, and Kuangyu Yen,

Lisa Prendergast and members of the Papamichos-Chronakis and Morillon's labs for

constructive discussions. This work has benefited from the facilities and expertise of the NGS

platform of Institute Curie, supported by the Agence Nationale de la Recherche (ANR-10-

EQPX-03, ANR10-INBS-09-08) and the Canceropôle Ile-de-France.

References

1.   Porrua, O. & Libri, D. RNA quality control in the nucleus: the Angels' share of RNA.
Biochim Biophys Acta **1829**, 604-11 (2013).
2.   Pelechano, V., Chavez, S. & Perez-Ortin, J.E. A complete set of nascent transcription
rates for yeast genes. *PLoS One* **5**, e15442 (2010).
3.   Kim, T.S. et al. RNA polymerase mapping during stress responses reveals widespread
nonproductive transcription in yeast. *Genome Biol* **11**, R75 (2010).
4.   Tuck, A.C. & Tollervey, D. A transcriptome-wide atlas of RNP composition reveals
diverse classes of mRNAs and lncRNAs. *Cell* **154**, 996-1009 (2013).
5.   Bresson, S. & Tollervey, D. Surveillance-ready transcription: nuclear RNA decay as a
default fate. *Open Biol* **8**(2018).
6.   Steurer, B. et al. Live-cell analysis of endogenous GFP-RPB1 uncovers rapid turnover
of initiating and promoter-paused RNA Polymerase II. *Proc Natl Acad Sci U S A* **115**,
E4368-E4376 (2018).
7.   Darzacq, X. et al. In vivo dynamics of RNA polymerase II transcription. *Nat Struct Mol*
Biol **14**, 796-806 (2007).
8.   Steinmetz, E.J., Conrad, N.K., Brow, D.A. & Corden, J.L. RNA-binding protein Nrd1
directs poly(A)-independent 3'-end formation of RNA polymerase II transcripts.
Nature **413**, 327-31 (2001).
9.   Kilchert, C., Wittmann, S. & Vasiljeva, L. The regulation and functions of the nuclear
RNA exosome complex. *Nat Rev Mol Cell Biol* **17**, 227-39 (2016).
10. Jensen, T.H., Jacquier, A. & Libri, D. Dealing with pervasive transcription. *Mol Cell* **52**,
473-84 (2013).
11. Mayer, A. et al. CTD tyrosine phosphorylation impairs termination factor recruitment
to RNA polymerase II. *Science* **336**, 1723-5 (2012).
12. Gonzalez-Aguilera, C. et al. Nab2 functions in the metabolism of RNA driven by
polymerases II and III. *Mol Biol Cell* **22**, 2729-40 (2011).
13. Yu, M.C. et al. Arginine methyltransferase affects interactions and recruitment of
mRNA processing and export factors. *Genes Dev* **18**, 2024-35 (2004).
14. Bresson, S., Tuck, A., Staneva, D. & Tollervey, D. Nuclear RNA Decay Pathways Aid
Rapid Remodeling of Gene Expression in Yeast. *Mol Cell* **65**, 787-800 e5 (2017).
15. Webb, S., Hector, R.D., Kudla, G. & Granneman, S. PAR-CLIP data indicate that Nrd1-
Nab3-dependent transcription termination regulates expression of hundreds of
protein coding genes in yeast. *Genome Biol* **15**, R8 (2014).
16. Schulz, D. et al. Transcriptome surveillance by selective termination of noncoding
RNA synthesis. *Cell* **155**, 1075-87 (2013).
17. Kapranov, P. et al. RNA maps reveal new RNA classes and a possible function for
pervasive transcription. *Science* **316**, 1484-8 (2007).
18. Yu, D., Ma, X., Zuo, Z., Wang, H. & Meng, Y. Classification of Transcription Boundary-
Associated RNAs (TBARs) in Animals and Plants. *Front Genet* **9**, 168 (2018).
19. Chiu, A.C. et al. Transcriptional Pause Sites Delineate Stable Nucleosome-Associated
Premature Polyadenylation Suppressed by U1 snRNP. *Mol Cell* **69**, 648-663 e7
(2018).
20. Erickson, B., Sheridan, R.M., Cortazar, M. & Bentley, D.L. Dynamic turnover of
paused Pol II complexes at human promoters. *Genes Dev* (2018).

21. Krebs, A.R. et al. Genome-wide Single-Molecule Footprinting Reveals High RNA
Polymerase II Turnover at Paused Promoters. *Mol Cell* **67**, 411-422 e4 (2017).
22. Gressel, S. et al. CDK9-dependent RNA polymerase II pausing controls transcription
initiation. *Elife* **6**(2017).
23. Shao, W. & Zeitlinger, J. Paused RNA polymerase II inhibits new transcriptional
initiation. *Nat Genet* **49**, 1045-1051 (2017).
24. Chen, F.X., Smith, E.R. & Shilatifard, A. Born to run: control of transcription
elongation by RNA polymerase II. *Nat Rev Mol Cell Biol* (2018).
25. Taft, R.J., Kaplan, C.D., Simons, C. & Mattick, J.S. Evolution, biogenesis and function
of promoter-associated RNAs. *Cell Cycle* **8**, 2332-8 (2009).
26. Whitehouse, I., Rando, O.J., Delrow, J. & Tsukiyama, T. Chromatin remodelling at
promoters suppresses antisense transcription. *Nature* **450**, 1031-5 (2007).
27. Boeger, H., Griesenbeck, J. & Kornberg, R.D. Nucleosome retention and the
stochastic nature of promoter chromatin remodeling for transcription. *Cell* **133**, 716-
26 (2008).
28. Smolle, M. et al. Chromatin remodelers Isw1 and Chd1 maintain chromatin structure
during transcription by preventing histone exchange. *Nat Struct Mol Biol* **19**, 884-92
(2012).
29. Morillon, A. et al. Isw1 chromatin remodeling ATPase coordinates transcription
elongation and termination by RNA polymerase II. *Cell* **115**, 425-35 (2003).
30. Babour, A. et al. The Chromatin Remodeler ISW1 Is a Quality Control Factor that
Surveys Nuclear mRNP Biogenesis. *Cell* **167**, 1201-1214 e15 (2016).
31. Poli, J., Gasser, S.M. & Papamichos-Chronakis, M. The INO80 remodeller in
transcription, replication and repair. *Philos Trans R Soc Lond B Biol Sci* **372**(2017).
32. Xue, Y. et al. Mot1, Ino80C, and NC2 Function Coordinately to Regulate Pervasive
Transcription in Yeast and Mammals. *Mol Cell* **67**, 594-607 e4 (2017).
33. Yen, K., Vinayachandran, V. & Pugh, B.F. SWR-C and INO80 chromatin remodelers
recognize nucleosome-free regions near +1 nucleosomes. *Cell* **154**, 1246-56 (2013).
34. Poli, J. et al. Mec1, INO80, and the PAF1 complex cooperate to limit transcription
replication conflicts through RNAPII removal during replication stress. *Genes Dev* **30**,
337-54 (2016).
35. Lafon, A. et al. INO80 Chromatin Remodeler Facilitates Release of RNA Polymerase II
from Chromatin for Ubiquitin-Mediated Proteasomal Degradation. *Mol Cell* **60**, 784-
96 (2015).
36. Tsvetanova, N.G., Klass, D.M., Salzman, J. & Brown, P.O. Proteome-wide search
reveals unexpected RNA-binding proteins in Saccharomyces cerevisiae. *PLoS One*
5(2010).
37. Edupuganti, R.R. et al. N(6)-methyladenosine (m(6)A) recruits and repels proteins to
regulate mRNA homeostasis. *Nat Struct Mol Biol* **24**, 870-878 (2017).
38. Churchman, L.S. & Weissman, J.S. Nascent transcript sequencing visualizes
transcription at nucleotide resolution. *Nature* **469**, 368-73 (2011).
39. Wissink, E.M., Vihervaara, A., Tippens, N.D. & Lis, J.T. Nascent RNA analyses: tracking
transcription and its regulation. *Nat Rev Genet* (2019).
40. Harlen, K.M. & Churchman, L.S. Subgenic Pol II interactomes identify region-specific
transcription elongation regulators. *Mol Syst Biol* **13**, 900 (2017).

41. Yen, K., Vinayachandran, V., Batta, K., Koerber, R.T. & Pugh, B.F. Genome-wide
nucleosome specificity and directionality of chromatin remodelers. *Cell* **149**, 1461-73
(2012).
42. Rhee, H.S. & Pugh, B.F. ChIP-exo method for identifying genomic location of DNA-
binding proteins with near-single-nucleotide accuracy. *Curr Protoc Mol Biol* **Chapter**
21, Unit 21 24 (2012).
43. Shaw, R.J. & Reines, D. Saccharomyces cerevisiae transcription elongation mutants
are defective in PUR5 induction in response to nucleotide depletion. *Mol Cell Biol* **20**,
7427-37 (2000).
44. Ehrensberger, A.H., Kelly, G.P. & Svejstrup, J.Q. Mechanistic interpretation of
promoter-proximal peaks and RNAPII density maps. *Cell* **154**, 713-5 (2013).
45. Baptista, T. et al. SAGA Is a General Cofactor for RNA Polymerase II Transcription.
Mol Cell **68**, 130-143 e5 (2017).
46. Timmers, H.T.M. & Tora, L. Transcript Buffering: A Balancing Act between mRNA
Synthesis and mRNA Degradation. *Mol Cell* **72**, 10-17 (2018).
47. Sun, M. et al. Comparative dynamic transcriptome analysis (cDTA) reveals mutual
feedback between mRNA synthesis and degradation. *Genome Res* **22**, 1350-9 (2012).
48. Malagon, F. et al. Mutations in the Saccharomyces cerevisiae RPB1 gene conferring
hypersensitivity to 6-azauracil. *Genetics* **172**, 2201-9 (2006).
49. Pelechano, V. & Perez-Ortin, J.E. The transcriptional inhibitor thiolutin blocks mRNA
degradation in yeast. *Yeast* **25**, 85-92 (2008).
50. Alpert, T., Straube, K., Carrillo Oesterreich, F., Herzel, L. & Neugebauer, K.M.
Widespread Transcriptional Readthrough Caused by Nab2 Depletion Leads to
Chimeric Transcripts with Retained Introns. *Cell Rep* **33**, 108324 (2020).
51. Aguilar, L.C. et al. Altered rRNA processing disrupts nuclear RNA homeostasis via
competition for the poly(A)-binding protein Nab2. *Nucleic Acids Res* **48**, 11675-11694
(2020).
52. Fasken, M.B., Laribee, R.N. & Corbett, A.H. Nab3 facilitates the function of the
TRAMP complex in RNA processing via recruitment of Rrp6 independent of Nrd1.
PLoS Genet **11**, e1005044 (2015).
53. Fox, M.J. & Mosley, A.L. Rrp6: Integrated roles in nuclear RNA metabolism and
transcription termination. *Wiley Interdiscip Rev RNA* **7**, 91-104 (2016).
54. Vasiljeva, L. & Buratowski, S. Nrd1 interacts with the nuclear exosome for 3'
processing of RNA polymerase II transcripts. *Mol Cell* **21**, 239-48 (2006).
55. Tudek, A. et al. Molecular basis for coordinating transcription termination with
noncoding RNA degradation. *Mol Cell* **55**, 467-81 (2014).
56. Holmes, R.K. et al. Loss of the Yeast SR Protein Npl3 Alters Gene Expression Due to
Transcription Readthrough. *PLoS Genet* **11**, e1005735 (2015).
57. Schwer, B., Mao, X. & Shuman, S. Accelerated mRNA decay in conditional mutants of
yeast mRNA capping enzyme. *Nucleic Acids Res* **26**, 2050-7 (1998).
58. Breslow, D.K. et al. A comprehensive strategy enabling high-resolution functional
analysis of the yeast genome. *Nat Methods* **5**, 711-8 (2008).
59. Arigo, J.T., Carroll, K.L., Ames, J.M. & Corden, J.L. Regulation of yeast NRD1
expression by premature transcription termination. *Mol Cell* **21**, 641-51 (2006).
60. Kim, M. et al. Distinct pathways for snoRNA and mRNA termination. *Mol Cell* **24**, 723-
734 (2006).

61.   Fox, M.J., Gao, H., Smith-Kinnaman, W.R., Liu, Y. & Mosley, A.L. The exosome
component Rrp6 is required for RNA polymerase II termination at specific targets of
the Nrd1-Nab3 pathway. *PLoS Genet* **11**, e1004999 (2015).
62.   van Dijk, E.L. et al. XUTs are a class of Xrn1-sensitive antisense regulatory non-coding
RNA in yeast. *Nature* **475**, 114-7 (2011).
63.   Albert, I. et al. Translational and rotational settings of H2A.Z nucleosomes across the
Saccharomyces cerevisiae genome. *Nature* **446**, 572-6 (2007).
64.   Zofall, M. et al. Histone H2A.Z cooperates with RNAi and heterochromatin factors to
suppress antisense RNAs. *Nature* **461**, 419-22 (2009).
65.   Wagner, E.J. et al. A genome-wide RNA interference screen reveals that variant
histones are necessary for replication-dependent histone pre-mRNA processing. *Mol*
Cell **28**, 692-9 (2007).
66.   Muniz, L. et al. Control of Gene Expression in Senescence through Transcriptional
Read-Through of Convergent Protein-Coding Genes. *Cell Rep* **21**, 2433-2446 (2017).
67.   Papamichos-Chronakis, M., Watanabe, S., Rando, O.J. & Peterson, C.L. Global
regulation of H2A.Z localization by the INO80 chromatin-remodeling enzyme is
essential for genome integrity. *Cell* **144**, 200-13 (2011).
68.   Eustermann, S. et al. Structural basis for ATP-dependent chromatin remodelling by
the INO80 complex. *Nature* **556**, 386-390 (2018).
69.   Brahma, S. et al. INO80 exchanges H2A.Z for H2A by translocating on DNA proximal
to histone dimers. *Nat Commun* **8**, 15616 (2017).
70.   Dheur, S., Nykamp, K.R., Viphakone, N., Swanson, M.S. & Minvielle-Sebastia, L. Yeast
mRNA Poly(A) tail length control can be reconstituted in vitro in the absence of
Pab1p-dependent Poly(A) nuclease activity. *J Biol Chem* **280**, 24532-8 (2005).
71.   Kuehner, J.N. & Brow, D.A. Regulation of a eukaryotic gene by GTP-dependent start
site selection and transcription attenuation. *Mol Cell* **31**, 201-11 (2008).
72.   Roth, K.M., Wolf, M.K., Rossi, M. & Butler, J.S. The nuclear exosome contributes to
autogenous control of NAB2 mRNA levels. *Mol Cell Biol* **25**, 1577-85 (2005).
73.   Rahl, P.B. et al. c-Myc regulates transcriptional pause release. *Cell* **141**, 432-45
(2010).
74.   Zhang, X. et al. Attenuation of RNA polymerase II pausing mitigates BRCA1-
associated R-loop accumulation and tumorigenesis. *Nat Commun* **8**, 15908 (2017).
75.   Aguilera, A. & Gomez-Gonzalez, B. Genome instability: a mechanistic view of its
causes and consequences. *Nat Rev Genet* **9**, 204-17 (2008).
76.   Kotsantis, P., Petermann, E. & Boulton, S.J. Mechanisms of Oncogene-Induced
Replication Stress: Jigsaw Falling into Place. *Cancer Discov* **8**, 537-555 (2018).
77.   Zhou, B. et al. INO80 governs superenhancer-mediated oncogenic transcription and
tumor growth in melanoma. *Genes Dev* **30**, 1440-53 (2016).
78.   Zhang, S. et al. INO80 is required for oncogenic transcription and tumor growth in
non-small cell lung cancer. *Oncogene* **36**, 1430-1439 (2017).
79.   Hemming, S.A. et al. RNA polymerase II subunit Rpb9 regulates transcription
elongation in vivo. *J Biol Chem* **275**, 35506-11 (2000).
80.   Riles, L., Shaw, R.J., Johnston, M. & Reines, D. Large-scale screening of yeast mutants
for sensitivity to the IMP dehydrogenase inhibitor 6-azauracil. *Yeast* **21**, 241-8
(2004).

81. Creamer, T.J. et al. Transcriptome-wide binding sites for components of the
Saccharomyces cerevisiae non-poly(A) termination pathway: Nrd1, Nab3, and Sen1.
PLoS Genet **7**, e1002329 (2011).
82. Jonsson, Z.O., Jha, S., Wohlschlegel, J.A. & Dutta, A. Rvb1p/Rvb2p recruit Arp5p and
assemble a functional Ino80 chromatin remodeling complex. *Mol Cell* **16**, 465-77
(2004).
83. Wery, M. et al. Nonsense-Mediated Decay Restricts LncRNA Levels in Yeast Unless
Blocked by Double-Stranded RNA Structure. *Mol Cell* **61**, 379-392 (2016).
84. Wery, M. et al. Native elongating transcript sequencing reveals global anti-
correlation between sense and antisense nascent transcription in fission yeast. *RNA*
24, 196-208 (2018).
85. Baptista, T. & Devys, D. Saccharomyces cerevisiae Metabolic Labeling with 4-
thiouracil and the Quantification of Newly Synthesized mRNA As a Proxy for RNA
Polymerase II Activity. *J Vis Exp* (2018).
86. Papamichos-Chronakis, M. & Peterson, C.L. The Ino80 chromatin-remodeling enzyme
regulates replisome function and stability. *Nat Struct Mol Biol* **15**, 338-45 (2008).
87. Rhee, H.S., Bataille, A.R., Zhang, L. & Pugh, B.F. Subnucleosomal structures and
nucleosome asymmetry across a genome. *Cell* **159**, 1377-88 (2014).
596

Fig. 1. Accumulation of nascent mRNA transcripts on chromatin proximally to

promoters and defective transcriptional elongation in the absence of INO80.

a, Metagene analysis for NET-seq density profiles averaged across scaled mRNA genes (n=

5432) in WT and *ino80Δ*. Tags were aligned to both Transcription Start Site (TSS) and

Polyadenylation Site (PAS) for each gene. n=2. **b,** Top: Schematic representation and

formula for calculating the Traveling Ratio (TR) from NET-seq mRNA densities. Bottom:

Cumulative distribution plot of Traveling Ratio for mRNA genes from a (n=5432). Genes

with TR>2 (WT: n=354, *ino80Δ*: n=314) are not shown. ****p < 0.0XXX with K-S test. **c,**

Scatterplot analysis comparing Ino80[33] and Rpb1 ChIP-exo densities between the

transcription start site and polyadenylation site (TSS to PAS) in mRNA yeast genes

(n=5798). Pearson's correlation coefficient (r) and p-value are indicated. **d,** Cell proliferation

analysis for WT and *ino80Δ* strains grown exponentially in SC-ura and in SC-ura containing

50 µg/ml 6-azauracil (6-AU) liquid media for the indicated time. Cell density was measured

at OD^{620} and plotted on a $\log_2$ scale. Doubling time was calculated from OD^{620}: 0.2 to OD^{620}:

0.4 for all strains in both conditions. Fold Change ratio (FC) of doubling time in 6-AU over

SC-ura for each strain is shown. n=4.

Fig. 3. mRNA synthesis and decay rates are reduced in the absence of INO80.

a, b, Scatterplot analysis for fold changes between WT and *ino80Δ* in newly synthesized

mRNA (a) and steady-state mRNA (b) levels (n=5012 genes) plotted on a $\log_2$ scale.

Thresholds of 1.5-fold change and 0.05 p values were considered for significant change

(green dots). n=3. **c,** cDTA profiles for *ino80Δ*. For all analysed genes, changes in synthesis

rates (SR) were plotted against the changes in mRNA decay rates (DR). Fold-change in SR

and DR were calculated between *ino80Δ* and WT and plotted on a $\log_2$ scale. 90% of genes

are contained within the outer contour. Yellow and red dots correspond to 60% of genes. **d,**

RT-qPCR analysis for *NRD1* and *RNA1* RNAs was conducted in WT and *ino80Δ* before (T_0)

and after addition of thiolutin in YPD. RNA abundance at each time-point was normalized

626 over the respective abundance of the *SCR1* RNA. RNA before thiolutin treatment was set at

627 100% in both strains. Remaining RNA was calculated as the amount of normalized RNA in

628 the indicated time-points relative to the normalized RNA at T_0. Values represent the average

629 from at least three independent experiments. Bars, standard errors.

Fig. 4. Loading of the RNA surveillance and termination factors Mtr4, Nab3 and Nab2

on promoter-proximal mRNA transcripts.

a, Formula for calculating Transcript Instability (TI) as the ratio of the averaged CRAC

densities of Nab3[56], Mtr4 or Nab2[4] to CRAC densities of Rpb1[56] for mRNA genes. **b,**

Schematic table for clustering mRNA genes according to the relative TR in WT (TR[WT]). The

number of genes and the level of promoter-proximal pausing are indicated for each cluster. **c,**

Boxplot analysis comparing TIs calculated for the promoter-proximal and gene body regions

of mRNA genes in the four clusters defined by their relative TR[WT]. p-values were calculated

by Wilcoxon rank-sum test. ****, p<0.0001. ***, p<0.001. **, p<0.01. ns, non-significant.

Fig. 5. Genetic and physical interactions of INO80 with the RNA quality control pathway.

a, 5-fold serial dilution of exponentially grown cells from the indicated strains were plated onto YPD and incubated at 30^{o}C (permissive temperature) or 37^{o}C (non-permissive temperature) for 4-5 days. **b,** 5-fold serial dilution of exponentially grown cells from the indicated strains were plated onto both SC-ura and SC-ura + 50μg/ml 6-Azauracil (6-AU) and incubated at 30^{o}C for 4-5 days. **c,** RT-qPCR analysis for *INO1* RNA was conducted in the indicated strains grown exponentially in SC media (0h) or SC without inositol for 14h. *INO1* RNA was normalized over *ACT1* RNA. Bars, standard errors (n=3). Asterisks indicate statistical significance of the indicated change as calculated by Tukey's multiple comparisons test. ns: non-significant. **d,** Nucleic acid-free lysates from cells co-expressing Nrd1-HA with Ino80 either untagged or tagged with FLAG epitope were subjected to FLAG-IP. Inputs and IP samples were immunoblotted for FLAG, Arp5, HA. Immunoblot against Pgk1 served as a control. **E,** Nucleic acid-free lysates from cells expressing Nab2-HA were subjected to either

656   HA-IP (Nab2-HA) or IgG-IP (mock). Inputs and IP samples were immunoblotted for Arp5

657   and HA.

a

WT
ino80Δ
SC-leu

WT
ino80Δ
0.25mM CuSO4

b

$CUP1/ACT^*$ Fold Change over WT

nrd1-5 2.8 *
arp8Δ 3.5 **
ino80Δ 3.0 *

c

RNA-seq density (log$_2$)

— WT, 5'-TTS-3' RNA
— ino80Δ, 5'-TTS-3' RNA
--- WT, total RNA
--- ino80Δ, total RNA

541800   541500   541300   Chr.IV, - strand

TSS TTS

SNR47

d

$$\text{Read-through Index (RI)} = \frac{\text{RNA-seq}^{(5'\text{-TTS-3'})} \text{ density}_{after\ TTS}}{\text{RNA-seq}^{(5'\text{-TTS-3'})} \text{ density}_{before\ TTS}}$$

RI

****

WT   ino80Δ

RI

ns

ns   *

WT   rrp6Δ   xrn1Δ

e

NRD1 mRNA

m⁷G — AAAAAA

m⁷G — AAAAAA
— AAAAAA

— AAAAAA

NRD1 5'
3'RACE product

NRD1 3'
3'RACE product

3' RACE NRD1 5'/NRD1 3' Fold Change over WT (log2)

rrp6Δ 0.35
ino80Δ -1.12 *
arp8Δ -0.90 *

Fig. 6. NNS-dependent transcriptional termination is defective in *ino80Δ*.

a, 5-fold serial dilution of cells from the indicated strains were plated onto both SC-leu and

SC-leu + 0.25mM CuSO4 and incubated at 30°C for 3 (SC-leu) and 7 days (CuSO4). **b,** RT-

qPCR analysis was conducted in WT, *nrd1-5*, *arp8Δ* and *ino80Δ* cells transformed with

either *ACT-CUP* or *ACT-SNR^{TTS}-CUP* plasmids. The *CUP1/ACT** ratio from the *ACT-*

SNR^{TTS}-CUP plasmid was normalized over the same ratio from the control *ACT-CUP*

plasmid (Extended Fig. 4c) to correct for differential transcription across all strains. Read-

through at the *SNR13^{TTS}* for each indicated strain is shown as Fold Change Ratio over WT.

Bars, standard errors from three independent experiments. p-values were calculated by t-test

after testing for normal distribution by Shapiro-Wilk test. *, p<0.05. **, p<0.01. **c,** Snapshot

of RNA-seq densities in WT and *ino80Δ* across the *SNR47* snoRNA gene. Solid lines, density

of reads encompassing the transcription termination site (5'-TTS-3') of the *SNR47* gene.

Broken lines, total RNA-seq density. Reads encompassing the TTS were selected from the

total RNA-seq reads as containing 5' upstream the *SNR47* TTS and 3' at or after the *SNR47*

TTS. Genomic coordinates of Chromosome IV around the *SNR47* gene (coordinates: 541700

to 541602) are shown. The closest downstream gene is *YDR042C* (TSS: Chromosome IV,

541203). n=4. **d,** Upper panel, formula for calculating the Read-through Index (RI) from

RNA-seq densities across snoRNA genes that encompass transcription termination sites (5'-

TTS-3'). RNA-seq reads with 5' upstream the TTS and 3' at or after the TTS were selected

from the total RNA-seq read pool and the ratio of the RNA-seq read density after TTS to the

RNA-seq read density before TTS in the selected reads was calculated. Lower panel, boxplot

analysis comparing RI from the indicated strains across snoRNA genes. p-values were

calculated by Wilcoxon matched-pairs rank test after testing for normal distribution. ****,

p<0.0001. *, p<0.05. ns, non-significant. **e,** Top: Scheme illustrating nested 3'RACE for

termination events at the promoter-proximal (5') and PAS (3') regions of the *NRD1* gene

(*NRD1 5'*, dark blue). Bottom: 3'RACE analysis for *NRD1* termination events was conducted

in WT and the indicated mutant strains. The relative enrichment of polyadenylated promoter-

proximal to PAS-terminated transcripts (5'/3') was calculated for all strains and the fold

change in each mutant compared to WT was plotted on a $\log_2$ scale. n=3. p-values were

calculated by t-test after testing for normal distribution by Shapiro-Wilk test. *, p<0.05.

Fig. 7. INO80 promotes the co-transcriptional recruitment of Nab2 to chromatin and its association with H2A.Z.

a, Chromatin immunoprecipitation for Nab2-HA and Rpb1 in WT and *ino80Δ* cells. Fold enrichment of each protein at the promoter-proximal region of *NRD1* gene (*NRD1 5'*) (upper panel, shaded in dark blue) was calculated relative to the *GAL1* promoter after correction for input DNA levels. ChIP ratio at the promoter-proximal region of *NRD1* gene (*NRD1 5'*) was calculated as Fold enrichment of Nab2-HA over Fold enrichment of Rpb1. Bars, standard

errors from four independent experiments. p-values were calculated by t-test after testing for

normal distribution by Shapiro-Wilk test. *, p<0.05. **b,** RNA immunoprecipitation for Nab2-

HA in WT and *ino80Δ* cells. Ratio of binding of Nab2 to the promoter-proximal *NRD1* RNA

(upper panel, *NRD1 5'*, dark blue) over binding to the 3' region of *NRD1* RNA (upper panel,

NRD1 3', light blue) after correction for input (total) RNA levels in the indicated strains.

Bars, standard errors from four independent experiments. p-values were calculated by t-test

after testing for normal distribution by Shapiro-Wilk test. ***, p<0.001. **c,** Cell proliferation

analysis for the indicated strains transformed with the *ACT-SNR^{TTS}-CUP* plasmid grown

exponentially in SC-ura (left panel) and in SC-ura liquid media containing 0.4mM CuSO$_4$ (6-

AU) (right panel) for the indicated time. Cell density was measured at OD^{620} and is shown on

a log$_2$ scale. **d,** Nested 3'RACE analysis for termination events at the *NRD1* gene was

conducted in WT and *htz1Δ*. Analysis was conducted as in Fig. 6e. n=4. p-values were

calculated by t-test after testing for normal distribution by Shapiro-Wilk test. **, p<0.01. **e,**

Chromatin immunoprecipitation for Nab2-HA and Rpb1 in WT and *htz1Δ* cells. Analysis was

conducted as in (a). Bars, standard errors from three independent experiments. p-values were

calculated by t-test after testing for normal distribution by Shapiro-Wilk test. **, p<0.01. **f,**

Representative H2A.Z immunoprecipitation experiment. Nucleic acid-free lysates from WT

and *ino80Δ* cells expressing Nab2-HA were subjected to H2A.Z-IP. Inputs and IP samples

were immunoblotted for H2A.Z and HA. Nab2-HA enrichment in H2A.Z pulldowns is

calculated as HA/H2A.Z IP ratio, corrected for the relative Nab2-HA abundance in the input

samples. Nab2-HA enrichment in WT is set arbitrarily to 1.0. All analysis was conducted

from the same SDS-page gel.

Extended Data Fig. 1. NET-seq and 6-AU cell growth analysis in the absence of INO80.

a, Heatmap of Pearson's correlation coefficients (R) for all pairwise combinations of NET-seq reads in independent biological replicates. **b,** Rpb1-FLAG is efficiently immunoprecipitated from yeast cells during NET-seq analysis. Lysates from WT and *ino80Δ* cells expressing *RPB1-FLAG* were subjected to FLAG immunoprecipitation using FLAG beads. Rpb1-FLAG was subsequently eluted from the beads with FLAG peptide. Input,

eluate and unbound samples were immunoblotted against FLAG epitope and total Rpb1.

Immunoblotting against Pgk1 serves as loading control. All samples were resolved in the

same SDS-page gel. **c,** Scatterplot analysis comparing the changes in NET-seq densities

between WT and *ino80Δ* for sense transcripts and their relative antisense transcripts over

mRNA gene regions, plotted on a $\log_2$ scale. Pearson's correlation coefficient (r) is indicated.

d, Cell proliferation analysis for the indicated strains grown exponentially in SC-ura (top

panel) and in SC-ura containing 50µg/ml 6-azauracil (6-AU) (bottom panel) liquid media for

the indicated time. Cell density was measured at OD^{620} and shown on a $\log_2$ scale. Shading

surrounding each line denotes standard error from four independent experiments. The mutant

strain deleted for the RNA Polymerase II subunit Rpb9 (*rpb9Δ*) has reported transcriptional

elongation defects in 6-AU[79] and was used as control. Doubling time from OD^{620} 0.2 to

OD^{620} 0.4 was calculated for all strains in both conditions. Fold Change ratio (FC) of

doubling time in 6-AU over SC-ura is shown (bottom panel). **e,** 5-fold serial dilution of cells

from the indicated strains were plated onto both SC-ura and SC-ura + 50µg/ml 6-Azauracil

(6-AU) and incubated at 30°C for 4-5 days. Genetic backgrounds (LS20, W303) are

indicated. The mutant strain deleted for the Dst1/TFIIS transcription elongation factor (*dst1Δ*)

was used as control[80].

Extended Data Fig. 3. Ectopic expression of INO80 rescues the nascent mRNA synthesis

defect of *ino80Δ* cells.

a, Scatterplot analysis for fold changes in newly synthesized mRNA between *ino80Δ* cells

expressing a WT allele of *INO80* from the pRS416 plasmid (*pINO80*) and WT cells

transformed with the empty vector pRS416 (EV) plotted on a $\log_2$ scale. Analysis as in

Fig.3a. **b,** Fold changes in mRNA decay rates between *ino80Δ* and WT for the indicated

genes as calculated by cDTA analysis. **c,** RT-qPCR analysis for *HPT1* RNA was conducted

in WT and *ino80Δ* before (T_0) and after addition of thiolutin in YPD as in Fig. 3d.

a

- Ino80-FLAG

- Arp5

- Pcf11-HA

Input   IP: FLAG

Extended Data Fig. 4. INO80 does not interact with Pcf11.

a, Nucleic acid-free lysates from cells co-expressing Pcf11-HA with Ino80 either untagged or tagged with FLAG epitope were subjected to FLAG-IP. Inputs and IP samples were immunoblotted for FLAG, Arp5 and HA. Samples were resolved in the same SDS-page gel.

Extended Data Fig. 5. Defective NNS-dependent transcription termination in *ino80Δ*

cells.

a, Schematic representation of the *ACT-CUP* (pGAC24) and *ACT-SNR^TTS-CUP* (pGAC24-

SNR13^TTS) reporter systems for transcription termination reported in[8]. The regions amplified

by RT-qPCR for *ACT** and *CUP1* are underlined. The region of the *SNR13* gene between

+125 to +232bp from TSS that was cloned in the *ACT-SNR13^TTS-CUP* plasmid[8] is shaded in

blue (*SNR13**). The expected phenotype of cells transformed with either the *ACT-CUP* or the

ACT-SNR13^TTS-CUP plasmid is shown. **b,** Snapshot of Nrd1 PAR-CLIP density profile

from[81] and Ino80 ChIP-exo density profile from [33] across the *SNR13* snoRNA gene. Tags

were aligned relative to TSS and plotted on a log_2 scale. The region between +125 to +232bp

from TSS cloned in the *ACT-SNR13^TTS-CUP* plasmid described in (a)[8] is highlighted in blue

(*SNR13**). **c,** RT-qPCR analysis for the *CUP1*/*ACT** ratio was conducted in the indicated

strains transformed with either the *ACT-CUP* (left panel) or *ACT-SNR^TTS-CUP* (right panel)

plasmids. Bars, standard errors from three independent experiments. Asterisks indicate

statistical significance of the change in the respective mutant compared to the WT as

calculated by t-test. ns: non-significant. *, $p<0.05$. ***, $p<0.001$. **d,** RT-qPCR analysis was

conducted in both WT and the *ino80-td* degron-inducible strain (*ino80-td*) [82] transformed

with the *ACT-CUP* control plasmid described in a. Galactose-induced activation for 3 hours

at 37°C (+) and glucose-induced suppression (–) of the *UBR1* N-recognin in the two strains is

indicated. Bars, standard errors from four independent experiments. p-values were calculated

by t-test after testing for normal distribution by Shapiro-Wilk test. ns, non-significant. **e,**

Snapshot representing RNA-seq densities in WT and *ino80Δ* across the ribosomal protein

coding gene YNL096C. Solid lines, density of reads encompassing the transcription

termination site (5'-TTS-3') as described in Fig. 6c. Analysis was performed around the

polyadenylation site (PAS). Broken lines, total RNA-seq density. Genomic coordinates of

Chromosome XVI around the *YNL096C* gene are shown. The closest gene downstream of

YNL096C is the *PHO23* mRNA gene (TSS: Chromosome XIV, 442509). **f,** Snapshot

representing ChIP-exo densities of Ino80 from [41] across the *NRD1* gene. **g,** Snapshot

representing NET-seq densities in WT and *ino80Δ* across the *NRD1* gene. **h,** Representative

image of nested 3'RACE analysis of termination events at the *NRD1* 5' and 3' regions

according to the scheme in Figure 6e and at the 3' of *ACT1* gene in WT and *rrp6Δ* on an

agarose gel. Boxes indicate the most frequent termination sites. Ladder is 100bp. *ACT1*

serves as loading control.

Extended Data Fig. 6. Co-transcriptional recruitment of Nab2 to chromatin depends on

INO80 and H2A.Z.

a, Snapshot representing CRAC densities for Nab2 from [4] across the *NRD1* gene. **b,**

Chromatin immunoprecipitation for Rpb1 and Nab2-HA in WT and *ino80Δ* cells. Fold

enrichment analysis at the promoter-proximal region of *NRD1* gene (*NRD1 5'*) (upper panel,

shaded in dark blue) was conducted as Fig. 7a. **c,** Chromatin immunoprecipitation for Nab2-

HA and Rpb1 in WT and *ino80Δ* cells. Fold enrichment of each protein at the TTS region of

SNR13 gene was calculated relative to the *GAL1* promoter after correction for input DNA

levels. ChIP ratio at the TTS region of *SNR13* gene was calculated as Fold enrichment of

Nab2-HA over Fold enrichment of Rpb1. Bars, standard errors from four independent

experiments. p-values were calculated by t-test after testing for normal distribution by

Shapiro-Wilk test. *, $p<0.05$. **d,** Nucleic acid-free lysates from WT and *ino80Δ* cells both

expressing Nab2-HA were subjected to HA-IP. Inputs and IP samples were immunoblotted

for Rpb1 and HA. Mock, WT protein lysate subjected to immunoprecipitation with a control

IgG antibody. **e,** Metagene analysis for ChIP-exo density profiles of histone H3 in WT and

ino80Δ aligned at the A nucleotide of the ATG sites in the TSS to TSS+200bp (Promoter-

proximal) and the TSS+200 to PAS (Gene body) regions. n=Number of ATG sites in each

region. **f,** Cell proliferation analysis for the indicated strains transformed with the *ACT-CUP*

plasmid grown exponentially in SC-ura (left panel) and in SC-ura liquid media containing

0.4mM $CuSO_4$ (6-AU) (right panel) for the indicated time. Cell density was measured at

OD^{620} and is shown on a $\log_2$ scale. **g,** Chromatin immunoprecipitation for Rpb1 and Nab2-

HA in WT and *htz1Δ* cells. Fold enrichment analysis at the promoter-proximal region of

NRD1 gene (*NRD1 5'*) was conducted as Fig. 7a. **h,** Nucleic acid-free lysates from WT cells

expressing Nab2-HA were subjected to immunoprecipitation with either an anti-H2A.Z

antibody (+) or a non-specific IgG antibody as a control (–), then washed in the presence of

300mM NaCl. Inputs and IP samples were immunoblotted for HA and H2A.Z. All analysis

was conducted from the same SDS-page gel. **I,** Nucleic acid-free lysates from cells

expressing Pcf11-HA were subjected to immunoprecipitation with either an anti-H2A.Z

antibody (+) or a non-specific IgG antibody as a control (–). Inputs and IP samples were

immunoblotted for HA, Arp5 and H2A.Z. The western for Pcf11-HA was over-exposed in

order to detect the background levels of Pcf11 binding to the beads. All analysis was

conducted from the same SDS-page gel.

 Extended Data Table 1 - Yeast strains used in this study

Name	Genotype	Source
YM50	LS20 [*MatΔ ade2 can1r cyh2r lys S URA3-52 trp1Δ his3-Δ200GAL:HO leu 2Δ*]	Sandell & Zakian, *Cell* 1993
YM85	LS20 *ino80Δ::KANMX*	Papamichos-Chronakis & Peterson, *Nat Struct Mol Biol* 2008
YM133	W303 *ubr1Δ::HIS3-GAL-Myc-UBR1*	Jónsson *et al.*, *Mol Cell* 2004
YM135	W303 *ubr1Δ::HIS3-GAL-Myc-UBR1 ino80Δ::URA3-HA-ino80-td*	Jónsson *et al.*, *Mol Cell* 2004
YM143	LS20 *htz1Δ::hisG-ura3Δ*	Papamichos-Chronakis *et al.*, *Cell* 2011
YM181	LS20 *arp8Δ::KANMX*	Papamichos-Chronakis *et al.*, *Cell* 2011
YM198	LS20 *hda1Δ::HPHMX*	This study
YM199	LS20 *arp8Δ::KANMX hda1Δ::HPHMX*	This study
YM311	W303 [*MATa ade2-1 can1-100 his3-11 leu2,3,112 trp1-1 ura3-1*]	Verma *et al.*, *Mol Cell* 2011
YM334	W303 *arp8Δ::KANMX*	Lafon *et al.*, *Mol Cell* 2015
YM467	LS20 *dst1Δ::KANMX*	This study
YM468	LS20 *rpb9Δ::KANMX*	This study
YM498	LS20 *rco1Δ::HPHMX*	This study
YM499	LS20 *arp8Δ::KANMX rco1Δ::HPHMX*	This study
YM753	BY4741 [*MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*] *NRD1-3XHA::HIS3*	This study
YM801	46a [*MATa ura3 his3 trp1 lys2 ade2 leu2 cup1Δ*]	Steinmetz & Brow, *Mol Cell Biol* 1996
YM802	46a *nrd1-5*	Steinmetz & Brow, *Mol Cell Biol* 1996
YM838	LS20 *arp5Δ::KANMX*	This study
YM850	BY4741 *PCF11-3XHA::HIS3*	This study
YM885	LS20 *RPB1-3XFLAG::KANMX*	This study

YM900	BY4741 *PCF11-3XHA::HIS3 INO80-5XFLAG::NATMX*	This study
YM901	BY4741 *NRD1-3XHA::HIS3 INO80-5XFLAG::NATMX*	This study
YM903	LS20 *RPB1-3XFLAG::KANMX ino80Δ::HPHMX*	This study
YM1002	46a *arp8Δ::HPHMX*	This study
YM1036	46a *nrd1-5 arp8Δ::HPHMX*	This study
YM1085	LS20 *xrn1Δ::KANMX*	This study
YM1139	LS20 *rrp6Δ::KANMX*	This study
YM1141	46a *rco1Δ::KANMX*	This study
YM1147	46a *nrd1-5 arp8Δ::HPHMX rco1Δ::KANMX*	This study
YM1197	BY4741 *abd1-DAmP::KANMX*	Breslow *et al*, *Nat Methods* 2008
YM1325	LS20 *NAB2-3XHA::KANMX*	This study
YM1331	LS20 *NAB2-3XHA::KANMX ino80Δ::HPHMX*	This study
YM1332	LS20 *NAB2-3XHA::KANMX ino80Δ::HPHMX*	This study
YM1452	LS20 *NAB2-3XHA::KANMX htz1Δ::HPHMX*	This study

Extended Data Table 2 – Expression of genes encoding for RNA surveillance proteins is

similar in WT, *ino80Δ* and *arp8Δ* cells. RNA abundance for the listed genes was measured

by spiked-in total RNA-seq analysis in the indicated strains. Values represent average from

two biological replicates for each experiment. The RNA-seq experiment for WT[1] and

ino80Δ was conducted independently from WT[2] and arp8*Δ*. Values represent average from

two biological replicates for each experiment.

Pathway/complex	Gene	WT[1]	*ino80Δ*	Fold Change	WT[2]	*arp8Δ*	Fold Change
NNS	*NRD1*	0.61	0.67	**1.1**	0.43	0.37	**0.8**
	NAB3	0.54	0.44	**0.8**	0.30	0.30	**1.0**
	SEN1	0.46	0.58	**1.3**	0.29	0.25	**0.8**
Exosome	*RRP6*	0.68	0.54	**0.8**	0.46	0.46	**1.0**
	DIS3/RRP44	0.82	0.73	**0.9**	0.64	0.62	**1.0**
TRAMP	*MTR4*	0.97	0.74	**0.8**	0.87	0.89	**1.0**
	TRF4	0.45	0.48	**1.1**	0.28	0.32	**1.1**
	AIR1	0.38	0.30	**0.8**	0.29	0.29	**1.0**
	AIR2	0.21	0.22	**1.1**	0.11	0.15	**1.4**
Cleavage and Polyadenylation	*HRP1*	1.21	1.01	**0.8**	1.04	0.83	**0.8**
	PCF11	0.26	0.24	**0.9**	0.19	0.18	**1.0**
	NAB2	0.70	0.59	**0.8**	0.51	0.54	**1.1**
	PAB1	8.95	6.85	**0.8**	4.53	3.89	**0.9**
Capping	*CEG1*	0.42	0.44	**1.1**	0.19	0.22	**1.1**
	CET1	0.27	0.26	**1.0**	0.17	0.19	**1.1**
Early mRNA processing	*CBC1*	0.96	0.74	**0.8**	0.58	0.68	**1.2**
	GBP2	1.43	1.09	**0.8**	0.83	0.74	**0.9**
	THO2	0.48	0.51	**1.0**	0.27	0.31	**1.2**
Export	*MEX67*	0.52	0.50	**1.0**	0.32	0.34	**1.1**
Cytoplasmic decay	*XRN1*	1.87	1.76	**0.9**	0.91	0.91	**1.0**
	SKI2	0.40	0.40	**1.0**	0.28	0.31	**1.1**

837 **Extended Data Table 3 – Readthrough index (RI) of budding yeast *SNR* snoRNA genes**

838 **as measured by total RNA-seq from the indicated strains.** The RNA-seq experiment for

839 WT[(1)] and *ino80Δ* was conducted independently from WT[(2)], *xrn1Δ*, *rrp6Δ*. Values represent

840 average from two biological replicates for each experiment.

	WT[(1)]	*ino80Δ*	WT[(2)]	*xrn1Δ*	*rrp6Δ*
NME1	0.05	0.10	0.09	0.12	0.06
SNR11	0.18	0.17	0.26	0.23	0.24
SNR128	0.29	0.36	0.36	0.38	0.29
SNR13	0.15	0.28	0.44	0.36	0.25
SNR161	0.20	0.21	0.30	0.20	0.15
SNR189	0.22	0.23	0.47	0.39	0.33
SNR190	0.01	0.01	0.01	0.01	0.01
SNR3	0.90	1.05	0.91	0.87	0.68
SNR30	0.11	0.16	0.25	0.24	0.51
SNR31	0.18	0.15	0.17	0.17	0.13
SNR32	0.64	0.63	0.73	0.83	0.67
SNR33	0.41	0.39	0.47	0.37	0.51
SNR34	0.49	0.55	0.55	0.41	0.29
SNR35	0.17	0.20	0.27	0.17	0.17
SNR36	0.36	0.38	0.33	0.29	0.25
SNR37	0.32	0.37	0.46	0.42	0.72
SNR39B	0.41	0.57	0.50	0.41	0.93
SNR4	0.17	0.18	0.18	0.15	0.34
SNR40	0.61	0.70	0.62	0.70	0.53
SNR41	0.23	0.22	0.22	0.31	0.34
SNR42	0.42	0.37	0.98	0.59	0.95
SNR43	0.23	0.24	0.35	0.40	0.73
SNR45	1.15	1.04	0.99	0.84	1.57
SNR46	0.37	0.58	0.52	1.06	0.20
SNR47	0.56	1.21	0.52	0.56	1.61
SNR48	0.59	0.71	0.75	0.76	0.93
SNR49	0.53	0.62	0.43	0.41	0.21
SNR5	0.42	0.44	0.53	0.45	0.57
SNR50	1.30	1.08	1.00	0.97	1.76
SNR51	0.79	1.30	1.03	0.76	1.45
SNR52	0.47	0.51	0.52	0.42	0.60
SNR53	0.54	0.79	0.52	0.65	0.74
SNR55	0.39	0.57	0.42	0.55	0.28
SNR56	0.40	0.68	0.46	0.52	2.72
SNR57	0.66	0.64	0.60	0.62	0.71
SNR58	1.01	0.87	1.14	0.69	1.07

SNR61	1.57	2.83	1.19	0.85	2.76
SNR62	1.78	2.03	1.66	1.51	3.55
SNR63	0.40	0.52	0.66	0.43	0.55
SNR64	0.33	0.49	0.22	0.23	0.49
SNR65	0.74	0.95	1.09	0.30	3.17
SNR66	1.23	2.42	1.07	1.33	1.74
SNR67	1.07	1.01	1.08	1.14	1.44
SNR68	0.36	0.75	0.82	0.60	1.09
SNR69	0.68	0.88	0.75	0.80	0.70
SNR70	1.56	1.72	2.16	2.34	1.28
SNR71	1.38	2.61	1.41	1.39	3.46
SNR72	0.39	0.62	0.33	0.37	0.42
SNR74	1.10	1.29	1.05	1.53	0.47
SNR75	2.10	2.46	1.72	2.14	1.37
SNR76	0.57	0.52	0.46	0.44	0.30
SNR77	1.07	1.27	1.07	1.13	0.62
SNR78	3.67	4.40	4.26	5.35	4.26
SNR79	1.73	1.63	1.65	1.23	2.45
SNR8	0.27	0.27	0.21	0.22	0.27
SNR80	0.06	0.17	0.14	0.13	0.23
SNR81	0.15	0.28	0.23	0.19	0.12
SNR82	0.66	0.77	0.92	0.89	0.94
SNR83	0.35	0.46	0.46	0.30	1.03
SNR84	0.58	0.77	0.60	0.81	0.26
SNR85	0.02	0.02	0.03	0.02	0.08
SNR86	0.41	0.40	0.25	0.26	0.17
SNR87	0.93	1.16	1.04	0.90	1.38
SNR9	0.28	0.32	0.38	0.36	0.27

METHODS

Bioinformatics Tools and genome annotations

Unless otherwise indicated, version of the tools used for all bioinformatics analyses are as

follows:

Samtools version 1.10

Bedtools version 2.29.2

Bowtie2 version 2.4.1

Hisat2 version 2.2.0

R version 3.6.2

BamCoverage version 3.5.0

Fastqc version 0.11.9

Cutadapt version 2.10

Trim_galore 0.6.4

MarkDuplicates version 2.23.1

FeatureCounts version 2.0.0

The *Saccharomyces cerevisiae* reference genome (S288C version R64-2-1) was downloaded

from Saccharomyces Genome Database (https://www.yeastgenome.org/). The annotation

used in the analysis for mRNA genes was produced by Antonin Morillon's lab, and based on

CAGE-seq. This also includes the following annotations: XUT genes from [83], SUT and CUT

genes from XUT et al (2009) and NUT genes from [16]. Annotations for other transcript types

(snRNA, snoRNA, etc) were downloaded from Saccharomyces Genome Database.

The *Schizosaccharomyces pombe* reference genome and annotation (ASM294v2.47) were

downloaded from pombase (www.pombase.org). Only protein coding genes were used in this

analysis.

NET-seq analyses

NET-seq was performed from yeast strains containing Rpb1-FLAG (3XFLAG), alone or

along with deletion of *INO80*. All strains appeared fully functional, with identical Ser2 and

Ser5 phosphorylation levels on Rpb1 C-terminal domain and identical growth rates to the

respective untagged strains (data not shown). NET-Seq libraries were constructed from

biological duplicates of YM903 (*ino80Δ*, *RPB1-FLAG*) and YM885 (WT, *RPB1-FLAG*) cells

and sequenced as previously described[84]. Reads shorter than 10 nt were filtered out. Reads

were mapped with Bowtie2 using default parameters. Duplicates reads were marked using

Picard MarkDuplicates. Coverage files (bigWig) were produced by custom script. Reads

locations from the alignment files (bam) were turned into bed files using bedtools bamtobed

(with « -split » option), ignoring duplicates reads and reads with mapping quality below 30,

piled up along the genome and turned into bigWig files using a python script. FeatureCounts

was used for quantification with options « -O --fraction s 2 --ignoreDup ». Normalization

factors were computed using DESeq2 method, using the whole count matrix. Travelling

Ratio was computed as: (read density on gene body) / (read density on TSS), with gene body

the region between TSS+200 nt to TTS and TSS the region from TSS to TSS + 200 nt. Read

density was computed as : (# reads on region / region length) *1000.

4tU labeling and purification of newly synthesized RNAs

RNA labeling and purification were performed as previously described with some minor

adjustments[45,85]. Briefly, 20 mL of wildtype and mutant S. cerevisiae cells were grown at

30°C to an OD600 ≈0.8 in YPD. Newly synthesized (ns) RNAs were labelled for 6 min with

4-thiouracil (4tU, Sigma-Aldrich) added to the yeast cultures to a final concentration of 5

mM. Similarly, S. pombe RNAs were labeled for 6 min in YES medium at 32°C. Cells were

pelleted directly after labeling and flash frozen in liquid nitrogen. All experiments were

performed in biological triplicates. Prior to RNA extraction S. cerevisiae and S. pombe cells

were mixed in a ratio of 3:1 and total RNA was extracted with Ribopure Yeast Kit (Ambion,

Life Technologies) according to the manufacturer's guidelines. 150 μg of total RNA were

biotinylated for 3 hours at room temperature using 200 μL of 1 mg/mL EZ-link HPDP-

Biotin (Pierce) with 100 μL of biotinylation buffer (100 mM Tris- HCl pH 7.5, 10 mM

EDTA) and 100 μL of DMSO adjusted to a final volume of 1 mL with DEPC-treated RNase-

free water (Sigma-Aldrich). After biotinylation, excess biotin was removed by mixing 1 mL

of chloroform to the reaction and the liquid phases were separated by centrifugation for 5 min

at 4°C and max speed. The aqueous phase was recovered and precipitated with isopropanol

(1/10 volumes 5M NaCl and 2.5 volumes isopropanol). The recovered RNA was resuspended

in 100 μL DEPC-treated RNase-free water and incubated for 10 min at 65°C followed by 5

min on ice. nsRNAs were bound to 100 μL of μMACS streptavidin microbeads (Miltenyi

Biotec) for 90 min at room temperature and purification was performed using the μMACS

streptavidin starting kit (Miltenyi Biotec). Columns were first equilibrated with 1 mL of

washing buffer (100 mM Tris-HCl at pH 7.5, 10 mM EDTA, 1M NaCl, 0.1% Tween20).

Samples were loaded and the flow-through was reapplied 2 times to the columns. The

columns were washed five times with increasing volumes of washing buffer (600, 700, 800,

900, and 1000 μL). Ultimately, labeled RNAs were eluted twice with 200 μL of 100mM

DTT. nsRNAs were precipitated overnight in 1/10 volume of 3 M NaOAc, 3 volumes of

100% ethanol and 20 μg of RNA-grade glycogen. The RNAs were recovered by

centrifugation and RNA pellets were wash with ice-cold 70% ethanol and resuspended in 20

μL of DEPC-treated RNase-free water (Sigma-Aldrich). Samples were stored at −80°C until

further use. Differential gene expression analysis was performed using DESeq2 1.16.1

Bioconductor R package on Saccharomyces cerevisiae counts normalized with size factors

computed by the median-of-ratios method proposed by Anders and Huber on

Schizosaccharomyces pombe counts (Anders and Huber 2010). cDTA analysis was

performed as described (Schwalb et al. 2012).

Total RNA-seq

Yeast strains were processed in duplicate. ERCC RNA Spike-in control mixes (Ambion,

catalogue #4456740) were added to 1µg of total RNA according to manufacturer. Ribosomal

RNAs were depleted from total RNA using the RiboMinus™ Eukaryote v2 Kit (Life

Technologies). Depletion efficiency and quality control of rRNA-depleted RNA was assessed

by analysis in RNA Pico 6000 chip for 2100 Bioanalyzer (Agilent). Total RNA-seq libraries

were prepared from 50ng of rRNA-depleted RNA using the TruSeq® Stranded Total RNA

Sample Preparation Kit (Illumina). Paired-end sequencing (2x50 nt) was performed on a

HiSeq 2500 sequencer giving between 65 and 84 million reads. Reads were mapped using

version 2.0.6 of TopHat, with a tolerance of 3 mismatches and a maximum size for introns

2Kb. Tags densities were normalized on spike-in controls for all subsequent analyses.

ChIP-exo

The ChIP-exo assay has been previously described (Rhee HS, Pugh BF. ChIP-exo method for

identifying genomic location of DNA-binding proteins with near-single-nucleotide accuracy.

Curr Protoc Mol Biol. 2012;Chapter 21:Unit 21). In brief, yeast strains were grown to

exponential phase in yeast extract peptone (YP) + 2% dextrose (30°C to $OD600\ nm = 0.8$),

then subjected to 1% formaldehyde crosslinking for 15 min at 25 °C. After quenched with

125mM final concentration of glycine for 5 min, cells were harvested and washed. Sonicated

chromatin was prepared by standard methods. Standard ChIP methods were used for histone

H3 (antibody Ab1791, Abcam) followed by lambda exonuclease treatment and library

construction. Libraries were sequenced on an Illumina 2000 sequencer. Reads were uniquely

mapped to the reference genome (sacCer3 build) using version 2.1.0 of Bowtie2 with a

tolerance of 1 mismatch in seed alignment. Data were normalized on the total number of

uniquely mapped reads, for each sample.

Cell proliferation analysis in liquid media

Liquid growth rate analysis was conducted using the CG-12 robot (S&P Robotics Inc.). Yeast cells transformed with a URA3-containing plasmid (pRS416) and grown exponentially in SC-ura at 30°C were diluted to OD620 0.1 in either SC-ura or SC-ura + 50µg/ml 6-AU and four 150µl samples from each strain were plated onto 96-well transparent plates covered with breathable films. Plates were incubated at 30°C for several days and OD620 was measured in the CG-12 robot (S&P Robotics Inc.) every hour with a shaking of 1000rpm for 1 minute before each read. Liquid growth rate analysis was not possible in the *nrd1-5* and associated strains due to flocculation. All cell proliferation experiments were repeated at least twice.

Growth rates analysis on agar plates

Growth of yeast strains was compared by plating 5-fold serial dilutions on SC / SC-inositol, SC-ura / SC-ura + 50µg/ml 6-azauracil (6-AU, Sigma), SC-leu / SC-leu + 0.4mM CuSO4 (Sigma) or SC-leu / SC-leu + 0.25 CuSO4 and incubating the plates at 30°C for several days. All plating experiments were performed at least three times.

RT-qPCR

Total RNA was extracted either with RNeasy Mini Kit (Qiagen) or by classical acid phenol-chloroform extraction. RNAs abundance was measured with SuperScript® III Platinum® SYBR® Green One-Step qRT-PCR Kit (Invitrogen) using specific primers for the targets of interest in a 10µl reaction volume, on Applied Biosystem Step-One Plus machines. RNAs abundance of each sample was compared with standard curves analysis. Abundance of *CUP1* from transcriptional read-through analyses using the *ACT-CUP* reporter system was compared to *ACT** RNA specifically transcribed from the same plasmid (Extended Data Fig.5a). Abundance of *NRD1* promoter-proximal RNA (*NRD1* 5') was compared to levels of *NRD1* promoter-distal RNA (*NRD1* 3').

Co-immunoprecipitation

Benzonase-treated yeast cell extracts were incubated with 20µl Anti-FLAG Affinity Gel

(Sigma Aldrich) in 10mM Tris-Cl pH 7.5, 150mM NaCl, 0.05mM EDTA, 0.5% NP40, 1X

Protease Inhibitor Cocktail (Roche) and 0.01mM PMSF for 3h. Beads were washed twice in

10mM Tris-Cl pH 7.5, 150mM NaCl, 0.05mM EDTA, 1X Protease Inhibitor Cocktail

(Roche) and 0.01mM PMSF and resuspended in SDS sample buffer. Western blot analyses

were performed using Monoclonal anti-FLAG M2 antibody (Sigma F3165), Anti-HA tag

antibody (Abcam, ab9110) and Anti-Arp5 antibody (Abcam, 12099). Images were acquired

using Syngene G:BOX. The experiment was reproduced three times. Immunoprecipitations

for H2A.Z and HA-tag were performed similarly on benzonase-treated yeast cell extract

using 36µl Protein-A sepharose beads coupled with 6µl of appropriate antibody (Anti-

H2A.Z, Active Motif 39647; Anti-HA tag, Abcam ab9110). Control immunoprecipitations

(mock) were performed with a non-specific antibody (Anti-Rabbit IgG, Promega). Western

blot analyses were performed using the following antibodies: Anti-HA, Anti-Arp5, Anti-

H2A.Z and Anti-Rpb1 y-80 (Santa Cruz Biotechnology, sc-25758). After the

immunoprecipitation for H2A.Z shown in Figure S6G, beads were washed twice in 10mM

Tris-Cl pH 7.5, 150mM NaCl, 0.05mM EDTA, 1X Protease Inhibitor Cocktail (Roche) and

0.01mM PMSF, then washed a third time in the same buffer containing 300mM NaCl and

finally resuspended in SDS sample buffer All experiments were reproduced three times.

Ino80 inducible degradation

The assay for inducible degradation of Ino80[82] was performed as previously described[35,86]

Briefly, a degron tag was fused to INO80 (Ino80-td) in cells carrying the UBR1 gene, that

encodes for the N-recognin Ubr1, under the control of the GAL1 promoter, and cells were

transformed with the *ACT-CUP* plasmid reporter system (Extended Data Fig. 5a). Cells were

grown in SC-leu media containing raffinose as carbon source at 25°C until OD600~0.6-0.7,

then shifted to SC-leu media containing galactose to induce *UBR1* expression. Subsequently,

complete Ino80-td protein degradation by the Ubr1 pathway was induced by 3 hours of heat

shock at 37°C. Cells were then immediately harvested by centrifugation at 4°C, 3500rpm for

3 minutes, then washed twice with ice-cold water and snap-frozen in liquid nitrogen for RNA

analysis. Yeast cells carrying the GAL1-driven UBR1 gene along with an unmodified INO80

genes were use as control. Furthermore, both (GAL1-UBR1) and (GAL1-UBR1 Ino80-td)

cells were grown in SC-leu media containing glucose as a negative control for UBR1

expression (Extended Data Fig. 5d, grey bars). RNA analysis was conducted by RT-qPCR as

described above.

ChIP-qPCR

Chromatin immunoprecipitation was performed as previously described in (Papamichos-

Chronakis M, Watanabe S, Rando OJ, Peterson CL. Global regulation of H2A.Z localization

by the INO80 chromatin-remodeling enzyme is essential for genome integrity. Cell.

2011;144(2):200-13.). Sonication was performed on a Bioruptor® Pico sonication device

(Diagenode) with ten 30-sec pulses with a 30-sec break between pulses. Immunoprecipitation

was carried out with Anti-HA tag antibody (Abcam ab9110) and Anti-Rpb1 y-80 antibody

(Santa Cruz Biotechnology, sc-25758) in parallel. The recovered DNA was subjected to

quantitative real-time PCR using Platinum® SYBR® Green qPCR SuperMix-UDG Kit

(Invitrogen) with specific primers for the targets of interest in 10µl reaction volume, on

Applied Biosystem Step-One Plus machines.

RNA-IP

RNA immunoprecipitation was performed as described in (Selth LA, Gilbert C, Svejstrup JQ.

RNA immunoprecipitation to determine RNA-protein associations in vivo. Cold Spring Harb

Protoc. 2009;2009(6):pdb prot5234.). Sonication was performed on a Bioruptor® Plus

sonication device (Diagenode) with two 15-sec pulses at 50% amplitude with a 1-min break

between pulses. Immunoprecipitation was carried out with Anti-HA tag antibody (Abcam

ab9110). The recovered RNA was analysed by RT-qPCR as described above.

De novo analysis of public ChIP-exo and CRAC data

NAB3 and POLII CRAC data were taken from GEO dataset GSE70191 and processed as

described in [56]. NAB2 and MTR CRAC data were taken from GEO dataset GSE46742 and

processed as described in [4]. ChIP-exo H3 data were downloaded from NCBI SRA dataset

SRA059355 and processed as described in [87].