Analysis of an experimental transition in individuality challenges the need to assign traits to levels

Caroline J. Rose[1,2], Katrin Hammerschmidt[1,3] & Paul B. Rainey[1,4,5]

[1]New Zealand Institute for Advanced Study, Massey University, Auckland, New Zealand.

[2]Present address: Centre d'Écologie Fonctionnelle et Évolutive (CEFE), CNRS, Montpellier, France.

[3]Present address: Institute of General Microbiology, Kiel University, Kiel, Germany

[4]Present address: Department of Microbial Population Biology, Max Planck Institute for Evolutionary Biology, Plön 24306, Germany and [5]Laboratoire de Génétique de l'Evolution, Chemistry, Biology and Innovation (CBI) UMR8231, ESPCI Paris, CNRS, PSL Research University, Paris, France

Author ORCIDs

Caroline J Rose https://orcid.org/0000-0001-7133-035

Katrin Hammerschmidt https://orcid.org/0000-0003-0172-8995

Paul B Rainey https://orcid.org/0000-0003-0879-5795

Correspondence and requests for materials should be addressed to CJR carojrose@gmail.com

Keywords: Group Selection – Emergence – Individuality – Multicellularity

Abstract

A major evolutionary transition in individuality, at any level of the biological hierarchy, occurs when groups become capable of participating in Darwinian processes as units of selection. Identifying transitions in individuality can be problematic because apparent selection at one level of the biological hierarchy may be a by-product of selection occurring at another level. Here we discuss various approaches to this 'levels-of-selection' problem, drawing attention to the futility of designating traits to hierarchical levels prior to determining the level of selection. While individuals can exhibit traits that are maintained by selection at any level in the biological hierarchy, terms synonymous with 'collective (group) trait' or 'particle trait' are often applied prematurely. We argue that traits should not be assigned *a priori* to a hierarchical level, but may respond to selection at different levels depending on environmental conditions. This point is illustrated by an experiment in which groups of the bacterium *Pseudomonas fluorescens* reproduced via life cycles involving soma- and germline-like phases. The rate of transition between the two cell types was a focus of selection, and might be regarded as a property of groups, cells, or even genes. Philosophical analyses reveal that this trait evolved to be higher in one treatment as a result of its emergent relation to group fitness (contributing to the evolution of individuality), and lower in another treatment because of cell-level selection. An *a priori* designation of transition rate as either a 'group trait' or a 'cell trait' may have masked the true level of selection.

Introduction

Evolutionary transitions in individuality are central to the emergence of biological complexity. During the transition from single cells to multicellular organisms, groups of cells – on multiple occasions (Baldauf 2003) – have emerged as 'Darwinian individuals' capable of evolving by natural selection – as units of selection in their own right. The causes of these transitions are unclear. Nonetheless, philosophical and theoretical literature has provided insight into the fundamental requirements for evolutionary transitions to take place (Buss 1987; Maynard Smith and Szathmáry 1995; Michod 1999; Bonner 2000; Godfrey-Smith 2009; Mikhailov et al. 2009; Libby and Rainey 2013a; Shelton and Michod 2014; Black et al. 2020). Recently, the question has also received interest from the field of experimental evolution (*e.g.* Ratcliff et al. 2012, 2013; Koschwanez et al. 2013; Hammerschmidt et al. 2014; Herron et al. 2019; Rose et al. 2020), which focuses more on the environmental and ecological circumstances that can establish conditions promoting transitions to multicellularity. Here we attempt to bridge the gap between the fields of philosophy of biology and experimental biology by application of philosophical frameworks to experimental data. In so doing we highlight the importance of careful consideration of the level(s) of selection prior to drawing conclusions based on intuition.

Levels of Selection

The view that the evolution of multicellularity necessitates a transition in individuality is based on a 'multilevel' perspective (Heisler and Damuth 1987; Damuth and Heisler 1988), which maintains that life is hierarchically organised, with 'higher' level units comprised of groups of particles from the level below. For example, eusocial colonies are made up of multicellular organisms, which in turn are integrated groups of cells. The very existence of a biological hierarchy is evidence of transitions from groups to Darwinian individuals. It

follows that during the evolutionary history of the hierarchy, groups emerged as

units that evolve by natural selection – that is, entities that can acquire

adaptations.

The essence of a transition in individuality lies in the capacity for higher-level

reproduction, a prerequisite for Darwinian evolution. While group reproduction

is necessary, it is not sufficient to ensure that natural selection operates at the

level of the group. The archetypal 'recipe' for a trait to evolve by natural

selection at a particular level of the biological hierarchy is that the trait under

selection must be heritable, and it must covary with fitness (reproductive

success) *at that level* (Lewontin 1970). However, covariance is a statistical

concept and not a causal one. There *may* be a causal (selective) influence of a

given trait on the fitness of a group if the trait covaries with group fitness, but

any such covariance may also be misleading (Okasha 2006). Therefore,

considering the possibility of natural selection independently at each level is

inadequate because an apparent Darwinian process at one level can be a side

effect of selection operating at another level – a 'cross-level by-product' (Sober

and Lewontin 1982; Okasha 2006; Shelton and Michod 2014). Okasha (2006,

p.78-79) frames the 'levels-of-selection problem' as follows:

"When is a character[1]-fitness covariance indicative of direct selection

at the level in question, and when is it a by-product of direct selection

at another hierarchical level?"

Attempting to understand the evolutionary origin of the biological hierarchy

naturally leads to a multilevel view of evolution by natural selection, however

alternative reductionist perspectives are also recognised. The supervenience

argument, for example, maintains that genuine group-level selection is

impossible because all properties of groups depend on those of their constituent

---

[1] A semantic note: The term 'trait' is used throughout the paper, however 'character' is on occasion used interchangeably with 'trait' when adopting the language of key reference material. 'Group' and 'collective' are also used interchangeably in this manner, as is the term 'particle', which is used by some references to refer to the units that comprise groups.

particles. The traits of the underlying particles ultimately determine both the
fitness and character of groups; the whole 'supervenes' on its parts (summarised
in Okasha 2012). The argument then follows that there can never be a causal
link between a group's character and its fitness because *all* causality is at the
particle level. The supervenience argument against group-level selection renders
the levels-of-selection problem irrelevant because it makes cross-level by-
products ubiquitous.

Perspectives that reduce all evolution to lower levels are not practical
frameworks for investigating evolutionary transitions because of limited
explanatory power. For example, the supervenience argument is unable to
account for how the biological hierarchy itself came into existence and must
logically reject the notion of a hierarchical structure in biological systems. Even
the reductionist perspectives of Dawkins (1976, 1982) and others, which
maintain that genic selection can account for all evolution, are not congruent
with the supervenience argument because genes themselves are groups of lower
level molecules whose evolution requires explanation. Okasha (2006) reasons
that reductionist arguments only show that a trait-fitness covariance at the
higher level must be a side effect of *"some lower level causal processes or other,*
but not necessarily lower level *selection"*. Certainly everything in existence can
be explained by its constituent particles, however reducibility to lower-level
selection is what is important for understanding evolutionary origins. The
underlying particle traits on which group fitness supervenes do not necessarily
include particle *fitness*, so it does not follow that group fitness always has a
lower-level *selective* explanation. Therefore, Okasha suggests that the concept
of a cross-level by-product should be restricted to a trait-fitness covariance at
one level that results from *selection* operating at another level.

In our view, if the level of *selection* is what matters, then characters or traits
themselves need not be assigned to any particular level. A trait may appear
intuitively to belong to a particular level, but other cases are less clear. For
example, the trait 'height' is necessarily the height of some entity that exists at a
particular level, e.g. the height of a tree, or a termite mound. On the other hand,

a bistable switch between two cell phenotypes might be viewed as a property of
cells, genes, or even a group of cells (manifesting as the proportion of the
different cell phenotypes in the population). Even seemingly clear-cut cases may
be less obvious when informed by the level of selection rather than by intuition.
The height of a termite colony, for example, may not play a role in selection
between colonies, but could be a by-product of selection acting on individual
termites within a colony for some trait that can only be empirically measured as
the height of the mound. The distinction between group trait and particle trait is
unnecessary and can lead to debates that detract from the real issue of the
selective cause of the trait in question. It makes no sense (and in fact invokes
circular logic) to utilize the terms 'group trait' or 'particle trait' prior to
determining the level(s) at which the trait was selected. If a given trait is not
assigned to a particular hierarchical level, discussions over whether it is
reducible to particle traits are irrelevant. The relevant factor in determining the
level of selection is the level at which a given *undesignated* trait is causally
associated with fitness.

The levels-of-selection problem is related to G.C. Williams' (1966) distinction
between 'group adaptation' and 'fortuitous group benefit'. In his well-known
thought experiment, Williams contrasts herds of deer that run at different
speeds. Herds made up of faster deer grow and 'reproduce' (split into new
herds) more quickly because slower deer are more susceptible to predation.
Consequently, selection for fast deer within herds leads to increasingly faster
herds of deer – a fortuitous group benefit. The fitness of the herds (the rate at
which they reproduce) covaries with the average running speed of the herd,
however this covariance is a cross-level by-product of within-herd selection for
faster deer - not a group adaptation, or the outcome of herd-level selection. To
assign deer speed as an adaptation of the herd would be to conclude that faster
deer were selected *because* of the effect of this adaptation on the reproduction
rate of herds. While herds indeed benefit from being comprised of faster deer,
causality cannot be attributed to between-herd selection. A fast individual deer
does not benefit the other deer within its herd, rather faster deer were selected

because of within-herd competition to evade predation; causality is at the lower
level.

The cross-level by-product is conveniently detected in the hypothetical example
above because Williams provides two factual statements that are necessary to
conclude that the evolution of fast deer is due solely to selection between deer:
1) faster individual deer are better able to evade predators, and 2) herd
reproduction occurs as a result of herds reaching a critical size (given by the
number of deer within a herd). If these were the only two biological facts then it
is straightforward to conclude that the relationship between herd speed and
herd reproduction is caused 'from below' by deer-level selection. However, if
these two facts were not given, the true selective cause of the evolution of fast
deer would be unknowable. Moreover, were *more* biological information
provided, the conclusion reached may be altered. Let us imagine that access to
more data revealed a trade-off between deer speed and deer reproduction, such
that while fast deer are more likely to survive in the presence of predators, slow
deer produce more offspring. In this scenario, deer may evolve to be faster or
slower depending on the density of predators in the environment. Let us make
the assumption that in a low predator density environment, there is nevertheless
selection for fast deer. In this case it would be dubious to conclude that
increased deer speed resulted solely from within-herd selection. Let us further
imagine that herd size is not the sole determinant of herd reproduction, rather
new 'offspring' herds are founded after an act of aggression, and such an act is
more likely to occur in herds with a higher proportion of fast deer. It is now
plausible that these fast herds are maintained, at least in part, by between-herd
selection.

The levels-of-selection problem is not often addressed adequately (if at all) in
experimental evolution literature. For example Ratcliff et al. (2012) evolved
'multicellular organisms' from the unicellular yeast *S. cerevisiae* by selection for
rapid settling cellular clumps in liquid media. The result was evolution of a
clustering 'snowflake' phenotype. These groups of cells reproduce via
multicellular propagules that separate from the parent snowflake by apoptosis

(cell death) of a single cell at the break point. The selective regime favoured
larger snowflakes and higher rates of apoptosis – resulting in higher rates of
group reproduction. If snowflakes vary in traits that affect their fitness, and if this
variation is heritable, then it is possible that these traits evolved in response to
between-snowflake selection. However, to claim that group-level selection –
and a transition in individuality - has occurred, it is necessary to show that the
observed adaptations not only improve the fitness of the snowflake, but that this
improvement is not a by-product of selection acting at the cell-level – a
fortuitous group benefit. In this experiment, cell fitness also increased and was
correlated with rate of apoptosis. By forming snowflakes that produce a higher
proportion of altruistic cell types, recipient cells can grow faster than their non-
cooperative ancestors. While it is possible that group-level selection may have
contributed to the increased rate of apoptosis, it is also possible that apoptosis
increased due to cell-level selection. Despite positive correlations between
apoptosis rate and both cell and snowflake reproduction, Ratcliff et al. assigned
the apoptosis trait to the snowflake level.

It is possible to identify an evolutionary transition in individuality by
determining the level(s) of selection. Groups evolve as higher-level Darwinian
individuals when their evolutionary success can no longer be fully accounted
for by the evolutionary success (*i.e.,* fitness) of their constituent particles. In
other words, a primordial multicellular group constitutes a unit of selection
once group functionality, rather than size, becomes the main determinant of
fitness (Okasha 2006). As illustrated by the examples above, detecting the level
of selection can be problematic because there exists a 'grey area' of mid-
transition groups (Godfrey-Smith 2009). In such groups it is unclear whether
certain properties are evolving due to selection at the group level, the cell level,
or both. We cannot simply assign an evolving trait to a particular level of
selection based on assumed benefits to this level. Furthermore, the emergence
and evolution of Darwinian properties themselves (*i.e.,* reproduction and
heritable variation) at the higher level requires a selective explanation. It is not
sufficient to simply transpose the existence of Darwinian properties, such as

variation, from the lower level to the higher (Bourrat 2015); the possibility of selection acting upon this variation is dependent on the mode of group reproduction (Libby and Rainey 2013b; De Monte and Rainey 2014; Hammerschmidt et al. 2014; Rose et al. 2020).

Hammerschmidt et al. (2014) reported the results of a selection experiment involving lineages of the bacterium *P. fluorescens* that differed in the mode of group reproduction. Groups reproduced via a single-celled bottleneck in two selection regimes, however in one regime this bottleneck was an alternative cell type (a 'germ' cell). Crucially, while all groups exhibited the hallmarks of natural selection (reproduction, variation and heredity), improvements in group fitness became 'decoupled' from the fitness of constituent cells only for groups that reproduced via a germline.

This experiment illustrates some difficulties with a simplistic interpretation of the process of natural selection at multiple levels. Recognition of a transition in individuality calls for careful consideration of the possibility of cross-level by-products. Various methods have been proposed for determining the hierarchical level(s) at which the association between a trait and fitness is causal, rather than merely statistical. Here we describe several approaches to tackling the levels-of-selection problem and apply them to the *P. fluorescens* populations. We demonstrate that a key trait evolved in the experiment due to selection at different levels depending on the environment.

The Experiment

The Hammerschmidt et al. (2014) experiment tracked the evolution of groups of the bacterium *P. fluorescens,* which had previously been shown to evolve between mat-forming (soma-like) cells and dispersing (germline-like) states via negative frequency-dependent selection (Rainey and Rainey 2003). When cultured in a static environment, rare cooperative "wrinkly spreader" (WS) mutants invade populations of planktonic "smooth" (SM) cells by forming a

surface-colonizing soma-like mat that allows cells to access oxygen. Within the

mat, SM "cheating" germline-like cells rise in frequency when rare due to

advantages that stem from not producing costly cell-cell adhesive.

The frequency-dependent interactions between SM and WS cell types were

utilized to implement a multicellular life cycle in which the WS mats were

analogous to soma, and SM cells functioned as a germline. This facilitated

differential group death and birth, which in turn allowed selection to operate

between groups (Fig 1a). After ten life cycle generations, group fitness (relative

number of group offspring produced) increased while cell fitness decreased; the

fitness of the higher and lower levels became decoupled – a signature of a

transition in individuality. Enhanced fitness of evolved groups was attributable

to a property selected at the group-level, namely, the capacity to transition

through the germ and soma phases of the life cycle (termed 'transition rate'),

and was not explained by improvements in individual cell proliferation - the

fitness of the SM 'propagules' and the total carrying capacity were both

reduced.

In parallel to this "cheat-embracing" selective regime, a "cheat-purging"

experiment was conducted, in which groups reproduced via a single WS

propagule cell (Fig 1b). While group fitness increased in both regimes, in the

cheat-purging regime this was correlated with increases in all indicators of cell

fitness. The following sections contrast the results of the cheat-embracing and

cheat-purging ecologies in light of various perspectives arising from the levels-

of-selection debate.

285 **Emergence**

One approach to the levels-of-selection problem is to distinguish between an

emergent trait – a property of a group that cannot be qualitatively or

quantitatively reduced to the sum of its parts (Corning 2002), and *non-emergent*

(or 'aggregate') traits. How such a distinction is made is not always

straightforward because all properties of a group can ultimately be explained by
the underlying particles (discussed above). Nevertheless, the 'emergent
character requirement' (Vrba and Eldredge 1984; Vrba 1989) posits that genuine
group-level selection, that is not reducible to selection at the particle level, can
only operate on emergent traits. Okasha (2006) objects to this rule because an
emergent trait may be a *result* of selection operating at the group-level, rather
than a pre-condition of it. The capacity for group reproduction, which is
necessary for group-level selection, is itself a derived, and arguably emergent,
trait (Griesemer 2000; Rainey 2007; Rainey and Kerr 2010). It is also possible
for an emergent trait to increase in frequency in the absence of group-level
selection if it is associated with traits that increase particle fitness, *i.e.*, an
emergent trait may inadvertently result from particle-level selection. It is
therefore generally accepted that the emergent/aggregate distinction is not
useful for solving the levels-of-selection problem (Damuth and Heisler 1988;
Okasha 2006, 2012; Godfrey-Smith 2009).

Stephen Jay Gould argued that while a given trait need not be emergent to
evolve at the level of the group, emergent *fitness* is what matters for group-level
selection (Lloyd and Gould 1993; Gould 2002). It is true that groups must
acquire the capacity for reproduction to allow the possibility of group-level
selection and a measure of group fitness. However, this potential for group-level
selection does not rule out cross-level by-products. Group fitness emerged
(increased) in both the cheat-purging and cheat-embracing selective regimes in
Hammerschmidt et al. (2014) however in the cheat-purging regime this was
likely due to a cross-level by-product because the emergent group fitness was
correlated with increased cell fitness.

Heisler and Damuth (1987) suggest that emergence of the *relation* between a
trait and group-level fitness is of fundamental importance for group-level
selection. A trait-fitness relation at a higher level is emergent if it cannot be
accounted for by a trait-fitness relation at a lower level. Okasha's (2006)
formulation of the levels-of-selection problem essentially recapitulates the

emergent relation requirement. It is possible to identify an emergent relation
when there is no positive association between particle and group fitness (Fig 2).

A relationship exists between cell fitness and group fitness in the *P. fluorescens*
groups that evolved under the cheat-purging regime (Hammerschmidt et al.
2014). This (probably causal) link presents no difficulty in determining the level
of selection, however, because the trait in question (transition rate) correlates
only with parameters of cell fitness, not group fitness (Fig 3a). It is clear
therefore, that the transition rate significantly decreased because of selection
acting at the cell level, and the increased group fitness can be attributed to the
correlation between group and cell fitness parameters – group fitness
supervenes on particle fitness. This is in contrast to the cheat-embracing regime,
in which no correlation exists between cell and group fitness in the evolved
populations thus the positive relationship between the increased transition rate
and group fitness is causal - an emergent relation (Fig 3b).

In Williams' (1966) example, however, there is a positive association between
deer fitness and herd fitness, so it is impossible to determine whether the
emergent relation between speed and herd fitness is causal from correlations
alone – there would be a red line between all three points in Figs 2 and 3 (the
same is true for the yeast snowflake example).

Contextual Analysis

Heisler and Damuth (1987) devised a method for detecting emergent relations
by analysing trait-fitness covariance from phenotypic data. Contextual Analysis
is a generalisation of the selection analysis developed by Lande and Arnold
(1983). This approach treats the levels-of-selection problem as a special case of
selection on correlated traits, where the dependence of fitness on correlated
traits is captured with a standard linear regression model. The covariance of
group fitness (Y) with a (so-called) 'group' trait (Z) could be non-zero (*i.e.,*
Cov(Y,Z) $\neq$ 0) in the absence of a causal link between Z and Y if Z correlates

with another trait that *does* affect fitness. There is a cross-level by-product if *Y* is
causally determined by the average fitness of its constituent particles, *W*. The
trait-group fitness covariance can be partitioned into two components: direct
group-level selection on a trait, *β₁Var(Z)*, and the by-product of particle-level
selection, *β₂Cov(W,Z)*:

$$Cov(Y,Z) = \beta_1 Var(Z) + \beta_2 Cov(W,Z) \qquad (1)$$

where β_1 is a measure of the direct effect of *Z* on *Y*, controlling for *W*, and β_2
measures the effect of *W* on *Y*, controlling for *Z*.

No relationship exists between transition rate and cell fitness in the cheat-
embracing *P. fluorescens* populations (Hammerschmidt et al. 2014). Contextual
analysis is therefore not necessary to partition the change in transition rate due
to cell- and group-level selection, because *Cov(W,Z) = 0*. The strength of
group-level selection for increased transition rate is equal to the covariance
between group fitness and transition rate:

$$Cov(Y,Z) = \beta_1 Var(Z) + 0 \qquad (2)$$

$$Cov(Y,Z) = \beta_1 Var(Z) \qquad (3)$$

365 As no relationship exists between transition rate and group fitness in the evolved
366 populations from the cheat-purging regime (Hammerschmidt et al. 2014), there
367 is no need to discriminate between levels of selection – decreased transition rate
368 was selected entirely at the lower level:

$$Cov(Y,Z) = 0 \qquad (4)$$

These results corroborate the idea that a trait need not be assigned to a
particular hierarchical level. The transition rate between the two cell types
might be regarded as a property of groups, cells, or even genes, often depending
on the inclinations of the investigator. Our view is that transition rate should not
be associated with one level because the direction of its evolution can change
due to selection operating at multiple levels in differing ecological contexts

(Rose et al. 2020). This trait evolved to be higher in the cheat-embracing
selective regime as a result of its influence on group fitness, and lower in the
cheat-purging regime because of selection at the cell-level. Does it then follow
that transition rate is a group trait in one ecological context and a cell trait in
another? To reiterate, the level of *selection* is the focus during an evolutionary
transition in individuality. An *a priori* designation of transition rate as either a
'group trait' or a 'cell trait' may have masked the true level of selection, or
worse, led to false conclusions.

385 **Additivity**

Lloyd (1989) contributed the 'additivity criterion' to the levels-of-selection
debate - a notion that was first put forward by Wimsatt (1980). The additivity
criterion posits that true group-level selection can only occur when the variance
in group fitness does not relate additively (linearly) to particle composition.
When there is a linear relationship between group fitness and particle
composition, then differences in group fitness can be fully explained by
selection at the particle level. A shift in the level of selection occurs when there
is a departure from additivity, *i.e,.* there is a non-linear relationship between the
particle composition of groups and fitness. The additivity criterion for group-
level selection is similar to the criterion of emergent relations: a 'new' non-
linear relation between a trait and fitness emerges at the level of the group.
However the additional specification that this relation be non-linear adds a
crucial layer of information that the emergent relation requirement alone lacks.
To illustrate, the emergent relations requirement could not assign selective
causality to the evolution of fast deer in the Williams example (discussed
above), however the stipulation by Williams that herds reproduce as a function
of herd size results in an additive relationship between herd fitness and their
composition, rendering the selection of fast deer to be at the lower (deer) level.

The cheat-embracing selective regime in the *P. fluorescens* populations clearly
results in a non-linear relationship between group fitness and particle
composition. SM 'germline' cell types are required at a high enough frequency
to be consistently detected during the regime, however a WS mat must also be
present to avoid extinction. Therefore, group fitness will be highest at
intermediate frequencies of SM cells, and zero if the proportion of SM cells is
either 0% or 100%. This non-additive relationship between cell composition
and group fitness is not pronounced in the experimental data (published in
Hammerschmidt et al. (2014) but re-analysed here) because the wildtype
transition rate is so low that the hypothetical 'high SM/low group fitness' groups
do not exist in the ancestral populations, and were not selected during the
experiment due to their low group fitness (Fig 4a). Nevertheless, a comparison
of the fit of linear and quadratic models to the data confirms the non-additive
relationship between the proportion of SM cells within a group and fitness.
Akaike's Information Criterion (Akaike 1973), which compares the quality of
statistical models by giving a relative estimate of the information lost by
overfitting a given model, indicated that the quadratic model was 99.99% and
99.88% more likely to explain the data than the linear model in the ancestral
and evolved groups respectively (Supplementary Table). This non-additive, and
therefore emergent, relationship between the proportion of SM cells (determined
by the transition rate trait) and group fitness is indicative of selection operating
at the level of the group.

In contrast, there is an additive relationship between the composition of groups
(as determined by transition rate) and their fitnesses in the cheat-purging
selective regime. Compared to the quadratic model, the linear model has a
70.54% and 69.94% probability of correctly fitting the ancestral and evolved
data respectively. The two ancestral lines that had both low group fitness and
low proportion of SM cells did not exhibit visible mats because of the invasion
of a third cell type, the "fuzzy spreader" (described in Ferguson et al. 2013).
These groups were therefore extinguished after one generation and the
proportion of SM alone became a reliable predictor of group fitness. In contrast

to the non-linear relationship between the fitness of a group and its composition

in the cheat-embracing regime, the additive dependence of a group's fitness on

its composition reveals lower-level causality in the cheat-purging regime. Non-

additivity in the cheat-embracing life cycle led to a transition in individuality

that was not possible for groups in the cheat-purging regime, whose fitnesses

depended entirely on particle composition.

Conclusion

The results from the experiments presented in Hammerschmidt et al. (2014) and

scrutinized here in further detail demonstrate that a major evolutionary

transition to multicellularity is possible when groups of cells reproduce via a

two-phase life cycle involving a germline-like cell. Conflict resolution evolved

in groups that reproduced without a germline (in the form of reduced transition

rate to SM 'germline-like' cell types), however these groups did not evolve

individuality. In contrast, during the course of the evolution experiment, a

causal relation between the transition rate trait and group fitness emerged in

groups that reproduced via a germline-like cell. This emergent relation, in

addition to a non-additive relation between the composition of the groups and

group fitness, demonstrates that the levels-of-selection problem was resolved,

shifting selection to the higher level and revealing new biological individuals

free to participate in an open ended manner in Darwinian evolution.

Importantly, while the transition rate trait evolved to be lower in one

experimental regime due to selection acting at the cell-level, the same trait

increased due to group-level selection in the other experimental setup,

challenging the common practice of assigning traits to levels prior to rigorous

levels-of-selection analyses.

References

Akaike H (1973) Information theory and an extension of the maximum likelihood principle. In: Petrov BN, Csaki F (eds) 2nd International Symposium on Information Theory. Akademiai Kiado, Budapest, pp 267–281

Baldauf SL (2003) The deep roots of eukaryotes. Science 300:1703–1706. https://doi.org/10.1126/science.1085544

Black AJ, Bourrat P, Rainey PB (2020) Ecological scaffolding and the evolution of individuality. Nat Ecol Evol 4:426–436. https://doi.org/10.1038/s41559-019-1086-9

Bonner JT (2000) First Signals: The Evolution of Multicellular Development. Princeton University Press, Princeton

Bourrat P (2015) Levels of Selection Are Artefacts of Different Fitness Temporal Measures. Ratio 28:40–50. https://doi.org/10.1111/rati.12053

Buss LW (1987) The Evolution of Individuality. Princeton University Press, Princeton

Corning PA (2002) The re-emergence of "emergence": A venerable concept in search of a theory. Complexity 7:18–30. https://doi.org/10.1002/cplx.10043

Damuth J, Heisler IL (1988) Alternative formulations of multilevel selection. Biol Philos 3:407–430. https://doi.org/10.1007/bf00647962

Dawkins R (1976) The Selfish Gene. Oxford University Press, Oxford

Dawkins R (1982) The Extended Phenotype: The Gene as the Unit of Selection. Oxford University Press, Oxford

De Monte S, Rainey PB (2014) Nascent multicellular life and the emergence of individuality. J Biosci. https://doi.org/10.1007/s12038-014-9420-5

Ferguson GC, Bertels F, Rainey PB (2013) Adaptive divergence in experimental
populations of *Pseudomonas fluorescens*. V. Insight into the niche specialist
"Fuzzy Spreader" compels revision of the model *Pseudomonas* radiation.
Genetics 195:1319–1335. https://doi.org/10.1534/genetics.113.154948

Godfrey-Smith P (2009) Darwinian Populations and Natural Selection. Oxford
University Press, Oxford

Gould SJ (2002) The Structure of Evolutionary Theory. Belknap Press,
Cambridge, MA

Griesemer J (2000) The units of evolutionary transition. Selection 1:67–80.
https://doi.org/10.1556/Select.1.2000.1-3.7

Hammerschmidt K, Rose CJ, Kerr B, Rainey PB (2014) Life cycles, fitness
decoupling and the evolution of multicellularity. Nature 515:75–79.
https://doi.org/10.1038/nature13884

Heisler IL, Damuth J (1987) A method for analyzing selection in hierarchically
structured populations. Am Nat 130:582–602

Herron MD, Borin JM, Boswell JC, et al (2019) De novo origins of
multicellularity in response to predation. Sci Rep 9:1–9.
https://doi.org/10.1038/s41598-019-39558-8

Koschwanez JH, Foster KR, Murray AW (2013) Improved use of a public good
selects for the evolution of undifferentiated multicellularity. eLife 2:e00367.
https://doi.org/10.7554/eLife.00367

Lande R, Arnold SJ (1983) The measurement of selection on correlated
characters. Evolution 37:1210–1226. https://doi.org/10.2307/2408842

Lewontin RC (1970) The units of selection. Annu Rev Ecol Evol S 1:1–18

Libby E, Rainey PB (2013a) A conceptual framework for the evolutionary origins
of multicellularity. Phys Biol 10:035001

Libby E, Rainey PB (2013b) Eco-evolutionary feedback and the tuning of proto-
developmental life cycles. PLoS ONE 8:e82274

Lloyd EA (1989) A structural approach to defining units of selection. Philos Sci
56:395–418. https://doi.org/10.2307/187992

Lloyd EA, Gould SJ (1993) Species selection on variability. Proc Natl Acad Sci
USA 90:595–599. https://doi.org/10.1073/pnas.90.2.595

Maynard Smith J, Szathmáry E (1995) The Major Transitions in Evolution.
Oxford University Press, Oxford

Michod RE (1999) Darwinian Dynamics, Evolutionary Transitions in Fitness and
Individuality. Princeton University Press, Princeton, N.J.

Mikhailov KV, Konstantinova AV, Nikitin MA, et al (2009) The origin of
Metazoa: a transition from temporal to spatial cell differentiation. Bioessays
31:758–768. https://doi.org/10.1002/bies.200800214

Okasha S (2006) Evolution and the Levels of Selection. Oxford University Press,
Oxford

Okasha S (2012) Emergence, hierarchy and top-down causation in evolutionary
biology. Interface Focus 2:49–54. https://doi.org/10.1098/rsfs.2011.0046

Rainey PB (2007) Unity from conflict. Nature 446:616–616

Rainey PB, Kerr B (2010) Cheats as first propagules: A new hypothesis for the
evolution of individuality during the transition from single cells to
multicellularity. Bioessays 32:872–880. https://doi.org/Doi
10.1002/Bies.201000039

Rainey PB, Rainey K (2003) Evolution of cooperation and conflict in
experimental bacterial populations. Nature 425:72–74

Ratcliff WC, Denison RF, Borrello M, Travisano M (2012) Experimental

evolution of multicellularity. Proc Natl Acad Sci USA 109:1595–1600.

https://doi.org/10.1073/pnas.1115323109

Ratcliff WC, Herron MD, Howell K, et al (2013) Experimental evolution of an

alternating uni- and multicellular life cycle in *Chlamydomonas reinhardtii*. Nat

Commun 4:1–7. https://doi.org/10.1038/ncomms3742

Rose CJ, Hammerschmidt K, Rainey PB (2020) Meta-population structure and

the evolutionary transition to multicellularity. bioRxiv 407163.

https://doi.org/10.1101/407163

Shelton DE, Michod RE (2014) Group selection and group adaptation during a

major evolutionary transition: insights from the evolution of multicellularity in

the volvocine algae. Biol Theory 1–18

Sober E, Lewontin RC (1982) Artifact, cause and genic selection. Philos Sci

49:157–180

Vrba E (1989) Levels of selection and sorting with special reference to the

species level. Oxford surveys in evolutionary biology 6:111–168

Vrba E, Eldredge N (1984) Individuals, hierarchies and processes; towards a

more complete evolutionary theory. Paleobiology 10:146–171

Williams GC (1966) Adaptation and Natural Selection. Princeton University

Press, Princeton, NJ

Wimsatt WC (1980) The units of selection and the structure of the multi-level

genome. PSA: Proceedings of the Biennial Meeting of the Philosophy of Science

Association 1980:122–183. https://doi.org/10.2307/192589

Declarations

Funding

The work was directly supported by the Marsden Fund Council from
government funding administered by the Royal Society of New Zealand, and in
part by grant RFP-12-20.

Conflicts of interest/Competing interests

The authors declare no competing financial interests.

Availability of data and material

Request for materials should be addressed to the corresponding author.

Code Availability

Not applicable

Fig 1. Simplified schematic contrasting the two experimental manipulations carried out by Hammerschmidt et al. (2014). a) In the "cheat-embracing" regime, groups of bacteria reproduced via a germline-like cell: a phenotypically different, non-cooperative cell type. b) In the "cheat-purging" regime, groups of bacteria reproduced via a single-celled bottleneck of the same cell type. Cell fitness is the total cell density, while group fitness is the relative ability of groups to beget offspring groups.

Fig 2. Emergent Relations. When there is no positive correlation between particle and group fitness, a trait-fitness relation at the group-level (a) is emergent; it cannot a by-product of a trait-fitness relation at the particle-level (b).

Fig 3. Detection of Emergent Relations in the Hammerschmidt et al. (2014) experiment. a) In the cheat-purging regime, the transition rate trait correlates with cell fitness and not group fitness: transition rate decreased and group fitness increased as a result of cell-level selection. b) In the cheat-embracing regime, there is no correlation between cell fitness and group fitness: there is an emergent relation between transition rate and group fitness. Small red and blue arrows represent an increase or decrease in the parameters indicated relative to ancestral lines. Long lines represent positive (red) or negative (blue) correlations, and long arrows indicate a likely causal relationship.

Fig 4. Relationship between particle composition and group fitness in the cheat-
embracing (a) and cheat-purging (b) regimes. Ancestral = grey, evolved = black.
Error bars are SEM, based on N=3.

Supplementary Table

Comparison of the fit of linear and quadratic models of the relationship between
the proportion of SM cells and group fitness in the cheat-embracing (CE) and
cheat-purging (CP) regimes using the Akaike's Information Criterion.

Prob correct: Probability that this model is the most likely to have generated the
data.

Parameters	Difference in AICc	N	Prob Correct	df	R^2
CE ancestral	151.5	360			
Linear model			**<0.01%**	358	0.712
Quadratic model			**>99.99%**	357	0.812
CE evolved	13.41	336			
Linear model			**0.12%**	334	0.128
Quadratic model			**99.88%**	333	0.167
CP ancestral	-1.747	360			
Linear model			**70.54%**	358	0.291
Quadratic model			**29.46%**	357	0.292
CP evolved	-1.689	360			
Linear model			**69.94%**	358	0.141
Quadratic model			**30.06%**	357	0.142