Polygenic adaptation and negative selection across traits, years and environments in a long-lived plant species (*Pinus pinaster* Ait., Pinaceae)

Marina de Miguel[†], Isabel Rodríguez-Quilón[‡], Myriam Heuertz[†], Agathe Hurel[†], Delphine Grivet[‡], Juan-Pablo Jaramillo-Correa[*], Giovanni G. Vendramin[§], Christophe Plomion[†], Juan Majada[††], Ricardo Alía[‡], Andrew J. Eckert[‡‡], Santiago C. González-Martínez[†§§]

[†]INRAE, Univ. Bordeaux, BIOGECO, F-33610 Cestas, France

[‡]Department of Forest Ecology and Genetics, Forest Research Centre, INIA, Carretera de la Coruña km 7.5, 28040 Madrid, Spain

[*]Department of Evolutionary Ecology, Institute of Ecology, Universidad Nacional Autónoma de México, AP 70-275 México City, CDMX 04510, Mexico

[§]Institute of Biosciences and Bioresources, Division of Florence, National Research Council, 50019 Sesto Fiorentino (FI), Italy

[††]Sección Forestal, SERIDA, Finca Experimental ''La Mata'', 33820 Grado, Principado de Asturias, Spain

[‡‡]Department of Biology, Virginia Commonwealth University, Richmond, VA 23284, USA

[§§]Corresponding author

Running title: Polygenic adaptation in maritime pine

Keywords: heritability, local adaptation, maritime pine, polygenicity, natural selection

Corresponding author:

Santiago C. González-Martínez

UMR 1202 INRAE – Univ. Bordeaux

69 route d'Arcachon, F-33610 Cestas, France

Tel: +33 (0) 5 35 38 53 20

santiago.gonzalez-martinez@inrae.fr

Abstract

A decade of association studies in multiple organisms suggests that most complex traits are polygenic; that is, they have a genetic architecture determined by numerous loci distributed across the genome, each with small effect-size. Thus, determining the degree of polygenicity and its variation across traits, environments and years is useful to understand the genetic basis of phenotypic variation. In this study, we applied multilocus approaches to estimate the degree of polygenicity of fitness-related traits in a long-lived plant (*Pinus pinaster* Ait., maritime pine) and to analyze how polygenicity changes across environments and years. To do so, we evaluated five categories of fitness-related traits (survival, height, phenology-related, functional, and biotic-stress response traits) in a clonal common garden network, planted in contrasted environments (over 12,500 trees). First, most of the analyzed traits showed evidence of local adaptation based on Q_{ST}-F_{ST} comparisons. Second, we observed a remarkably stable degree of polygenicity, averaging 6% (range of 0-27%), across traits, environments and years. As previously suggested for humans, some of these traits showed also evidence of negative selection, which could explain, at least partially, the high degree of polygenicity. The observed genetic architecture of fitness-related traits in maritime pine supports the polygenic adaptation model. Because polygenic adaptation can occur rapidly, our study suggests that current predictions on the capacity of natural forest tree populations to adapt to new environments should be revised, which is of special relevance in the current context of climate change.

Introduction

Population adaptive responses to environmental changes depend on the genetic architecture of fitness-related traits (Hayward and Sella 2019). Although not initially conceived for the study of adaptation, genome-wide association studies (GWAS) have provided essential information to understand the genetic basis of complex traits. The implementation of GWAS allowed the identification of genetic variants affecting fitness-related traits, their allele frequencies, the magnitudes of their effects, and their interactions with one another and the environment. Examples exist for humans (reviewed by Visscher *et al.* 2017), other animals (e.g. Sharma *et al.* 2015; Pitchers *et al.* 2019) and plants (e.g. González-Martínez *et al.* 2006; Huang and Han 2014; Alonso-Blanco *et al.* 2016). Surprisingly, in some species, such as humans and forest trees (Resende *et al.* 2012; Lind *et al.* 2018), the genetic variants associated with phenotypic variation accounted for only small fractions of trait heritability, as estimated through pedigree analysis, causing the so-called 'missing heritability' paradox (Maher 2008). Several explanations have been provided to solve this paradox (Manolio *et al.* 2009; Brachi *et al.* 2011; Björkegren *et al.* 2015; Pallares 2019). In particular, different sources of evidence point to polygenicity, i.e. trait architecture determined by a large number of variants, each with a small effect-size, as a potential reason for the low levels of heritability explained by current GWAS, which would thus be unpowered to detect most causal variants (Yang *et al.* 2010; Shi *et al.* 2016; Boyle *et al.* 2017).

The study of adaptation has traditionally been addressed from contrasting research paradigms (Höllinger *et al.* 2019). While quantitative genetic approaches view adaptation as the result of changes in allele frequencies at an idealized infinite number of loci, each with infinitesimal effects on fitness (Fisher 1918), population genetic approaches made more emphasis in the detection of selective sweeps, where new beneficial mutations rapidly become fixed at a small number of loci (Smith and Haigh 1974). The hypothesis that natural selection (mostly) acts

through subtle allele frequency shifts on standing genetic variation at numerous loci

distributed across the genome has been suggested in previous evolutionary studies (Orr and

Coyne 1992; e.g. McKay and Latta 2002; Le Corre and Kremer 2003). Nevertheless, it was

Pritchard *et al.* (2010) who first brought together population and quantitative genetic theory

with conclusions from GWAS to formulate a new model for the study of adaptation – the

polygenic adaptation model. Under this model, some genes may harbor new mutations that

have been fixed by natural selection, but the most common pattern would be the genome-wide

increase of favored alleles, without the fixation of most causative variants. Thus, the expected

genome-wide footprint resulting from natural selection would not be that of a classical hard

sweep, but would rather involve a large number of causal variants, each with subtle allele

frequency changes (Pritchard *et al.* 2010; Pritchard and Rienzo 2010; Hermisson and

Pennings 2017).

Several new methods have been developed to detect the genetic signatures of natural selection

under the polygenic adaptation model (Guan and Stephens 2011; Turchin *et al.* 2012; Daub *et*

al. 2013; Berg and Coop 2014; Field *et al.* 2016; Zeng *et al.* 2018; Edge and Coop 2019;

Speidel *et al.* 2019; Lloyd-Jones *et al.* 2019). Applications using these methods, however,

were often unknowingly biased by subtle patterns of population structure (Liu *et al.* 2018;

Josephs *et al.* 2019; Rosenberg *et al.* 2019; Berg *et al.* 2019a; Sohail *et al.* 2019).

Nonetheless, even considering the inflation of polygenic signals due to unrecognized

population structure, mounting evidence over the last decade using a variety of

methodologies, supports the polygenic adaptation model in a diversity of organisms, such as

humans (Hancock *et al.* 2010a, b; Daub *et al.* 2013; Shi *et al.* 2016; Zeng *et al.* 2018;

Gnecchi-Ruscone *et al.* 2018; Berg *et al.* 2019b), insects (Friedline *et al.* 2019), molluscs

(Bernatchez *et al.* 2019), model plants (He *et al.* 2016), crops (Josephs *et al.* 2019; Wisser *et*

al. 2019), and forest trees (Lind *et al.* 2017; De La Torre *et al.* 2019). However, there are still

multiple open questions regarding the degree of polygenicity at adaptive traits, the distribution

of effect sizes the involved loci, and how the genetic architecture changes under varying

selective forces, especially for non-model species (Lind *et al.* 2018).

Following the expectations of the polygenic adaptation model, the heritability of complex

traits is often associated with loci that are widespread across the whole genome, also

including SNPs located in genes and pathways that do not show a clear functional connection

to the trait of interest (Boyle *et al.* 2017). The omnigenic model, formulated by Boyle *et al.*

(2017), provides a plausible hypothesis to explain this. The interconnection of gene regulatory

networks implies that the vast majority of expressed genes likely influence the functions of

core genes directly linked to fitness-related traits. Nevertheless, the study of polygenic

adaptation at the pathway level is useful to identify gene sets of special relevance for

adaptation (Daub *et al.* 2013). In particular, polygenic adaptation at pathway level has been

proved especially useful to study non-model species, as reliance on selective sweep models

often led to poor inferences (Hämälä *et al.* 2020). For example, Mayol *et al.* (2020) detected

signatures of polygenic selection at the pathway level in English yew (*Taxus baccata* L.)

Taxus baccata and identified negative selection as an important mechanism driving the

pathway-level signal. Similarly, negative selection has been identified as a pervasive

mechanism determining the polygenic architecture of fitness-related traits in humans (Zeng *et*

al. 2018; O'Connor *et al.* 2019). In particular, negative selection has been proposed to favor

polygenicity in complex traits by removing large-effect variants, because of their deleterious

effects, while small-effect variants would remain unaffected; a process named 'flattening', as

the genetic signal is 'flattened' relative to the underlying biology (O'Connor *et al.* 2019).

Despite theoretical advances and the development of new methods to study polygenic

adaptation, the addressed questions remain constrained by the specific life-history traits of a

few model species. Maritime pine (*Pinus pinaster* Ait.) is an ideal case study to investigate

polygenic adaptation in an ecologically and economically important group of species, the

forest trees: it is a long-lived plant inhabiting nearly undomesticated random-mating

populations with high genetic diversity (González-Martínez *et al.* 2002; Jaramillo-Correa *et*

al. 2015). It expanded from several isolated glacial refugia, and it is now distributed across

the western Mediterranean Basin and the European Atlantic front in scattered populations

under contrasting environments (Bucci *et al.* 2007). In addition, an artificial clonal

propagation program in maritime pine allowed us to estimate precisely variance components

and investigate selective forces driving trait evolution under contrasting environments.

Specifically, we i) tested the hypothesis that most complex adaptive traits in a long-lived plant

are polygenic, providing a first estimate of the degree polygenicity in a forest tree, and ii) that

their genetic architecture is mostly driven by negative selection. We then iii) investigated how

these patterns change with time and through environmental settings, which is of special

relevance for long-lived organisms, such as forest trees.

Materials and Methods

Clonal common garden network (CLONAPIN)

We studied phenotypic variation in a clonal common garden network (CLONAPIN)

composed of five sites covering the natural environmental range of maritime pine, from harsh

Mediterranean climates to mild Atlantic ones. Common gardens comprise trees from 36

populations, sampled across the species natural distribution and covering the six previously

identified gene pools for this species (Jaramillo-Correa *et al.* 2015; Supplemental Figure S1).

Open-pollinated seeds were collected in natural stands, and germinated in a nursery; then one

seedling per open-pollinated family was selected and vegetatively propagated by cuttings

(following Majada *et al.* 2011). A total of 535 genotypes (clones) belonging to 35 populations

were used to establish four of the clonal common gardens (three sites in Spain: Cabada,

Cáceres and Madrid; and one in Portugal: Fundão; Table 1) with eight ramets per clone set in

a randomized complete block design (*N*=4,273 trees). These clonal common gardens were

planted in 2010. In 2011, a fifth common garden was established in Pierroton (France),

comprising 443 clones from all 36 populations (*N*=3,434 trees). Common gardens in Cabada,

Fundão and Pierroton are located in the Atlantic region, with high annual rainfall and mild

temperatures. Common gardens in Cáceres and Madrid are located in continental areas under

Mediterranean influence, with large seasonal temperature oscillations and a marked summer

drought. In addition, clay soils in Cáceres hampered plant growth and diminished survival

(Table 1).

Phenotypic evaluation

A total of 28 phenotypic trait-environment combinations were evaluated in this study.

Assayed phenotypic traits were classified into five groups: survival, height, phenology-

related, functional, and biotic-stress response (see Supplemental Table S1 for an exhaustive

list of the measured traits). In brief, tree survival and height were evaluated in the five

common gardens (including different years in Pierroton, with measures taken in 2013, 2015,

and 2018). Phenology-related traits were evaluated in the Atlantic sites only (Cabada, Fundão

and Pierroton), including different years of evaluation in Pierroton (2015 and 2017). In

Cabada and Fundão, growth phenology was estimated through the presence of polycyclism

(i.e. the ability for a plant to produce several flushes in the same growing season; (Girard *et*

al. 2011)) and a phenology growth index (1):

172
$$\text{Phenology Growth Index} = \frac{spring\ \text{growth}}{total\ \text{growth}} = \frac{(tree\ \text{height}\ may_n - tree\ \text{height}\ dec_{n-1})}{(tree\ \text{height}\ dec_n - tree\ \text{height}\ dec_{n-1})} \qquad (1)$$

where *may* and *dec* correspond to the months May and December of the year n and the year n-

1, respectively.

In Pierroton, phenology of bud burst was estimated through a scale ranging from 0 to 5 (0:

bud with no winter elongation, 1: elongation of the bud, 2: emergence of brachyblasts, 3:

brachyblasts begin to separate, 4: elongating needles, 5: total elongation of the needles) (see

Hurel *et al.* 2019). The first Julian day at each stage (S1 to S5) was scored for each tree.

Julian days were converted into accumulated degree-days (with base temperature 0°C) from

the first day of the year, to take into account the between-year variability in temperature. The

number of degree-days between stages 1 and 4 defines the duration of bud burst. Daily mean

temperatures to calculate accumulated degree-days were downloaded from the nearest

climatic station (located just a few hundred meters from the common garden, station

33122004 of the INRAE Agroclim database: https://www6.paca.inrae.fr/agroclim/Les-outils).

Functional traits, including nitrogen and carbon content and isotopic composition ($\delta^{15}N$ and

$\delta^{13}C$, respectively), as well as specific leaf area (SLA, a measure of leaf area per unit of dry

mass), were evaluated in the common garden located in Fundão (Portugal). A bulk of five

needles positioned 10 cm below the upper part of the shoot to avoid sampling bias (Warren *et*

al. 2001) were sampled and prepared in a standard way for analysis (Brendel 2001).

Determination of carbon content and isotopic composition was performed with a mass

spectrometer at the University of Colorado isotope laboratory. Raw values were corrected by

their position in the plate according to the standards, and this value was used for the

subsequent analysis. SLA is an estimation of the compromise among light capture, CO_2

assimilation, and the restrictions imposed by water loss through transpiration (Sefton *et al.*

2002). Low SLA suggests high leaf construction cost, and thus higher stress tolerance (Díaz *et*

al. 2016). Thus, this key leaf trait is also associated with fitness components, such as tree

survival (Greenwood *et al.* 2017). Given that there is a positive relationship between $\delta^{13}C$ and

water use efficiency (Farquhar and Richards 1984), $\delta^{13}C$ has been widely used as a surrogate

to study tree adaptation to water-limiting environments (e.g. Aranda *et al.* 2010; Walker *et al.*

2015). Similarly, $\delta^{15}N$ is an indirect index related to the nitrogen cycle (Craine *et al.* 2015).

Assessment of biotic-stress response in a high number of trees is logistically complex.

Therefore, it was evaluated only for a subset of clones (see Supplemental Table S1) in the

Pierroton common garden (France). This common garden was selected because of the

importance of the Landes region in maritime pine breeding for wood production. Biotic-stress

response was evaluated based on susceptibility to two major pine pathogens, *Diplodia sapinea*

and *Armillaria ostoyae*, as well as the incidence of the defoliator pest, *Thaumetopoea*

pityocampa (pine processionary moth) (see Hurel *et al.* 2019 for details). *D. sapinea* causes

several diseases in conifers, which may be exacerbated under climate change and compromise

pine forest health (Desprez-Loustau *et al.* 2006). Susceptibility to *D. sapinea* was evaluated

following controlled inoculation as the lesion extent around the inoculated site (hereafter

referred as necrosis) and a scalar notation of needle discoloration (0: no discoloration, 2: some

needles around the necrosis were discolored, and 3: all needles around the necrosis were

discolored). *A. ostoyae* is a conifer root pathogen causing growth cessation and eventually

death (Heinzelmann *et al.* 2019). To evaluate the incidence of this pathogen, *A. ostoyae*

mycelium culture was prepared in plastic jars. The level of humidity observed in the plastic

jar was visually scored as dry, medium or humid. Susceptibility to *A. ostoyae* was assessed

after controlled inoculation as the lesion length in the sapwood (i.e. wood browning, hereafter

also referred as necrosis). We accounted for the potential influence of variation in humidity on

wood browning by including the level of humidity in the jar as a covariate for *A. ostoyae*

susceptibility analysis (see below). Finally, the pine processionary moth is an insect that

rapidly defoliates pines leading to forest decline (Jacquet *et al.* 2013). The presence or

absence of pine processionary moth nests in the tree crowns was assessed in March 2018.

DNA extraction and SNP genotyping

Needles were collected from one ramet per clone in the Cabada common garden (N=523).

Genomic DNA was extracted using the Invisorb® DNA Plant HTS 96 Kit/C kit (Invitek

GmbH, Berlin, Germany). A 9k *Illumina Infinium* SNP array developed by Plomion *et al.*

(2016b) was used for genotyping. This array was constructed using previously identified and

newly *in silico*-developed SNPs, either from randomly screened EST sequences or

specifically detected at candidate genes for adaptation to biotic and abiotic factors (see

Plomion *et al.* 2016b for further details). Genotyped SNPs covered all 12 chromosomes of *P.*

pinaster according to previous linkage mapping (Plomion *et al.* 2016b). For this study, 6,100

SNPs were finally retained following standard filtering (GenTrain score > 0.35, GenCall50

score > 0.15 and Call frequency > 0.85) and removal of SNPs with uncertain clustering

patterns (visual inspection using *GenomeStudio v. 2.0*). Individuals with more than 15%

missing data were also removed. This resulted in 5,165 polymorphic SNPs that were included

in the estimation of molecular population differentiation (F_{ST}) and the polygenic association

study.

Quantitative genetics analysis

Genetic components of the phenotypic variance were estimated using Generalized Linear

Mixed-Effects Models (GLMM) fitted in a Bayesian framework using Markov chain Monte

Carlo (MCMC) methods. The model, described in equation (2), was implemented for those

phenotypic traits evaluated at multiple sites of the CLONAPIN common garden network (see

Supplemental Table S1). To estimate the genetic control of the genotype-by-environment

(G×E) interactions, the model described in equation (3) was fitted for those traits measured at

all sites of the CLONAPIN common garden network (i.e. height and survival).

$$y_{ijkl} = \mu + S_i + S(B)_{ij} + P_k + P(C)_{kl} + S_i * C_l + \varepsilon_{ijkl} \qquad (2)$$

$$y_{ijkl} = \mu + S_i + S(B)_{ij} + P_k + P(C)_{kl} + S_i * P_k + S_i * C_l + \varepsilon_{ijkl} \qquad (3)$$

where, for a given trait y, μ denotes the overall phenotypic mean, S_i refers to the fixed effect

of site i, B_j represents the random effect of experimental block j nested within site i, P_k is the

random effect of population k, C denotes the random effect of clone l nested within

population k, and ε is the residual effect.

Simplified models with or without covariates represented by equations (4) and (5) were

implemented for phenotypic traits measured in just one site of the CLONAPIN common

garden network (see Supplemental Table S1).

$$y_{ijk} = \mu + B_i + P_j + P(C)_{jk} + \varepsilon_{ijk} \qquad (4)$$

$$y_{ijk} = \mu + B_i + cov + P_j + P(C)_{jk} + \varepsilon_{ijk} \qquad (5)$$

Where, for a given trait y, μ denotes the overall phenotypic mean, B_i represents the fixed

effect of experimental block i, P_j is the random effect of population j, C denotes the random

effect of clone k nested within population j, and ε is the residual effect. In the model

represented by equation (5), *cov* represents a covariate implemented when modeling the

presence of pine processionary moth nests (i.e. an estimate of tree height) and necrosis caused

by *A. ostoyae* (i.e., level of humidity in the experimental jar).

All models were fitted with the R package *MCMCglmm* (Hadfield 2010). Phenotypic traits

showing Gaussian distributions where modeled using the identity link function, while

phenotypic traits exhibiting a binomial distribution (survival, polycyclism, *D. sapinea* needle

discoloration, and presence or absence of pine processionary moth) were modeled either with

logit or *probit* link functions (see Supplemental Table S2 for an exhaustive list of model

parameter specifications). Multivariate-normal prior distributions with mean centered at zero

and large variance matrices (10^8) were used for fixed effects. For ordinal traits, a Gelman

prior for the variance of fixed effects was set, as suggested by Gelman et al. (2008). Inverse

Wishart non-informative priors were used for the variances and covariances of random

effects, with the matrix parameter V set to 1, and a parameter n set to 0.002 (Hadfield 2010).

Parameter expanded priors were used to improve the chain convergence and mixing, as

suggested by Gelman (2006) for models with near-zero variance components. Priors with a

larger degree of belief parameter (n set to 1), specifying that a large proportion of the

variation is under genetic control (as suggested by Wilson *et al.* 2010) did not change the

results (data not shown). Models were run for at least 550,000 iterations, including a burn-in

of 50,000 iterations and a thinning interval of 100 iterations. Four chains per model were run

to test for parameter convergence. The potential scale reduction factor (psrf) was consistently

below 1.02 for all the models (Supplemental Table S2) (Gelman and Rubin 1992).

Variance components were then used to compute broad-sense heritability (H^2) as (6):

282
$$H^2 = \frac{\sigma^2_{clone}}{(\sigma^2_{clone} + \sigma^2_e)} \qquad (6)$$

where σ^2_{clone} is the variance among clones within populations and σ^2_e the residual variance.

For estimating broad-sense heritability for traits following a binomial distribution, we

included an extra term in the denominator ($+ \pi^2/3$) to account for implicit *logit* link function

variance; similarly, we added one to the denominator to account for the *probit* link function

(Nakagawa and Schielzeth 2010).

The GLMMs described above were used to estimate genetic values using Best Linear

Unbiased Predictors (BLUPs) (Henderson 1973; Robinson 1991). The genetic value of each

clone was defined as the population BLUP plus the clone BLUP. BLUPs for G×E, were

obtained from equation (3) and calculated following equation (7).

292
$$G \times E\ BLUP = \left(\frac{\sum BLUP_{pop\ atl}}{N\ atl} - \frac{\sum BLUP_{pop\ med}}{N\ med}\right) + \left(\frac{\sum BLUP_{clone\ atl}}{N\ atl} - \frac{\sum BLUP_{clone\ med}}{N\ med}\right) \qquad (7)$$

where $BLUP_{pop}$ *atl* is the population BLUP in sites under Atlantic climate (Cabada, Fundão,

and Pierroton), $BLUP_{pop}$ *med* the population BLUP in sites under Mediterranean climate

(Cáceres and Madrid), $BLUP_{clone}$ *atl* the clone BLUP in sites under Atlantic climate,

$BLUP_{clone}$ *med,* the clone BLUP in sites under Mediterranean climate, *N atl* the number of

sites under Atlantic climate, and *N med* the number of sites under Mediterranean climate.

Parameter estimates from quantitative genetics analyses are presented as the mode of the

posterior distribution; 95% credible intervals were computed as the highest density region of

each posterior parameter distribution.

Q_{ST}-F_{ST} comparison

Molecular population differentiation (F_{ST}) was estimated according to Weir and Cockerham

(1984) using the 5,165 SNPs from the *Illumina Infinium* SNP array and the diveRsity R

package (Keenan *et al.* 2013). The 95% confidence interval of the global F_{ST} estimate was

computed by bootstrapping across loci (1,000 bootstrap iterations). Quantitative genetic

differentiation among populations was calculated following Spitze (1993) using the variance

components estimated from the previously described models (equations 2-5):

$$Q_{ST} = \frac{\sigma^2_{pop}}{\sigma^2_{pop} + 2\sigma^2_{clone}} \qquad (8)$$

where σ^2_{pop} is the variance among populations, and σ^2_{clone} is the variance among clones within

populations. Quantitative (Q_{ST}) and molecular (F_{ST}) genetic differentiation among populations

were considered significantly different when Q_{ST} and F_{ST} posterior distributions had non-

overlapping 95% confidence intervals.

Polygenicity across traits, years and environments

Polygenicity was evaluated as the proportion of SNPs with non-zero effects on phenotypic

traits. First, we conducted posterior inference via model averaging and subset selection

(VSR), as implemented in piMASS software (Guan and Stephens 2011). This method allows

to identify combinations of SNPs likely affecting a phenotype and to estimate the proportion

of trait variance explained by the SNPs in the data set. Hereafter, we referred to this quantity

as the genetic explained variance (*GEV*), which, in this study, represents the BLUP variance

explained by SNP additive effects. Second, we used the Bayesian mixed linear model (MLM)

framework developed by Zeng *et al.* (2018) as implemented in CGTB 2.0 software. This last

model simultaneously estimates: i) SNP-based heritability (considering the SNPs with non-

zero effects on the trait), hereafter referred as *GEV*, analogously to VSR estimates, ii)

polygenicity (as defined above), and iii) the relationship between SNP effect-size and minor

allele frequency (*S*, a common indicative of negative selection). When negative selection is

operating, *S* is expected to be negative, as most new mutations are deleterious and high-effect

SNPs are kept at low frequencies. Estimates with 95% credible intervals of parameter

posterior distributions not overlapping zero were considered as significant. Prior to these

analyses, neutral population genetic structure was accounted for by running linear models

relating the genetic values for each trait (with site and block effects removed) to the admixture

coefficients for each clone (*Q*-scores) obtained using a STRUCTURE run for *K*=6 based on

neutral markers (see Jaramillo-Correa *et al.* 2015 for further details). From this linear model,

we extracted the normalized residuals for each trait, as recommended in piMASS manual.

Analyses were run separately for different traits, years, and environments (see Supplemental

Table S1). VSR models were run for 2,000,000 iterations with a burn-in of 100,000 iterations

and a thinning interval of 100. After several preliminary runs, the maximum number of SNPs

included in VSR models was fixed to 2,000 (i.e. maximum allowed polygenicity of ~40%).

MLM models were run for 500,100 iterations, including a burn-in of 100 iterations, and a

thinning interval of 10 iterations. Parameter estimates from both VSRs and MLMs were

presented as the median of the posterior distribution, instead of the mode, for better handling

of bimodal distributions (Supplemental Figure S2). The 95% credible intervals were

computed as the highest density region of the posterior parameter distribution, as above.

Annotation and gene function enrichment at pathway level

The transcripts containing the 5,165 polymorphic SNPs were downloaded from SustainPine

v.3.0 database (Canales *et al.* 2014). DNA sequences were translated with BioEdit v. 7.2.6

(Hall 1999) and submitted to BlastKOALA (Kanehisa *et al.* 2016) for annotation and

functional characterization using InterPro annotations, GO terms, and KEGG pathway

identification. Annotations were compared with those available at SustainPine, and

conflicting cases were examined individually by privileging similarity to genes correctly

identified in other conifers or forest trees. Contigs with no clear annotations (e.g. hypothetical

or unknown proteins, or unsolved conflicting annotations) were removed from the database.

For the retained contigs, the top-two KEGG terms were used for assignation to one or more

specific metabolic pathways/modules based on KEGG orthology. Genes for which no hit with

KEGG database was found, were assigned to metabolic pathways/modules based on the

InterPro annotation. We privileged metabolic pathways/modules that could be unequivocally

assigned to a given phenotypic response (e.g. circadian rhythm to bud phenology or pathogen

interaction to biotic stress response) or linked to various stress responses (e.g. DNA

recombination and repair, ubiquitin system or transcription factor machinery to survival and

biotic stress response). In total, seventeen pathways/modules were retained containing a total

of 628 (19.7% out of 3,194) genes, with 1,233 polymorphic SNPs (Supplemental Table S3).

For enrichment tests using polysel (Daub *et al.* 2013), the seventeen pathways/modules were

defined as gene sets. First, we computed two statistics at the gene level (i.e. objStat in polysel)

based on the per-SNP estimates obtained from the VSR implemented in piMASS: the

maximum, over all SNPs included in a gene, of the Rao-Backwellized posterior probability of

inclusion *maxpostrb*, and the maximum of the absolute value of Rao-Backwellized effect size

maxabsbetarb. To account for a weak correlation of these statistics with the number of SNPs

per gene, we used the AssignBins and RescaleBins functions in polysel, which automatically

assigns gene scores (objStat) into bins defined from the number of SNPs per gene. We then

rescaled scores within bins and computed the sum(objStat) of each statistic over all genes per

gene set. Since the sum(objStat) for random gene sets (sizes n = 10, 50, 250 genes) was not

normally distributed, we built empirical null distributions by randomly sampling gene sets of

the same size as the sets to be tested. Then, we performed one-sided tests evaluating whether

the observed sum(objStat) was smaller than the 5^{th} or larger than the 95^{th} percentile of the

sum(obStat) null distribution. Higher-tail significant results for *maxpostrb* indicate gene sets

enriched with higher overall probability of being selected during the VSR procedure

implemented in piMASS. Higher-tail significant results for *maxabsbetarb* points to gene sets

enriched with higher overall SNP effect-sizes. Contrarily, lower-tail significant results for

both statistics suggests conserved gene sets, containing genes with smaller overall probability

of inclusion or SNP effect-size estimates. We report *p*-values based on this comparison, as

well as *q*-values from a False Discovery Rate (FDR) approach implemented in the R package

qvalue (R Core Team 2019). The level of connection between gene sets was weak with only

four genes associated with more than one gene set (633 gene – gene set combinations for 628

genes). For this reason, we did not assess enrichment for pruned gene sets (see Daub *et al.*

2013).

Results

Broad-sense heritability and genetic differentiation among populations

All traits had low to moderate estimates of broad-sense heritability (Supplemental Table S1),

with the exception of nitrogen and carbon amount that did not show genetic variation.

Consequently, polygenic association methods failed to converge for these two traits and they

were excluded from further analyses. H^2 ranged from 0.32 for bud burst measured in 2015 to

zero for survival in the French Atlantic environment in 2013. Interestingly, survival showed

significant estimates of H^2 only in the sites under (harsher) Mediterranean climate. The

highest H^2 estimates were observed for phenology-related traits followed by tree height. H^2 for

a given trait varied across environments (e.g. height, survival and phenology-related traits) but

showed little variation across years (Supplemental Table S1).

The global F_{ST} was 0.112 (95% confidence interval: 0.090 - 0.141). All groups of phenotypic

traits, excepting survival, had at least one trait with statistically higher Q_{ST} than F_{ST}

(Supplemental Table S1). The highest Q_{ST} was obtained for susceptibility to *D. sapinea*

infection measured as necrosis length, followed by $\delta^{13}C$ and tree height, which also showed

similar Q_{ST} values across environments and tree ages (Figure 1).

Genetic architecture (polygenicity) of adaptive traits

Polygenicity estimates were consistent between the VSR and MLM methods (Supplemental

Table S4). Both methods showed substantial polygenic control for most of the phenotypic

traits, with an average of 6% (0-27%) of the genotyped SNPs having non-zero effects.

Significant polygenicity was found in all five trait categories for at least one trait (Figures 2

and 3; Supplemental Table S4). Polygenicity for height was stable across environments and

years, when measured multiple times under the same environment (i.e. in the French Atlantic

common garden) (Figure 3). Along the same line, polygenicity for phenology-related traits

and tree survival also remained stable across environments, although 95% credible intervals

overlapped zero in some cases. The low polygenicity values observed for survival in the

French Atlantic common garden are probably a consequence of the low levels of phenotypic

variability in this site, with almost no mortality (97.12% of planted trees were alive at the

evaluation time, Supplemental Table S1). Polygenicity was heterogeneous for biotic-stress

response and functional traits (Figure 2). For instance, susceptibility to *D. sapinea* was more

polygenic than to *A. ostoyae* or than incidence of pine processionary moth. For functional

traits, SLA and δ^{15}N showed the highest levels of polygenicity, while δ^{13}C showed a

considerably lower proportion of SNPs with non-zero effects.

In addition, *GEV* was consistent between methods, although VSR tended to render higher

values (Supplemental Table S4). On average *GEV* was 0.38 across traits, with a minimum of

0.018 for survival in the French Atlantic environment in 2018, and a maximum of 0.99 for *D.*

sapinea necrosis. *GEV* estimated with the VSR method for the G×E component on tree height

(considering Atlantic versus Mediterranean environments) was low but significant (median =

0.238, 95% credible interval = 0.043 - 0.409), indicating some SNPs with significant effects

on growth plasticity. However, this result could not be confirmed with the MLM method.

Moreover, *GEV* for the G×E component on tree survival was not significant with any model.

Polygenicity and *GEV* were positively and consistently correlated for both VSR and MLM

models (Figure 4). This positive correlation suggested that SNP-based heritability is mainly

determined by genetic variants with similarly small effects, and that differences in

polygenicity across traits are mostly accounting for differences in explained genetic variance,

rather than the distribution of SNP effect-size (Supplemental Figures S2 and S3).

Evidence of negative selection

The correlation between SNP effect-size and minor allele frequency (MAF), *S*, was used to

identify the type and mode of natural selection acting upon phenotypic traits. Out of the 28

assayed traits, we were able to estimate *S* through the MLM method for 19 of them. Estimates

ranged from -1.68 (bud burst in 2017) to 0.55 (tree survival in French Atlantic environment),

but only seven traits from four out of five trait categories (survival, height, phenology-related,

and functional traits) were significant (Figure 5). No significant effect was observed for any

trait belonging to the biotic-stress response category. Remarkably, all seven significant

estimates of S were negative (ranging from -1.68 for bud burst in 2017 to -0.99 for survival in

the Iberian Atlantic environment).

Estimates of S for height were consistent across years and environments. However, S

estimated for tree survival was only significant in the Iberian Atlantic environment. For

phenology-related traits, S was significant only for bud burst measured in 2017 (Figure 5).

These results contrast with the consistent level of polygenicity for all these traits across years

and environments. Interestingly, our results suggest a stable polygenic architecture, but an

environment and year-dependent impact of negative selection at some traits.

Gene function enrichment at pathway level

Tests for gene function enrichment at the pathway level provided significant results for

survival in the Iberian Atlantic environment, phenology-related and biotic-stress response

traits, and height in the French Atlantic and Mediterranean environments. Genes coding for

transcription factors showed higher probability of being included in the VSR models

(*maxpostprb)* and higher estimated SNP effect-sizes (*maxabsbetarb)* for survival in the

Iberian Atlantic environment (Table 2). Two gene sets associated to bud burst in 2015 showed

signals of polygenic selection: *monolignol biosynthesis*, which had high overall values of both

maxpostprb and *maxabsbetarb*, and *glycan metabolism*, which showed low overall

maxabsbetarb estimates (Table 2). Furthermore, phenology growth index was associated with

enrichment for genes related to *cell growth and death*, *DNA recombination and repair* and *UV*

response, which mostly have low *maxabsbetarb* values (Table 2). *D. sapinea* susceptibility

was associated with enrichment of genes, with high overall *maxabsbetarb* and *maxpostprb,* in

the *ubiquitin system* for *D. sapinea* necrosis*,* and in the *signal transduction* and *flavonoid*

biosynthesis gene sets for *D. sapinea* needle discoloration (Table 2). Interestingly, height was

enriched for genes from different pathways when measured in contrasting environments. For

instance, in the French Atlantic environment genes coding for *transcription factors* showed

high *maxabsbetarb* and *maxpostprb*, while genes within the *cytoskeleton* pathway showed

overall low *maxabsbetarb* values in the Mediterranean environment.

Discussion

Unraveling the genetic architecture of adaptive traits is challenging because of the difficulty

to identify variants with small effect-sizes using GWAS. Here, we addressed this challenge

obtaining precise phenotypic information (over 12,500 trees were evaluated) for an extensive

number of fitness-related traits measured on clonal replicates. Specifically, we tested if a high

proportion of the genetic variance of fitness-related traits in a long-lived forest tree (maritime

pine) can be explained by a large number of small size-effect variants, in line with the

polygenic adaptation model. We also tested whether negative selection is pervasive for such

polygenic traits. Our results showed patterns of local adaptation for most of the analyzed

traits, highlighting its relationship with fitness, and also revealed a high and remarkably stable

degree of polygenicity, across traits, years, and environments. Moreover, using two

complementary multilocus approaches we accounted for a considerable proportion of the

heritability estimated for these highly polygenic traits, and identified negative selection as a

key driver of local adaptation.

Evidence of local adaptation in maritime pine

All phenotypic categories presented significant within-population genetic variation (i.e.

broad-sense heritability), and were consequently susceptible to respond to natural selection

(Visscher *et al.* 2008). Estimates of heritability were consistent with previous results for these

traits in forest trees (reviewed by Lind *et al.* 2018). In addition, our results were consistent

with adaptive differentiation ($Q_{ST} > F_{ST}$) for 11 out of 26 analyzed traits, involving four out of

the five trait categories (no evidence for survival traits). These results are in accordance with

reports of pervasive local adaptation in forest trees (Savolainen *et al.* 2007, 2013; Alberto *et*

al. 2013; Lind *et al.* 2018).

The stability of Q_{ST} estimates for height across environments and years highlights the strength

of directional selection for height in this species; a trait that can thus be used for the

delimitation of conservation and management units (Rodríguez-Quilón *et al.* 2016).

Contrarily, phenology-related traits showed contrasting estimates of Q_{ST} depending on the

environment and year of measurement. This result highlights that the evolutionary forces

driving population genetic differences in phenology-related traits are environmentally and

temporally-dependent, which can slow-down attaining phenotypic optima under rapidly

changing climates. Polygenic adaptation could be specially relevant for these traits because it

can produce rapid phenotypic changes, as it only requires small adjustments in allele

frequencies in the contributing loci rather than selective sweeps on new mutations (Jain and

Stephan 2017; Dayan *et al.* 2019; Wisser *et al.* 2019).

Unexpectedly, survival, a trait directly related with a component of fitness (i.e. viability), did

not show evidence of local adaptation in maritime pine. The low levels of phenotypic

variability observed for survival in this study may explain these results. Future studies should

focus on quantitative evaluations of survival (e.g. adding a time-frame, such as time until

death or order of dead trees) to better gather the complexity of this trait, and be able to discern

genetic differences among populations. The strong selective pressure in the Mediterranean

region exacerbated genetic differences in survival among clones and resulted in slightly

higher estimates of heritability (similarly to Gaspar *et al.* 2013). Additionally, we observed

significant phenotypic plasticity for height and survival, the two traits measured in all five

experimental sites. While our results hinted a heritable component for plasticity, this question

still deserves further investigation to elucidate the importance of phenotypic plasticity in the

adaptive response of maritime pine to changing environmental conditions (Alía *et al.* 2014;

Vizcaíno-Palomar *et al.* 2019).

Two traits in particular had remarkably high levels of adaptive genetic differentiation among

populations, δ^{13}C and *D. sapinea* necrosis (Figure 1), but genetic variation within populations

was low, compromising their adaptive potential. These traits deserve special attention because

of the implication of water-use efficiency in drought resistance (reviewed by Plomion *et al.*

2016a) and the new pathogenic outbreaks of *D. sapinea* expected on maritime pine

plantations fostered by climate change (Fabre *et al.* 2011; Brodde *et al.* 2019). In contrast to

our findings, a lack of adaptive genetic differentiation for δ^{13}C was previously reported for

maritime pine by Lamy *et al.* (2011), as well as for broad-leaved trees (Torres-Ruiz *et al.*

2019). Although this disagreement may be influenced by the much larger number of

populations we analyzed (see Whitlock and Guillaume 2009) as compared to Lamy *et al.*

(2011), we cannot rule out discrepancies due to the estimation of total genetic variance in our

study (i.e. based on clones), instead of additive genetic variance. Nevertheless, non-additive

genetic effects in maritime pine traits related to drought resistance have been reported to be of

little importance (Gaspar *et al.* 2013), and they should not have affected our estimates.

Genetic architecture (polygenicity) of fitness-related traits

Most traits assessed had a considerable degree of polygenicity, ranging between 4-15%,

which is on the same order of magnitude as for humans (Zeng *et al.* 2018). Polygenicity was

relatively similar across all analyzed traits and therefore did not depend on the level of genetic

control, as estimated by heritability through quantitative genetic analysis. Mei *et al*. (2018)

observations in humans predicted different genetic architectures as a function of genome size.

Surprisingly, although the maritime pine genome is more than seven times larger than that of

humans (De La Torre *et al.* 2014), we found similar estimates of polygenicity between both

species. The distributions of SNP effect-sizes showed that hundreds of SNPs with near-zero

effect-size contributed together to shape phenotypic differences among clones. This highly

polygenic architecture could be explained by the omnigenic model (Boyle *et al.* 2017).

Indeed, as in humans, we expect high biological complexity and interconnectivity of gene

expression networks in forest trees, resulting in the association of virtually all expressed genes

in relevant tissues with the observed phenotypes (Wray *et al.* 2018). However, this

explanation would not account for the lack of high effect-size SNPs in our data set composed

mostly of SNPs from candidate genes (see below).

The implementation of polygenic adaptation studies outside of humans is slowly emerging

(Csilléry *et al.* 2014; He *et al.* 2016; Lind *et al.* 2017; Barghi *et al.* 2019; Friedline *et al.* 2019;

Wisser *et al.* 2019), providing increased evidence that polygenic adaptation in complex traits

may be pervasive (Sella and Barton 2019). As a result, new evolutionary questions relevant

for different organisms are arising. For instance, in forest trees, for which local adaptation is

frequently observed (Savolainen *et al.* 2007, 2013; Alberto *et al.* 2013; this study), the

contribution of alleles with small effect-size and selection coefficients (and therefore more

prompted to be swamped by gene flow) to shaping local adaptation is a question that remains

open (Yeaman 2015). Another fundamental question, in particular for conifers, is the role of

genetic redundancy. It has been suggested that genetic redundancy favors polygenic

adaptation and speed up the achievement of phenotypic optima through multiple genetic

pathways leading to similar phenotypes (Höllinger *et al.* 2019; Barghi *et al.* 2019).

Unraveling this relationship in conifers, whose genomes are characterized by a high number

of paralogs (Diaz-Sala *et al.* 2013), may shed new light about how rapidly these taxa can

adapt to environmental changes. Moreover, the influence of genome size in the genetic

architecture of fitness-related traits, as well as the relationship between heritability and

polygenicity, deserve further investigation including a better coverage of conifer genomes, as

well as improved knowledge of non-coding regions (Mackay *et al.* 2012).

Recent studies in human height (a classic example of polygenic adaption) have suggested that

detecting polygenicity may be affected by subtle biases in GWAS caused by population

structure (Berg *et al.* 2019a; Sohail *et al.* 2019). In our study, the clonal common garden

network allowed separating the genetic and the environmental effect on phenotypes to

identify which traits are contributing to adaptation. In addition, we corrected the BLUPs

estimates for the effect of neutral population genetic structure. In this sense, our work

highlights the potential of combining precise estimation of the genetic effect on phenotypes

with multi-locus genotype-phenotype association models to elucidate the mechanisms that

allow the maintenance of genetic variation in adaptive traits, especially in species with

complex demographic history. Undoubtedly, next steps to decipher polygenic adaptation in

species with varied life-history traits should implement upcoming polygenic association

methods that directly correct for population stratification (e.g. Josephs *et al.* 2019).

Performance of polygenic adaptation approaches (VSR and MLM)

We evaluated the performance of polygenic approaches (VSR and MLM) through the

comparison of SNP-based genetic variance estimates, *GEV*. Despite some slight differences,

notably for biotic-stress response traits that were limited by low sample sizes, both methods

were robust and provided consistent estimations. The large proportion of the genetic variance

explained by SNP-based models, usually higher than 50%, suggests that, by adopting a

polygenic analytical model, we were able to account for a significant part of the heritability

inferred through pedigree-based analysis, even when using a modest number of SNPs. It is

worth noting that the performance of polygenic models did not depend on the estimated

degree of heritability, as evidenced by the absence of correlation between GEV and H^2 ($\rho =$

0.04 for VSR, $\rho = -0.05$ for MLM, $p > 0.05$ in both cases). For instance, polygenic models

allowed to explain around 45% of the broad-sense heritability, also for low-heritable traits,

such as survival in Mediterranean sites, polycyclism, and SLA. GEV can be interpreted as an

analogous of the SNP-based heritability, with the particularity that GEV refers to proportion

of the variance in genetic values, rather than on the phenotypic values that are explained by

associated SNPs (see Materials and Methods for further details). SNP-based heritability is

becoming a fundamental parameter in quantitative genetics because it can yield insights into

the 'missing heritability' of complex traits (Hou *et al.* 2019). In this sense, our study shows

that polygenic approaches can be a promising strategy to account for a significant part of this

missing heritability that is commonly observed in GWAS in forest trees (reviewed by Hall *et*

al. 2016; Lind *et al.* 2018).

However, insights provided by SNP-based estimations of GEV should be interpreted with

caution. First, because maritime pine has a huge genome size (around 28 Gbp; Grotkopp *et al.*

2004; Zonneveld 2012) and a rapid decay in linkage disequilibrium (Neale and Savolainen

2004), a larger number of genotyped SNPs should be needed to obtain a good genomic

coverage. And second, because rare variants are usually difficult to incorporate in genotyping

platforms, such as the one used in our study. Such rare variants may indeed account for an

important proportion of the heritability in complex traits (Young 2019). Even though further

investigations are needed to draw stronger conclusions, robust and consistent estimates of

polygenicity across methods were fostered herein by a precise phenotypic evaluation in a

large number of individuals (over 12,500 trees).

Stability of polygenicity estimates across environments and years

The temporal and spatial heterogeneity of selection can impact the evolution of the genetic

architecture underlying adaptation (Sella and Barton 2019). Monitoring the patterns of genetic

architecture not only across environments but also across years is an important issue in long-

lived forest trees that may experience changing selection pressures along their lifetimes. In

this sense, our study is not only a validation of the polygenic adaptation model in a new

organism, but a contribution to improving our understanding of adaptation. Surprisingly, the

estimated degree of polygenicity remained stable across environments for all trait categories,

especially tree height. Additionally, we observed highly stable genetic architectures for

height, phenology, and survival across years. For the case of tree height, polygenicity was

highly stable for three time-point measures along a time-span of 6 years, comprising seedling

and juvenile stages, during which trees are more vulnerable and selection pressure are more

pronounced (Leck *et al.* 2008). However, analysis of gene function enrichment (see below)

suggests that different genetic pathways could be underlying phenotypic variation in

contrasting environments. Moreover, differences in gene expression may also underlie

adaptation under different environments and years (Mähler *et al.* 2017; Hämälä *et al.* 2020) .

The role of negative selection in polygenic adaptation

All significant correlations between SNP effect-size and MAF were negative (for tree height,

bud burst and SLA), suggesting a genetic architecture modeled, at least partially, by the action

of negative selection, i.e. SNPs with large effects are rare because they mostly have

deleterious effects and are thus selected against (O'Connor *et al.* 2019). The MLM method

did not allow elucidating whether negative estimates of S were the consequence of an

enrichment of trait-increasing or trait-decreasing alleles (Zeng *et al.* 2018), but it certainly

suggests that these traits have been under some form of negative selection. The effect of

purifying selection is widespread in model plant genomes (Wright and Andolfatto 2008), and

it has been largely evidenced in trees (Krutovsky and Neale 2005; Palmé *et al.* 2009; Eckert *et*

al. 2013; De La Torre *et al.* 2017; Grivet *et al.* 2017). Indeed, negative selection, and its

variation across populations and through time, has been pointed out as a main cause for

maintaining polygenicity (Zeng *et al.* 2018; O'Connor *et al.* 2019). Thus, negative selection

may also explain, at least partially, the degree of polygenicity observed for fitness-related

traits in maritime pine (but see below), as well as the absence of large effect-size SNPs in

previous association studies for this species (Lepoittevin *et al.* 2012; Budde *et al.* 2014; Hurel

et al. 2019).

Nevertheless, strikingly, the negative selection patterns observed across environments and

years did not mimic the trend observed for polygenicity. That is, negative selection was

consistently inferred for height, but its strength changed across environments and years for

survival and phenology-related traits. This uncoupling between negative selection and

polygenicity may result from the fact that our limited coverage of maritime pine genome did

not account for (most) rare variants, which can considerably affect S estimates (Zeng *et al.*

2018). In addition, polygenic adaptation generally results in highly stochastic genetic

responses driven by non-predictable changes in allele frequencies (Zhang *et al.* 2013).

Finally, we detected signals of gene enrichment for 10 pathways that had higher values of

maximum SNP effect-size or higher posterior probability of being included in the polygenic

models: height in the French Atlantic environment and survival in the Iberian Atlantic

environment were enriched for genes coding for *transcription factors*, bud burst in 2015 for

genes within the *monolignol biosynthesis* pathway, and *D. sapinea* susceptibility (considering

both the induced necrosis and needle discoloration) for genes within the *ubiquitin system*,

signal transduction and *flavonoid biosynthesis* pathways. Assuming that evolution of these

pathways is driven by negative selection, these patterns could be interpreted as a consequence

of the accumulation of (slightly) deleterious alleles, resulting in higher proportions of SNPs

with non-zero effect-size on these phenotypic traits. This higher tolerance to retain deleterious

mutations could be explained by a high genetic redundancy (Nowak *et al.* 1997; Krakauer and

Nowak 1999). Otherwise, if we were to assume a higher impact of positive than negative

selection, the observed patterns would imply an accumulation of beneficial mutations in these

pathways, which is a hypothesis worth exploring using sequence-based neutrality tests in

future studies.

Another five pathways were enriched in lower effect-sizes alleles: genes involved in

cytoskeleton were linked with height in the Mediterranean environment, those in the *glycan*

metabolism pathway were associated with bud burst in 2015, and those for *cell growth and*

death, *DNA recombination and repair*, and *UV response* were associated with phenology

growth index. These pathways perform general functions and could be constituted by

functionally important genes. In this case, the observed patterns suggest higher genetic

constraints on these functionally important genes, for which negative selection should be

highly efficient (Wright and Andolfatto 2008). Interestingly, our results suggest that even for

stable estimates of polygenicity, different gene pathways could underlie polygenic adaptation

for height in contrasting environments. Finally, although our gene enrichment analysis

revealed some pathways with stronger evidence for polygenic adaptation, we cannot discard

the influence of other (non-studied) gene pathways, as pointed by the omnigenic theory

(Boyle *et al.* 2017).

Conclusions

The study of genetic adaptation is currently facing new challenges. The advancement of

GWAS relies on the development of methods able to detect causal variants of small effect-

size, or at low allele frequencies. Our study, adopting a polygenic adaptation model on well-

characterized maritime pine clones planted in contrasted environments, contributed to a better

understanding of the heritability of complex adaptive traits in long-lived organisms, and its

underlying genetic architecture. Our results showed that most complex adaptive traits are

polygenic, with several of them showing also signatures of negative selection. The degree of

polygenicity was similar for traits spanning different functional categories, and this genetic

architecture was considerably stable over time and across environments. Current models for

predicting population trajectories in forest trees under climate change are based on

identification of outlier SNPs with relatively large effects on phenotypes and/or strong

correlation with climate variables (e.g. Jaramillo-Correa *et al.* 2015; Rellstab *et al.* 2016; Lu

et al. 2019). Because polygenic adaptation can take place rapidly (see, for example, Jain and

Stephan 2017), current prediction models are probably underestimating the capacity of natural

forest tree populations to adapt to new environments. Thus, adopting a polygenic adaptation

perspective could significantly improve prediction accuracy, and provide new scenarios to

inform forest conservation and reforestation programs (Valladares *et al.* 2014; Fady *et al.*

2016). Also, a better understanding of the genetic architecture of economically valuable

polygenic traits can improve genomic-assisted breeding, and allow building better genomic

selection models (Grattapaglia *et al.* 2018).

Acknowledgements

We thank A. Saldaña, F. del Caño, E. Ballesteros and D. Barba (INIA) and the 'Unité

Expérimentale Forêt Pierroton' (UEFP, INRAE; doi: 10.15454/1.5483264699193726E12) for

field assistance. Data used in this research are part of the Spanish Network of Genetic Trials

(GENFORED, http://www.genfored.es). We thank all persons and institutions linked to the

establishment and maintenance of field trials used in this study. Thanks are extended to

Antoine Kremer, Martin Lascoux and Outi Savolainen for valuable insights and discussions

on models of local adaptation in forest trees.

Funding

This study was funded by the Spanish Ministry of Economy and Competitiveness through

projects RTA2010-00120-C02-02 (CLONAPIN), CGL2011-30182-C02-01 (AdapCon) and

AGL2012-40151-C03-02 (FENOPIN). The study was also supported by the 'Initiative

d'Excellence (IdEx) de l'Université de Bordeaux - Chaires d'installation 2015' (EcoGenPin)

and the European Union's Horizon 2020 research and innovation programme under grant

agreement No 773383 (B4EST).

Author contributions

Mde-M collected field data, carried out the statistical analyses and drafted the manuscript.

JM, RA and CP designed and established the common gardens, and helped with field data

collection. IR-Q, DG, CP, GGV and SCG-M contributed to the SNP assay design and

molecular laboratory work. AE, RA and SCG-M conceived and designed the study. IR-Q and

AH collected field data and helped with the statistical analyses. JPJ-C identified gene

pathways and defined gene-sets. MH, SCG-M, DG, JPJ-C and GGV contributed to the

statistical analysis of genomic data. SCG-M coordinated the study. All authors contributed to

manuscript discussion and review, and gave final approval for publication.

Supplemental material

Table S1. Phenotypic data summary and quantitative genetic analysis. V_g stands for

genetic variance (posterior mean of the variance explained by clone effect), H^2 stands for

broad-sense heritability and Q_{ST} for genetic differentiation among populations (posterior mode

and 95% credible interval are presented).

Table S2. MCMCglmm Bayesian model parametrization. Psrf stands for the Gelman-

Rubin potential scale reduction factor criterion, a measure of model convergence. Good

convergence of models is expected for psrf < 1.02.

Table S3. List of genes included in the 17 gene sets considered for gene function

enrichment at pathway level. Annotation based on KEGG: Kyoto Encyclopedia of Genes

and Genomes (https://www.genome.jp/kegg/) is also provided. *Annotation* label indicates

genes for which no hit with KEGG database was found and thus were assigned to metabolic

pathways/modules based on the InterPro annotation.

Table S4. Number of non-zero effect-size SNPs (*nbnon-zero*) and genetic explained

variance (*GEV*) estimated using Bayesian variable selection regression (VSR), as

implemented in piMASS software, and the Bayesian linear mixed model, MLM,

implemented in GCTB software. For MLM, the coefficients of correlation between SNP

effect-size and minor allele frequency (S) are also provided. The parameters are presented as

the posterior median and 95% credible intervals. Estimates not overlapping zero are marked

in bold. NA: models that did not converge.

Figure S1. Sampled maritime pine populations (circles) and common garden sites (other

symbols). Neutral gene pools (identified in Jaramillo-Correa *et al.* 2015) outline the species

natural distribution range in different colors.

Figure S2. Posterior distribution of the number of non-zero size-effect SNPs for 26 traits

belonging to five categories: survival, height, phenology-related, functional, and biotic-

stress response traits. The number of non-zero size-effect SNPs was estimated through two

Bayesian methods: posterior inference via model averaging and subset selection (VSR), as

implemented in the software piMASS (Guan and Stephens 2011), and the Mixed Linear

Model (MLM) implemented in the software CGTB (Zeng *et al.* 2018). The posterior median

is indicated with a dashed line.

Figure S3. Posterior distribution of SNP effect-sizes for 26 traits belonging to five

categories: survival, height, phenology-related, functional, and biotic-stress response

traits. SNP effect-size was estimated through two Bayesian methods: posterior inference via

model averaging and subset selection (VSR), as implemented in the software piMASS (Guan

and Stephens 2011), and the Mixed Linear Model (MLM) implemented in the software CGTB

(Zeng *et al.* 2018).

Literature cited

Alberto F. J., S. N. Aitken, R. Alía, S. C. González□Martínez, H. Hänninen, *et al.*, 2013 Potential for evolutionary responses to climate change – evidence from tree populations. Glob Chang Biol 19: 1645–1661. https://doi.org/10.1111/gcb.12181

Alía R., R. Chambel, E. Notivol, J. Climent, and S. C. González-Martínez, 2014 Environment-dependent microevolution in a Mediterranean pine (*Pinus pinaster* Aiton). BMC Evol Biol 14: 200. https://doi.org/10.1186/s12862-014-0200-5

Alonso-Blanco C., J. Andrade, C. Becker, F. Bemm, J. Bergelson, *et al.*, 2016 1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana. Cell 166: 481–491. https://doi.org/10.1016/j.cell.2016.05.063

Aranda I., R. Alía, U. Ortega, Â. K. Dantas, and J. Majada, 2010 Intra-specific variability in biomass partitioning and carbon isotopic discrimination under moderate drought stress in seedlings from four *Pinus pinaster* populations. Tree Genet Genomes 6: 169–178. https://doi.org/10.1007/s11295-009-0238-5

Barghi N., R. Tobler, V. Nolte, A. M. Jakšić, F. Mallard, *et al.*, 2019 Genetic redundancy fuels polygenic adaptation in *Drosophila*. PLoS Biol. 17: e3000128. https://doi.org/10.1371/journal.pbio.3000128

Berg J. J., and G. Coop, 2014 A Population Genetic Signal of Polygenic Adaptation. PLoS Genetics 10: e1004412. https://doi.org/10.1371/journal.pgen.1004412

Berg J. J., A. Harpak, N. Sinnott-Armstrong, A. M. Joergensen, H. Mostafavi, *et al.*, 2019a Reduced signal for polygenic adaptation of height in UK Biobank. eLife 8: e39725. https://doi.org/10.7554/eLife.39725

Berg J. J., X. Zhang, and G. Coop, 2019b Polygenic Adaptation has Impacted Multiple Anthropometric Traits. bioRxiv 167551. https://doi.org/10.1101/167551

Bernatchez S., A. Xuereb, M. Laporte, L. Benestan, R. Steeves, *et al.*, 2019 Seascape genomics of eastern oyster (*Crassostrea virginica*) along the Atlantic coast of Canada. Evol. Appl. 12: 587–609. https://doi.org/10.1111/eva.12741

Björkegren J. L. M., J. C. Kovacic, J. T. Dudley, and E. E. Schadt, 2015 Genome-Wide Significant Loci: How Important Are They? J Am Coll Cardiol 65: 830–845. https://doi.org/10.1016/j.jacc.2014.12.033

Boyle E. A., Y. I. Li, and J. K. Pritchard, 2017 An Expanded View of Complex Traits: From Polygenic to Omnigenic. Cell 169: 1177–1186. https://doi.org/10.1016/j.cell.2017.05.038

Brachi B., G. P. Morris, and J. O. Borevitz, 2011 Genome-wide association studies in plants: the missing heritability is in the field. Genome Biol. 12: 232. https://doi.org/10.1186/gb-2011-12-10-232

Brendel O., 2001 Does bulk-needle $\delta13C$ reflect short-term discrimination? Ann. For. Sci. 58: 135–141.

Brodde L., K. Adamson, J. Julio Camarero, C. Castaño, R. Drenkhan, *et al.*, 2019 Diplodia

Tip Blight on Its Way to the North: Drivers of Disease Emergence in Northern Europe. Front
Plant Sci 9: 1818. https://doi.org/10.3389/fpls.2018.01818

Bucci G., S. C. González⬜Martínez, G. L. Provost, C. Plomion, M. M. Ribeiro, *et al.*, 2007
Range-wide phylogeography and gene zones in *Pinus pinaster* Ait. revealed by chloroplast
microsatellite markers. Mol Ecol 16: 2137–2153. https://doi.org/10.1111/j.1365-
294X.2007.03275.x

Budde K. B., M. Heuertz, A. Hernández-Serrano, J. G. Pausas, G. G. Vendramin, *et al.*, 2014
In situ genetic association for serotiny, a fire-related trait, in Mediterranean maritime pine
(*Pinus pinaster*). New Phytol 201: 230–241. https://doi.org/10.1111/nph.12483

Canales J., R. Bautista, P. Label, J. Gómez-Maldonado, I. Lesur, *et al.*, 2014 De novo
assembly of maritime pine transcriptome: implications for forest breeding and biotechnology.
Plant Biotechnol. J. 12: 286–99. https://doi.org/10.1111/pbi.12136

Craine J. M., E. N. J. Brookshire, M. D. Cramer, N. J. Hasselquist, K. Koba, *et al.*, 2015
Ecological interpretations of nitrogen isotope ratios of terrestrial plants and soils. Plant Soil
396: 1–26. https://doi.org/10.1007/s11104-015-2542-1

Csilléry K., H. Lalagüe, G. G. Vendramin, S. C. González-Martínez, B. Fady, *et al.*, 2014
Detecting short spatial scale local adaptation and epistatic selection in climate-related
candidate genes in European beech (*Fagus sylvatica*) populations. Mol Ecol 23: 4696–4708.
https://doi.org/10.1111/mec.12902

Daub J. T., T. Hofer, E. Cutivet, I. Dupanloup, L. Quintana-Murci, *et al.*, 2013 Evidence for
Polygenic Adaptation to Pathogens in the Human Genome. Mol Biol Evol 30: 1544–1558.
https://doi.org/10.1093/molbev/mst080

Dayan D. I., X. Du, T. Z. Baris, D. N. Wagner, D. L. Crawford, *et al.*, 2019 Population
genomics of rapid evolution in natural populations: polygenic selection in response to power
station thermal effluents. BMC Evol. Biol. 19: 61. https://doi.org/10.1186/s12862-019-1392-5

De La Torre A. R., I. Birol, J. Bousquet, P. K. Ingvarsson, S. Jansson, *et al.*, 2014 Insights
into conifer giga-genomes. Plant Physiol. 166: 1724–32.
https://doi.org/10.1104/pp.114.248708

De La Torre A. R., Z. Li, Y. Van de Peer, and P. K. Ingvarsson, 2017 Contrasting Rates of
Molecular Evolution and Patterns of Selection among Gymnosperms and Flowering Plants.
Mol. Biol. Evol. 34: 1363–1377. https://doi.org/10.1093/molbev/msx069

De La Torre A. R., B. Wilhite, and D. B. Neale, 2019 Environmental Genome-Wide
Association Reveals Climate Adaptation Is Shaped by Subtle to Moderate Allele Frequency
Shifts in Loblolly Pine. Genome Biol Evol 11: 2976–2989.
https://doi.org/10.1093/gbe/evz220

Desprez-Loustau M.-L., B. Marçais, L.-M. Nageleisen, D. Piou, and A. Vannini, 2006
Interactive effects of drought and pathogens in forest trees. Ann. For. Sci. 63: 597–612.
https://doi.org/10.1051/forest:2006040

Díaz S., J. Kattge, J. H. C. Cornelissen, I. J. Wright, S. Lavorel, *et al.*, 2016 The global
spectrum of plant form and function. Nature 529: 167–171.

https://doi.org/10.1038/nature16489

Diaz-Sala C., J. A. Cabezas, B. Fernández de Simón, D. Abarca, M. Á. Guevara, *et al.*, 2013
The uniqueness of conifers, pp. 67–96 in *From Plant Genomics to Plant Biotechnology*,
edited by Poltronieri P., Burbulis N., Fogher C. Woodhead Publishing, Cambridge.

Eckert A. J., A. D. Bower, K. D. Jermstad, J. L. Wegrzyn, B. J. Knaus, *et al.*, 2013 Multilocus
analyses reveal little evidence for lineage-wide adaptive evolution within major clades of soft
pines (*Pinus* subgenus *Strobus*). Mol Ecol 22: 5635–5650. https://doi.org/10.1111/mec.12514

Edge M. D., and G. Coop, 2019 Reconstructing the History of Polygenic Scores Using
Coalescent Trees. Genetics 211: 235–262. https://doi.org/10.1534/genetics.118.301687

Fabre B., D. Piou, M.-L. Desprez☐Loustau, and B. Marçais, 2011 Can the emergence of pine
Diplodia shoot blight in France be explained by changes in pathogen pressure linked to
climate change? Glob Chang Biol 17: 3218–3227. https://doi.org/10.1111/j.1365-
2486.2011.02428.x

Fady B., J. Cottrell, L. Ackzell, R. Alía, B. Muys, *et al.*, 2016 Forests and global change:
what can genetics contribute to the major forest management and policy challenges of the
twenty-first century? Reg Environ Change 16: 927–939. https://doi.org/10.1007/s10113-015-
0843-9

Farquhar G. D., and R. A. Richards, 1984 Isotopic Composition of Plant Carbon Correlates
With Water-Use Efficiency of Wheat Genotypes. Functional Plant Biol. 11: 539–552.
https://doi.org/10.1071/pp9840539

Field Y., E. Boyle, N. Telis, Z. Gao, K. J. Gaulton, *et al.*, 2016 Detection of human adaptation
during the past 2000 years. Science 354: 760–764. https://doi.org/10.1126/science.aah5114

Fisher R. A., 1918 The correlation between relatives on the supposition of Mendelian
inheritance. Transaction Royal Society Edinburgh 52: 399–433.

Fréjaville T., and M. Benito Garzón, 2018 The EuMedClim Database: Yearly Climate Data
(1901–2014) of 1 km Resolution Grids for Europe and the Mediterranean Basin. Front. Ecol.
Evol. 6: 31. https://doi.org/10.3389/fevo.2018.00031

Friedline C. J., T. M. Faske, B. M. Lind, E. M. Hobson, D. Parry, *et al.*, 2019 Evolutionary
genomics of gypsy moth populations sampled along a latitudinal gradient. Mol. Ecol. 28:
2206–2223. https://doi.org/10.1111/mec.15069

Gaspar M. J., T. Velasco, I. Feito, R. Alía, and J. Majada, 2013 Genetic Variation of Drought
Tolerance in *Pinus pinaster* at Three Hierarchical Levels: A Comparison of Induced Osmotic
Stress and Field Testing. PLOS ONE 8: e79094.
https://doi.org/10.1371/journal.pone.0079094

Gelman A., and D. B. Rubin, 1992 Inference from iterative simulation using multiple
sequences. Stat Sci 7: 457–511.

Gelman A., 2006 Prior distributions for variance parameters in hierarchical models (comment
on article by Browne and Draper). Bayesian Anal 1: 515–534. https://doi.org/10.1214/06-
BA117A

Gelman A., D. A. van Dyk, Z. Huang, and Boscardin, 2008 Using Redundant
Parameterizations to Fit Hierarchical Models. J Comput Graph Stat 17: 95–122.

Girard F., M. Vennetier, S. Ouarmim, Y. Caraglio, and L. Misson, 2011 Polycyclism, a
fundamental tree growth process, decline with recent climate change: the example of *Pinus*
halepensis Mill. in Mediterranean France. Trees 25: 311–322. https://doi.org/10.1007/s00468-
010-0507-9

Gnecchi-Ruscone G. A., P. Abondio, S. De Fanti, S. Sarno, M. G. Sherpa, *et al.*, 2018
Evidence of Polygenic Adaptation to High Altitude from Tibetan and Sherpa Genomes.
Genome Biol Evol 10: 2919–2930. https://doi.org/10.1093/gbe/evy233

González-Martínez S. C., R. Alía, and L. Gil, 2002 Population genetic structure in a
Mediterranean pine (*Pinus pinaster* Ait.): a comparison of allozyme markers and quantitative
traits. Heredity 89: 199–206. https://doi.org/10.1038/sj.hdy.6800114

González-Martínez S. C., K. V. Krutovsky, and D. B. Neale, 2006 Forest-tree population
genomics and adaptive evolution. New Phytol 170: 227–238.

Grattapaglia D., O. B. Silva-Junior, R. T. Resende, E. P. Cappa, B. S. F. Müller, *et al.*, 2018
Quantitative Genetics and Genomics Converge to Accelerate Forest Tree Breeding. Front.
Plant Sci. 9: 1693. https://doi.org/10.3389/fpls.2018.01693

Greenwood S., P. Ruiz☐Benito, J. Martínez☐Vilalta, F. Lloret, T. Kitzberger, *et al.*, 2017
Tree mortality across biomes is promoted by drought intensity, lower wood density and
higher specific leaf area. Ecol Lett 20: 539–553. https://doi.org/10.1111/ele.12748

Grivet D., K. Avia, A. Vaattovaara, A. J. Eckert, D. B. Neale, *et al.*, 2017 High rate of
adaptive evolution in two widespread European pines. Mol Ecol 26: 6857–6870.
https://doi.org/10.1111/mec.14402

Grotkopp E., M. Rejmánek, M. J. Sanderson, and T. L. Rost, 2004 Evolution of Genome Size
in Pines (*Pinus*) and Its Life-History Correlates: Supertree Analyses. Evolution 58: 1705.
https://doi.org/10.1554/03-545

Guan Y., and M. Stephens, 2011 Bayesian variable selection regression for genome-wide
association studies and other large-scale problems. Ann Appl Stat 5: 1780–1815.
https://doi.org/10.1214/11-AOAS455

Hadfield J. D., 2010 MCMC Methods for Multi-Response Generalized Linear Mixed Models:
The **MCMCglmm** *R* Package. Journal of Statistical Software 33.
https://doi.org/10.18637/jss.v033.i02

Hall T., 1999 BioEdit: a user-friendly biological sequence alignment editor and analysis
program for Window 95/98/NT. Nucleic Acids Symposium Series 41: 95–98.

Hall D., H. R. Hallingbäck, and H. X. Wu, 2016 Estimation of number and size of QTL
effects in forest tree traits. Tree Genet Genomes 12: 110. https://doi.org/10.1007/s11295-016-
1073-0

Hämälä T., M. J. Guiltinan, J. H. Marden, S. N. Maximova, C. W. dePamphilis, *et al.*, 2020
Gene Expression Modularity Reveals Footprints of Polygenic Adaptation in *Theobroma*
cacao. Mol Biol Evol 37: 110–123. https://doi.org/10.1093/molbev/msz206

Hancock A. M., D. B. Witonsky, E. Ehler, G. Alkorta-Aranburu, C. Beall, *et al.*, 2010a Human adaptations to diet, subsistence, and ecoregion are due to subtle shifts in allele frequency. PNAS 107: 8924–8930. https://doi.org/10.1073/pnas.0914625107

Hancock A. M., G. Alkorta-Aranburu, D. B. Witonsky, and A. Di Rienzo, 2010b Adaptations to new environments in humans: the role of subtle allele frequency shifts. Phil. Trans. R. Soc. B 365: 2459–2468. https://doi.org/10.1098/rstb.2010.0032

Hayward L. K., and G. Sella, 2019 Polygenic adaptation after a sudden change in environment. bioRxiv. https://doi.org/10.1101/792952

He F., A. L. Arce, G. Schmitz, M. Koornneef, P. Novikova, *et al.*, 2016 The Footprint of Polygenic Adaptation on Stress-Responsive Cis-Regulatory Divergence in the *Arabidopsis* Genus. Mol Biol Evol 33: 2088–2101. https://doi.org/10.1093/molbev/msw096

Heinzelmann R., C. Dutech, T. Tsykun, F. Labbé, J.-P. Soularue, *et al.*, 2019 Latest advances and future perspectives in *Armillaria* research. Can. J. For. Res. 41: 1–23. https://doi.org/10.1080/07060661.2018.1558284

Henderson C. R., 1973 Sire evaluation and genetic trends. J Anim Sci 1973: 10–41. https://doi.org/10.1093/ansci/1973.Symposium.10

Hermisson J., and P. S. Pennings, 2017 Soft sweeps and beyond: understanding the patterns and probabilities of selection footprints under rapid adaptation. Methods. Ecol. Evol 8: 700–716. https://doi.org/10.1111/2041-210X.12808

Höllinger I., P. S. Pennings, and J. Hermisson, 2019 Polygenic adaptation: From sweeps to subtle frequency shifts. PLOS Genetics 15: e1008035.

Hou K., K. S. Burch, A. Majumdar, H. Shi, N. Mancuso, *et al.*, 2019 Accurate estimation of SNP-heritability from biobank-scale data irrespective of genetic architecture. Nat Genet 51: 1244–1251. https://doi.org/10.1038/s41588-019-0465-0

Huang X., and B. Han, 2014 Natural Variations and Genome-Wide Association Studies in Crop Plants. Annu. Rev. Plant Biol 65: 531–551. https://doi.org/10.1146/annurev-arplant-050213-035715

Hurel A., M. de Miguel, C. Dutech, M.-L. Desprez-Loustau, C. Plomion, *et al.*, 2019 Genetic basis of susceptibility to *Diplodia sapinea* and *Armillaria ostoyae* in maritime pine. bioRxiv 699389. https://doi.org/10.1101/699389

Jacquet J.-S., A. Bosc, A. P. O'Grady, and H. Jactel, 2013 Pine growth response to processionary moth defoliation across a 40-year chronosequence. Forest Ecol Manag 293: 29–38. https://doi.org/10.1016/j.foreco.2012.12.003

Jain K., and W. Stephan, 2017 Rapid Adaptation of a Polygenic Trait After a Sudden Environmental Shift. Genetics 206: 389–406. https://doi.org/10.1534/genetics.116.196972

Jaramillo-Correa J.-P., I. Rodríguez-Quilón, D. Grivet, C. Lepoittevin, F. Sebastiani, *et al.*, 2015 Molecular Proxies for Climate Maladaptation in a Long-Lived Tree (*Pinus pinaster* Aiton, Pinaceae). Genetics 199: 793–807. https://doi.org/10.1534/genetics.114.173252

Josephs E. B., J. J. Berg, J. Ross-Ibarra, and G. Coop, 2019 Detecting adaptive differentiation

in structured populations with genomic data and common gardens. Genetics 211: 989–1004.
https://doi.org/10.1101/368506

Kanehisa M., Y. Sato, and K. Morishima, 2016 BlastKOALA and GhostKOALA: KEGG
Tools for Functional Characterization of Genome and Metagenome Sequences. J. Mol. Biol.
428: 726–731. https://doi.org/10.1016/j.jmb.2015.11.006

Keenan K., P. McGinnity, T. F. Cross, W. W. Crozier, and P. A. Prodöhl, 2013 diveRsity: An
R package for the estimation and exploration of population genetics parameters and their
associated errors. Methods. Ecol. Evol 4: 782–788. https://doi.org/10.1111/2041-210X.12067

Krakauer D. C., and M. A. Nowak, 1999 Evolutionary preservation of redundant duplicated
genes. Semin. Cell Dev. Biol. 10: 555–559. https://doi.org/10.1006/scdb.1999.0337

Krutovsky K. V., and D. B. Neale, 2005 Nucleotide Diversity and Linkage Disequilibrium in
Cold-Hardiness- and Wood Quality-Related Candidate Genes in Douglas Fir. Genetics 171:
2029–2041. https://doi.org/10.1534/genetics.105.044420

Lamy J.-B., L. Bouffier, R. Burlett, C. Plomion, H. Cochard, *et al.*, 2011 Uniform Selection
as a Primary Force Reducing Population Genetic Differentiation of Cavitation Resistance
across a Species Range. PLoS ONE 6: e23476. https://doi.org/10.1371/journal.pone.0023476

Le Corre V., and A. Kremer, 2003 Genetic Variability at Neutral Markers, Quantitative Trait
Loci and Trait in a Subdivided Population Under Selection. Genetics 164: 1205–1219.

Leck M. A., V. T. Parker, and R. L. Simpson, 2008 *Seedling Ecology and Evolution.*
Cambridge University Press.

Lepoittevin C., L. Harvengt, C. Plomion, and P. Garnier-Géré, 2012 Association mapping for
growth, straightness and wood chemistry traits in the *Pinus pinaster* Aquitaine breeding
population. Tree Genet Genomes 8: 113–126. https://doi.org/10.1007/s11295-011-0426-y

Lind B. M., C. J. Friedline, J. L. Wegrzyn, P. E. Maloney, D. R. Vogler, *et al.*, 2017 Water
availability drives signatures of local adaptation in whitebark pine (*Pinus albicaulis* Engelm.)
across fine spatial scales of the Lake Tahoe Basin, USA. Mol Ecol 26: 3168–3185.
https://doi.org/10.1111/mec.14106

Lind B. M., M. Menon, C. E. Bolte, T. M. Faske, and A. J. Eckert, 2018 The genomics of
local adaptation in trees: are we out of the woods yet? Tree Genet Genomes 14: 29.
https://doi.org/10.1007/s11295-017-1224-y

Liu X., P.-R. Loh, L. J. O'Connor, S. Gazal, A. Schoech, *et al.*, 2018 Quantification of
genetic components of population differentiation in UK Biobank traits reveals signals of
polygenic selection. bioRxiv 357483. https://doi.org/10.1101/357483

Lloyd-Jones L. R., J. Zeng, J. Sidorenko, L. Yengo, G. Moser, *et al.*, 2019 Improved
polygenic prediction by Bayesian multiple regression on summary statistics. Nat Commun 10:
1–11. https://doi.org/10.1038/s41467-019-12653-0

Lu M., K. V. Krutovsky, and C. A. Loopstra, 2019 Predicting adaptive genetic variation of
loblolly pine (*Pinus taeda* L.) populations under projected future climates based on
multivariate models. J Hered 110: 857–865. https://doi.org/10.1093/jhered/esz065

Mackay J., J. F. D. Dean, C. Plomion, D. G. Peterson, F. M. Cánovas, *et al.*, 2012 Towards
decoding the conifer giga-genome. Plant Mol Biol 80: 555–69.
https://doi.org/10.1007/s11103-012-9961-7

Maher B., 2008 Personal genomes: The case of the missing heritability. Nature 456: 18–21.
https://doi.org/doi:10.1038/456018a

Mähler N., J. Wang, B. K. Terebieniec, P. K. Ingvarsson, N. R. Street, *et al.*, 2017 Gene co-
expression network connectivity is an important determinant of selective constraint. PLOS
Genet 13: e1006402. https://doi.org/10.1371/journal.pgen.1006402

Majada J., C. Martínez-Alonso, I. Feito, A. Kidelman, I. Aranda, *et al.*, 2011 Mini-cuttings:
an effective technique for the propagation of *Pinus pinaster* Ait. New Forests 41: 399–412.
https://doi.org/10.1007/s11056-010-9232-x

Manolio T. A., F. S. Collins, N. J. Cox, D. B. Goldstein, L. A. Hindorff, *et al.*, 2009 Finding
the missing heritability of complex diseases. Nature 461: 747–753.
https://doi.org/10.1038/nature08494

Mayol M., M. Riba, S. Cavers, D. Grivet, L. Vincenot, *et al.*, 2020 A multiscale approach to
detect selection in nonmodel tree species: Widespread adaptation despite population decline
in *Taxus baccata* L. Evol. Appl. 0. https://doi.org/10.1111/eva.12838

McKay J. K., and R. G. Latta, 2002 Adaptive population divergence: markers, QTL and traits.
Trends Ecol. Evol 17: 285–291. https://doi.org/10.1016/S0169-5347(02)02478-3

Mei W., M. G. Stetter, D. J. Gates, M. C. Stitzer, and J. Ross□Ibarra, 2018 Adaptation in
plant genomes: Bigger is different. Am. J. Bot 105: 16–19. https://doi.org/10.1002/ajb2.1002

Nakagawa S., and H. Schielzeth, 2010 Repeatability for Gaussian and non-Gaussian data: a
practical guide for biologists. Biol. Rev. 85: 935–956. https://doi.org/10.1111/j.1469-
185X.2010.00141.x

Neale D. B., and O. Savolainen, 2004 Association genetics of complex traits in conifers.
Trends Plant Sci. 9: 325–330. https://doi.org/10.1016/j.tplants.2004.05.006

Nowak M. A., M. C. Boerlijst, J. Cooke, and J. M. Smith, 1997 Evolution of genetic
redundancy. Nature 388: 167–171. https://doi.org/10.1038/40618

O'Connor L. J., A. P. Schoech, F. Hormozdiari, S. Gazal, N. Patterson, *et al.*, 2019 Extreme
Polygenicity of Complex Traits Is Explained by Negative Selection. Am J hum genet 105:
456–476. https://doi.org/10.1016/j.ajhg.2019.07.003

Orr H. A., and J. A. Coyne, 1992 The Genetics of Adaptation: A Reassessment. Am Nat 140:
725–742. https://doi.org/10.1086/285437

Pallares L. F., 2019 Searching for solutions to the missing heritability problem. eLife 8:
e53018. https://doi.org/10.7554/eLife.53018

Palmé A. E., T. Pyhäjärvi, W. Wachowiak, and O. Savolainen, 2009 Selection on Nuclear
Genes in a Pinus Phylogeny. Mol Biol Evol 26: 893–905.
https://doi.org/10.1093/molbev/msp010

Pitchers W., J. Nye, E. J. Márquez, A. Kowalski, I. Dworkin, *et al.*, 2019 A Multivariate

Genome-Wide Association Study of Wing Shape in *Drosophila melanogaster*. Genetics 211:
1429–1447. https://doi.org/10.1534/genetics.118.301342

Plomion C., J. Bartholomé, L. Bouffier, O. Brendel, H. Cochard, *et al.*, 2016a Understanding
the genetic bases of adaptation to soil water deficit in trees through the examination of water
use efficiency and cavitation resistance: maritime pine as a case study. JPH 3: e008.
https://doi.org/10.20870/jph.2016.e008

Plomion C., J. Bartholomé, I. Lesur, C. Boury, I. Rodríguez-Quilón, *et al.*, 2016b High-
density SNP assay development for genetic analysis in maritime pine (*Pinus pinaster*). Mol
Ecol Res 16: 574–587. https://doi.org/10.1111/1755-0998.12464

Pritchard J. K., J. K. Pickrell, and G. Coop, 2010 The Genetics of Human Adaptation: Hard
Sweeps, Soft Sweeps, and Polygenic Adaptation. Curr Biol 20: R208–R215.
https://doi.org/10.1016/j.cub.2009.11.055

Pritchard J. K., and A. D. Rienzo, 2010 Adaptation – not by sweeps alone. Nat Rev Genet 11:
665–667. https://doi.org/10.1038/nrg2880

R Core Team, 2019 *R: A language and environment for statistical computing*. R Foundation
for Statistical Computing, Vienna, Austria.

Rellstab C., S. Zoller, L. Walthert, I. Lesur, A. R. Pluess, *et al.*, 2016 Signatures of local
adaptation in candidate genes of oaks (*Quercus* spp.) with respect to present and future
climatic conditions. Mol Ecol 25: 5907–5924. https://doi.org/10.1111/mec.13889

Resende M. D. V., M. F. R. Resende Jr, C. P. Sansaloni, C. D. Petroli, A. A. Missiaggia, *et*
al., 2012 Genomic selection for growth and wood quality in Eucalyptus: capturing the missing
heritability and accelerating breeding for complex traits in forest trees. New Phytol 116–128.
https://doi.org/10.1111/j.1469-8137.2011.04038.x@10.1002

Robinson G. K., 1991 That BLUP is a Good Thing: The Estimation of Random Effects.
Statist. Sci. 6: 15–32. https://doi.org/10.1214/ss/1177011926

Rodríguez-Quilón I., L. Santos-del-Blanco, M. J. Serra-Varela, J. Koskela, S. C. González-
Martínez, *et al.*, 2016 Capturing neutral and adaptive genetic diversity for conservation in a
highly structured tree species. Ecol App 26: 2254–2266.

Rosenberg N. A., M. D. Edge, J. K. Pritchard, and M. W. Feldman, 2019 Interpreting
polygenic scores, polygenic adaptation, and human phenotypic differences. Evol. Med. Public
Health. 26–34. https://doi.org/10.1093/emph/eoy036

Savolainen O., T. Pyhäjärvi, and T. Knürr, 2007 Gene Flow and Local Adaptation in Trees.
Ann.   Rev.   Ecol.   Evol.   Syst.   38:   595–619.
https://doi.org/10.1146/annurev.ecolsys.38.091206.095646

Savolainen O., M. Lascoux, and J. Merilä, 2013 Ecological genomics of local adaptation. Nat
Rev Genet 14: 807–820. https://doi.org/10.1038/nrg3522

Sefton C. A., K. Montagu, B. J. Atwell, and J. P. Conroy, 2002 Anatomical variation in
juvenile eucalypt leaves accounts for differences in specific leaf area and CO_2 assimilation
rates. Aust. J. Bot. 50: 301–310. https://doi.org/10.1071/bt01059

Sella G., and N. H. Barton, 2019 Thinking About the Evolution of Complex Traits in the Era
of Genome-Wide Association Studies. Annu. Rev. Genom. Hum. Genet. 20: 461–493.
https://doi.org/10.1146/annurev-genom-083115-022316

Sharma A., J. S. Lee, C. G. Dang, P. Sudrajad, H. C. Kim, *et al.*, 2015 Stories and Challenges
of Genome Wide Association Studies in Livestock — A Review. Asian-Australas J Anim Sci
28: 1371–1379. https://doi.org/10.5713/ajas.14.0715

Shi H., G. Kichaev, and B. Pasaniuc, 2016 Contrasting the Genetic Architecture of 30
Complex Traits from Summary Association Data. Am J hum genet 99: 139–153.
https://doi.org/10.1016/j.ajhg.2016.05.013

Smith J. M., and J. Haigh, 1974 The hitch-hiking effect of a favourable gene. Genet. Res. 23:
23–35. https://doi.org/10.1017/S0016672300014634

Sohail M., R. M. Maier, A. Ganna, A. Bloemendal, A. R. Martin, *et al.*, 2019 Polygenic
adaptation on height is overestimated due to uncorrected stratification in genome-wide
association studies. eLife 8: e39702. https://doi.org/10.7554/eLife.39702

Speidel L., M. Forest, S. Shi, and S. R. Myers, 2019 A method for genome-wide genealogy
estimation for thousands of samples. Nat Genet 51: 1321–1329.
https://doi.org/10.1038/s41588-019-0484-x

Spitze K., 1993 Population Structurein Daphnia obtusa: Quantitative Genetic and Allozymic
Variation. Genetics 135: 367–374.

Torres-Ruiz J. M., A. Kremer, M. R. Carins-Murphy, T. J. Brodribb, L. J. Lamarque, *et al.*,
2019 Genetic differentiation in functional traits among European sessile oak populations. Tree
Physiol 39: 1736–1749. https://doi.org/10.1093/treephys/tpz090

Turchin M. C., C. W. Chiang, C. D. Palmer, S. Sankararaman, D. Reich, *et al.*, 2012 Evidence
of widespread selection on standing variation in Europe at height-associated SNPs. Nat Genet
44: 1015–1019. https://doi.org/10.1038/ng.2368

Valladares F., S. Matesanz, F. Guilhaumon, M. B. Araújo, L. Balaguer, *et al.*, 2014 The
effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under
climate change. Ecol Lett 17: 1351–1364. https://doi.org/10.1111/ele.12348

Visscher P. M., W. G. Hill, and N. R. Wray, 2008 Heritability in the genomics era —
concepts and misconceptions. Nat Rev Genet 9: 255–266. https://doi.org/10.1038/nrg2322

Visscher P. M., N. R. Wray, Q. Zhang, P. Sklar, M. I. McCarthy, *et al.*, 2017 10 Years of
GWAS Discovery: Biology, Function, and Translation. Am J hum genet 101: 5–22.
https://doi.org/10.1016/j.ajhg.2017.06.005

Vizcaíno-Palomar N., B. Fady, R. Alía, A. Raffin, S. Mutke, *et al.*, 2019 Patterns of
phenotypic plasticity among populations of three Mediterranean pine species and implications
for evolutionary responses to climate change. bioRxiv 716084.
https://doi.org/10.1101/716084

Walker X. J., M. C. Mack, and J. F. Johnstone, 2015 Stable carbon isotope analysis reveals
widespread drought stress in boreal black spruce forests. Glob Chang Biol 21: 3102–3113.
https://doi.org/10.1111/gcb.12893

Warren C. R., J. F. McGrath, and M. A. Adams, 2001 Water availability and carbon isotope
discrimination in conifers. Oecologia 127: 476–486. https://doi.org/10.1007/s004420000609

Weir B. S., and C. C. Cockerham, 1984 ESTIMATING F-STATISTICS FOR THE
ANALYSIS OF POPULATION STRUCTURE. Evol 38: 1358–1370.
https://doi.org/10.1111/j.1558-5646.1984.tb05657.x

Whitlock M. C., and F. Guillaume, 2009 Testing for Spatially Divergent Selection:
Comparing QST to FST. Genetics 183: 1055–1063.
https://doi.org/10.1534/genetics.108.099812

Wilson A. J., D. Réale, M. N. Clements, M. M. Morrissey, E. Postma, *et al.*, 2010 An
ecologist's guide to the animal model. J Anim Ecol 79: 13–26. https://doi.org/10.1111/j.1365-
2656.2009.01639.x

Wisser R. J., Z. Fang, J. B. Holland, J. E. C. Teixeira, J. Dougherty, *et al.*, 2019 The Genomic
Basis for Short-Term Evolution of Environmental Adaptation in Maize. Genetics 213: 1479–
1494. https://doi.org/10.1534/genetics.119.302780

Wray N. R., C. Wijmenga, P. F. Sullivan, J. Yang, and P. M. Visscher, 2018 Common
Disease Is More Complex Than Implied by the Core Gene Omnigenic Model. Cell 173:
1573–1580. https://doi.org/10.1016/j.cell.2018.05.051

Wright S. I., and P. Andolfatto, 2008 The Impact of Natural Selection on the Genome:
Emerging Patterns in Drosophila and Arabidopsis. Ann. Rev. Ecol. Evol. Syst. 39: 193–213.
https://doi.org/10.1146/annurev.ecolsys.39.110707.173342

Yang J., B. Benyamin, B. P. McEvoy, S. Gordon, A. K. Henders, *et al.*, 2010 Common SNPs
explain a large proportion of the heritability for human height. Nat Genet 42: 565–569.
https://doi.org/10.1038/ng.608

Yeaman S., 2015 Local Adaptation by Alleles of Small Effect. Am Nat 186: S74–S89.
https://doi.org/10.1086/682405

Young A. I., 2019 Solving the missing heritability problem, (J. Flint, Ed.). PLoS Genet 15:
e1008222. https://doi.org/10.1371/journal.pgen.1008222

Zeng J., R. de Vlaming, Y. Wu, M. R. Robinson, L. R. Lloyd-Jones, *et al.*, 2018 Signatures of
negative selection in the genetic architecture of human complex traits. Nat Genet 50: 746–
753. https://doi.org/10.1038/s41588-018-0101-4

Zhang G., L. J. Muglia, R. Chakraborty, J. M. Akey, and S. M. Williams, 2013 Signatures of
natural selection on genetic variants affecting complex human traits. Appl Transl Genom 2:
78–94. https://doi.org/10.1016/j.atg.2013.10.002

Zonneveld B. J. M., 2012 Conifer genome sizes of 172 species, covering 64 of 67 genera,
range from 8 to 72 picogram. Nor. J. Bot. 30: 490–502. https://doi.org/10.1111/j.1756-
1051.2012.01516.x

Site	Country	Coordinates	Environment	Plantation year	N trees (clones)	Annual precipitation (mm)	Summer precipitation (mm)	Annual mean temperature (ºC)	Annual temperature range (ºC)	Soil type
Cabada	Spain	43°25'17" N 06°32'38" W	Iberian Atlantic	2010	4,272 (535)	890	126	12.9	24.0	Cambisol
Fundão	Portugal	40°06'38" N 07°28'58" W		2010	4,272 (535)	1122	58	14.0	26.9	Cambisol
Pierroton	France	44°44'42" N 00°47'04" W	French Atlantic	2011	3,434 (443)	933	199	13.8	26.7	Arenosol
Madrid	Spain	40°30'47" N 03°18'44" W	Mediterranean	2010	4,272 (535)	378	35	14.8	32.8	Arenosol
Cáceres	Spain	40°02'24" N 05°22'19" W		2010	4,272 (535)	374	21	16.7	32.6	Fluvisol

Table 1. CLONAPIN common garden network (5 sites). Climatic data correspond to the mean of each parameter for the period 2005-2014 obtained from the EuMedClim database (Fréjaville and Benito Garzón 2018).

Trait	Environment	Gene set	Statistic tested	Sign of enrichment	*p*-value	*q*-value (<0.10)
Height	French Atlantic	*Transcription factor*	*maxabsbetarb*	Higher	0.003	0.05
			maxpostprb	Higher	0.004	0.07
	Mediterranean	*Cytoskeleton*	*maxabsbetarb*	Lower	0.003	0.06
Survival	Iberian Atlantic	*Transcription factor*	*maxabsbetarb*	Higher	0.001	0.01
			maxpostprb	Higher	<0.001	0.005
Bud burst 2015	French Atlantic	*Monolignol biosynthesis*	*maxabsbetarb*	Higher	0.003	0.05
		Monolignol biosynthesis	*maxpostprb*	Higher	0.005	0.08
		Glycan metabolism	*maxabsbetarb*	Lower	0.040	0.09
Phenology growth index	Iberian Atlantic	*Cell growth and death*	*maxabsbetarb*	Lower	0.010	0.03
		DNA recomb and repair	*maxabsbetarb*	Lower	0.008	0.03
		UV response	*maxabsbetarb*	Lower	0.005	0.03
D. sapinea necrosis	French Atlantic	*Ubiquitin system*	*maxabsbetarb*	Higher	0.002	0.04
			maxpostprb	Higher	0.003	0.06
D. sapinea discoloration	French Atlantic	*Signal transduction*	*maxabsbetarb*	Higher	0.004	0.08
			maxpostprb	Higher	0.003	0.06
		Flavonoid biosynthesis	*maxpostprb*	Higher	0.007	0.07

Table 2. Gene sets with gene function enrichment at pathway/module level. Two statistics obtained from the VSR method were tested: the maximum of any SNP per gene of the Rao-Backwellized posterior probability of inclusion (*maxpostprb*) and the maximum of any SNP per gene of the absolute value of the Rao-Backwellized effect-size (*maxabsbetarb*). Sign of enrichment refers to two-tailed null hypothesis testing.

Figures

Figure 1. Comparison of Q_{ST} and F_{ST} estimates across traits, environments and years. A) Q_{ST} for a selection of traits belonging to five categories: survival, height, phenology-related traits, functional traits and biotic-stress response (see Supplemental Table S1 for all traits). B) Q_{ST} for height estimated in three different environments: Mediterranean, Iberian Atlantic, and French Atlantic, and a global Q_{ST} for the three environments together. In the French Atlantic common garden, height was measured in three different years: 2013, 2015 and 2018. Global F_{ST} estimate is presented by a red line surrounded by the 95% confidence intervals computed by bootstrapping.

Figure 2. Polygenicity estimated from Bayesian mixed linear models (MLMs) for a selection of traits (see Supplemental Table S4 for all traits). Polygenicity was estimated as the proportion of non-zero size-effect SNPs. Posterior median and 95% credible intervals are presented.

Figure 3. Polygenicity estimated from Bayesian mixed linear models (MLMs) across environments and years. A) Variation of polygenicity across environments. B) Temporal variation of polygenicity. Polygenicity was estimated as the proportion of non-zero size-effect SNPs. Posterior median and 95% credible intervals are presented.

Figure 4. Correlation between polygenicity (proportion of non-zero size-effect SNPs) and

***GEV* (explained genetic variance).** A) MLM method implemented in CGTB software. B)

VSR method implemented in piMASS software. Each point represents the posterior median.

Figure 5. Correlation between SNP effect-size and Minor Allele Frequency (MAF). The coefficient of correlation between SNP effect-size and MAF (*S*) was estimated through the MLM method. The posterior distribution of *S* (median and 95% credible intervals) are presented.