1 **Title: Merkel Cell Polyomavirus Small Tumor Antigen Activates Matrix Metallopeptidase-9**
2 **Gene Expression for Cell Migration and Invasion**
3

4 **Authors**
5   Nnenna Nwogu[1,2], Luz E. Ortiz[1,2], Adrian Whitehouse[3], Hyun Jin Kwun[1,2],*
6
7 **Affiliations**
8   [1]Department of Microbiology and Immunology, Penn State University College of
9   Medicine, Hershey, PA, USA
10
11   [2]Penn State Cancer Institute, Hershey, PA, USA
12
13   [3]School of Molecular and Cellular Biology and Astbury Centre for Structural Molecular
14   Biology, Faculty of Biological Sciences, University of Leeds, Leeds, United Kingdom.
15
16
17
*Correspondence: Hyun Jin Kwun
Department of Microbiology and Immunology, Penn State University College of
Medicine, Hershey, PA 17033, USA
E-mail addresses: hxk479@psu.edu
Phone: (717) 512-7241
Fax: (717) 623-7715

41 Competing interests: The authors declare no competing interests.

42
43
44

Keywords: Merkel cell carcinoma, Merkel cell polyomavirus, small tumor antigen, Matrix
Metallopeptidase-9, FBW7, Cell migration, Cell invasion

ABSTRACT

Merkel cell polyomavirus (MCV) small T antigen (sT) is the main oncoprotein for the

development of Merkel cell carcinoma (MCC). MCC is a rare, clinically aggressive

neuroendocrine tumor of the skin with a high propensity for local, regional, and distant spread.

The dysregulation of matrix metalloproteinase-9 (MMP-9) has been implicated in multiple

essential roles in the development of various malignant tumor cell invasion and metastasis.

Previously, MCV sT was shown to induce the migratory and invasive phenotype of MCC cells

through the transcriptional activation of the Sheddase molecule, ADAM 10 (A disintegrin and

metalloprotease domain-containing protein 10). In this study, we show that MCV sT protein

stimulates differential expression of epithelial–mesenchymal transition (EMT) associated genes,

including MMP-9 and Snail. This effect is dependent on the presence of the large T stabilization

domain (LSD), which is known to be responsible for cell transformation through targeting of

promiscuous E3 ligases, including FBW7, a known MMP-9 and Snail regulator. Chemical

treatments of MMP-9 markedly inhibited sT-induced cell migration and invasion. These results

suggest that MCV sT contributes to the activation of MMP-9 as a result of FBW7 targeting, and

increases the invasive potential of cells, which can be used for targeted therapeutic intervention.

IMPORTANCE

Merkel cell carcinoma (MCC) is the most aggressive cutaneous tumor without clearly defined

treatment. Although MCC has a high propensity for metastasis, little is known about the

underlying mechanisms that drive MCC invasion and metastatic progression. MMP-9 has shown

to play a detrimental role in many metastatic human cancers, including melanoma and other non-

melanoma skin cancers. Our study shows that MCV sT-mediated MMP-9 activation is driven

through the LSD, a known E3 ligase targeting domain, in MCC. MMP-9 may serve as the

biochemical culprit to target and develop a novel approach for the treatment of metastatic MCC.

INTRODUCTION

Merkel cell carcinoma (MCC) is a rare skin cancer of neuroendocrine origin with a high propensity to metastasize (1). Although the incidence rate of MCC is lower than melanoma, it is highly aggressive with an estimated mortality rate of 33%-46%; hence it is significantly more lethal than malignant melanoma (2). Merkel cell polyomavirus (MCV) is the etiological agent of MCC. The majority of MCC cases are associated with MCV as observed by monoclonal integration of the MCV genome in the tumor DNA (3). As a classic polyomavirus, the genomic organization of MCV is similar to other known human polyomaviruses. MCV expresses small and large tumor antigens (sT and LT), which are essential for viral replication and pathogenesis. MCC tumor-derived MCV LT sequences integrated into MCC genomes contain mutations prematurely truncating the C-terminal growth inhibitory domain (4), while MCV sT remains intact.

MCV sT has been shown to mediate multiple oncogenic mechanisms that contribute to MCC development. Inhibition of SCF (Skp1, Cullin, F-box containing complex) E3 ligases by MCV sT appears to induce several viral and cellular oncoprotein activation, leading to enhanced MCV replication and cell proliferation (5, 6). Aberrant activation of oncogenic potential in MCV sT expressing cells also promoted the malignant phenotypes that are involved in genomic instability such as centrosome amplification, aneuploidy, and micronuclei formation (7). This oncogenic activity of MCV sT requires the LT stabilization domain (LSD), a unique and disordered domain of MCV sT, which is known to interact with SCF E3 ligase complexes (5, 7). Although the exact mechanism by which MCV sT targets E3 ligases is yet to be elucidated, it is clear that the LSD plays a significant role in the distinctive transforming activities induced by MCV sT in vitro and in vivo (5-7).

The E3 ubiquitin ligase F-box and WD repeat domain containing 7 (FBW7) functions as a putative tumor suppressor and an evolutionarily conserved substrate receptor of SCF ubiquitin ligase complex and plays vital roles in cell proliferation and cell migration (8). In various cancers, including renal cancer (9, 10), gastric cancer (11) and hepatocellular carcinoma (12), FBW7 inhibition promotes metastasis and epithelial–mesenchymal transition (EMT) by upregulating matrix metalloproteinase expression, specifically MMP-2, MMP-9, and MMP-13.

Matrix metalloproteinases (MMPs) are a zinc-dependent family of proteolytic enzymes that

participate in the degradation of the extracellular matrix (ECM). Dysregulation of these proteases

has been observed in multiple cancers where enhanced expression of certain MMP proteins

contribute to cell migration, invasion, and angiogenesis (13, 14). Specifically, MMP-9 has been

linked to multiple hallmarks of cancer, including but not limited to metastasis, invasion,

immunological surveillance, and angiogenesis (15). MMP-9, also known as 92 kDa type IV

collagenase (16), plays a vital role in the degradation of elastin and partially hydrolyzed collagen

that is essential for maintaining epithelial structural integrity. Various studies have shown that

human tumor virus-associated oncoproteins play a critical role in metastasis and EMT-related

mechanisms. Hepatitis B virus (HBV)-encoded X protein (HBx) (17), Kaposi's sarcoma-

associated herpesvirus (KSHV) K1 (18), and Epstein-Barr virus (EBV) latent membrane protein

1 (LMP-1) proteins (19) are known to upregulate MMP-9 expression, thereby contributing to

invasiveness and metastasis, key hallmarks of cancer (20).

MCV sT stimulates cell motility by inducing microtubule destabilization (21), actin

rearrangement (22) and cell dissociation by disruption of cell junctions (23). Interrogation of

previously published quantitative proteomic datasets of MCV sT-expressing cells (21) indicates

that MCV sT activated expression of Snail, a transcription factor that enhances mesenchymal

genes, and MMP-9. In contrast, MCV sT significantly downregulated genes related to cell

adhesion molecules, suggesting the potential function of MCV sT in the regulation of EMT.

MMP-9 and Snail activation by MCV sT was strictly dependent on the presence of the LSD,

which resulted in the enhancement of cell migration in mouse fibroblast cells and human cancer

cell lines. Our findings indicate that MCV sT targeting of cellular E3 ligases may play a role in

MCV sT-induced cell migration and invasion in MCC. Notably, chemical treatment with MMP-9

inhibitors resulted in significant inhibition of MCV sT-induced cell migration and invasion. This

suggests that MMP-9 protein may be a viable target for novel therapeutic intervention for

disseminated MCC.

RESULTS

MCV sT expression induces differential expression of proteins associated with EMT.

Recent studies have highlighted the involvement of MCV sT in the highly migratory and cell dissociation phenotypes of MCC, elucidating its highly multifunctional roles in MCC (21-23). Previously described SILAC (stable isotope labeling by amino acids in cell culture)-based quantitative proteomics data (21) was further interrogated to assess the alterations in the host cell proteome upon expression of MCV sT in a HEK293 derived cell line (i293-sT) **(Fig. 1A)** (21). These results highlighted an alteration in proteins associated with enhancement of cell migration (microtubule-associated cytoskeletal organization) and cell adhesion as previously reported and the basement membrane proteins, a specialized form of the ECM. Specifically, the quantitative proteomic analysis showed an almost two-fold decrease in Collagen alpha-2(IV) chain (COL4A2) and Laminin subunit gamma 1(LAMC1), two essential components of the basement membrane. The basement membrane is crucial for epithelial structural integrity. It is comprised of a network of glycoproteins and proteoglycans such as Type IV collagen and laminin and provides a barrier from invasion by tumor cells (24). These results suggest that MCV sT plays a role in the basement membrane degradation, an essential process for the metastatic invasion of tumor cells into the circulatory system to occur in MCC. Aligned with these observations, transcriptome analysis has suggested that specific markers associated with EMT are increased upon MCV sT expression (25).

To validate the potential regulation of EMT-related markers by MCV sT, RT-qPCR was performed. Changes in mRNA levels of classic EMT markers were assessed in MCC13 cells overexpressing a control of MCV sT construct. Upon MCV sT expression, a significant downregulation in epithelial markers E-cadherin, Zonula occludens-1 (ZO-1) and Occludin was observed **(Fig. 1B)**. Conversely, mesenchymal markers Slug, Snail, ZEB1, ZEB2, MMP-3, and MMP-9 were upregulated upon MCV sT expression. These results infer the possibility of MCV sT inducing an EMT, which contributes to the metastatic potential of MCV-associated MCC.

MCV sT inhibition of FBW7 contributes to migratory phenotype.

An essential requirement of metastasis involves the dissemination of tumor cells to various organs from the primary tumor (26). Multiple studies have demonstrated that loss of FBW7

promotes cell invasion and migration in numerous cancers through modulation of EMT-related

cellular factors such as MMP-9 and Snail (27-29), which are upregulated transcriptionally upon

MCV sT expression **(Fig. 1B).** The MCV sT LSD region is known to bind and inhibit the FBW7

(5, 7). As a result of this inhibition, FBW7 oncogenic substrates are stabilized in MCV sT-

expressing cells, which may contribute to MCV sT-induced migratory phenotype. Although

MCV sT and FBW7 interaction has been characterized in vitro by co-immunoprecipitation in

over-expressing cell, the in vitro techniques do not identify whether this interaction occurs with

endogenous proteins and, therefore, may not reflect the native behavior of their endogenous

counterparts. In situ Proximity ligation assay (PLA) can detect interactions with high specificity

and sensitivity due to the coupling of antibody recognition and DNA amplification, which

provides a technical advantage over other protein-protein interaction assays often plagued with

long preparation times and extensive troubleshooting. For that reason, we utilized PLA combined

with flow cytometry to revalidate this interaction (30, 31). As shown in **Fig. 2A**, quantification

of wild type MCV sT interaction with FBW7 resulted in high-intensity PLA signal comparative

to our positive control c-Myc, a well-known FBW7 substrate (32). This interaction was markedly

diminished by expression of sT_{LSDm} **(Fig. 2A)**, an LSD alanine mutant of MCV sT, consistent

with the finding from a previous report (5).

To determine whether MCV sT targeting of FBW7 contributes to sT-induced migratory

phenotype, a scratch assay was performed comparing vector control, MCV sT_{WT}, and MCV

sT_{LSDm} in U2OS cells. Images of the scratch area were recorded at time point 0 and 24 h post

scratch. Compared to empty vector negative control, MCV sT greatly enhanced the motility and

migration of U2OS cells, consistent with previous studies **(Fig. 2B)** (21-23, 33). In contrast,

MCV sT_{LSDm} did not show a significant increase in cell migration. Over the 24 h period of the

assay, we see no significant positive or negative effect on cell number confirmed by a viability

assay, indicating that the resulting phenotype is specific to cell migration **(Fig. 2B)**. Similarly,

enhanced cell migration was readily detected with wild type sT in NIH3T3 mouse fibroblast

(Fig. 2C) and MCC13 (MCV negative MCC cell line) **(Fig. S1B)**, while this phenotype is not

induced by sT_{LSDm}. This suggests that sT targeting of FBW7 may be involved in the MCV sT-

induced cell migratory phenotype.

MCV sT inhibition of FBW7 prevents turnover of MMP-9.

As shown in **Fig. 1B**, MCV sT induces MMP-9, an essential protein associated with the FBW7-

EMT axis in human cancers (34). To confirm MCV sT induction of MMP-9 expression, a variety

of cell lines, 293, COS-7, MCC13, and U2OS cells were transfected with a vector control and

MCV sT plasmids. MMP-9 gene expression is primarily regulated transcriptionally, resulting in

low basal levels of these proteases in normal physiology (35). RT-qPCR results showed that

MCV sT expression significantly increased MMP-9 transcript levels in all cell lines tested (**Fig.**

3A).

We posited that MCV sT targeting of FBW7 plays a role in promoting the migratory potential of

MCC by preventing MMP-9 protein turnover. Both transcriptional and post-transcriptional levels

of MMP-9 were assessed in the presence of the MCV sT_{WT} or MCV sT_{LSDm} in U2OS and

MCC13 cell lines (**Fig. 3B**). Results showed that MCV sT significantly induced the upregulation

of MMP-9 transcripts when analyzed by RT-qPCR in both cell lines, which was not observed

upon mutation of the MCV sT LSD (**Fig. 3B**). Additionally, we performed immunoblot analysis

to evaluate the effect of sT on MMP-9 protein levels (**Fig. 3C and 3D**). Studies have shown

MMP-9 exists in several forms; a monomeric pro- (∼92 kDa), a disulfide-bonded homodimeric

(∼220 kDa) and multiple active forms (∼67 and 82 kDa) (36). The active and dimeric forms of

MMP-9 play a role in the invasive and migratory phenotypes of cancer cells (37, 38). MCV sT_{WT}

expression induced the upregulation of the dimer, monomer, and active forms of MMP-9.

However, mutation of the LSD prevented MCV sT-mediated upregulation of MMP-9 as

expression levels remained comparable with the control, suggesting that MCV sT-mediated

upregulation of MMP-9 is LSD-dependent (**Fig. 3**).

MMP-9 inhibition impedes MCV sT-induced cell migration.

We next sought to determine if MMP-9 inhibition would have an impact on MCV sT induced

motile and migratory potentials. The migratory phenotype of U2OS cells transfected with vector

control, sT_{WT} and sT_{LSDm} was assessed using a scratch assay in the absence or presence of non-

cytotoxic concentrations of MMP9-I and MMP9-II inhibitors (**Fig. 4A and Fig. S2A**). MMP-9

inhibition resulted in a significant decrease in the distance traveled by MCV sT expressing cells

(**Fig. 4A and 4B**), confirming that the MMP-9 is a critical migratory factor that is regulated by

MCV sT. Incubation of both inhibitors showed a slight decrease in the motility of vector control

cells, implying that any changes observed in migratory rates of MCV sT expression cells are not

due to changes in cell viability or cytotoxicity. Both inhibitors showed a minor impact on the

motility of sT_{LSDm} expressing cells, comparable to vector control cells.

MMP-9 is essential for cell motility and migration in MCC.

To demonstrate that MMP-9 is vital for cell motility and migration in metastatic MCC, a

transwell migration assay was performed using MCV-positive MCC cell lines. This assay

quantified the migration ability of MCC cells towards a chemoattractant across a permeable

chamber. MCV-positive MCC cell lines, MKL-1 and MS-1 (**Fig. 4C**), were incubated in the

absence or presence of the MMP9-I and MMP9-II inhibitors at non-toxic concentrations (**Fig.**

S2B). After treatment, cells were allowed to migrate for 48 h and the total numbers of migrated

cells were measured by cell counting Kit-8 assay. Results showed the migration of MCV-

positive MCC cell lines was significantly reduced (~40 to 50%) upon incubation of both MMP-9

inhibitors in comparison to the untreated control, suggesting that MMP-9 expression contributes

to the migratory capacity of MCV-positive MCC (**Fig. 4C**). Together, these results indicate that

MMP-9 is required for MCV sT-mediated cell migration enhancement in MCC.

MCV sT invasive phenotype is LSD-dependent.

The invasiveness of epithelial cancers is a multi-step process and a key hallmark involves the

degradation of the basement membrane. Type IV collagen is a major component in most

basement membranes. Multiple studies have correlated overexpression of MMPs with not only

an enhancement of cell migration and metastasis, but also the invasiveness of cancer cells (13,

15). In particular, MMP-9 is a key protease associated with the degradation of ECM components,

including type IV collagen and laminin, which in turn facilitates invasion of tumors into the

circulatory system and promotes metastasis. To test if enzymatic activation of MMP-9 is

regulated by sT LSD, we evaluated the effect of MCV sT on MMP-9 substrate collagen IV, by

immunofluorescence staining in U2OS cells. Our results demonstrate that collagen IV expression

in MCV sT_{WT} expressing cells is significantly reduced in comparison to vector control cells,

potentially due to MMP-9 activation induced by MCV sT. In contrast, MCV sT_{LSDm} expressing

cells did not show a decrease in collagen IV expression (**Fig. 5A, Fig. S3**). Moreover, our

regression analysis revealed that collagen IV expression levels are highly correlated with MCV

sT_{WT} or sT_{LSDm} expression levels (**Fig. 5B**). To further validate the effect of MCV sT LSD on

collagen IV degradation, we performed an invasion assay using collagen pre-coated inserts in

U2OS and MCC13 cells. sT_{WT} induced 4 to 5-fold increases in collagen invasion compared to

either vector control or sT_{LSDm} (**Fig. 5C**), inferring that MCV sT induces not only cell migration,

but also cancer cell invasion through the LSD.

MCV sT activates expression of EMT regulator, Snail.

A positive regulatory loop has been identified between MMP-9 and Snail. siRNA mediated

inhibition of MMP-9 significantly reduces expression of Snail, and conversely, knockdown of

Snail, a transcription factor of MMP-9, suppresses expression of MMP-9 (39). FBW7 abrogation

of Snail protein also inhibits MMP-9 expression (40). Interestingly, MCV sT induced Snail

expression in our initial transcriptional analysis (**Fig. 1B**). Since both MMP-9 and Snail are vital

mediators of EMT, we assessed the effect of sT LSD on transcriptional and protein levels of

Snail in U2OS cells. RT-qPCR results showed that MCV sT_{WT} upregulated mRNA levels of

Snail, while this transcriptional change was not observed upon mutation of the LSD (**Fig. 6A**).

Similar to RT-qPCR data, our results demonstrated a significant increase in Snail protein levels

upon MCV sT expression in an LSD-dependent manner (**Fig. 6B**).

DISCUSSION

Metastasis is the endpoint of a series of biological processes by which a tumor cell detaches from

the primary tumor and disseminates to a distant site through the circulatory system and

establishes a secondary tumor (41). Oncogenic viruses often modulate the EMT axis via

regulating E-cadherin repression (42-44), fibroblast growth factor (FGF) ligand modulation (44,

45), cadherin switching (46, 47), induction of transcription factors such as TWIST (48, 49) and

Snail (42, 50), and MMP-9 upregulation (17-19). These cellular targets can regulate cancer cell

migration and invasion; therefore, they could be exploited for therapeutic strategies in virus-

induced metastatic cancers.

Multiple F box proteins can function as tumor suppressors by negatively regulating oncoproteins,

and various studies have focused on elucidating this mechanism in tumorigenesis and EMT

progression (34). In this report, we show that MCV sT LSD inhibition of FBW7 promotes the
upregulation of MMP-9 and contributes to MCV sT-mediated cell migration and invasion. The
mechanism by which FBW7 regulates MMP-9 expression is currently unclear, although many
studies have shown MMP-9 expression is directly and indirectly regulated by FBW7 substrates
such as XBP1, Notch1, and Snail (12, 40, 51). Snail is known to be a substrate of both FBW7
(40) and β-TrCP (52), another major SCF E3 ligase that MCV sT targets through the LSD (7).
As previously shown, Snail gene expression induces the loss of epithelial markers and the gain of
mesenchymal markers, as well as promoting changes in cell motility and invasive properties
(53). Our study initially focused on an MMP-9 specific metastatic progression induced by MCV
sT due to limited availability of Snail inhibitors. However, the distressed proteome balance in
EMT molecules induced by MCV sT might be triggered by Snail activation through MCV sT
targeting multiple E3 ligases, which requires further investigation.
302
While the detailed regulatory mechanisms and specificity of sT function in transcription
modulation remain unclear, studies have shown that MCV sT mediates cellular
transcriptome/chromatin remodeling (25, 54) which may alter transcriptional activity and gene
expression. Consistent with our data, Berrios et al. also reported that MCV sT downregulates
extracellular matrix organization and cell adhesion molecules in their transcriptome analysis
(25). We have demonstrated that MCV sT specifically activates both mRNA and protein levels
of the EMT-related cellular proteins MMP-9 and Snail through the LSD; however, our results do
not rule out the possibility that this effect is potentially modulated by multiple mechanisms in
MCC. Nonetheless, it is clear that MCV sT LSD plays a critical role in regulating metastasis-
initiating capacity in MCC that could be a potential target for therapeutic interventions.

The underlying mechanism for the high propensity of MCC tumors to metastasize is yet to be
elucidated. Because of the rare and aggressive nature of metastatic MCC and the lack of standard
chemotherapy, there are no prospective studies of outcomes following treatment of distant
metastatic MCC. Since recent FDA approvals of Avelumab and Pembrolizumab represent the
only approved treatment option for metastatic MCC (55, 56), it is necessary to evaluate the
preclinical anticancer activity of efficient and economical chemotherapeutics through
retrospective analysis for both MCV-negative and positive MCC patients. MCV-negative MCC

tumors patients are more likely to present with advanced disease than patients with virus-positive

tumors (66.7% vs. 48.3%) (57). However, targeting the signaling pathways implicated in

regulating tumor invasion could be an effective therapeutic protocol for both types of metastatic

MCC treatment.

Our study is the first approach to investigate the therapeutic potential of matrix metalloproteinase

in MCC. MCV sT specifically activates the EMT-related cellular proteins MMP-9 through the

LSD, which we targeted by commercially available inhibitors, and revealed a potential secondary

treatment for distant metastatic MCC.

MATERIALS AND METHODS

Cells. 293, U2OS, and COS-7 cells were maintained in Dulbecco's modified Eagle's medium

(DMEM) containing 10% fetal bovine serum (FBS) (Seradigm). MCC13, MS-1, and MKL-1

cell lines were maintained in RPMI 1640 medium supplemented with 10% FBS (Seradigm).

NIH3T3 cells were maintained in DMEM with 10% bovine calf serum (Seradigm).

Plasmids, transfection and transduction. Plasmids for vector control, codon-optimized cDNA

constructs for sT_{WT} and sT_{LSDm} have been previously described (5). HA-Fbw7 and Flag-cMyc

(32) plasmids were kindly provided by Dr. Nakayama (Kyushu University, Japan).

FBW7ΔDF(d231-324) was generated by overlapping PCR using primers listed in Table S2. For

sT protein expression, cells were transfected using Lipofectamine 3000 (Invitrogen) or

jetOPTIMUS (Polyplus Transfection) according to the manufacturer's protocol. For lentiviral

transduction, codon-optimized cDNAs for MCV sT_{WT}, MCV $sT_{LSDm,}$ (5) and H-RasV12 were

inserted into pLVX empty vector. Plasmids used for this study were listed in **Table S1**

(Supplemental material). For lentivirus production, 293FT (Invitrogen) cells were used for

induction according to the manufacturer's instructions. Cells were selected with puromycin (3

μg/ml) after infection for one week.

Reverse transcription-quantitative polymerase chain reaction (RT-qPCR). RNA was

extracted using Monarch Total RNA miniprep kit (New England Biolabs), as per the

manufacturer's instruction. 250 ng of RNA was used as a template in each reaction with iTaq

Universal One-Step RT-qPCR Kit (Bio-Rad) or Luna Universal One-Step RT-qPCR Kit (New

England Biolabs). Primer sequences used are described in **Table S2**. With GAPDH as an internal

control, quantitative analysis was performed using the comparative $\Delta\Delta$Ct method.

Quantitative immunoblotting (IB) and antibodies. Cells were lysed in IP buffer (50 mM Tris-

HCl (pH 8.0), 150 mM NaCl, 1% TritonX-100, 1 mM PMSF, 1 mM benzamidine) and sonicated

whole cell lysates were used for direct immunoblotting. Primary antibodies were incubated

overnight at 4°C, followed by 1 h secondary antibody incubation at RT. All signals were

detected using quantitative Infrared (IR) secondary antibodies (IRDye 800CW goat anti-mouse,

800CW goat anti-rabbit, 680LT goat anti-rabbit IgG, 680LT goat anti-mouse IgG) (LI-COR).

Signal intensities were analyzed using a laser-scanning imaging system, Odyssey CLX (LI-

COR). Antibodies used for this study are listed in **Table S3**. Protein levels were quantitated and

normalized by control, α-Tubulin, or β-Actin, using an Odyssey LI-COR IR imaging system.

SILAC data analysis. The previously published SILAC-based quantitative proteomic data set

analyzing host cell proteome changes upon MCV sT expression (21) and further interrogated

using the Database for Annotation, Visualization and Integrated Discovery (DAVID) v6.7 (51).

For quantitative analysis, a 2.0-fold cutoff was chosen as a basis for investigating potential

proteome changes (50).

Proximity ligation assay (PLA) Flow cytometry. PLA was performed using a Duolink assay

kit (Sigma-Aldrich) according to the manufacturer's instructions. To evaluate MCV sT and

FBW7 interaction, HA-FBW7ΔDF(d231-324) was co-expressed with sT_{WT} or sT_{LSDm}.

FBW7ΔDF was also co-expressed with c-Myc, a known FBW7 substrate, as a positive control

(32). Primary antibodies were utilized at optimized concentrations with HA-Tag (C29F4)

(1:500), c-Myc (9E10) (1:500), and 2T2 (1:500) (Millipore). Cells were analyzed by flow

cytometry on a 16-color BD LSR Fortessa. The acquired data were analyzed using FlowJo

software (Tree Star, Ashland, OR, USA).

Scratch wound-healing assay. Cells were seeded into the Poly-L-Lysine-coated 6-well plates
and transfected with either empty vector or sT_{WT} or sT_{LSDm} plasmids. Because MCV sT promotes
serum-independent cell growth (58), a serum starvation condition was not considered for our
scratch assay to exclude cell proliferation effect by sT. After 48 h, a scratch was created by
scraping the monolayer using a p1000 pipette tip. The migration of cells toward the scratch was
observed over a 24 h period, and images were taken every 8 h under a REVOLVE4 fluorescent
microscope (Echo Laboratories). Inhibitor-based scratch assays were incubated for 24 h prior to
transfection with 0.1 and 1 μM of 9-I and 9-II inhibitors respectively.

Transwell cell migration assay. Cells grown in DMEM with 10% FBS were trypsinized and
resuspended in DMEM. 1×10^5 cells were gently added to the transwell insert (8 μm, Greiner
Bio-One). DMEM with 10% FBS was added to the bottom of the lower chamber (24-well plate).
The cells were incubated in the culture incubator at 37 °C plus 5% CO2 for the indicated time.
The cells migrated from the insert to the well through the filter. The filter was fixed with 4%
paraformaldehyde in PBS for 10 min, then stained with 1% Crystal Violet in 2% ethanol for 20
min for NIH3T3 cells and MCC cells were counted using a Cell Counting Kit-8 (CCK-8)
(Sigma-Aldrich). The stained cells on the lower side were counted under a microscope from 5
different randomly selected views. All conditions were the same for assays performed in
triplicate.

Immunofluorescence. U2OS cells grown on glass coverslips were transfected with empty vector
or sT wild type or LSD mutant expression constructs. After 48 h, cells were fixed in 1:1
methanol/acetone at -20^oC, permeabilized, and blocked in PBS with 5% BSA and 0.3 M glycine
for 1 h. Cells were labeled with the appropriate primary antibodies and then incubated with the
appropriate Alexa Fluor-conjugated secondary antibody. Cells were analyzed with a
REVOLVE4 fluorescent microscope (Echo Laboratories).

Collagen invasion assay. U2OS and MCC13 cells were transfected with wild type and LSD
mutant sT constructs for 48 h, followed by overnight serum starvation. 1×10^6 cells
resuspending in serum-free media in each condition were seeded in a 24-well cell invasion plate
containing polymerized collagen-coated membrane inserts. The collagen inserts had a pore size

of 8 μm (Chemicon QCM Collagen Cell Invasion Assay, ECM551). Complete medium was used

as a chemoattractant in the lower chamber and cells were left to incubate for 72 h. Cells/media

were carefully aspirated by pipetting any residual suspension in the transwell insert. Inserts were

transferred to a clean well and were carefully stained with 400 μL cell staining solution at room

temperature for 20 minutes, followed by a gentle wash in deionized water. While slightly damp,

unattached cells were removed cautiously by cotton swabs from the collagen inserts and allowed

to dry at room temperature for 15 minutes. Dried inserts were transferred to clean wells

containing 200 μL of extraction buffer and incubated for 15 minutes at room temperature.

Following the extraction incubation, 100 μL of the extraction solution was pipetted into 96 well

plates, and optical density was measured at 560 nm.

Cell Proliferation Assay. U2OS transfected cells (vector control, MCV sT$_{WT}$ and MCV sT$_{LSDm}$

plasmids) were seeded in 96 well plates (1×10^4 cells/well) 48 h post-transfection. Cell

proliferation was monitored using a WST-8 based assay Cell Counting Kit-8 (CCK-8) according

to the manufacturer's protocol. OD values were divided by the OD value of day 0 for

normalization.

Chemical inhibitors. MMP-9 inhibitors-I and II (EMD Millipore) were used at 0.1 to 0.2 μM

and 1 to 2 μM, respectively. Cell toxicity was measured using a Cell Counting Kit-8 (CCK-8)

(Sigma-Aldrich) according to the manufacturer's protocol.

Statistical analysis. Statistical significance between two groups was determined using one- or

two-tailed student's t-tests in GraphPad Prism (GraphPad Software, Inc., La Jolla, CA, USA).

The difference was considered significant when $p < 0.05$ for multiple testing. *, **, *** = p-

value < 0.01, 0.005 and 0.001, respectively.

Acknowledgments

We thank Dr. Patrick S Moore and Dr. Yuan Chang for kind sharing of MCV-related reagents.

H.J.K. was supported in part by an Institutional Research Grant, IRG-17-175-04 from the

American Cancer Society, and by the Pennsylvania Department of Health Tobacco CURE

Funds. N.N. was supported by training grant T32 CA060395 from the National Cancer Institute,

National Institutes of Health.

References

1. Becker JC, Stang A, DeCaprio JA, Cerroni L, Lebbé C, Veness M, Nghiem P. 2017. Merkel cell carcinoma. Nat Rev Dis Primers 3:17077.

2. Miller RW, Rabkin CS. 1999. Merkel cell carcinoma and melanoma: Etiological similarities and differences (vol 8, pg 153, 1999). Cancer Epidemiology Biomarkers & Prevention 8:485-485.

3. Feng H, Shuda M, Chang Y, Moore PS. 2008. Clonal integration of a polyomavirus in human Merkel cell carcinoma. Science 319:1096-100.

4. Shuda M, Feng HC, Kwun HJ, Rosen ST, Gjoerup O, Moore PS, Chang Y. 2008. T antigen mutations are a human tumor-specific signature for Merkel cell polyomavirus. Proceedings of the National Academy of Sciences of the United States of America 105:16272-16277.

5. Kwun HJ, Shuda M, Feng H, Camacho CJ, Moore PS, Chang Y. 2013. Merkel Cell Polyomavirus Small T Antigen Controls Viral Replication and Oncoprotein Expression by Targeting the Cellular Ubiquitin Ligase SCFFbw7. Cell Host & Microbe 14:125-135.

6. Verhaegen ME, Mangelberger D, Harms PW, Vozheiko TD, Weick JW, Wilbert DM, Saunders TL, Ermilov AN, Bichakjian CK, Johnson TM, Imperiale MJ, Dlugosz AA. 2015. Merkel cell polyomavirus small T antigen is oncogenic in transgenic mice. J Invest Dermatol 135:1415-1424.

7. Kwun HJ, Wendzicki JA, Shuda Y, Moore PS, Chang Y. 2017. Merkel cell polyomavirus small T antigen induces genome instability by E3 ubiquitin ligase targeting. Oncogene 36:6784-6792.

8. Ang XL, Wade Harper J. 2005. SCF-mediated protein degradation and cell cycle control. Oncogene 24:2860-70.

9. Cai YK, Zhang M, Qiu XF, Wang BW, Fu Y, Zeng J, Bai J, Yang GS. 2017. Upregulation of FBXW7 Suppresses Renal Cancer Metastasis and Epithelial Mesenchymal Transition. Disease Markers.

10. He H, Dai J, Xu Z, He W, Wang X, Zhu Y, Wang H. 2018. Fbxw7 regulates renal cell carcinoma migration and invasion via suppression of the epithelial-mesenchymal transition. Oncol Lett 15:3694-3702.

11. Calcagno DQ, Freitas VM, Leal MF, De Souza CRT, Demachki S, Montenegro R, Assumpcao PP, Khayat AS, Smith MDC, dos Santos A, Burbano RR. 2013. MYC, FBXW7 and TP53 copy number variation and expression in Gastric Cancer. Bmc Gastroenterology 13.

12. Wang X, Zhang J, Zhou L, Sun W, Zheng ZG, Lu P, Gao Y, Yang XS, Zhang ZC, Tao KS, Dou KF. 2015. Fbxw7 regulates hepatocellular carcinoma migration and invasion via Notch1 signaling pathway. International Journal of Oncology 47:231-243.

13. Gialeli C, Theocharis AD, Karamanos NK. 2011. Roles of matrix metalloproteinases in cancer progression and their pharmacological targeting. Febs Journal 278:16-27.

14. Nagase H, Woessner JF. 1999. Matrix metalloproteinases. Journal of Biological Chemistry 274:21491-21494.

15. Huang H. 2018. Matrix Metalloproteinase-9 (MMP-9) as a Cancer Biomarker and MMP-9 Biosensors: Recent Advances. Sensors 18.

487   16.   Wilhelm SM, Collier IE, Marmer BL, Eisen AZ, Grant GA, Goldberg GI. 1989. SV40-
488   transformed human lung fibroblasts secrete a 92-kDa type IV collagenase which is
489   identical to that secreted by normal human macrophages. J Biol Chem 264:17213-21.
490   17.   Chung TW, Lee YC, Kim CH. 2004. Hepatitis B viral HBx induces matrix
491   metalloproteinase-9 gene expression through activation of ERK and PI-3K/AKT
492   pathways: involvement of invasive potential. FASEB J 18:1123-5.
493   18.   Wang L, Wakisaka N, Tomlinson CC, DeWire SM, Krall S, Pagano JS, Damania B.
494   2004. The Kaposi's sarcoma-associated herpesvirus (KSHV/HHV-8) K1 protein induces
495   expression of angiogenic and invasion factors. Cancer Research 64:2774-2781.
496   19.   Yoshizaki T, Sato H, Furukawa M, Pagano JS. 1998. The expression of matrix
497   metalloproteinase 9 is enhanced by Epstein-Barr virus latent membrane protein 1.
498   Proceedings of the National Academy of Sciences of the United States of America
499   95:3621-3626.
500   20.   Hanahan D, Weinberg RA. 2011. Hallmarks of Cancer: The Next Generation. Cell
501   144:646-674.
502   21.   Knight LM, Stakaityte G, Wood JJ, Abdul-Sada H, Griffiths DA, Howell GJ, Wheat R,
503   Blair GE, Steven NM, Macdonald A, Blackbourn DJ, Whitehouse A. 2015. Merkel cell
504   polyomavirus small T antigen mediates microtubule destabilization to promote cell
505   motility and migration. J Virol 89:35-47.
506   22.   Stakaityte G, Nwogu N, Dobson SJ, Knight LM, Wasson CW, Salguero FJ, Blackbourn
507   DJ, Blair GE, Mankouri J, Macdonald A, Whitehouse A. 2018. Merkel Cell
508   Polyomavirus Small T Antigen Drives Cell Motility via Rho-GTPase-Induced
509   Filopodium Formation. Journal of Virology 92.
510   23.   Nwogu N, Boyne JR, Dobson SJ, Poterlowicz K, Blair GE, Macdonald A, Mankouri J,
511   Whitehouse A. 2018. Cellular sheddases are induced by Merkel cell polyomavirus small
512   tumour antigen to mediate cell dissociation and invasiveness. Plos Pathogens 14.
513   24.   Horejs CM. 2016. Basement membrane fragments in the context of the epithelial-to-
514   mesenchymal transition. European Journal of Cell Biology 95:427-440.
515   25.   Berrios C, Padi M, Keibler MA, Park DE, Molla V, Cheng JW, Lee SM, Stephanopoulos
516   G, Quackenbush J, DeCaprio JA. 2016. Merkel Cell Polyomavirus Small T Antigen
517   Promotes Pro-Glycolytic Metabolic Perturbations Required for Transformation. Plos
518   Pathogens 12:21.
519   26.   Bravo-Cordero JJ, Hodgson L, Condeelis J. 2012. Directed cell invasion and migration
520   during metastasis. Current Opinion in Cell Biology 24:277-283.
521   27.   Sailo BL, Banik K, Girisa S, Bordoloi D, Fan L, Halim CE, Wang H, Kumar AP, Zheng
522   DL, Mao XL, Sethi G, Kunnumakkara AB. 2019. FBXW7 in Cancer: What Has Been
523   Unraveled Thus Far? Cancers 11:31.
524   28.   Zhang Y, Zhang XX, Ye MX, Jing PY, Xiong J, Han ZP, Kong J, Li MY, Lai XF, Chang
525   N, Zhang J. 2018. FBW7 loss promotes epithelial-to-mesenchymal transition in non-
526   small cell lung cancer through the stabilization of Snail protein. Cancer Letters 419:75-
527   83.
528   29.   Lamouille S, Xu J, Derynck R. 2014. Molecular mechanisms of epithelial-mesenchymal
529   transition. Nature Reviews Molecular Cell Biology 15:178-196.
530   30.   Zhu XZ, Zelmer A, Wellmann S. 2017. Visualization of Protein-protein Interaction in
531   Nuclear and Cytoplasmic Fractions by Co-immunoprecipitation and In Situ Proximity
532   Ligation Assay. Jove-Journal of Visualized Experiments.

31.   Andersen SS, Hvid M, Pedersen FS, Deleuran B. 2013. Proximity ligation assay
combined with flow cytometry is a powerful tool for the detection of cytokine receptor
dimerization. Cytokine 64:54-57.
32.   Yada M, Hatakeyama S, Kamura T, Nishiyama M, Tsunematsu R, Imaki H, Ishida N,
Okumura F, Nakayama K, Nakayama KI. 2004. Phosphorylation-dependent degradation
of c-Myc is mediated by the F-box protein Fbw7. EMBO J 23:2116-25.
33.   Stakaityte G, Nwogu N, Lippiat JD, Blair GE, Poterlowicz K, Boyne JR, Macdonald A,
Mankouri J, Whitehouse A. 2018. The cellular chloride channels CLIC1 and CLIC4
contribute to virus-mediated cell motility. Journal of Biological Chemistry 293:4582-
4590.
34.   Song YZ, Lin M, Liu Y, Wang ZW, Zhu XQ. 2019. Emerging role of F-box proteins in
the regulation of epithelial-mesenchymal transition and stem cells in human cancers.
Stem Cell Research & Therapy 10.
35.   Fanjul-Fernandez M, Folgueras AR, Cabrera S, Lopez-Otin C. 2010. Matrix
metalloproteinases: Evolution, gene regulation and functional analysis in mouse models.
Biochimica Et Biophysica Acta-Molecular Cell Research 1803:3-19.
36.   Olson MW, Bernardo MM, Pietila M, Gervasi DC, Toth M, Kotra LP, Massova I,
Mobashery S, Fridman R. 2000. Characterization of the monomeric and dimeric forms of
latent and active matrix metalloproteinase-9 - Differential rates for activation by
stromelysin 1. Journal of Biological Chemistry 275:2661-2668.
37.   Dufour A, Zucker S, Sampson NS, Kuscu C, Cao JA. 2010. Role of Matrix
Metalloproteinase-9 Dimers in Cell Migration DESIGN OF INHIBITORY PEPTIDES.
Journal of Biological Chemistry 285:35944-35956.
38.   Roomi MW, Kalinovsky T, Rath M, Niedzwiecki A. 2014. Effect of a nutrient mixture
on matrix metalloproteinase-9 dimers in various human cancer cell lines. Int J Oncol
44:986-92.
39.   Lin CY, Tsai PH, Kandaswami CC, Lee PP, Huang CJ, Hwang JJ, Lee MT. 2011. Matrix
metalloproteinase-9 cooperates with transcription factor Snail to induce epithelial-
mesenchymal transition. Cancer Science 102:815-827.
40.   Fu Q, Lu Z, Fu X, Ma S, Lu X. 2019. MicroRNA 27b promotes cardiac fibrosis by
targeting the FBW7/Snail pathway. Aging (Albany NY) 11:11865-11879.
41.   Valastyan S, Weinberg RA. 2011. Tumor Metastasis: Molecular Insights and Evolving
Paradigms. Cell 147:275-292.
42.   Morris MA, Laverick L, Wei WB, Davis AM, O'Neill S, Wood L, Wright J, Dawson
CW, Young LS. 2018. The EBV-Encoded Oncoprotein, LMP1, Induces an Epithelial-to-
Mesenchymal Transition (EMT) via Its CTAR1 Domain through Integrin-Mediated
ERK-MAPK Signalling. Cancers 10.
43.   Liu J, Lian Z, Han S, Waye MMY, Wang H, Wu MC, Wu K, Ding J, Arbuthnot P, Kew
M, Fan D, Feitelson MA. 2006. Downregulation of E-cadherin by hepatitis B virus X
antigen in hepatocellullar carcinoma. Oncogene 25:1008-1017.
44.   Cheng YM, Chou CY, Hsu YC, Chen MJ, Wing LYC. 2012. The role of human
papillomavirus type 16 E6/E7 oncoproteins in cervical epithelial-mesenchymal transition
and carcinogenesis. Oncology Letters 3:667-671.
45.   Wakisaka N, Murono S, Yoshizaki T, Furukawa M, Pagano JS. 2002. Epstein-barr virus
latent membrane protein 1 induces and causes release of fibroblast growth factor-2.
Cancer Res 62:6337-44.

46.   Shair KHY, Schnegg CI, Raab-Traub N. 2009. Epstein-Barr Virus Latent Membrane
Protein-1 Effects on Junctional Plakoglobin and Induction of a Cadherin Switch. Cancer
Research 69:5734-5742.
47.   Hu DX, Zhou JS, Wang FF, Shi HY, Li Y, Li BH. 2015. HPV-16 E6/E7 promotes cell
migration and invasion in cervical cancer via regulating cadherin switch in vitro and in
vivo. Archives of Gynecology and Obstetrics 292:1345-1354.
48.   Horikawa T, Yang J, Kondo S, Yoshizaki T, Joab I, Furukawa N, Pagano JS. 2007. Twist
and epithelial-mesenchymal transition are induced by the EBV oncoprotein latent
membrane protein 1 and are associated with metastatic nasopharyngeal carcinoma.
Cancer Research 67:1970-1978.
49.   Chen X, Bode AM, Dong ZG, Cao Y. 2016. The epithelial-mesenchymal transition
(EMT) is regulated by oncoviruses in cancer. Faseb Journal 30:3001-3010.
50.   Horikawa T, Yoshizaki T, Kondo S, Furukawa M, Kaizaki Y, Pagano JS. 2011. Epstein-
Barr Virus latent membrane protein 1 induces Snail and epithelial-mesenchymal
transition in metastatic nasopharyngeal carcinoma. British Journal of Cancer 104:1160-
1167.
51.   Chae U, Lee H, Kim B, Jung H, Kim BM, Lee AH, Lee DS, Min SH. 2019. A negative
feedback loop between XBP1 and Fbw7 regulates cancer development. Oncogenesis
8:12.
52.   Zhou BHP, Deng J, Xia WY, Xu JH, Li YM, Gunduz M, Hung MC. 2004. Dual
regulation of Snail by GSK-3 beta-mediated phosphorylation in control of epithelial-
mesenchymal transition. Nature Cell Biology 6:931-+.
53.   Przybylo JA, Radisky DC. 2007. Matrix metalloproteinase-induced epithelial-
mesenchymal transition: Tumor progression at Snail's pace. International Journal of
Biochemistry & Cell Biology 39:1082-1088.
54.   Cheng JW, Park DE, Berrios C, White EA, Arora R, Yoon R, Branigan T, Xiao TF,
Westerling T, Federation A, Zeid R, Strober B, Swanson SK, Florens L, Bradner JE,
Brown M, Howley PM, Padi M, Washburn MP, DeCaprio JA. 2017. Merkel cell
polyomavirus recruits MYCL to the EP400 complex to promote oncogenesis. Plos
Pathogens 13:31.
55.   Joseph J, Zobniw C, Davis J, Anderson J, Trinh V. 2018. Avelumab: A Review of Its
Application in Metastatic Merkel Cell Carcinoma. Annals of Pharmacotherapy 52:928-
935.
56.   Shirley M. 2018. Avelumab: A Review in Metastatic Merkel Cell Carcinoma. Targeted
Oncology 13:409-416.
57.   Moshiri AS, Doumani R, Yelistratova L, Blom A, Lachance K, Shinohara MM, Delaney
M, Chang O, McArdle S, Thomas H, Asgari MM, Huang ML, Schwartz SM, Nghiem P.
2017. Polyomavirus-Negative Merkel Cell Carcinoma: A More Aggressive Subtype
Based on Analysis of 282 Cases Using Multimodal Tumor Virus Detection. Journal of
Investigative Dermatology 137:819-827.
58.   Shuda M, Kwun HJ, Feng HC, Chang Y, Moore PS. 2011. Human Merkel cell
polyomavirus small T antigen is an oncoprotein targeting the 4E-BP1 translation
regulator. Journal of Clinical Investigation 121:3623-3634.
622
623

 FIGURE LEGENDS

FIGURE 1

MCV sT leads to differential expression of proteins associated with epithelial to

mesenchymal transition (EMT). (A) Quantitative proteomics analysis illustrating differential

expression of EMT associated proteins upon MCV sT expression. Proteins associated with cell

adhesion and structural integrity of the extracellular matrix are downregulated upon MCV sT

expression. While expression of proteins which encourage cell migration by reorganization of

the actin network and microtubule destabilization are upregulated. **(B)** MCV sT regulates EMT-

associated gene expression. MCC13 cells were transfected with control or MCV sT expression

plasmids. While epithelial markers were downregulated, mesenchymal markers were

significantly upregulated upon MCV sT expression. Cellular RNA was extracted using a trizol

reagent and transcript levels were analyzed by RT-qPCR using the comparative $\Delta\Delta$Ct method (n

= 3).

FIGURE 2

MCV sT induces cell motility in an LSD-dependent manner. (A) Validation of MCV sT and

FBW7 interaction. To confirm the interaction of HA-FBW7 and MCV sT LSD domain, a PLA-

flow cytometric analysis was carried out. Wild type MCV sT displayed an interaction with

FBW7 comparative to the positive control interaction of c-Myc with FBW7, while mutation of

sT LSD greatly ablated sT interaction with FBW7. Primary antibodies were utilized at optimized

concentrations with HA-Tag (C29F4) (1:500), c-Myc (9E10) (1:500), and 2T2 (1:500). Protein

expression was evaluated by immunoblot analysis in **Fig. S1A**. **(B)** MCV sT induced-cell

migration is LSD-dependent. Scratch assay. Poly-L-lysine-coated 6-well plates were seeded with

U2OS cells and transfected with either a vector control, sT_{WT} and sT_{LSDm} (Fbw7 binding mutant)

plasmids. Migration of cells toward the scratch was observed over a 24 h period, and images

were taken using a REVOLVE4 fluorescent microscope (Echo Laboratories). Scratch assays

were performed in triplicate and measured using Fiji Image J analysis software. Differences

between means (*p* value) were analyzed using a t-test with GraphPad Prism software. Protein

expression was detected by immunoblot analysis to validate successful transfection using 2T2

antibody for sT antigens and α-Tubulin, respectively. No significant differences in cell

proliferation were observed between cells expressing MCV sT within 24 h, indicating that cell

proliferation does not interfere with the measurement of sT- induced cell migration. **(C)** MCV sT

promotes rodent fibroblast cell migration. NIH3T3 cells stably expressing an empty vector, H-

RasV12, sT_{WT} and sT_{LSDm} were trypsinized and $2x10^5$ cells were used for transwell migration

and scratch assay. H-RasV12 was used as a positive control. The experiments were performed

two times, and the results were reproducible. The graph indicates the fold difference of migrated

cells relative to the vector control sample. Protein expression was determined by

immunoblotting.

FIGURE 3

MCV sT activates Matrix metalloproteinase 9 (MMP-9). (A) MCV sT expression results in

upregulation of MMP-9 mRNA levels. Various cell lines (293, COS-7, MCC13 and U2OS) were

transfected with either empty vector or MCV sT_{WT} expressing plasmids to measure MMP-9

mRNA levels. After 48 h, total RNA was isolated and analyzed by RT-qPCR. **(B)** MCV sT

upregulates MMP-9 transcription through the LSD. U2OS and MCC13 cells transfected with

empty vector control, MCV sT_{WT} and MCV sT_{LSDm} expressing plasmids. Transcript levels of

MMP-9 were analyzed using the comparative $\Delta\Delta Ct$ method. (n = 3). Differences between means

(*p* value) were analyzed using a t-test with GraphPad Prism software. **(C)** MCV sT upregulates

MMP-9 protein expression through the LSD. U2OS cells were transfected with empty vector,

sT_{WT} and sT_{LSDm} expression plasmids. After 48 h, immunoblot analysis was performed to

analyze expression of MMP-9, sT and α-tubulin (Ci). Densitometry quantification of

immunoblots was carried out using the Image studio software and is shown as a fold change

relative to the loading control α-tubulin (Cii). Data analyzed using three biological replicates per

experiment (n = 3). **(D)** MCV sT reproducibly activates MMP-9 expression in MCC13.

FIGURE 4

MMP-9 inhibition impedes MCV sT-induced cell migration. (A) MCV sT promotes MMP-9-

induced cell migration. Scratch assay. Poly-L-lysine-coated 6-well plates were seeded with

U2OS cells and incubated with specific MMP-9 inhibitors at predetermined concentrations. Cells

were transfected with either a vector control, sT_{WT} and sT_{LSDm} plasmids. After 48 h, a scratch

was created and migration of cells toward the scratch was observed over a 24 h period. The size

of the wound was measured at 0 and 24 h and presented as the fold change in (**B**). Scratch assays

were performed in triplicate. (**C**) MMP-9 is required for MCC migration. MCV positive MCC

cell lines, MKL-1 and MS-1, were incubated with DMSO or the MMP-9 inhibitors 9-I (0.2 µM)

and 9-II (2 µM); 9-I (0.1 µM) and 9-II (1 µM), respectively. Cells were then transferred into

transwell inserts and allowed to migrate for 48 h. Migrated cells were measured using cell

counting kit-8 (CCK-8). Data analyzed using three biological replicates per experiment, n = 3.

Differences between means (*p* value) were analyzed using a t-test with GraphPad Prism

software.

FIGURE 5

MCV sT LSD induces collagen degradation. (**A**) MCV sT decreases collagen IV expression.

U2OS cells were transfected with empty vector control, MCV sT_{WT} and MCV sT_{LSDm} plasmids.

Cells were fixed at 48 h post transfection and endogenous collagen IV levels were measured by

indirect immunofluorescence using a specific antibody. MCV sT expression was detected with

2T2 antigen antibody. Nuclear counterstain (DAPI-Blue), MCV sT_{WT} and MCV sT_{LSDm} (Green),

and collagen IV(Red). (**B**) sT regulates collagen IV expression through the LSD. Mean

Florescence intensity of collagen IV in wild type (Bi) and LSD mutant sT-expressing cells (Bii)

was analyzed using Fiji Image J software. The calculated values were plotted for regression

analysis using Prism software. (**C**) sT induces cell invasion through the LSD. (Ci) Collagen

invasion assay. Serum starved sT-expressing U2OS cell were seeded on the precoated collagen

inserts and incubated for 72 h, then labeled with a cell staining solution for 20 min. Upon

extraction of the cell staining solution, absorbance at OD560 was measured. Data analyzed using

three replicates per experiment; the experiments were performed two times. The results were

reproducible and differences between means (*p* value) were analyzed using a t-test with

GraphPad Prism software. (Cii) Expression of sT. Protein expression levels of wild type and

mutant MCV sT were detected by immunoblot analysis to validate successful transfection.

Quantitative infrared fluorescence immunoblotting was performed using a 2T2 antibody for sT

antigens and α-Tubulin as an equal loading control.

FIGURE 6

MCV sT LSD induces Snail expression. (A) MCV sT activates the transcription of Snail in an

LSD-dependent manner. U2OS cells were transfected with empty vector control, sT_{WT} and

sT_{LSDm} expressing plasmids. Cellular RNA was extracted at 48 h post transfection and transcript

levels were analyzed using the comparative $\Delta\Delta Ct$ method (n = 3). **(B)** Snail protein expression is

induced by MCV sT through the LSD. Immunoblot analysis was performed on the cellular

lysates and analyzed using Snail specific antibody. α-Tubulin was used as a measure of equal

loading and the 2T2 antibody was used to confirm MCV sT wild type and mutant expression.

FIG 1

A

B

FIG 2

A

Bi

Bii

C

FIG 3

FIG 4

A

	DMSO	9-I	9-II
Vector			
sT$_{WT}$			
sT$_{LSDm}$			

Time(h)   0   24   0   24   0   24

B

Fold difference -Distance Travelled (Normalized to Untreated vector)

Vector
sT$_{WT}$
sT$_{LSDm}$

****
****
*
*
ns
**
ns

DMSO 9-I 9-II   DMSO 9-I 9-II   DMSO 9-I 9-II

C

Fold Change (Migrated cells)

DMSO
9-I
9-II

****
****
ns
****
****
ns

MKL-1   MS-1

FIG 5

FIG 6

A

ns

****   ****

mRNA fold difference normalized to GAPDH (SNAIL)

Vector
sT$_{WT}$
sT$_{LSDm}$

U2OS

B

Vector   sT$_{WT}$ sT$_{LSDm}$

Snail

MCV sT

α–Tubulin