1   **Title:** Strong effect of *Penicillium roqueforti* populations on volatile and metabolic
2   compounds responsible for aromas, flavor and texture in blue cheeses
3
4   **Running title:** Impact of mold population on blue cheeses
5

6   **Authors:**
7   Thibault CARON[1,4], Mélanie LE PIVER[4], Anne-Claire PÉRON[3], Pascale LIEBEN[3], René
8   LAVIGNE[2], Sammy BRUNEL[4], Daniel ROUEYRE[4], Michel PLACE[4], Pascal
9   BONNARME[3], Tatiana GIRAUD[1]*, Antoine BRANCA[1]*, Sophie LANDAUD[3]*,
10   Christophe CHASSARD[2]*
11
1: Ecologie Systematique Evolution, Université Paris Saclay, CNRS, AgroParisTech, 91400
Orsay, France
2: Université Clermont Auvergne, INRAE, Vetagro Sup, UMRF, 20 Côte de Reyne, 15000
Aurillac, France
3: Université Paris-Saclay, INRAE, AgroParisTech, UMR SayFood, 78850 Thiverval-
Grignon, France
4: Laboratoire Interprofessionnel de Production – SAS L.I.P., 34 rue de Salers, 15 000
Aurillac, France
20
21   *joint senior authors
22
23   **Contact information:** Thibault Caron thibaultcaron864@gmail.com

24
25   **Keywords:** Roquefort cheese, fungi, *Penicillium*, domestication, volatile compounds
26

Acknowledgments: We thank Béatrice DESSERRE, Céline DELBES and Cécile CALLON
for advice and technical assistance in microbiology, Sébastien THEIL for technical support
for metabarcoding analyses, Patricia LE THUAUT, Manon SURIN and Brigitte POLLET for
technical support for metabolomic analyses, Sara PARISOT for milk delivery and quality,
Pierre CONCHON for technical support for image analysis, LIAL-MC for the various
reference measurements in physicochemistry, Christophe LACROIX and Alfonso DIE for the
determination of short-chain fatty acids in fermentation supernatants.
This study was funded by the LIP SAS, ANRT (*Association Nationale Recherche
Technologie*), the ERC Genomefun 309403 starting grant, the BLUE ERC Proof-of-Concept
grant, the Fondation Louis D grant (French Academy of Sciences) and the ANR-19-CE20-
0002-02 Fungadapt ANR grant.

Abstract

Studies of food microorganism domestication can provide important insight into adaptation mechanisms and lead to commercial applications. *Penicillium roqueforti* is a fungus with four genetically differentiated populations, two of which were independently domesticated for blue cheese-making, with the other two populations thriving in other environments. Most blue cheeses are made with strains from a single *P. roqueforti* population, whereas Roquefort cheeses are inoculated with strains from a second population. We made blue cheeses in accordance with the production specifications for Roquefort-type cheeses, inoculating each cheese with a single *P. roqueforti* strain, using a total of three strains from each of the four populations. We investigated differences between the cheeses made with the strains from the four *P. roqueforti* populations, in terms of the induced flora, the proportion of blue color, water activity and the identity and abundance of aqueous and organic metabolites as proxies for proteolysis and lipolysis as well as volatile compounds responsible for flavor and aroma. We found that the population-of-origin of the *P. roqueforti* strains used for inoculation had a minor impact on bacterial diversity and no effect on the abundance of the main microorganism. The cheeses produced with *P. roqueforti* strains from cheese populations had a higher percentage of blue area and a higher abundance of the volatile compounds typical of blue cheeses, such as methyl ketones and secondary alcohols. In particular, the Roquefort strains produced higher amounts of these aromatic compounds, partly due to more efficient proteolysis and lipolysis. The Roquefort strains also led to cheeses with a lower water availability, an important feature for preventing spoilage in blue cheeses, which is subject to controls for the sale of Roquefort cheese. The typical appearance and flavors of blue cheeses thus result from human selection on *P. roqueforti,* leading to the acquisition of specific features by the two cheese populations. These findings have important implications for our understanding of adaptation and domestication, and for cheese improvement.

Domestication is an evolutionary process that has been studied by many biologists since

Darwin. Indeed, domestication is an excellent model for understanding adaptation, because it

results from strong and recent selection on traits that are often known and of interest to

humans (Larson *et al.*, 2014). Furthermore, studies of domestication frequently have

important implications for the improvement of cultivated organisms. Domesticated fungi

have been less studied than crops in this respect, despite being excellent models in this field

(Giraud *et al.*, 2017; Gladieux *et al.*, 2014). Most fungi can be cultured in Petri dishes, can

remain alive for decades when stored in freezers and are propagated asexually. All these

features facilitate experimentation. Fungal metabolism results in the production of various

compounds of interest, including fuel alcohols, enzymes and antibiotics (Bigelis, 2001). The

most ancient and frequent human use of fungi is for fermentation, to preserve and mature

food. For example, the yeast *Saccharomyces cerevisiae* is used in the production of bread,

wine and beer, and the filamentous fungus *Aspergillus oryzae* is used for the production of

soy sauce and sake (Dupont *et al.*, 2017). These models have provided important insight into

mechanisms of adaptation and domestication (Almeida *et al.*, 2014; Baker *et al.*, 2015;

Gallone *et al.*, 2016; Gibbons *et al.*, 2012; Gonçalves *et al.*, 2016; Libkind *et al.*, 2011; Sicard

et al., 2011).

The *Penicillium* genus contains more than 300 species, several of which are used by humans.

For example, penicillin was discovered in *P. rubens,* and two other species, *P. nalgiovense*

and *P. salami,* are used in the production of dry-cured meat (Fleming, 1929; Ludemann *et al.*,

2010, Perrone *et al.*, 2015). For centuries, *Penicillium roqueforti* (Thom) has been used in the

maturation of all blue cheeses worldwide (Coton *et al.*, 2020a; Labbe *et al.*, 2004, 2009;

Vabre, 2015). This fungus generates cheeses with a blue-veined appearance, by producing

melanized conidia within cavities in the cheese in which oxygen is readily available (Moreau,

1980). *Penicillium roqueforti* is also found in non-cheese environments (Pitt *et al.*, 2009;

Ropars *et al.*, 2012), and four genetically differentiated clusters of individuals (i.e.,

populations) have been identified in *P. roqueforti*. Two populations are used for

cheesemaking, whereas the other two populations thrive in silage, lumber or spoiled food

(Dumas *et al.*, 2020; Gillot *et al.*, 2015; Ropars *et al.*, 2014). Genomic and experimental

approaches have provided compelling evidence for the domestication of cheese *P. roqueforti*

populations (Cheeseman *et al.*, 2014; Dumas *et al.*, 2020; Gillot *et al.*, 2015, 2017; Ropars *et*

al., 2015, 2016b, 2017) which have been reviewed (Coton *et al.*, 2020b; Ropars *et al.*,

2020a). Indeed, the populations of *P. roqueforti* used to make blue cheeses display the

characteristic features of domesticated organisms: genetic and phenotypic differences relative

to non-cheese populations, with, in particular, traits beneficial for cheese production, such as

faster growth on cheese medium (Dumas *et al.*, 2020; Gillot *et al.*, 2015; Ropars *et al.*, 2014,

2016), but also lower fertility and lower fitness in nutrient-poor conditions (Ropars *et al.*,

2015, 2016). Both cheese populations have lower levels of genetic diversity than the two non-

cheese populations, indicating an occurrence of bottlenecks (Dumas *et al.*, 2020), which

typically occur during domestication. The two cheese populations are genetically and

phenotypically differentiated from each other, suggesting that they result from independent

domestication events (Dumas *et al.*, 2020). One of the cheese populations, the non-Roquefort

population, is a clonal lineage with a very low level of genetic diversity used to produce most

types of blue cheeses worldwide. The second cheese population, the Roquefort population, is

genetically more diverse and contains all the strains used to produce blue cheeses from the

emblematic Roquefort protected designation of origin (PDO) (Dumas *et al.*, 2020). *In vitro*

tests have shown that the non-Roquefort population displays faster tributyrin degradation (*i.e.*

a certain type of lipolysis) and higher salt tolerance, faster *in vitro* growth on cheese medium,

better exclusion of competitors than the Roquefort population (Dumas *et al.*, 2020; Ropars *et*

al., 2014, 2015), and an absence of toxic mycophenolic acid production (Gillot *et al.*, 2017).

The specific features of the Roquefort population may result from the constraints of the PDO,

requiring the use of local strains and at least 90 days of maturation, and preventing the use of

strains from the non-Roquefort population better suited to modern modes of production

(Dumas *et al.*, 2020). Genomic footprints of domestication (i.e., of adaptive genetic changes)

have been identified in the two *P. roqueforti* populations used for cheesemaking. Indeed, it

has been suggested that horizontally transferred genes found only in the non-Roquefort

population are involved in the production of an antifungal peptide and in lactose catabolism

(Cheeseman *et al.*, 2014; Ropars *et al.*, 2014, 2015). The effects of positive selection have

been detected in genes with predicted functions in flavor compound production, in each of

the cheese populations (Dumas *et al.*, 2020).

Thus, the four *P. roqueforti* populations probably harbor multiple specific traits, leading to

the generation of cheeses with different physicochemical properties and flavors, although this

has yet to be tested. Assessments of the effect of the population-of-origin of the *P. roqueforti*

strain used on the features of the cheese will i) provide important fundamental knowledge

about the trait under selection for cheesemaking and adaptation to the cheese environment, ii)

provide a basis for the elucidation of other genomic changes and iii) be crucial to

improvements in strain use and improvement. *Penicillium roqueforti* is used as a secondary

starter for flavor production, mostly through proteolysis (*i.e.* casein catabolism) and lipolysis

during ripening (Moreau, 1980). The main characteristic feature of blue cheeses, and of

Roquefort PDO cheeses in particular, is their intense, spicy flavors (Kinsella *et al.*, 1976;

Rothe *et al.*, 1982). The specific volatile and metabolic compounds responsible for these

flavors are generated principally by lipolysis in blue cheeses (Cerning *et al.*, 1987; Collins *et*

al., 2003) and their intensity varies among *P. roqueforti* strains (Dumas *et al.*, 2020; Gillot *et*

al., 2017; Larsen *et al.*, 1999). The fatty acids released by lipolysis are the precursors of

aldehydes, alcohols, acids, lactones and methyl ketones, which provide the moldy aromas

typical of blue cheeses (Collins *et al.*, 2003). *Penicillium roqueforti* degrades most proteins,

but the efficiency of proteolysis differs between strains (Cerning *et al.*, 1987; Dumas *et al.*,

2020; Gillot *et al.*, 2017; Larsen *et al.*, 1998). The resulting peptides contribute to flavors,

and their degradation into amino acids further influences cheese aroma and the growth of

other microorganisms (McSweeney *et al.*, 2000; Williams *et al.*, 2004). *Penicillium*

roqueforti also contributes to lactate degradation, which is necessary for deacidification and

promotes the development of less acid-tolerant microorganisms (McSweeney *et al.*, 2017).

Through these effects, and by producing secondary metabolites with antimicrobial properties,

P. roqueforti may also affect the microbial composition of the cheese (Kopp *et al.*, 1979;

Vallone *et al.*, 2014). Another parameter potentially affected by *P. roqueforti* populations and

restricting the occurrence of spoiler microorganisms is the lack of free water, (i.e., low water

activity, Aw), which is subject to strong controls for the sale of Roquefort cheese and is

affected by the degree of proteolysis (Ardö *et al.*, 2017). The *P. roqueforti* population may,

thus, also have an indirect effect on the features of the cheese, through various effects on

beneficial or undesirable contaminants.

The differences between *P. roqueforti* populations have, to date, been studied only *in vitro* or

in very rudimentary cheese models. Here, our objective was to assess the effect of the *P.*

roqueforti population-of-origin of the strain used for inoculation on the features of blue

cheeses. We focused on several features considered important for cheese quality. Given the

evidence from previous studies that cheese *P. roqueforti* populations have been domesticated,

any differences between the cheeses produced with cheese and non-cheese populations,

and/or between the two cheese populations can probably be considered to reflect selection by

humans for the production of good cheeses, either on standing variation in the ancestral *P.*

roqueforti population or for *de novo* mutations. We therefore produced blue cheeses in

conditions very similar to those used in industrial Roquefort PDO production, using, in

particular, milk from the local "Lacaune" breed, with strains from the four *P. roqueforti*

populations. We compared several important cheese features between the four populations: i)

physicochemical features, relating to texture and biochemical composition, ii) cheese

microbiota composition and abundance, which may have effects on several cheese features,

iii) the blue area as a proportion of cheese slices, which is important for the blue-veined

appearance of the cheese and is dependent on the growth and sporulation of *P. roqueforti* in

cheese cavities, and iv) the types and abundance of the metabolic and volatile compounds

produced, which influence flavor and aroma. We investigated the differences in these features

between the cheeses produced with strains from the four *P. roqueforti* populations (Roquefort

cheese, non-Roquefort cheese, silage and lumber/food spoiler populations). We also

investigated the possible differences between cheeses made with cheese and non-cheese

populations, and between the Roquefort and non-Roquefort cheese populations. Assessments

of the traits differing between cheese and non-cheese *P. roqueforti* populations, and between

the two cheese populations, and investigations of whether the cheese populations are more

suitable for cheese-making, are of fundamental importance for understanding the

domestication of cheese fungi, through the identification of traits subjected to selection.

Furthermore, the identification of differences between *P. roqueforti* populations of relevance

for cheese making has many applied consequences for the cheese industry, in terms of strain

choice for different kinds of blue cheeses, paving the way for the improvement of mold

strains by generating progenies from crosses of the two cheese populations, and for the

choice of traits for measurement and selection in offspring.

Materials and Methods

The Materials and Methods are described in more details in the Supplementary Methods.

Experimental design and fungal strains: We made cheeses with three different strains from

each of the four *P. roqueforti* populations (Figure 1, a strain is defined here as a haploid

individual obtained by monospore isolation), according to population assignments obtained in

a previous study based on molecular markers (Dumas *et al.*, 2020). For cheese production

(Figure 1B), we used a single strain per cheese for inoculation, using a total of three different

strains from each of the four *P. roqueforti* populations (Figure 1A). Due to the limited

production capacity of the experimental facility, we were unable to make all the cheeses at

the same time. We therefore split production into three assays, each including one strain from

each of the four populations (Figure 1A). For each strain in each assay, we created three

production replicates, with two cheeses per strain in each replicate, to ensure the production

of sufficient material for sampling. In total, we produced 72 cheeses (4 strains * 2 cheeses * 3

replicates * 3 assays, Figure 1A). The assays were performed sequentially from February to

April. The effect of the seasonal change in milk composition was therefore confounded with

the strain effect within the population, hereafter referred to as the "assay effect". The three

replicates within each assay were also set up at different times, a few days apart, and thus

with different batches of raw milk (Figure 1A). LCP (*Laboratoire de Cryptogamie de Paris*)

strains were obtained from the *Muséum d'Histoire Naturelle de Paris*, France, and BRFM

(Biological Resource Fungi Marseille') strains were obtained from the *Centre International*

de Ressources Microbiennes de Marseille, France.

Cheesemaking: The cheesemaking protocol was typical of the procedures used by the main

producers of Roquefort cheese and complied with the Roquefort PDO specifications, except

that the ripening process took place in artificial cellars in the INRA facilities at Aurillac and

with strains from different *P. roqueforti* populations. Briefly, we slowly mixed about 35 L of

milk in each vat, and heated it to 32.5°C. We then added 20 mg.L^{-1} mesophilic starter culture

containing *Leuconostoc* spp., *Lactococcus lactis* subsp. *cremoris*, *L. lactis* subsp. *lactis*, and

L. lactis subsp. *lactis* biovar *diacetylactis* (FD-DVS CHN-11, CHR HANSEN, Saint-

Germain-lès-Arpajon, France), 25 mg.L^{-1} gas-producing *Leuconostoc mesenteroïdes* subsp.

mesenteroïdes (CHOOZIT LM 57 LYO 20 DCU, DANISCO, Paris, France) and about 3.6 x

10^6 CFU.L^{-1} *P. roqueforti* spores (SAS LIP, Aurillac, France). The mixture was homogenized

for five minutes, and we then added 0.25 ml of active chymosin at a concentration of 520

mg.L^{-1} (Laboratoires Humeau, La Chapelle-sur-Erdre, France). The resulting curd was cut

into 1.5 cm^3 cubes, left to rest for five minutes, and allowed to ferment for 50 minutes. The

curd was drained and placed in perforated cylinders of 20x10 cm at 26°C/85% humidity. The

cheeses were turned five times after molding. On the second day, the cheeses were unmolded.

On the third day, the cheeses were rubbed with sterilized coarse salt and transferred to

ripening cellars at 11°C / 95% humidity. On the fifth day, the cheeses were resalted in the

same way. On the seventh day, the cheeses were pricked. They were left to ripen until day 20,

when they were wrapped in sterile aluminum foil and allowed to mature at -2°C until day

180.

Cheese samples were collected on days 0, 9, 20, 90 and 180. These time points are referred to

hereafter as "stages". Day 0 corresponded to raw milk for microbiology analysis and sowed

curd for metabolic and volatile compound analyses. Days 9 and 20 corresponded to half-way

through and the end of the ripening period, respectively. Days 90 and 180 (i.e. 3 and 6

months) corresponded to the minimum maturation times for the Roquefort PDO and a typical

maturation period for sale, respectively.

Microbial and metabarcoding analyses: We estimated the concentrations of various

microbial communities in the initial unpasteurized milk and at various stages of cheese

maturation, to determine whether the *P. roqueforti* population affected the cheese microbiota

(for more information, see the Supplementary Methods). We counted the total number of

aerobic mesophilic bacteria on plate count agar (PCA; Nelson, 1940). Mesophilic lactic acid

bacteria were counted on Man Rogosa Sharpe agar (MRS; De Man *et al.*, 1960) and

thermophilic lactic acid bacteria were counted on M17 agar (Terzaghi, 1975). Dextran-

positive *Leuconostoc* spp. were counted on Mayeux, Sandine and Elliker medium (MSE;

Mayeux *et al.*, 1962). Molds and yeasts were counted on oxytetracycline gelose agar medium

(OGA; Mossel *et al.*, 1970). Gram-positive catalase-positive bacteria were counted on

cheese-ripening bacterial medium (CRBM; Denis *et al.*, 2001). Enterobacteria were counted

on violet red bile glucose (VRBG; Mossel *et al.*, 1978). We further investigated the possible

effects of the *P. roqueforti* population on the identity and relative abundance of other

microorganisms in cheeses, by performing metabarcoding analyses on our experimental

cheeses at 9 and 20 days (during the ripening period), by extracting DNA, and amplifying

and sequencing the V3-V4 region of the 16S rDNA gene. Amplicons were sequenced with

Illumina Miseq technology. The mean read depth of the 71 samples (36 at 9 and 20 days of

ripening minus one lost) was 28539 reads. Amplicon data from high-throughput sequencing

were analyzed with Find Rapidly OTUs in Galaxy Solution (FROGS) v3.0 (Escudié *et al.*,

2018). For each OTU, taxonomic assignment was determined with the Silva-132

(https://www.arb-silva.de/) and 16S rDNA RefSeq databases

(https://blast.ncbi.nlm.nih.gov/Blast.cgi). Four widely used diversity parameters were

calculated from OTU compositions: two for alpha diversity (Shannon index, Simpson index)

and the Bray-Curtis dissimilarity for beta diversity.

Blue area: We estimated the percentage area of the cheese that was blue, on fresh inner

cheese slices (Figure 3B). This percentage is dependent on the formation of cavities within

the cheese, the growth of *P. roqueforti* within the cavities and its sporulation, as the blue

color is due to the melanin present in *P. roqueforti* spores. We cut all the 20-, 90- and 180-

day cheeses in half and took three photographs of each fresh slice with a Canon PowerShot

SX410 IS (JPEG format, 5152 x 3864 pixels, 100 ISO, without flash). We analyzed these

images with imageJ 1.52n (Schneider *et al.*, 2012; Figure S2): (i) the brightness and contrast

of the raw images were standardized with a stack contrast adjustment plugin, using a

reference image (Figure S2A,B) (ii) rectangular selection was used to crop the images

manually to ensure that they contained only the cheese slices (Figure S2C), (iii) the red

channel was segmented with a grayscale threshold of 102 to distinguish colonized cavities

from the inner white parts of the cheese and empty cavities (Figure S2D), (iv) the number of

dark pixels in the cavities was counted and divided by the total number of pixels for the entire

cheese slice, to determine the percentage blue area of the slice. Within the dark patches in the

cheese cavities colonized by the fungus, all pixels were considered dark (script at

https://gitlab.com/snippets/1945218).

Physicochemistry: We performed standard physicochemical measurements on the cheeses.

We measured dry matter content, fat content as a proportion of dry matter, the moisture

content of the defatted cheese, total, soluble and non-protein nitrogen contents, chloride and

salt content, water activity and pH at various stages of maturation, according to reference

methods (for more information, see the Supplementary Methods). We measured glucose,

lactose, lactate, acetate and butyrate concentrations in the cheeses on days 9 and 20, by high-

performance liquid chromatography (HPLC, for more information, see the Supplementary

Methods).

Metabolic and volatile compounds: We investigated possible differences in proteolytic and

lipolytic activities between the four populations, by UHPLC-MS after two extraction

procedures (in water and an organic solvent). We analyzed the abundance of free fatty acids

and residual glycerides in 90-day cheeses, by coupling a global extraction (accelerated

solvent extraction with hexane-isobutanol) with UHPLC-MS analysis in the positive

(triglycerides) and negative (fatty acids) ionization modes (for more information, see the

Supplementary Methods). We investigated the identity and abundance of volatile flavor and

aroma compounds, using a dynamic headspace system (DHS) with a Gerstel MPS

autosampler (Mülheim an der Ruhr, Germany) and gas chromatography-mass spectrometry

analysis with an Agilent 7890B GC system coupled to an Agilent 5977B quadrupole mass

spectrometer (Santa Clara, United States). Statistical analyses were performed with R

software (http://www.r-project.org/). Further details about the materials and methods are

provided in the Supplementary Methods.

Results

***Penicillium roqueforti* population-of-origin influences bacterial diversity slightly, but**

has no effect on the abundance of the main microorganisms in the cheese. We

investigated whether the population-of-origin of the *P. roqueforti* strains used for inoculation

affected the composition of the cheese microbiota, by estimating the densities of key

microbial communities through colony counts (CFU/g) on various specific culture media and

a metabarcoding approach based on 16S sequencing targeting the bacteria in cheeses at

several stages of maturation. Based on microbial counts on plate, we found no significant

effect of *P. roqueforti* population on the abundance of any of the counted microorganisms,

including molds (i.e. *P. roqueforti)*, at any sampled stage of maturation (Supplementary

Figure 1; Supplementary Table 1A).

The metabarcoding approach targeting bacteria identified mostly sequences from the

Lactococcus and *Leuconostoc* spp. starters (about 58 and 40%, respectively, Figure 2A,

Supplementary Table 1B), which are responsible for acidification and cavity formation,

respectively, in the cheese. The remaining sequences corresponded to 12 bacterial genera

frequently found in raw milk cheeses, such as *Lactobacillus*, *Staphylococcus* and

Arthrobacter (<1%, Supplementary Table 1B). However, the large predominance of starters

made it impossible to obtain sufficient data for other bacteria to assess differences in the

abundance of particular bacteria between cheeses made with strains from the four *P.*

roqueforti populations (Supplementary Table 1C). We also targeted eukaryotes, based on ITS

sequences, in a few samples. Most sequences were assigned to *P. roqueforti* spp. (89%) and,

to a lesser extent, *Candida* spp. (10%, data not shown), so no further analyses were

performed on the remaining low-abundance sequences. We estimated three OTU (operational

taxonomic unit) diversity parameters based on bacterial barcode sequence abundances, to

measure OTU richness and/or evenness. Bray-Curtis dissimilarity showed that cheeses made

with strains from the same *P. roqueforti* population were no more similar than those made

with strains from different *P. roqueforti* populations. However, we found a significant effect

of *P. roqueforti* population, in addition to a stage effect, on the Shannon and Simpson

diversity indices (Figures 2B and 2C). Cheeses made with strains from the cheese *P.*

roqueforti populations tended to have a higher bacterial OTU diversity, particularly at nine

days of maturation and for the Roquefort population (Figures 2B and 2C), although the post-

hoc analyses were not powerful enough to detect significant pairwise differences

(Supplementary Table 1C). There may also be undetected differences at species level or for

low-abundance microorganisms that might nevertheless have substantial effects. However,

even if this were the case, it would constitute an indirect effect of the *P. roqueforti*

population, as this was the only difference during our cheesemaking process.

Higher proportion of blue area in cheeses produced with cheese *P. roqueforti* populations. The percentage blue area in cheese cavities was determined on freshly cut cheese slices (Figure 3B). The blue veins result from the formation of cavities in the cheese, and the growth and sporulation of *P. roqueforti* in these cavities. The percentage blue area was significantly higher in cheeses produced with cheese population strains than in those produced with non-cheese population strains (Figure 3A and 3B; Supplementary Table 1D). We also found a significant decrease in the percentage blue area from 20 to 180 days of maturation, for all populations except the Roquefort population, for which the percentage blue area remained relatively constant (Figure 3A; Supplementary Table 1D).

More efficient proteolysis and lipolysis by the Roquefort *P. roqueforti* population. We investigated the proteolysis and lipolysis efficiencies of the four populations in cheeses after 90 days of maturation. Both targeted and non-targeted chromatographic analyses showed that proteolytic efficiency was highest in the Roquefort *P. roqueforti* population. We performed the targeted analysis with standards for the principal 23 amino acids (Supplementary Table 2A). We found that eight amino acids discriminated significantly between cheeses made with the different *P. roqueforti* populations (Supplementary Table 1E, mainly ornithine, leucine, and alanine), 15 discriminated between the cheese and non-cheese populations (mainly spermidine, isoleucine, methionine, glutamic acid, citrulline, serine and leucine) and 14 distinguished between the Roquefort and non-Roquefort populations (Supplementary Figure 3A, mainly valine, leucine, isoleucine, serine and threonine). The cheeses made with strains from cheese populations, and from the Roquefort population, in particular, had a higher total amino-acid concentration (Supplementary Tables 1E and 2B).

We also assessed proteolytic activity in a non-targeted analysis (fingerprint approach) on

whole chromatograms (8,364 signals), which provided much more powerful discrimination

between metabolites. Each metabolite generates a signal specific to its mass-to-charge (m/z)

ratio at a given retention time. We obtained the largest number of aqueous signals, indicating

the most efficient proteolysis, for cheeses inoculated with strains from the Roquefort

population, followed by the lumber and non-Roquefort cheese populations, which were not

significantly different from each other, and proteolysis was least efficient for the silage

population (Figure 4; Supplementary Table 1F).

Lipolysis was also more efficient for the Roquefort population than for the other populations.

We investigated whether the *P. roqueforti* population influenced the abundance of free fatty

acids and residual glycerides, as a proxy for the efficiency of lipolysis, in 90-day cheeses,

with targeted and non-targeted chromatographic analyses in the positive and negative

ionization modes. We specifically targeted glycerides and free fatty acids. In the targeted

analysis, we identified seven free fatty acids and 20 triglycerides, and found that three free

fatty acids (stearic, oleic and linoleic acids) were significantly more concentrated in cheeses

made with Roquefort strains than in those made with strains from non-Roquefort populations

(Supplementary Table 1G). In the non-targeted analysis, we obtained 3,094 signals and

observed a higher abundance for organic signals specific to free fatty acids, indicating the

most efficient lipolysis, in cheeses made with strains from the Roquefort population, followed

by the lumber and non-Roquefort cheese populations, which were very similar to each other,

with lipolytic efficiency lowest for cheeses made with strains from the silage population

(Figure 5; Supplementary Table 1H). For residual glycerides, we obtained 8,472 signals, with

no significant difference between the populations (Supplementary Figure 4; Supplementary

Table 1I).

We performed the most relevant standard physicochemical measurements on raw milk and on

the cheeses at 9, 20, 90 and 180 days of maturation. As expected, we observed a maturation

stage effect for 11 of the 16 physicochemical parameters (Supplementary Table 1J). Non-

protein nitrogenous content was significantly higher in cheeses inoculated with strains from

cheese *P. roqueforti* populations than in cheeses inoculated with strains from the other

populations, consistent with the greater efficiency of proteolysis associated with these strains

(Supplementary Figure 5A). Cheese water activity differed significantly between the cheeses

made with strains from the four *P. roqueforti* populations (Supplementary figure 5B): it was

significantly lower for the Roquefort cheese population than for the non-Roquefort cheese

and silage populations (see statistics in the Supplementary Table 1J).

Strong influence of *P. roqueforti* population on volatile compound production. We

investigated the effect of *P. roqueforti* population on cheese volatile compounds in 90-day

cheeses. We focused on the GC-MS data for the 40 principal volatile compounds considered

to be most important for the aromatic quality of blue cheeses (Rothe *et al.*, 1982): 11 acids,

12 ketones, 10 esters, six alcohols and one aldehyde (Supplementary Table 3). We found that

P. roqueforti population strongly influenced the relative abundance of the compounds from

these aromatic families in the cheeses (Supplementary Table 1K; Figures 6 and 7, and see

below). In fact, the odors of the cheeses differed considerably (pers. obs.): the cheeses made

with strains from the cheese *P. roqueforti* populations smelled as good as typical ripened blue

cheeses, whereas those made with strains from non-cheese *P. roqueforti* populations had

unpleasant odors, similar to those of a wet mop (Supplementary Figure 6; personal

observation).

The abundance of acids, methyl ketones and secondary alcohols resulting from proteolysis

and lipolysis, and contributing to the typical flavor of blue cheese, was higher in cheeses

produced with strains from cheese populations than in those produced with strains from non-

cheese populations (Figure 6). These compounds were present at a particularly high

abundance in cheeses made with strains from the Roquefort population. Four of the 40

compounds analyzed were proteolysis by-products (primary alcohols: 3-methyl-butanal, 3-

methyl-butanol and isopropyl-alcohol, hereafter referred to as alcohols I, and 3-methyl-

butanoic acid, hereafter referred to as acid I; Supplementary Table 3). The abundance of

alcohols I was significantly higher in cheeses made with strains from cheese *P. roqueforti*

populations than in those made with strains from non-cheese populations, and the highest

values were obtained for the Roquefort population (mainly 3-methyl-butanol; Figure 6;

Supplementary Table 1K). Acid I (i.e. 3-methyl-butanoic acid) was also present in higher

abundance in cheeses made with strains from the Roquefort population than in other cheeses

(Figure 6). Two acids, by-products of glycolysis (acetic and propionic acid, hereafter referred

to as acids II), were present at higher abundance in cheeses made with strains from the

Roquefort and lumber/food spoiler *P. roqueforti* populations than in other cheeses (Figure 6;

Supplementary Tables 1K and 3). The other 35 aromatic compounds (i.e. acids from beta-

oxidation, hereafter referred to as acids III, ketones, secondary alcohols hereafter referred to

as alcohols II, and esters) were almost all direct or indirect by-products of lipolysis

(Supplementary Table 3). The abundance of acids III was higher in cheeses made with strains

from the Roquefort and lumber/food spoiler populations than in cheeses made with strains

from the non-Roquefort cheese population (mainly butanoic, pentanoic, hexanoic and

octanoic acid; Figure 6; Supplementary Table 1K). The levels of these compounds were

lowest in cheeses made with strains from the silage population. Esters and methyl ketones

(especially 2-pentanone and 2-heptanone) were present at higher abundance in cheeses made

with strains from cheese *P. roqueforti* populations (Supplementary Table 1K). Cheeses made

with strains from the Roquefort population contained the highest abundance of methyl

ketones, and these compounds were barely detectable in cheeses made from silage population

strains (mainly 2-heptanone, 2-pentanone and 2-nonanone; Figure 7A). The abundance of

alcohols II, particularly 2-heptanol, was also much higher in cheeses made with Roquefort

population strains than in other cheeses (Supplementary Table 1K; Figure 7B).

Discussion

Cheese *P. roqueforti* populations have been selected to produce better blue cheeses.

Measurements of multiple features of blue cheeses made under conditions resembling those

typically used in commercial Roquefort production revealed a strong influence of the

differentiated *P. roqueforti* populations on several aspects of cheese quality. The cheese

populations appeared the best adapted to cheesemaking, in terms of both the appearance and

volatile compound content of the resulting cheese. The differences between the four *P.*

roqueforti populations and the more appealing cheeses produced with strains from the cheese

populations suggest that humans have exerted selection for the production of better cheeses,

and this corresponds to domestication. Indeed, we found that cheese *P. roqueforti* strains

produced a higher percentage blue area on cheese slices, an important visual aspect of blue

cheeses. We also found that proteolysis and lipolysis were more efficient in cheeses made

with Roquefort population strains than in cheeses made with strains from the other *P.*

roqueforti populations, resulting in a higher abundance of desirable volatile compounds,

including alcohols and associated acids. Cheese water activity was lower in cheeses made

with strains from the Roquefort population, probably due to more efficient proteolysis (Ardö

et al., 2017). We found no significant difference in the identities and abundances of

microorganisms between the cheeses made with strains from the four *P. roqueforti*

populations. Some minor differences in bacterial alpha diversity were observed, however, and

the differences in all other measurements than diversity between cheeses probably reflected a

direct effect of the specific features of the *P. roqueforti* population, although minor indirect

effects involving the induction of more diverse bacterial communities by cheese *P. roqueforti*

strains may also have occurred. Overall, our findings strongly support the view that cheese *P.*

roqueforti populations have been selected by humans for better appearance and aroma. This

selection may have involved the choice of the most beneficial strains for making good

cheeses from standing variation, and/or the selection of *de novo* genetic changes. Previous

studies found footprints of genomic changes in cheese populations in the form of beneficial

horizontal gene transfers and positive selection (Dumas *et al.*, 2020; Ropars *et al.*, 2015).

Previous studies reported differences between *P. roqueforti* populations, in terms of growth,

lipolysis and proteolysis, but on synthetic media (Dumas *et al.*, 2020; Ropars *et al.*, 2015).

Here, using experimental cheeses made in commercial cheese production conditions, we

reveal important features specific to cheese *P. roqueforti* populations, and to the Roquefort

and non-Roquefort cheese populations. These findings are important in the context of

domestication, for understanding rapid adaptation and diversification, and future studies

based on quantitative trait mapping may be able to identify further genomic changes

responsible for the specific features of the populations, according to the contrasting

phenotypes revealed here. Progenies can indeed be obtained from crosses between strains

from different populations of *P. roqueforti* (Ropars *et al.*, 2015), and this could facilitate

strain improvement through recombination between the different populations. Our results are,

therefore, also important for improving blue cheese production.

The four *P. roqueforti* populations induce similar microbiotas, but water availability is
lower with cheese population strains, restricting the occurrence of spoiler
microorganisms. Based on microbiological counts, we found no significant differences in
abundance for any of the species monitored between cheeses made with strains from the four
populations of *P. roqueforti*. In particular, we found no significant difference in the
abundance of molds on Petri dishes. However, microbiological counts are known to provide
poor estimates of fungal biomass, especially for mycelium growth (Schnurer, 1993).

The abundance and identity of the microorganisms studied were globally similar to those in
four commercial Roquefort cheeses (personal information from C. Callon; Devoyod *et al.*,
1968) and closely related blue cheeses (Diezhandino *et al.*, 2015). The metabarcoding
approach suggested that the different *P. roqueforti* populations induced bacterial
communities of different levels of diversity. The cheese populations, and the Roquefort
population in particular, were associated with the highest level of alpha diversity. The
differences in cheese bacterial diversity, although minor, suggest that the differences between
cheeses made with strains from the four *P. roqueforti* populations may be due not only to a
direct effect of *P. roqueforti* population, but also to an indirect effect mediated by the
induction of bacterial communities of different diversities. The large predominance of
bacterial starters made it impossible to collect sufficient data for an assessment of the
differences in relative abundance between subdominant bacterial species on the basis of
metabarcoding. We also found a significant difference in water activity between cheeses
made with strains from different *P. roqueforti* populations, the lowest value obtained being
that for the Roquefort population. This may also reflect human selection, as low water
activity restricts the occurrence of spoiler microorganisms. This characteristic is, therefore,
subject to tight control in Roquefort cheeses for sale, particularly those for export.

Cheese *P. roqueforti* populations produce bluer cheeses. We found significantly higher percentage blue areas in cheese slices from cheeses made with cheese *P. roqueforti* strains than in those made from non-cheese strains, potentially reflecting greater *P. roqueforti* growth in cheese and/or a higher sporulation efficiency in cavities. The percentage blue area in cheese slices also depends on the formation of cavities in the cheese, as *P. roqueforti* can only sporulate in cavities in which oxygen is available. The cavities are mostly generated by the gas-producing bacterium *Leuconostoc mesenteroides*, the abundance of which did not differ between the cheeses made with strains from different *P. roqueforti* populations, suggesting a direct effect of *P. roqueforti* populations on the blueness of cheese slices. The significantly higher percentage blue area in slices of cheeses made with cheese *P. roqueforti* strains than in those made with non-cheese strains therefore probably reflects better cheese and cavity colonization and sporulation, probably due to selection on the basis of appearance. The percentage blue area decreased by the end of maturation, perhaps due to the death of the fungus. Only cheeses made with Roquefort strains retained a high percentage blue area at 90 days of maturation, again potentially reflecting selection in pre-industrial times, when Roquefort cheeses had to be stored for several months at cave temperature before sale. The minimum maturation time for Roquefort PDO remains 90 days, which is longer than for other blue cheeses. These findings contrast with a previous study showing that a non-Roquefort population colonized the cavities of model cheeses better than other populations (Dumas *et al.*, 2020); this discrepancy may reflect differences between studies in terms of the measurements used (total percentage blue area versus percentage blue area within cavities), the type of milk (ewe versus goat) or the mode of cheesemaking (rudimentary models versus commercial-like cheeses). Our findings are consistent with the presence of horizontally

transferred genes in cheese populations with predicted functions in fungal development,

including sporulation and hyphal growth (Dumas *et al.*, 2020).

Proteolysis and lipolysis are more efficient in the Roquefort *P. roqueforti* population.

Based on chemical analyses and powerful chromatographic discrimination methods, we

showed that the abundance of amino acids and small peptides (i.e., residual products of

proteolysis) was highest in cheeses made with Roquefort *P. roqueforti* strains. Thus, these

strains had the highest capacity for proteolysis, which is an important process in

cheesemaking. Indeed, proteolysis contributes to the development of cheese texture, flavors

and aromas (Andersen *et al.*, 2010; Ardö, 2006, 2017; McSweeney, 1997; Roudot-Algaron,

1996). Previous measurements of proteolytic activity in synthetic media detected significant

differences between *P. roqueforti* populations, but not between the two cheese populations

(Dumas *et al.*, 2020). We show here that experimental cheeses made with strains from the

Roquefort population have a higher content of residual products of proteolysis, a sign of more

advanced ripening.

We also found that lipolysis was more efficient in the cheeses made with strains from the

Roquefort *P. roqueforti* population. By contrast, previous studies in synthetic media found

that lipolysis was most efficient in the non-Roquefort population (Dumas *et al.*, 2020). The

discrepancy between these studies demonstrates the need for measurements in real cheeses

for the reliable assessment of metabolic activities. Lipolytic activity is known to affect cheese

texture and the production of volatile compounds affecting pungency (Alonso *et al.*, 1987;

González De Llano *et al.*, 1990, 1992; Martín *et al.*, 2016; Thierry *et al.*, 2017; Woo *et al.*,

1984). The more efficient proteolysis and lipolysis in the Roquefort *P. roqueforti* population

should have a strong impact on cheese texture and flavor. It therefore probably results from

selection to obtain better cheeses, i.e. from a domestication process, as previously reported

for other fungi (Almeida *et al.*, 2014; Baker *et al.*, 2015; Gallone *et al.*, 2016; Gibbons *et al.*,

2012; Gonçalves *et al.*, 2016; Libkind *et al.*, 2011; Sicard *et al.*, 2011). Roquefort cheeses are

widely considered to be the blue cheeses with the strongest aromas and flavors. The less

efficient lipolysis and proteolysis in the non-Roquefort population may result from more

recent selection for milder cheeses.

Cheese *P. roqueforti* populations produce cheeses with more abundant and diverse

volatile compounds. We found major differences between the cheeses made with strains

from different *P. roqueforti* populations, in terms of the volatile compounds resulting from

lipolysis and, to a lesser extent, also from proteolysis. Only four of the aromatic compounds

detected in our cheeses (3-methyl-butanal, 3-methyl-butanol, isopropyl-alcohol and 3-methyl-

butanoic acid) were by-products of casein proteolysis (McSweeney *et al.*, 2000), and the

concentrations of these molecules were significantly higher in cheeses made with Roquefort

P. roqueforti strains, consistent with the higher proteolysis efficiency and amino-acid

precursor (i.e. valine, leucine and isoleucine) concentrations of these strains. These

compounds produce fruity (banana), cheesy and alcoholic notes, which were probably

important selection criteria during the domestication of the Roquefort *P. roqueforti*

population. For the products of metabolic pathways leading from amino acids to alcohols

(Ehrlich pathway with aldehyde reduction) or acids (aldehyde oxidation; Ganesan *et al.*,

2017), the higher concentration of alcohols than of acids observed for all populations is

consistent with the general micro-aerobic conditions of blue cheese cavities.

Most of the aromatic compounds identified were direct or indirect by-products of lipolysis,

consistent with the known key role of lipolysis in the generation of typical blue cheese aroma

(Cerning *et al.*, 1987; Collins *et al.*, 2003). The aromatic compounds resulting from lipolysis

belonged to four chemical families (acids, methyl ketones, secondary alcohols and esters).

Methyl ketones were the most diverse and abundant for cheese *P. roqueforti* populations,

particularly for the Roquefort population, in which abundance was highest for 2-pentanone

and 2-heptanone; 2-heptanone underlies the characteristic "blue cheese" sensory descriptor

(Anderson *et al.*, 1966; González De Llano *et al.*, 1990, 1992; Moio *et al.*, 2000). In *P.*

roqueforti, methyl ketones with odd numbers of carbons are mostly produced by fatty-acid

beta-oxidation, whereas those with even numbers of carbons may be produced by the beta-

oxidation or autoxidation of fatty acids (Spinnler, 2011). These compounds are produced by

the decarboxylation of hexanoic acid and octanoic acid, respectively, which were the most

abundant acids found in our cheeses. This reaction is considered to be a form of

detoxification, because methyl ketones are less toxic than acids (Kinderlerer, 1993; Spinnler,

2011). Interestingly, this pathway appeared to be more active in the cheese *P. roqueforti*

populations, as methyl ketone levels were lower in cheeses made with lumber (four-fold

difference) and silage (10-fold lower) strains than in cheeses made with cheese population

strains. Methyl ketone concentrations were not directly associated with the concentrations of

their precursors (acids), the highest concentrations being found in the lumber and Roquefort

populations. The biosynthesis pathway producing methyl ketones must, therefore, be more

efficient in cheese populations, particularly the non-Roquefort population. The cheese *P.*

roqueforti populations were probably selected for their higher acid detoxification capacity, as

this produces aromatic compounds with a very positive impact on flavor (Spinnler, 2011).

The concentrations of secondary alcohols (resulting from the reduction of methyl ketones)

were also higher in cheeses produced by cheese *P. roqueforti* strains, particularly those of the

Roquefort population, for which they were seven times higher than for the non-Roquefort

cheese population and 20 times higher than for the silage/lumber populations; 2-heptanol was

the major alcoholic compound produced. The reduction of 2-heptanone to 2-heptanol occurs

specifically in anaerobic conditions and is much stronger in the Roquefort population; aerobic

conditions were similar for all the populations. The Roquefort *P. roqueforti* population may

also have been selected for this feature, as secondary alcohols provide "fruity notes", which

are associated with better aromatic quality (Spinnler, 2011). Methyl ketones may be reduced

to alcohols by an alcohol dehydrogenase, as occurs when aldehyde is reduced to alcohol via

the Ehrlich pathway. Alcohol dehydrogenase genes may thus have been targets of selection in

the Roquefort *P. roqueforti* population, although they were not detected as evolving under

positive selection in a previous study (Dumas *et al.*, 2020).

We also found higher levels of esters in cheeses made with cheese *P. roqueforti* populations.

Esters are produced principally by the esterification of ethanol with acids generated by beta-

oxidation. *Leuconostoc* starters can produce ethanol, and ester synthesis has also been

described as a detoxification mechanism (Mason *et al.*, 2000). These results further indicate

that cheese *P. roqueforti* populations, particularly the Roquefort population, have been

selected for acid detoxification capacity, leading to a large variety of less toxic aromatic

compounds with strong aromas and flavors.

Overall, the aromas of cheeses made with cheese *P. roqueforti* strains had more appealing

aromas, and this was particularly true for cheeses made with Roquefort strains. These aroma

properties probably reflect selection by humans. The cheeses made with silage and lumber

populations had a mildly unpleasant smell, whereas those made with cheese strains smelled

like typical blue cheeses, with cheeses made with Roquefort strains having the strongest

smell. This may reflect previously reported horizontal gene transfers in cheese populations,

involving genes with predicted functions in lipolysis or amino-acid catabolism, and the

positive selection of genes involved in aroma production (Dumas *et al.*, 2020). We compared

P. roqueforti populations between cheeses made following commercial modes of production,

which represents a major advance relative to previous studies based on experimental models

or synthetic media (Dumas *et al.*, 2020; Gillot *et al.*, 2017). We used unpasteurized ewe's

milk, in accordance with the requirements for Roquefort PDO production, which also affects

cheese aromas. In future studies, it would be interesting to determine whether the use of

pasteurized or unpasteurized ewe's milk or cow's milk leads to similar specific features of the

Roquefort versus non-Roquefort cheese *P. roqueforti* populations, as there may have been

selection during domestication, leading to an adaptation of the Roquefort population for the

catabolism of unpasteurized ewe's milk.

Conclusion. We show that the *P. roqueforti* population has a strong impact on cheese

quality, appearance and aroma. The populations used for cheesemaking led to bluer cheeses,

with better aromas, probably due to domestication involving the selection of multiple fungal

traits by humans seeking to make the best possible cheeses. French cheese producers have

been inoculating cheeses with *P. roqueforti* spores from moldy rye bread since the end of the

19[th] century (Labbe *et al.*, 2004, 2009; Vabre, 2015). This process made it possible for them

to re-inoculate with the strains producing the best cheeses, thereby applying a strong selection

pressure. The two cheese populations displayed a number of specific features, with the

Roquefort population notably producing more intense and specific aromas and flavors. The

selection of different fungal varieties for different usages has also been reported in the

fermenting yeast *Saccharomyces cerevisiae* (Gallone *et al.*, 2016; Legras *et al.*, 2018).

Previous studies on *P. roqueforti* detected recurrent changes in amino acids and horizontal

gene transfers in cheese populations, both of which facilitated rapid adaptation (Dumas *et al.*,

2020; Ropars *et al.*, 2015). Our findings provide greater insight into *P. roqueforti*

domestication and pave the way for strain improvement through the targeting of relevant

traits. A protocol inducing sexual selection has been developed in *P. roqueforti* (Ropars *et al.*,

2014), making it possible to perform crosses between strains from the two cheese

populations, each of which harbors very little genetic diversity (Dumas *et al.*, 2020), to

generate variability and to identify strains with high levels of performance. The results of this

study will facilitate the choice of the parental strains for crossing and of the most important

phenotypes to be measured in the offspring. Parental strains with strongly contrasting

phenotypes for the traits important for cheesemaking that we found to be differentiated

between populations (such as volatile compound production, lipolysis and proteolysis) should

be used, to maximize variability in the progeny. Other traits may also be worth investigating

in the future, to understand the changes that have occurred during the domestication of the

two cheese *P. roqueforti* populations, particularly as concerns toxin production in cheeses.

Previous *in vitro* studies have shown a lack of production of the toxic mycophenolic acid by

the non-Roquefort population due to a deletion in one of the genes of the synthesis pathway

(Gillot *et al.*, 2017). In the *P. camemberti* fungus, domestication has also led to a lack of

production of a mycotoxin due to a deletion in a gene (Ropars *et al.*, 2020b).

References

Almeida, M., Hébert, A., Abraham, A.-L., Rasmussen, S., Monnet, C., Pons, N., Delbès-
Paus, C., Loux, V., Batto, J.-M., Leonard, P., Kennedy, S., Ehrlich, S., Pop, M., Montel, M.-
C., Irlinger, F. & Renault, P. (2014). Construction of a dairy microbial genome catalog opens
new perspectives for the metagenomic analysis of dairy fermented products. *BMC Genomics*,
15, 1101.

Alonso, L., Juarez, M., Ramos, M., & Martin□Alvarez, P. J. (1987). Effects of changes
during ripening and frozen storage on the physicochemical and sensory characteristics of
Cabrales cheeses. *International Journal of Food Science & Technology*, *22*(5), 525–534.

Andersen, L. T., Ardö, Y., & Bredie, W. L. P. (2010). Study of taste-active compounds in the
water-soluble extract of mature Cheddar cheese. *International Dairy Journal*, *20*(8), 528–
536.

Andersson, A. F., Lindberg, M., Jakobsson, H., Bäckhed, F., Nyrén, P., & Engstrand, L.
(2008). Comparative analysis of human gut microbiota by barcoded pyrosequencing. *PLoS*
ONE, *3*(7), e2836.

Anderson, D. F., & Day, E. A. (1966). Quantitation, evaluation, and effect of certain
microorganisms on flavor components of blue cheese. *Journal of Agricultural and Food*
Chemistry, *14*(3), 241–245.

Ardö, Y. (2006). Flavour formation by amino acid catabolism. *Biotechnology Advances*,
24(2), 238–242.

Ardö, Y., McSweeney, P. L. H., Magboul, A. A. A., Upadhyay, V. K., & Fox, P. F. (2017).
Biochemistry of cheese ripening: proteolysis. In P. L. H. McSweeney, P. F. Fox, P. D. Cotter,
& D. W. Everett (Eds.), *Cheese: Chemistry, Physics and Microbiology* (Fourth Edi, pp. 445–
482). Elsevier Ltd.

Auguie, B. (2017). gridExtra: miscellaneous functions for "Grid" graphics. Retrieved from
https://cran.r-project.org/web/packages/gridExtra/index.html

Baker, E. P., Wang, B., Bellora, N., Peris, D., Hulfachor, A. B., Koshalek, J. A., Libkind, D.
& Hittinger, C. T.(2015). The genome sequence of *Saccharomyces eubayanus* and the
domestication of lager-brewing yeasts. *Molecular Biology and Evolution*, *32*(11), 2818–2831.

Bates, D., Mächloer, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects
models using lme4. *Journal of Statistical Software*, *67*(1), 1–48.

Bigelis, R. (2001). Fungal fermentation: industrial. *Encyclopedia of Life Sciences*, 1–8.

Bjørn-Helge, M., Wehrens, R., & Hovde Liland, K. (2018). *pls: partial least squares and*
principal component regression (R package version 2.7-0). Retrieved from https://cran.r-
project.org/package=pls

Cerning, J., Gripon, J.-C., Lamberet, G., & Lenoir, J. (1987). Les activités biochimiques des
Penicillium utilisés en fromagerie. *Le Lait*, *67*(1), 3–39.

Cheese and processed cheese: determination of chloride content potentiometric titration
method. International standards 4a. International Dairy Federation, Brussels, Belgium. 1988.
https://www.iso.org/standard/43922.html. ISO 5943:2006 [IDF 88:2006]

Cheese and processed cheese: determination of total solids content. International standards
4a. (1982). *International Dairy Federation, Brussels, Belgium*. Brussels, Belgium.
http://doi.org/ISO 5534:2004 [IDF 4:2004]

Cheeseman, K., Ropars, J., Renault, P., Dupont, J., Gouzy, J., Branca, A., *et al.*. (2014).
Multiple recent horizontal transfers of a large genomic region in cheese making fungi. *Nature*
Communications, 5, 2876.

Collins, Y. F., McSweeney, P. L. H., & Wilkinson, M. G. (2003). Lipolysis and free fatty
acid catabolism in cheese: a review of current knowledge. *International Dairy Journal*,
13(11), 841–866.

Coton, E., Coton, M., Hymery, N., Jany, J. L., Mounier, J., & Jany, J. L. (2020b). *Penicillium*
roqueforti: an overview of its genetics, physiology, metabolism and biotechnological
applications. *Fungal Biology Reviews*, https://doi.org/10.1016/j.fbr.2020.03.001.

Coton, E., Jany, J.-L., & Coton, M. (2020a). Yeasts and Molds: *Penicillium roqueforti*. In
Reference Module in Food Science (pp. 1–8). Elsevier.

De Man, J. C., Rogosa, M., & Sharpe, M. E. (1960). A medium used for the cultivation of
Lactobacilli. *Journal of Applied Bacteriology*, *23*, 130–135.

Denis, C., Gueguen, M., Henry, E., & Levert, D. (2001). New media for the numeration of
cheese surface bacteria. *Le Lait*, *81*(3), 365–379.

Devoyod, J. J., & Muller, M. (1969). La flore microbienne du fromage de Roquefort III. Les
streptocoques lactiques et les *Leuconostoc*; Influence de différents micro-organismes de
contamination. *Le Lait*, *487*, 369–399. Retrieved from https://doi.org/10.1051/lait:196948715

Diezhandino, I., Fernández, D., González, L., McSweeney, P. L. H., & Fresno, J. M. (2015).
Microbiological, physico-chemical and proteolytic changes in a Spanish blue cheese during
ripening (Valdeón cheese). *Food Chemistry*, *168*, 134–141.

Dumas, É., Feurtey, A., Rodríguez de la Vega, R. C., Le Prieur, S., Snirc, A., Coton, M.,
Thierry, A., Coton, E., Le Piver, M., Roueyre, D., Ropars, J., Branca, A. & Giraud, T. (2020).
Independent domestication events in the blue-cheese fungus *Penicillium roqueforti*.
Molecular Ecology, *33913*(January), 451773.

Dupont, J., Dequin, S., Giraud, T., Le Tacon, F., Masit, S., Ropars, J., Richard, F. & Selosse,
M.-A. (2017). Fungi as a source of food. *Microbiology Spectrum*, *5*(3), 1–22.

Duval, P., Chatelard-Chauvin, C., Gayard, C., Rifa, E., Bouchard, P., Hulin, S., Picque, D. &
Montel, M.-C. (2016). Microbial dynamics in industrial blue veined cheeses in different
packaging. *International Dairy Journal*, *56*, 198–207.

Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., & Knight, R. (2011). UCHIME
improves sensitivity and speed of chimera detection. *Bioinformatics*, *27*(16), 2194–2200.

Escudié, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., Maman, S., Hernandez-Raquet, G., Combes, S. & Pascal, G. (2018). FROGS: Find, Rapidly, OTUs with Galaxy Solution. *Bioinformatics*, *34*(8), 1287–1294.

Fleming, A. (1929). On the antibacterial action of cultures of a *Penicillium*, with special reference to their use in the isolation of *B. influenzæ*. *British Journal of Experimental Pathology*, *10*(3), 226–236.

Fox, J., & Weisberg, S. (2019). *An {R} Companion to applied Regression*. (T. Oaks, Ed.) (Third). Sage. Retrieved from https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Gallone, B., Steensels, J., Prahl, T., Soriaga, L., Saels, V., Herrera-Malaver, B., Merlevede A, Roncoroni M, Voordeckers K, Miraglia L., Teiling, C., Steffy, B., Taylor, M., Schwartz, A., Richardson, T., White, C., Baele, G., Maere, S., & Verstrepen, K. J. (2016). Domestication and divergence of *Saccharomyces cerevisiae* beer yeasts. *Cell*, *166*(6), 1397–1410.

Ganesan, B., & Weimer, B. C. (2017). Amino acid catabolism and its relationship to cheese flavor outcomes. In P. L. H. McSweeney, P. F. Fox, P. D. Cotter, & D. W. Everett (Eds.), *Cheese: Chemistry, Physics and Microbiology: Fourth Edition* (Fourth Edi, Vol. 1, pp. 483–516). Elsevier Ltd.

Gatto, L., & Lilley, K. S. (2012). Msnbase-an R/Bioconductor package for isobaric tagged mass spectrometry data visualization, processing and quantitation. *Bioinformatics*, *28*(2), 288–289.

Gibbons, J. G., Salichos, L., Slot, J. C., Rinker, D. C., McGary, K. L., King, J. G., Klich M.A., Tabb D.L., McDonald W.H. & Rokas, A. (2012). The evolutionary imprint of domestication on genome variation and function of the filamentous fungus *Aspergillus oryzae*. *Current Biology*, *22*(15), 1403–1409.

Gillot, G., Jany, J.-L., Coton, M., Le Floch, G., Debaets, S., Ropars, J., López-Villavicencio M., Dupont J., Branca A., Giraud T. & Coton, E. (2015). (Fifty shades of blue :) Insights into *Penicillium roqueforti* morphological and genetic diversity. *PLoS ONE*, *10*(6), e0129849.

Gillot, G., Jany, J.-L., Poirier, E., Maillard, M., Debaets, S., Thierry, A., Coton E. & Coton, M. (2017). Functional diversity within the *Penicillium roqueforti* species. *International Journal of Food Microbiology*, *241*, 141–150.

Giraud, T., Koskella, B., & Laine, A.-L. (2017). Introduction: microbial local adaptation: insights from natural populations, genomics and experimental evolution. *Molecular Ecology*, *26*(7), 1703–1710.

Gladieux, P., Ropars, J., Badouin, H., Branca, A., Aguileta, G., De Vienne, D. M., Rodríguez De La Vega R.C., Branco S. & Giraud, T. (2014). Fungal evolutionary genomics provides insight into the mechanisms of adaptive divergence in eukaryotes. *Molecular Ecology*, *23*(4), 753–773.

Gomri, G. (1946). Buffers in the range of pH 6.5 to 9.6. *Proceedings of Society for Experimental Biology and Medicine*, *6*(1), 33–34.

Gonçalves, M., Pontes, A., Almeida, P., Barbosa, R., Serra, M., Libkind, D., Hutzler M.,
Gonçalves P. & Sampaio, J. P. (2016). Distinct domestication trajectories in top-fermenting
beer yeasts and wine yeasts. *Current Biology*, *26*(20), 2750–2761.

González de Llano, D., Ramos, M., Polo, C., Sanz, J., & Martinez-Castro, I. (1990).
Evolution of the volatile components of an artisanal blue cheese during ripening. *Journal of*
Dairy Science, *73*(7), 1676–1683.

González de Llano, D., Ramos, M., Rodriguez, A., Montilla, A., & Juarez, M. (1992).
Microbiological and physicochemical characteristics of Gamonedo blue cheese during
ripening. *International Dairy Journal*, *2*(2), 121–135.

Hébert, A., Forquin-Gomez, M. P., Roux, A., Aubert, J., Junot, C., Heilier, J. F., Landaud S.,
Bonnarmeb P. & Beckerich, J. M. (2013). New insights into sulfur metabolism in yeasts as
revealed by studies of *Yarrowia lipolytica*. *Applied and Environmental Microbiology*, *79*(4),
1200–1211.

ISO. (2004). Determination of water activity NF ISO 21807. *International Standardisation*
Organisation. http://doi.org/ISO 21807:2004

Kinderlerer, J. L. (1993). Fungal strategies for detoxification of medium chain fatty acids.
International Biodeterioration and Biodegradation, *32*(1–3), 213–224.

Kinsella, J. E., & Hwang, D. H. (1976). Enzymes of *Penicillium roqueforti* involved in the
biosynthesis of cheese flavor. *C R C Critical Reviews in Food Science and Nutrition*, *8*(2),
191–228.

Kopp, B., & Rehm, H. J. (1979). Antimicrobial action of roquefortine. *European Journal of*
Applied Microbiology and Biotechnology, *6*, 397–401 (1979).

Kuhl, C., Tautenhahn, R., Böttcher, C., Larson, T. R., & Neumann, S. (2012). CAMERA: An
integrated strategy for compound spectra extraction and annotation of liquid
chromatography/mass spectrometry data sets. *Analytical Chemistry*, *84*(1), 283–289.

Labbe, M., & Serres, J. P. (2004). *Chroniques du Roquefort - De la préhistoire à l'aube*
industrielle. Grand Imprimeur, La Primaube, France.

Labbe, M., & Serres, J. P. (2009). *Chroniques du Roquefort: des Hommes, des entreprises,*
des marques, période moderne. Grand Imprimeur, La Primaube, France.

Larsen, M. D., & Jensen, K. (1999). The effects of environmental conditions on the lipolytic
activity of strains of *Penicillium roqueforti*. *International Journal of Food Microbiology*,
46(2), 159–166.

Larsen, M. D., Kristiansen, K. R., & Hansen, T. K. (1998). Characterization of the proteolytic
activity of starter cultures of *Penicillium roqueforti* for production of blue veined cheeses.
International Journal of Food Microbiology, *43*(3), 215–221.

Larson, G., & Fuller, D. Q. (2014). The evolution of animal domestication. *Annual Review of*
Ecology, Evolution, and Systematics, *45*, 115–136.

Lazuka, A., Auer, L., Bozonnet, S., Morgavi, D. P., O'Donohue, M., & Hernandez-Raquet,
G. (2015). Efficient anaerobic transformation of raw wheat straw by a robust cow rumen-
derived microbial consortium. *Bioresource Technology*, *196*, 241–249.

Le Boucher, C., Courant, F., Jeanson, S., Chereau, S., Maillard, M., Royer, A. L., Thierry A.,
Dervilly-Pinel G., Le Bizec B. & Lortal, S. (2013). First mass spectrometry metabolic
fingerprinting of bacterial metabolism in a model cheese. *Food Chemistry*, *141*(2), 1032–
1040.

Legras, J.-L., Galeote, V., Bigey, F., Camarasa, C., Marsit, S., Nidelet, T., Couloux A., Guy
J., Franco-Duarte R. Marcet-Houben, M. Gabaldon, T., Schuller, D., Sampaio, J. P. &
Dequin, S. (2018). Adaptation of *S. cerevisiae* to fermented food environments reveals
remarkable genome plasticity and the footprints of domestication. *Molecular Biology and*
Evolution, *35*(7), 1712–1727.

Libkind, D., Hittinger, C. T., Valeŕio, E., Gonçalves, C., Dover, J., Johnston, M., Gonçalves,
P. & Sampaio, J. P. (2011). Microbe domestication and the identification of the wild genetic
stock of lager-brewing yeast. *Proceedings of the National Academy of Sciences of the United*
States of America, *108*(35), 14539–14544.

Liu, Z., Lozupone, C., Hamady, M., Bushman, F. D., & Knight, R. (2007). Short
pyrosequencing reads suffice for accurate microbial community analysis. *Nucleic Acids*
Research, *35*(18), e120.

Ludemann, V., Greco, M., Rodríguez, M. P., Basílico, J. C., & Pardo, A. G. (2010). Conidial
production by *Penicillium nalgiovense* for use as starter cultures in dry fermented sausages
by solid state fermentation. *LWT - Food Science and Technology*, *43*(2), 315–318.

Mahé, F., Rognes, T., Quince, C., de Vargas, C., & Dunthorn, M. (2014). Swarm: robust and
fast clustering method for amplicon-based studies. *PeerJ*, *2*, e593.

Martín, J.-F., & Coton, M. (2016). Blue cheese: microbiota and fungal metabolites. In J.
Frias, C. Martinez-Villaluenga, & E. Peñas (Eds.), *Fermented Foods in Health and Disease*
Prevention (pp. 275–303). Elsevier Inc.

Mason, A. B., & Dufour, J. P. (2000). Alcohol acetyltransferases and the significance of ester
synthesis in yeast. *Yeast*, *16*(14), 1287–1298.

Mayeux, J. V, Sandine, W. E., & Elliker, P. R. (1962). A selective medium for detecting
Leuconostoc organisms in mixed-strain starter cultures. *Journal of Dairy Science*, *45*, 655–
656.

McSweeney, P. L. H. (1997). The flavour of milk and dairy products: III. Cheese: Taste.
International Journal of Dairy Technology, *50*(4), 123–128.

McSweeney, P. L. H., Fox, P. F., & Ciocia, F. (2017). Metabolism of residual lactose and of
lactate and citrate. In P. L. H. McSweeney, P. F. Fox, P. D. Cotter, & D. W. Everett (Eds.),
Cheese: Chemistry, Physics and Microbiology: Fourth Edition (Fourth Edi, pp. 411–421).
Elsevier Ltd.

McSweeney, P. L. H., & Sousa, M. J. (2000). Biochemical pathways for the production of
flavour compounds in cheeses during ripening: A review. *Le Lait*, *80*(3), 293–324.

Moio, L., Piombino, P., & Addeo, F. (2000). Odour-impact compounds of Gorgonzola
cheese. *Journal of Dairy Research*, *67*(2), 273–285.

Moreau, C. (1980). Le *Penicillium roqueforti*, morphologie, physiologie, intérêt en industrie
fromagère, mycotoxines. *Le Lait*, *60*(595–596), 254–271.

Mossel, D. A. A., Eelderink, I., Koopmans, M., & Van Rossem, F. (1978). Optimisation of a
MacConkey-type medium for the enumeration of *Enterobacteriaceae*. *Laboratory Practices*,
27, 1049–1050.

Mossel, D. A. A., Kleynen□Semmeling, A. M. C., Vincentie, H. M., Beerens, H., &
Catsaras, M. (1970). Oxytetracycline□Glucose□Yeast extract agar for selective enumeration
of moulds and yeasts in foods and clinical material. *Journal of Applied Bacteriology*, *33*(3),
454–457.

Nelson, F. E. (1940). The effect of the new standard milk agar on the plate count of dairy
products. *Journal of Bacteriology*, *39*(3), 263–272. Retrieved from
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC374570/

Oksanen, J., Blanchet, G. F., Firendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin,
P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M. H. H., Szoecs, E. & Wagner, H.
(2019). vegan: community ecology package. Retrieved from https://cran.r-
project.org/package=vegan

Perrone, G., Samson, R. A., Frisvad, J. C., Susca, A., Gunde-Cimerman, N., Epifani, F., &
Houbraken, J. A. M. P. (2015). *Penicillium salamii*, a new species occurring during seasoning
of dry-cured meat. *International Journal of Food Microbiology*, *193*, 91–98.

Pham, V. T., Lacroix, C., Braegger, C. P., & Chassard, C. (2016). Early colonization of
functional groups of microbes in the infant gut. *Environmental Microbiology*, *18*(7), 2246–
2258.

Pitt, J. I., & Hocking, A. D. (2009). *Fungi and food spoilage* (Springer). Springer, Boston,
MA.

Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile
open source tool for metagenomics. *PeerJ*, *4*, e2584.

Ropars J., Bennetot B., Caron T., Lo Y.C. and Giraud T. (2020a). The domestication of
Penicillium cheese fungi. *Comptes Rendus de l'Académie des Sciences - Biologie, 343*, 155-
176.

Ropars, J., Cruaud, C., Lacoste, S., & Dupont, J. (2012). A taxonomic and ecological
overview of cheese fungi. *International Journal of Food Microbiology*, *155*(3), 199–210.

Ropars J., Didiot E., Rodriguez de la Vega R., Bennetot B., Coton M., Coton E., Snirc A., Le
Prieur S. and Giraud T. (2020b). Domestication of the emblematic white cheese-making
fungus *Penicillium camemberti* and its diversification into two varieties *Current Biology, 30*,
4441-4453.

Ropars, J., Lo, Y.-C., Dumas, É., Snirc, A., Begerow, D., Rollnik, T., Lacoste, S., Dupont, J.,
Giraud, T. & López-Villavicencio, M. (2016). Fertility depression among cheese-making

Penicillium roqueforti strains suggests degeneration during domestication. *Evolution*, *70*(9),
2099–2109.

Ropars, J., López-Villavicencio, M., Dupont, J., Snirc, A., Gillot, G., Coton, M., Jany, J.-L.,
Coton, E. & Giraud, T. (2014). Induction of sexual reproduction and genetic diversity in the
cheese fungus *Penicillium roqueforti*. *Evolutionary Applications*, *7*(4), 433–441.

Ropars, J., López-Villavicencio, M., Snirc, A., Lacoste, S., & Giraud, T. (2017). Blue cheese-
making has shaped the population genetic structure of the mould *Penicillium roqueforti*.
PLoS ONE, *12*(3), e0171387.

Ropars, J., Rodríguez de la Vega, R. C., Gouzy, J., Dupont, J., Swennen, D., Dumas, É.,
Giraud, T. & Branca, A. (2016). Diversity and mechanisms of genomic adaptation in
Penicillium. In R. P. de Vries, I. Benoit Gelber, & A. Rordam (Eds.), *Aspergillus and*
Penicillium in the post-genomic era (HAL). Caister Academic Press.

Ropars, J., Rodríguez de la Vega, R. C., López-Villavicencio, M., Gouzy, J., Sallet, E.,
Dumas, É., Lacoste, S., Debuchy, R., Dupont, J., Branca, A. & Giraud, T. (2015). Adaptive
horizontal gene transfers between multiple cheese-associated fungi. *Current Biology*, *25*(19),
2562–2569.

Rothe, M., Engst, W., & Erhardt, V. (1982). Studies on characterization of blue cheese
flavour. *Die Nahrung*, *26*(7/8), 591–602.

Roudot-Algaron, F. (1996). Le goût des acides aminés, des peptides et des protéines:
Exemple de peptides sapides dans les hydrolysats de caséines. *Lait*, *76*(4), 313–348.

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years
of image analysis. *Nature Methods*, *9*(7), 671–675.

Schnurer, J. (1993). Comparison of methods for estimating the biomass of three food-borne
fungi with different growth patterns. *Applied and Environmental Microbiology*, *59*(2), 552–
555.

Searle, S. R., Speed, F. M., & Milliken, G. A. (1980). Population marginal means in the
linear model: An alternative to least squares means. *American Statistician*, *34*(4), 216–221.

Sicard, D., & Legras, J.-L. (2011). Bread, beer and wine: Yeast domestication in the
Saccharomyces sensu stricto complex. *Comptes Rendus - Biologies*, *334*(3), 229–236.

Smith, C. A., Want, E. J., O'Maille, G., Abagyan, R., & Siuzdak, G. (2006). XCMS:
Processing mass spectrometry data for metabolite profiling using nonlinear peak alignment,
matching, and identification. *Analytical Chemistry*, *78*(3), 779–787.

Snow, G. (2016). TeachingDemos: Demonstrations for teaching and learning. Retrieved from
https://cran.r-project.org/web/packages/TeachingDemos/index.html

Spinnler, H.-E. (2011). Rôle des lipides dans la perception olfactive des produits laitiers.
Sciences Des Aliments, *30*, 103–120.

Terzaghi, B. E., & Sandine, W. E. (1975). Improved medium for lactic streptococci and their
bacteriophages. *Applied Microbiology*, *29*(6), 807–813.

Thierry, A., Collins, Y. F., Abeijón Mukdsi, M. C., McSweeney, P. L. H., Wilkinson, M. G.,
& Spinnler, H.-E. (2017). Lipolysis and metabolism of fatty acids in cheese. In P. L. H.
McSweeney, P. F. Fox, P. D. Cotter, & D. W. Everett (Eds.), *Cheese: Chemistry, Physics and*
Microbiology: Fourth Edition (Fourth Edi, pp. 423–444). Elsevier Ltd.

Urbanek, S., & Horner, J. (2019). Cairo: R graphics device using Cairo graphics library for
creating high-quality bitmap (PNG, JPEG, TIFF), vector (PDF, SVG, PostScript) and display
(X11 and Win32) output. Retrieved from https://cran.r-
project.org/web/packages/Cairo/index.html

Vabre, S. (2015). *Le sacre du Roquefort*. Presses universitaires François Rabelais, Tours,
France.

Vallone, L., Giardini, A., & Soncini, G. (2014). Secondary metabolites from *Penicillium*
roqueforti, a starter for the production of Gorgonzola cheese. *Italian Journal of Food Safety*,
3(3), 173–177 (2118).

Van Den Boogaart, K. G., Tolosana-Delgado, R., & Bren, M. (2018). compositions: A
unified R package to analyze compositional data. *Computers & Geosciences*, *34*(4), 320–338.

Villanueva, R. A. M., Chen, Z. J., & Wickham, H. (2016). *ggplot2: Elegant graphics for data*
analysis using the grammar of graphics (2nd Editio). Springer-Verlag New York.

Wickham, H. (2011). The split-apply-combine strategy for data analysis. *Journal of*
Statistical Software, *40*(1), 1–29.

Wilke, C. O. (2019). cowplot: Streamlined plot theme and plot annotations for "ggplot2".
Retrieved from https://cran.r-project.org/web/packages/cowplot/index.html

Williams, A. G., Beattie, S. H., & Banks, J. M. (2004). Enzymes involved in flavour
formation by bacteria isolated from the smear population of surface-ripened cheese.
International Journal of Dairy Technology, *57*(1), 7–13.

Woo, A. H., & Lindsay, R. C. (1984). Concentrations of major free fatty acids and flavor
development in Italian cheese varieties. *Journal of Dairy Science*, *67*(5), 960–968.

Figure legends

Figure 1: Experimental cheesemaking. (A) Experimental design for cheesemaking, using one

strain per cheese, and three different strains from each of the four *Penicillium roqueforti*

populations (non-Roquefort cheese in blue, Roquefort cheese in purple, silage/food spoiler in

orange, lumber/food spoiler in green, the lineages of which are shown on the left). Each assay

(February, March, April) corresponded to a single strain from each of the four populations,

with three production replicates at different times, different batches of unpasteurized milk

and with two cheeses produced per strain in each replicate. The identities of the strains used

are indicated on the left of each assay, for each of the four *P. roqueforti* populations. (B)

Photograph of the experimental cheeses after 20 days of maturation.

Figure 2: (A) Relative abundance of the six main bacterial operational taxonomic units in

cheeses made with strains from the four *Penicillium roqueforti* populations (non-Roquefort

cheese in blue, Roquefort cheese in purple, silage/food spoiler in orange and lumber/food

spoiler in green) in cheeses at 9 (left) and 20 (right) days of maturation. (B & C) Mean

bacterial genus diversity; A: Shannon index, B: Inverse of Simpson index (= 1 - Simpson

index) for the operational taxonomic units detected by metabarcoding in 9-day cheeses (left)

and 20-day cheeses (right) made with strains from the four *Penicillium roqueforti* populations

(lumber/food spoiler in green, non-Roquefort cheese in blue, Roquefort cheese in purple and

silage/food spoiler in orange).

Figure 3: (A) Mean percentage blue area per cheese slice at 20, 90 and 180 days of

maturation, for cheeses made with strains from the four *Penicillium roqueforti* populations

(non-Roquefort cheese in blue, Roquefort cheese in purple, silage/food spoiler in orange and

lumber/food spoiler in green). Error bars indicate 95% confidence intervals. (B) Illustration

of the differences in mean percentage blue area per cheese slice at 180 days of maturation

between the four *Penicillium roqueforti* populations (non-Roquefort cheese in blue,

Roquefort cheese in purple, silage/food spoiler in orange and lumber/food spoiler in green).

Contrast and brightness have been standardized and the edges cropped.

Figure 4: Violin plot depicting the distribution of the sums of 3,864 non-targeted aqueous

signal peak areas, weighted by their mass-to-charge ratios ("m/z"), obtained in positive

ionization mode for 90-day cheeses made with strains from the four *Penicillium roqueforti*

populations (lumber/food spoiler in green, non-Roquefort cheese in blue, Roquefort cheese in

purple and silage/food spoiler in orange). Boxplots within violin plots represent the median

(center line), the 25th and 75th percentiles (box bounds), the 5th and 95th percentiles

(whiskers), the points being the outliers from these 95th and 5th percentiles. The red dots

depict the mean values.

1007

Figure 5: Violin plot depicting the distribution of the sums of 3,094 non-targeted organic

signal peak areas, weighted by their mass-to-charge ratios ("m/z"), obtained in negative

ionization mode for 90-day cheeses made with strains from the four *Penicillium roqueforti*

populations (lumber/food spoiler in green, non-Roquefort cheese in blue, Roquefort cheese in

purple and silage/food spoiler in orange). Boxplots within violin plots represent the median

(center line), the 25th and 75th percentiles (box bounds), the 5th and 95th percentiles

(whiskers), and the red dots depict the mean values.

1015

Figure 6: Volatile compound production (integrated peak areas from chromatograms in

arbitrary units) in 90-day cheeses inoculated with strains from the four *Penicillium roqueforti*

populations (non-Roquefort cheese in blue, Roquefort cheese in purple, silage/food spoiler in

orange and lumber/food spoiler in green). The areas for each family of compounds are the

sum of the integrated areas of the compounds belonging to the family concerned. Alcohols I

and II are derived from proteolysis and lipolysis, respectively. Acids I, II and III are derived

from proteolysis, glycolysis and lipolysis, respectively (Supplementary Table 3). The color of

the titles indicates the affiliation of the compounds to their families, as in Figure S6.

1024

1025 **Figure 7**: Integrated surface area (from chromatograms in arbitrary units) of the three most

1026 discriminant methyl ketones (A) and four secondary alcohols (B) for each assay (February,

1027 March, April) for the three strains of each *Penicillium roqueforti* population (lumber/food

1028 spoiler in green, non-Roquefort cheese in blue, Roquefort cheese in purple, silage/food

1029 spoiler in orange). Error bars represent standard deviations across cheese replicates.

1030

1031 **Figure legends for the Supplementary Material**

1032

Figure S1: Abundance of microorganisms (in log colony-forming units/g) of the eight types

monitored at various stages of cheese maturation (i.e. unpasteurized milk, 9, 20, 90 and 180

days), for each of the four *Penicillium roqueforti* populations used to inoculate the cheeses

(non-Roquefort cheese in blue, Roquefort cheese in purple, silage/food spoiler in orange and

lumber/food spoiler in green). Error bars represent standard deviations across assays.

Figure S2: Illustration of image processing for estimation of the percentage blue area on

cheese slices: (a) example of an unprocessed image of a cheese slice; (b) image after

brightness and contrast standardization; (c) image after cropping; (d) corresponding image

binarization with a grayscale of 102 on the red channel. White and black correspond to pixel

classification: in white, the inner part of the cheese and empty cavities; in black, cavities

filled with the fungus.

1044

Figure S3: Differences in aminoacid content between cheeses according to the population-of-

origin of the *Penicillium roqueforti* strains. A. Discrimination between 90-day cheeses made

with cheese (blue) and non-cheese (green) *P. roqueforti* populations (left), or Roquefort

cheese (purple) and non-Roquefort cheese (blue) *P. roqueforti* populations (right), based on

the abundance of the 23 identified amino acids present, according to an orthogonal signal-

corrected partial least squares (PLS) discriminant analysis. Vertical and horizontal axes

represent PLS1 and PLS 2 scores and gray arrows represent the relative contribution of

loadings of signals significantly discriminating the group considered in a *t*-test with jackknife

resampling. B. Abundance of molecules from particular classes detected in cheeses: mean

integrated peak area from chromatograms in arbitrary units (bars, left axis) and cumulative

percentage (line with dots, right axis) of aqueous extracts across all 90-day cheeses.

1056

Figure S4: Violin plot depicting the distribution of the sums of 8,472 non-targeted organic

signal peak areas, weighted by their mass-to-charge ratios ("m/z"), obtained in positive

ionization mode for 90-day cheeses made with strains from the four *Penicillium roqueforti*

populations (lumber/food spoiler in green, non-Roquefort cheese in blue, Roquefort cheese in

purple and silage/food spoiler in orange). Boxplots within violin plots represent the median

(center line), the 25th and 75th percentiles (box bounds), the 5th and 95th percentiles

(whiskers), and the red dots depict the mean values.

1064

1065 **Figure S5**: Non-protein nitrogen levels at 20, 90 and 180 days of maturation, and water

1066 activity at 90 and 180 days of maturation. Comparison of cheeses made with strains from

1067 different *Penicillium roqueforti* populations (non-Roquefort cheese in blue, Roquefort cheese

1068 in purple, silage/food spoiler in orange and lumber/food spoiler in green). Error bars indicate

1069   95% confidence intervals.

1070

Figure S6: Discrimination between 90-day cheeses inoculated with strains from the four

Penicillium roqueforti populations (non-Roquefort cheese in blue, Roquefort cheese in

purple, silage/food spoiler in yellow and lumber/food spoiler in green), based on the

abundance of 41 volatile compounds in an orthogonal signal-corrected partial least squares

(PLS) discriminant analysis. Vertical and horizontal axes represent the PLS1 and PLS2

variances, and arrows represent the relative contributions of compound odor loadings

significantly   discriminating   the   group   considered   (according   to

www.thegoodscentscompany.com) in a *t*-test with jackknife resampling. The odor colors

indicate the families in Figure 6 to which the associated compounds belong.

1080

Conflict of interest: TC, MLP, SB, DR and MP were employed by SAS LIP, which

produces starters for fermented food products, during the course of the study and therefore

declare a competing financial interest. None of the other authors has any conflict of interest to

declare. With the exception of TC employed by the LIP, the funders had no role in decisions

concerning design, data analysis and interpretation, or in the decision to submit the work for

publication.

Figure 1

A

Four *P. roqueforti* populations

assay	February			March			April		
production replicate	milk	milk	milk	milk	milk	milk	milk	milk	milk
strains									
BRFM 3021									
BRFM 3022									
BRFM 3023									

non Roquefort

LCP 6064
LCP 6040
LCP 6039

lumber / food spoiler

LCP 6043
LCP 5885
LCP 3914

silage / food spoiler

BRFM 3020
BRFM 3019
BRFM 3024

Roquefort

analyses

Biological counts — Bacterial metabarcoding — surface covered by blue — general physico-chemistry — organic, aqueous and volatiles extraction

B

Figure 2A

Penicillium roqueforti population

■ lumber/food spoiler   ■ non-Roquefort   ■ Roquefort   ■ silage/food spoiler

Figure 2B

Figure 2C

Figure 3A

Figure 3B

Non-cheese *P. roqueforti* populations	Cheese *P. roqueforti* populations
lumber/food spoiler (5.2 %)	non-Roquefort (10.2 %)
silage/food spoiler (5.5%)	Roquefort (11.8 %)

Figure 4

Figure 5

Figure 6

Figure 7