Title: Strong effect of *Penicillium roqueforti* populations on volatile and metabolic compounds responsible for aromas, flavour and texture in blue cheeses

Running title: Impact of mold population impact on blue cheeses

Authors:
Thibault CARON[1,4], Mélanie LE PIVER[4], Anne-Claire PÉRON[3], Pascale LIEBEN[3], René LAVIGNE[2], Sammy BRUNEL[4], Daniel ROUEYRE[4], Michel PLACE[4], Pascal BONNARME[3], Tatiana GIRAUD[1]*, Antoine BRANCA[1]*, Sophie LANDAUD[3]*, Christophe CHASSARD[2]*

1: Ecologie Systematique Evolution, Université Paris Saclay, CNRS, AgroParisTech, 91400 Orsay, France
2: Université Clermont Auvergne, INRAE, Vetagro Sup, UMRF, 20 Côte de Reyne, 15000 Aurillac, France
3: Université Paris-Saclay, INRAE, AgroParisTech, UMR SayFood, 78850 Thiverval-Grignon, France
4: Laboratoire Interprofessionnel de Production – SAS L.I.P., 34 rue de Salers, 15 000 Aurillac, France

*should be considered joint senior authors

Contact information: Antoine Branca antoine.branca@u-psud.fr

Keywords: Roquefort cheese, fungi, *Penicillium,* domestication, volatile compounds

Acknowledgments: We thank Béatrice DESSERRE, Céline DELBES and Cécile CALLON for advice and technical assistance in microbiology, Sébastien THEIL for the technical support of metabarcoding analyses, Patricia LE THUAUT, Manon SURIN and Brigitte POLLET for the technical support of metabolomic analyses, Sara PARISOT for milk delivery and quality, Pierre CONCHON for the technical support of image analysis, LIAL-MC for the various reference measurements in physico-chemistry, Christophe LACROIX and Alfonso DIE for the determination of short-chain fatty acids in fermentation supernatants. This study has been funded by the LIP SAS, ANRT (association nationale recherche technologie), by the ERC Genomefun 309403 Stg and Blue Proof of Concept grants, the Fondation Louis D grant (French Academy of Sciences) and the ANR-19-CE20-0002-02 Fungadapt ANR grant.

Abstract

Studies of food microorganism domestication can provide important insight into adaptation

mechanisms and lead to commercial applications. The *Penicillium roqueforti* fungus consists

of four genetically differentiated populations, two of which have been domesticated for blue

cheese-making, the other two thriving in other environments. Most blue cheeses are made

with strains from a single *P. roqueforti* population, whereas Roquefort cheeses are inoculated

with strains from a second population. We made blue cheeses in accordance with the

production specifications for Roquefort-type cheeses, inoculating each cheese with a single

P. roqueforti strain and using three strains from each of the four populations. The strain

population-of-origin had a minor impact on bacterial diversity and none on the main

microorganism abundance. The strains from cheese populations produced cheeses with

higher percentages of blue area and larger amounts of desired volatile compounds. In

particular, the Roquefort strains produced larger amounts of appealing aromatic compounds,

in part due to their greater efficiency of proteolysis and lipolysis. The typical appearance and

flavors of blue cheeses thus result from human selection on *P. roqueforti,* and the two cheese

populations have acquired specific features. This has important implications for our

understanding of adaptation and domestication, and for cheese improvement.

Domestication is an evolutionary process that has been studied by many biologists since

Darwin. Indeed, domestication is an excellent model for understanding adaptation, being the

result of strong and recent selection on traits that are often known and of interest for humans

(Larson *et al.,* 2014). In addition, studies of domestication frequently have important

implications for the improvement of cultivated organisms. However, domesticated fungi have

been much less studied than crops, despite being excellent models in this field (Gladieux *et*

al., 2014; Giraud *et al.,* 2017). Most fungi can be cultured in Petri dishes, can remain alive

for decades when stored in freezers and are propagated asexually. All these features facilitate

experiments. Fungal metabolism produces various compounds of interest, including fuels,

enzymes and antibiotics (Bigelis, 2001). The most ancient and frequent use of fungi by

humans is for fermentation, to preserve and mature food. For example, the yeast

Saccharomyces cerevisiae is used for bread, wine and beer fermentation, and the filamentous

fungus *Aspergillus oryzae* is used for soy sauce and sake fermentation (Dupont *et al.,* 2017).

These models have provided important insight into mechanisms of adaptation and

domestication (Almeida *et al.,* 2014; Baker *et al.,* 2015; Gallone *et al.,* 2016; Gibbons *et al.,*

2012; Gonçalves *et al.,* 2016; Libkind *et al.,* 2011; Sicard *et al.,* 2011).

The *Penicillium* genus contains more than 300 species, several of which are used by humans.

For example, penicillin was discovered in *P. rubens,* and *P. nalgiovense* and *P. salamii* are

used for the production of dry-cured meat (Fleming, 1929; Ludemann *et al.,* 2010, Perrone *et*

al., 2015). For centuries, *Penicillium roqueforti* (Thom) has been used in the maturation of all

the many varieties of blue cheese worldwide (Labbe *et al.,* 2004, 2009; Vabre, 2015). This

fungus generates the blue-veined appearance of these cheeses, by producing melanized spores

in cavities within the cheese in which oxygen is available (Moreau, 1980). *Penicillium*

roqueforti is also found in non-cheese environments (Pitt *et al.,* 2009; Ropars *et al.,* 2012),

and four genetically differentiated clusters of individuals (i.e., populations) have been

identified in *P. roqueforti*. Two populations are used for cheesemaking, whereas the other

two populations thrive in silage, lumber or spoiled food (Ropars *et al.,* 2014; Gillot *et al.,*

2015; Dumas *et al.,* 2020). Genomic and experimental approaches have provided compelling

evidence for the domestication of cheese *P. roqueforti* populations (Cheeseman *et al.,* 2014;

Ropars *et al.,* 2015, 2016 a & b, 2017; Gillot *et al.,* 2015, 2017; Dumas *et al.,* 2020). Indeed,

the populations of *P. roqueforti* used to make blue cheeses display the characteristic features

of domesticated organisms: genetic and phenotypic differences relative to non-cheese

populations, with, in particular, traits beneficial for cheese production, such as faster growth

on cheese medium (Ropars *et al.,* 2014, 2016; Gillot *et al.,* 2015; Dumas *et al.,* 2020), but

also lower fertility and lower fitness in nutrient-poor conditions (Ropars *et al.,* 2015, 2016).

Both cheese populations have lower levels of genetic diversity than the two non-cheese

populations, indicating an occurrence of bottlenecks (Dumas *et al.,* 2020), which typically

occur during domestication. The two cheese populations are genetically and phenotypically

differentiated from each other, suggesting that they result from independent domestication

events (Dumas *et al.,* 2020). One of the cheese populations, the non-Roquefort population, is

a clonal lineage with a very low level of genetic diversity, used to produce most types of blue

cheeses worldwide. The second cheese population, the Roquefort population, is genetically

more diverse and contains all the strains used to produce blue cheeses from the emblematic

Roquefort protected designation of origin (PDO) (Dumas *et al.,* 2020). *In vitro* tests showed

that the non-Roquefort population displayed faster tributyrin degradation (*i.e.* a certain type

of lipolysis) and a higher salt tolerance, faster *in vitro* growth on cheese medium and better

exclusion of competitors than the Roquefort population (Ropars *et al.,* 2014, 2015; Dumas *et*

al., 2020). The specific features of the Roquefort population may result from the constraints

of the PDO, requiring the use of local strains and at least 90 days of maturation, and

preventing the use of strains from the non-Roquefort population better suited to modern

modes of production (Dumas *et al.,* 2020). Genomic footprints of domestication (i.e., of

adaptive genetic changes) have also been identified in the two *P. roqueforti* populations used

for cheesemaking. Indeed, it has been suggested that horizontally transferred genes found

only in the non-Roquefort population are involved in the production of an antifungal peptide

and in lactose catabolism (Ropars *et al.,* 2014, 2015; Cheeseman *et al.,* 2014). The effects of

positive selection have been detected in genes with predicted functions in flavor compound

production, in each of the cheese populations (Dumas *et al.,* 2020).

Thus, the four *P. roqueforti* populations probably harbor multiple specific traits, leading to

the generation of cheeses with different physicochemical properties and flavors, although this

has yet to be tested. Assessments of the effect of the population-of-origin of the *P. roqueforti*

strain used on the features of the cheese will i) provide important fundamental knowledge

about the trait under selection for cheesemaking and adaptation to the cheese environment, ii)

provide a basis for the elucidation of other genomic changes and iii) be of crucial applied

importance for governing strain use and strain improvement. *Penicillium roqueforti* is used as

a secondary starter for flavor production, mostly through proteolysis (*i.e.* casein catabolism)

and lipolysis during ripening (Moreau, 1980). The main characteristic feature of blue cheeses,

and of Roquefort PDO cheeses in particular, is their intense, spicy flavors (Kinsella *et al.,*

1976; Rothe *et al.,* 1982). The specific volatile and metabolic compounds responsible for

these flavors are generated principally by lipolysis in blue cheeses (Cerning *et al.,* 1987;

Collins *et al.,* 2003), but their intensity varies between *P. roqueforti* strains (Larsen *et al.,*

1999; Dumas *et al.,* 2020). The fatty acids released by lipolysis are the precursors of

aldehydes, alcohols, acids, lactones and methyl ketones, which provide the moldy aromas

typical of blue cheeses (Collins *et al.,* 2003). *Penicillium roqueforti* degrades most proteins,

but proteolysis efficiency varies between strains (Cerning *et al.*, 1987; Larsen *et al.*, 1998;

Dumas *et al.*, 2020). The resulting peptides contribute to flavors, and their degradation into

amino acids further influences cheese aroma and the growth of other microorganisms

(Williams *et al.*, 2004; McSweeney *et al.*, 2000). *Penicillium roqueforti* also contributes to

lactate degradation, which is necessary for deacidification and promotes the development of

less acid-tolerant microorganisms (McSweeney *et al.*, 2017). Through these effects, and by

producing secondary metabolites with antimicrobial properties, *P. roqueforti* may also affect

the microbial composition of the cheese (Kopp *et al.*, 1979; Vallone *et al.*, 2014). Another

parameter potentially affected by *P. roqueforti* populations and restricting the occurrence of

spoiler microorganisms is the lack of free water, (i.e., a low water activity *aka* Aw), which is

heavily controlled for Roquefort cheese sales and is affected by the degree of proteolysis

(Ardö *et al.*, 2017). The *P. roqueforti* population may thus also have an indirect effect on the

features of the cheese, through various effects on beneficial or undesirable contaminants.

The differences between *P. roqueforti* populations have, to date, been studied only *in vitro* or

in very rudimentary cheese models. Here, our objective was to assess the effect of the *P.*

roqueforti population-of-origin of the inoculated strains on the features of blue cheeses

produced in conditions closely mimicking those of commercial Roquefort PDO cheese

production. We focused on several features considered important for cheese quality. Given

the evidence from previous studies that cheese *P. roqueforti* populations have been

domesticated, any differences between the cheeses produced with cheese and non-cheese

populations, and/or between the two cheese populations would probably reflect human

selection for the production of good cheeses, either on standing variation in the ancestral *P.*

roqueforti population or for *de novo* mutations. Identifying the differences between *P.*

roqueforti populations in terms of their properties for cheesemaking (e.g., ripening dynamics

and specific flavors) would improve our understanding of domestication and adaptation

processes, and might drive important applications and developments. We therefore produced

blue cheeses in conditions very similar to those used in industrial Roquefort PDO production,

using, in particular, milk from the local "Lacaune" breed, with strains from the four *P.*

roqueforti populations. We compared several important cheese features between the four

populations: i) physicochemical features, relating to texture and biochemical composition, ii)

cheese microbiota composition and abundance, which may have effects on several cheese

features, iii) the proportion of the blue area in cheese slices, which is important for the blue-

veined appearance of the cheese and is dependent on the growth and sporulation of *P.*

roqueforti in cheese cavities, and iv) the metabolic and volatile compounds produced and

their amounts, which influence flavor and aroma. We investigated the differences in these

features between the cheeses produced with strains from the four *P. roqueforti* populations

(Roquefort cheese, non-Roquefort cheese, silage, and lumber/food spoiler populations). We

also investigated the possible differences between cheeses made with cheese and non-cheese

populations, and between the Roquefort and non-Roquefort cheese populations. Assessments

of the traits differing between cheese and non-cheese *P. roqueforti* populations, and between

the two cheese populations, and investigations of whether the cheese populations are more

suitable for cheese-making, i) is of fundamental importance for understanding the

domestication of cheese fungi, through the identification of traits subjected to selection, and

ii) has many applied consequences for the cheese industry, in terms of strain choice for

different kinds of blue cheeses, paving the way for the improvement of mold strains by

generating progenies from crosses of the two cheese populations, and for the choice of traits

for measurement and selection in offsprings.

Materials and Methods

More details about the Materials and Methods are provided in the Supplementary Methods.

Cheesemaking: The cheesemaking protocol was typical of the procedures used by the main

producers of Roquefort cheese and complied with the Roquefort PDO specifications, except

that the ripening process took place in artificial cellars in the INRA facilities at Aurillac, with

strains from different *P. roqueforti* populations (Figure 1A; a strain is defined here as a

haploid individual obtained by monospore isolation). We made cheeses by inoculating a

single strain per cheese, using in total three different strains from each of the four *P.*

roqueforti populations (Figure 1A). The strains were assigned to these populations in a

previous study, on the basis of molecular markers (Dumas *et al.*, 2020). Due to the limited

production capacity of the experimental facility, it was not possible to make all the cheeses at

the same time. We therefore split cheese production into three assays, each including one

strain from each of the four populations (Figure 1A). For each strain in each assay, we

created three production replicates, with two cheeses per strain in each replicate, to ensure

that enough material was produced for sampling. In total, we produced 72 cheeses (4 strains *

2 cheeses * 3 replicates * 3 assays; figure 1B). The assays were performed sequentially from

February to April. The effect of the seasonal change in milk composition was therefore

confounded with the strain effect within the population, hereafter referred to as the "assay

effect". The three replicates within each assay were also set up at different times, a few days

apart, and thus with different batches of unpasteurized milk (Figure 1A).

Microbial analyses: We estimated the concentrations of various microorganism

communities in the initial unpasteurized milk and at various stages of cheese maturation (for

more information see the Supplementary Methods). We performed a metabarcoding analysis

on the experimental cheeses at 9 and 20 days of maturation, by sequencing the 16S DNA

fragment with Illumina Miseq technology and analyzing sequences with Find Rapidly OTUs

in Galaxy Solution (FROGS), v3.0 (Escudié *et al.*, 2018). For each OTU, taxonomic

assignment was determined with the Silva-132 (https://www.arb-silva.de/) and 16S rDNA

RefSeq databases (https://blast.ncbi.nlm.nih.gov/Blast.cgi).

Blue area: We estimated the percentage area of the cheese that was blue, on fresh inner

cheese slices, by analyzing images of the slices with ImageJ software and counting the

number of dark pixels.

Physicochemistry: We performed standard physicochemical measurements on the cheeses.

We measured dry matter content, fat over dry matter content, the moisture content of the

defatted cheese, total, soluble and non-protein nitrogen contents, chloride and salt content,

water activity and pH at various stages of maturation, according to reference methods (for

more information, see the Supplementary Methods). We measured glucose, lactose, lactate,

acetate and butyrate concentrations in the cheeses on days 9 and 20, by high-performance

liquid chromatography (HPLC, for more information, see the Supplementary Methods).

Metabolic and volatile compounds: We investigated possible differences in proteolytic and

lipolytic activities between the four populations, by UHPLC-MS after two extraction

procedures (water and an organic solvent). We analyzed the amounts of free fatty acids and

residual glycerides in 90-day cheeses, by coupling a global extraction (accelerated solvent

extraction with hexane-isobutanol) with UHPLC-MS analysis in the positive (triglycerides)

and negative (fatty acids) ionization modes (for more information, see the Supplementary

Methods). We investigated the identity and abundance of volatile flavor and aroma

compounds, using a dynamic headspace system (DHS) with a Gerstel MPS autosampler

(Mülheim an der Ruhr, Germany) and gas chromatography-mass spectrometry analysis with a

7890B Agilent GC system coupled to an Agilent 5977B quadrupole mass spectrometer (Santa

Clara, United States). Statistical analyses were performed with R software (http://www.r-

project.org/). Further details about the materials and methods are provided in the

Supplementary Methods.

Results

Design and cheesemaking. We made cheeses by inoculating a single strain per cheese and

using in total three different strains (biological replicates) from each of the four *P. roqueforti*

populations (Figure 1A). We divided the production into three assays, each including one

strain from each of the four populations (Figure 1A). For each strain in each assay, we

generated three production replicates at different times, with different batches of

unpasteurized milk, each production replicate encompassing two cheeses per strain. The

assays were performed sequentially from February to April. The effect of the seasonal change

in milk composition was therefore confounded with the strain effect, hereafter referred to as

the "assay effect". The experimental design did not, therefore, allow for the testing of a strain

effect, but it was possible to test for a population effect (the replicates being the three strains

used per population), which was our goal. The seasonal effect, if any, would blur the

population effect, so any differences between populations detected in this study can be

considered to be robust.

Penicillium roqueforti **population-of-origin influences the bacterial diversity of the**

cheese, but not the abundance of the main microorganisms. We investigated whether *P.*

roqueforti population-of-origin affected the composition of the cheese microbiota, by

estimating the densities of key microbial communities with colony counts (cfu/g) on various

specific culture media (for total aerobic mesophilic bacteria, mesophilic lactic acid bacteria,

thermophilic lactic acid bacteria, dextran-producing *Leuconostoc* spp., molds and yeasts,

Gram-positive catalase-positive bacteria and enterobacteria) and with a metabarcoding

approach based on 16S sequencing targeting the bacteria in cheeses at several stages of

maturation. The abundance and identity of the microorganisms studied (Supplementary

Figures 1A and 1B) were similar to those in four commercial Roquefort cheeses (Devoyod *et*

al., 1968; personal information from C. Callon) and closely related blue cheeses

(Diezhandino *et al.,* 2015). Based on microbial counts, we found no significant effect of *P.*

roqueforti population on the abundance of any of the counted microorganisms, including

molds (i.e., mainly *P. roqueforti),* at any stage of maturation (Supplementary Table 1A).

The metabarcoding approach targeting bacteria identified mostly sequences from the

Lactococcus and *Leuconostoc spp.* starters, which are responsible for acidification and cavity

formation in the cheese, respectively. The remaining sequences corresponded to 12 bacterial

genera frequently found in unpasteurized milk cheeses, such as *Lactobacillus,*

Staphylococcus and *Arthrobacter*. However, the large predominance of starters made it

impossible to obtain sufficient data for other bacteria to assess differences in the abundance

of particular bacteria between cheeses made with strains from the four *P. roqueforti*

populations (Supplementary Table 1B). We estimated three OTU (operational taxonomic

unit) diversity parameters based on bacterial barcode sequence abundances, to measure OTU

richness and/or evenness. Bray-Curtis dissimilarity showed that cheeses made with strains

from the same *P. roqueforti* population were no more similar than those made with strains

from different *P. roqueforti* populations. However, we found a significant effect of *P.*

roqueforti population, in addition to a stage effect, on the Shannon and Simpson diversity

indices. Cheeses made with strains from the cheese *P. roqueforti* populations tended to have a

higher bacterial OTU diversity, particularly at nine days of maturation and for the Roquefort

population (Supplementary figure 2A and 2B), although the post-hoc analyses were not

powerful enough to detect significant pairwise differences (Supplementary Table 1B). The

differences in cheese bacterial diversity, although minor, suggest that the differences between

cheeses made with strains from the four *P. roqueforti* populations may be due not only to a

direct effect of *P. roqueforti* population, but also to an indirect effect mediated by the

induction of bacterial communities of different diversities. There may also be undetected

differences at species level or for low-abundance microorganisms that might nevertheless

have substantial effects. However, even if this were the case, it would constitute an indirect

effect of the *P. roqueforti* population, as this was the only difference during our

cheesemaking process.

Higher proportion of blue area in cheeses produced with cheese *P. roqueforti*

populations. We estimated the percentage of the cheese area that was blue on fresh inner

cheese slices. The blue veins depend on the formation of cavities in the cheese, and the

growth and sporulation of *P. roqueforti* in these cavities. The percentage blue area was

significantly larger in cheeses produced with cheese population strains than in those produced

with non-cheese population strains (Figure 3; Supplementary Table 1C). We also found a

significant decrease in blue area from 20 to 180 days of maturation, for all populations except

the Roquefort population, for which the percentage blue area remained approximately

constant (Figure 3; Supplementary Table 1C).

More efficient proteolysis and lipolysis by the Roquefort *P. roqueforti* population. The *P.*

roqueforti strains used for cheesemaking are known to have high proteolytic and lipolytic

activities, which play a key role in cheese ripening. We therefore investigated the proteolysis

and lipolysis efficiencies of the four populations. Both targeted and non-targeted

chromatographic analyses showed that proteolysis efficiency was highest in the Roquefort *P.*

roqueforti population. We performed the targeted analysis with standards for the principal 23

amino acids (Supplementary Table 2A). We found that eight amino acids discriminated

significantly between cheeses made with the different *P. roqueforti* populations

(Supplementary Table 1D), 15 discriminated between the cheese and non-cheese populations

and 14 distinguished between the Roquefort and non-Roquefort populations (Supplementary

Figure 4A). The cheeses made with strains from cheese populations, and from the Roquefort

population, in particular, had a higher total amino-acid concentration (Supplementary Tables

1D and 2B).

We also assessed proteolysis activity in a non-targeted analysis (fingerprint approach) on

whole chromatograms (8,364 signals), which provided much more powerful discrimination

between metabolites. Each metabolite generates a signal specific to its mass-to-charge (m/z)

ratio at a given retention time. We obtained the largest number of aqueous signals, indicating

the most efficient proteolysis, in cheeses inoculated with strains from the Roquefort

population, followed by the lumber and non-Roquefort cheese populations, which were not

significantly different from each other, and proteolysis was least efficient for the silage

population (Figure 4; Supplementary Table 1E).

Lipolysis was also more efficient for the Roquefort population than for the other populations.

We investigated whether the *P. roqueforti* population influenced the amounts of free fatty

acids and residual glycerides, as a proxy for lipolysis efficiency, in 90-day cheeses, with

targeted and non-targeted chromatographic analyses in the positive and negative ionization

modes. We specifically targeted glycerides and free fatty acids. In the targeted analysis, we

identified seven free fatty acids and 20 triglycerides, and found that three free fatty acids

were significantly more concentrated in cheeses made with Roquefort strains than in those

made with strains from non-Roquefort populations (Supplementary Table 1F). In the non-

targeted analysis, we obtained 3,094 signals and observed higher amounts of organic signals

specific to free fatty acids, indicating the most efficient lipolysis, in cheeses made with strains

from the Roquefort population, followed by the lumber and non-Roquefort cheese

populations, which were very similar to each other, with lipolysis efficiency lowest for

cheeses made with strains from the silage population (Figure 5; Supplementary Table 1G).

For residual glycerides, we obtained 8,472 signals, with no significant difference between the

populations (Supplementary Figure 5; Supplementary Table 1H).

As expected, we observed a maturation stage effect for 11 of the 16 physicochemical

parameters (Supplementary Table 1I). Non-protein nitrogenous content was significantly

higher in cheeses inoculated with strains from cheese *P. roqueforti* populations than in

cheeses inoculated with strains from the other populations, consistent with the greater

efficiency of proteolysis associated with these strains (Supplementary Figure 6A). Cheese

water activity differed significantly between the cheeses made with strains from the four *P.*

roqueforti populations (Supplementary figure 6B): it was significantly lower for the

Roquefort cheese population than for the non-Roquefort cheese and silage populations

(Supplementary Table 1I).

Strong influence of *P. roqueforti* population on volatile compound production. We

investigated the effect of *P. roqueforti* population on cheese aroma and flavor, by

determining the relative abundance of the most relevant volatile compounds in 90-day

cheeses. We focused on the GC-MS data for the 40 principal volatile compounds considered

to be markers of the aromatic quality of blue cheeses (Rothe *et al.,* 1982): 11 acids, 12

ketones, 10 esters, six alcohols and one aldehyde (Supplementary Table 3). We found that *P.*

roqueforti population strongly influenced the amounts of the compounds from these aromatic

families in the cheeses (Supplementary Table 1J; Figures 6 and 7). Indeed, the odors of the

cheeses differed considerably: the cheeses made with strains from the cheese *P. roqueforti*

populations smelled as good as typical ripened blue cheeses, whereas those made with strains

from non-cheese *P. roqueforti* populations had unpleasant odors, similar to those of a wet

swab (Supplementary Figure 7; personal observation).

The amounts of acids, methyl ketones and secondary alcohols resulting from proteolysis and

lipolysis, and contributing to the typical flavor of blue cheese, were larger in cheeses

produced with strains from cheese populations than in those produced with strains from non-

cheese populations. These compounds were present in particularly large amounts in cheeses

made with strains from the Roquefort population. Four of the 40 compounds analyzed were

proteolysis by-products (primary alcohols: 3-methyl-butanal, 3-methyl-butanol and

isopropyl-alcohol, named hereafter alcohols I, and 3-methyl-butanoic acid, named hereafter

acid I; Supplementary Table 3). The abundance of alcohols I was significantly higher in

cheeses made with strains from cheese *P. roqueforti* populations than in those made with

strains from non-cheese populations, and the highest values were obtained for the Roquefort

population (Supplementary Table 1J). Acid I was also present in larger amounts in cheeses

made with strains from the Roquefort population than in other cheeses. Two acids, by-

products of glycolysis (named hereafter acids II), were present in larger amounts in cheeses

made with strains from the Roquefort and lumber/food spoiler *P. roqueforti* populations than

in other cheeses (Supplementary Tables 1J and 3). The other 35 aromatic compounds (i.e.

acids from beta-oxidation, named hereafter acids III, ketones, secondary alcohols named

hereafter alcohols II, and esters) were almost all direct or indirect by-products of lipolysis

(Supplementary Table 3). The abundance of acids III was higher in cheeses made with strains

from the Roquefort and lumber/food spoiler populations than in cheeses made with strains

from the non-Roquefort cheese population. The levels of these compounds were lowest in

cheeses made with strains from the silage population. Larger amounts of esters and methyl

ketones (especially 2-pentanone and 2-heptanone) were found in cheeses made with strains

from cheese *P. roqueforti* populations (Supplementary Table 1J). Cheeses made with strains

from the Roquefort population contained the largest amounts of methyl ketones, and these

compounds were barely detectable in cheeses made from silage population strains (Figure

7A). The levels of alcohols II, particularly 2-heptanol, were also much higher in cheeses

made with Roquefort population strains than in other cheeses (Supplementary Table 1J;

Figure 7B).

Discussion

Cheese *P. roqueforti* populations have been selected to produce better blue cheeses.

Measurements of multiple features of blue cheeses made under conditions resembling those

typically used in commercial Roquefort production revealed a strong influence of the

differentiated *P. roqueforti* populations on cheese quality, with the cheese populations

appearing the best adapted to cheesemaking, in terms of both the appearance and aromatic

quality of the resulting cheese. The differences between the four *P. roqueforti* populations

and the more appealing cheeses produced with strains from the cheese populations suggest

that humans have exerted selection for the production of better cheeses, either on standing

variation or on *de novo* mutations, and this corresponds to domestication. Indeed, we found

that cheese *P. roqueforti* strains produced a larger percentage blue area on cheese slices, an

important visual aspect of blue cheeses. We also found that proteolysis and lipolysis were

more efficient in cheeses made with Roquefort population strains than in cheeses made with

strains from the other *P. roqueforti* populations, resulting in the production of larger amounts

of desirable volatile compounds, including alcohols and associated acids. Cheese water

activity was lower in cheeses made with strains from the Roquefort population, probably due

to more efficient proteolysis (Ardö *et al.*, 2017). We found no significant difference in the

identities and abundances of microorganisms between the cheeses made with strains from the

four *P. roqueforti* populations. Some minor differences in species diversity were observed,

however, and the differences between cheeses probably reflected a direct effect of the

specific features of the *P. roqueforti* population, although minor indirect effects involving the

induction of more diverse bacterial communities by cheese *P. roqueforti* strains may also

have occurred. Overall, our findings strongly support the view that cheese *P. roqueforti*

populations have been selected by humans for better appearance and aroma. This selection

may have involved the choice of the most beneficial strains for making good cheeses from

standing variation, and/or the selection of *de novo* genetic changes. Previous studies found

footprints of genomic changes in cheese populations in the form of beneficial horizontal gene

transfers and positive selection (Dumas *et al.*, 2020; Ropars *et al.*, 2015).

Previous studies reported differences between *P. roqueforti* populations, in terms of growth,

lipolysis and proteolysis, but on synthetic media (Dumas *et al.*, 2020; Ropars *et al.*, 2015).

Here, using experimental cheeses made in commercial cheese production conditions, we

reveal important features specific to cheese *P. roqueforti* populations, and to the Roquefort

and non-Roquefort cheese populations. These findings are important in the context of

domestication, for understanding rapid adaptation and diversification, and future studies

based on quantitative trait mapping may be able to identify further genomic changes

responsible for the specific features of the populations, according to the contrasting

phenotypes revealed here. Progenies can indeed be obtained from crosses between strains

from different populations of *P. roqueforti* (Ropars *et al.*, 2015), and this could facilitate

strain improvement through recombination between the different populations. Our results are,

therefore, also important for improving blue cheese production.

The four *P. roqueforti* populations induce similar microbiotas, but water availability is lower with cheese population strains, restricting the occurrence of spoiler microorganisms. Based on microbiological counts, we found no significant differences in abundance for any of the species monitored between cheeses made with strains from the four populations of *P. roqueforti*. In particular, we found no significant difference in the abundance of molds on Petri dishes. However, microbiological counts are known to provide poor estimates of fungal biomass, especially for mycelium growth (Schnurer, 1993).

The metabarcoding approach suggested that the different *P. roqueforti* populations induced bacterial communities of different levels of diversity. The cheese populations, and the Roquefort population in particular, were associated with the highest level of diversity. The large predominance of bacterial starters made it impossible to collect sufficient data for an assessment of the differences in relative abundance between subdominant bacterial species on the basis of metabarcoding. We also found a significant difference in water activity between cheeses made with strains from the four *P. roqueforti* populations, the lowest value obtained being that for the Roquefort population. This may also reflect human selection, as low water activity restricts the occurrence of spoiler microorganisms, and is therefore highly controlled for Roquefort cheese sales, particularly those for export.

Cheese *P. roqueforti* populations produce bluer cheeses. We found significantly higher percentage blue areas in cheese slices from cheeses made with cheese *P. roqueforti* strains than in those made from non-cheese strains, potentially reflecting greater *P. roqueforti* growth in cheese and/or a higher sporulation efficiency in cavities. The percentage blue area in cheese slices also depends on the formation of cavities in the cheese, as *P. roqueforti* can

only sporulate in cavities in which oxygen is available. The cavities are mostly generated by

the gas-producing bacterium *Leuconostoc mesenteroides*, the abundance of which did not

differ between the cheeses made with strains from different *P. roqueforti* populations,

suggesting a direct effect of *P. roqueforti* populations on the blueness of cheese slices. The

significantly higher percentage blue area in slices of cheese made with cheese *P. roqueforti*

strains than in those made with non-cheese strains therefore probably reflects better cheese

and cavity colonization and sporulation, probably due to selection on the basis of appearance.

The percentage blue area decreased by the end of maturation, perhaps due to the death of the

fungus. Only cheeses made with Roquefort strains retained a high percentage blue area at 90

days of maturation, again potentially reflecting selection in pre-industrial times, when

Roquefort cheeses had to be stored for several months at cave temperature before sale. The

minimum maturation time for Roquefort PDO remains 90 days, which is longer than for other

blue cheeses. These findings contrast with a previous study showing that non-Roquefort

population colonized the cavities of model cheeses better than other populations (Dumas *et*

al., 2020); this discrepancy may reflect differences between studies in terms of the

measurements used (total percentage blue area versus percentage blue area within cavities),

the type of milk (ewe versus goat) or the mode of cheesemaking (rudimentary models versus

commercial-like cheeses). Our findings are consistent with the presence of horizontally

transferred genes in cheese populations with predicted functions in fungal development,

including sporulation and hyphal growth (Dumas *et al.*, 2020).

Proteolysis and lipolysis are more efficient in the Roquefort *P. roqueforti* population.

Based on chemical analyses and powerful chromatographic discrimination methods, we

showed that the abundance of amino acids and small peptides (i.e., residual products of

proteolysis) was highest in cheeses made with Roquefort *P. roqueforti* strains. Thus, these

strains had the highest capacity for proteolysis, which is an important process in

cheesemaking. Indeed, proteolysis contributes to the development of cheese texture, flavors

and aromas (Ardö *et al.*, 2017; Andersen *et al.*, 2010; McSweeney, 1997; Roudot-Algaron,

1996; Ardö, 2006). Previous measurements of proteolytic activity in synthetic media detected

significant differences between *P. roqueforti* populations, but not between the two cheese

populations (Dumas *et al.*, 2020). We show here that experimental cheeses made with strains

from the Roquefort population have a higher content of residual products of proteolysis, a

sign of more advanced ripening.

We also found that lipolysis was more efficient in the cheeses made with strains from the

Roquefort *P. roqueforti* population. By contrast, previous studies in synthetic media found

that lipolysis was most efficient in the non-Roquefort population (Dumas *et al.*, 2020). The

discrepancy between these studies demonstrates the need for measurements in real cheeses

for the reliable assessment of metabolic activities. Lipolytic activity is known to affect cheese

texture and the production of volatile compounds affecting pungency (Alonso *et al.*, 1987;

González De Llano *et al.*, 1990, 1992; Martín *et al.*, 2016; Thierry *et al.*, 2017; Woo *et al.*,

1984). The more efficient proteolysis and lipolysis in the Roquefort *P. roqueforti* population

should have a strong impact on cheese texture and flavor. It therefore probably results from

selection to obtain better cheeses, i.e. from a domestication process, as previously reported

for other fungi (Almeida *et al.*, 2014; Baker *et al.*, 2015; Gallone *et al.*, 2016; Gibbons *et al.*,

2012; Gonçalves *et al.*, 2016; Libkind *et al.*, 2011; Sicard *et al.*, 2011). Roquefort cheeses are

widely considered to be the blue cheeses with the strongest aromas and flavours; the less

efficient lipolysis and proteolysis in the non-Roquefort population may result from more

recent selection for milder cheeses.

Cheese *P. roqueforti* populations produce cheeses with better flavor and aromas. We

found major differences between the cheeses made with strains from different *P. roqueforti*

populations, in terms of the volatile compounds resulting from lipolysis and, to a lesser

extent, also from proteolysis. Only four of the aromatic compounds detected in our cheeses

(3-methyl-butanal, 3-methyl-butanol, isopropyl-alcohol and 3-methyl-butanoic acid) were by-

products of casein proteolysis (McSweeney *et al.,* 2000), and the concentrations of these

molecules were significantly higher in cheeses made with Roquefort *P. roqueforti* strains,

consistent with the higher proteolysis efficiency and amino-acid precursor (i.e. valine, leucine

and isoleucine) concentrations of these strains. These compounds produce fruity (banana),

cheesy and alcoholic notes, which were probably important selection criteria during the

domestication of the Roquefort *P. roqueforti* population. For the products of metabolic

pathways leading from amino acids to alcohols (Ehrlich pathway with aldehyde reduction) or

acids (aldehyde oxidation; Ganesan *et al.,* 2017), the higher concentration of alcohols than of

acids observed for all populations is consistent with the general micro-aerobic conditions of

blue cheese cavities.

Most of the aromatic compounds identified were direct or indirect by-products of lipolysis,

consistent with the known key role of lipolysis in the generation of typical blue cheese aroma

(Cerning *et al.,* 1987; Collins *et al.,* 2003). The aromatic compounds resulting from lipolysis

belonged to four chemical families (acids, methyl ketones, secondary alcohols and esters).

Methyl ketones were the most diverse and abundant for cheese *P. roqueforti* populations,

particularly for the Roquefort population, in which 2-pentanone and 2-heptanone were

present in the largest amounts; 2-heptanone underlies the characteristic "blue cheese" sensory

descriptor (González De Llano *et al.,* 1990, 1992; Moio *et al.,* 2000; Anderson *et al.,* 1966).

In *P. roqueforti,* methyl ketones with odd numbers of carbons are mostly produced by fatty-

acid beta-oxidation, whereas those with even numbers of carbons may be produced by the

beta-oxidation or autoxidation of fatty acids (Spinnler, 2011). These compounds are produced

by the decarboxylation of hexanoic acid and octanoic acid, respectively, which were the most

abundant acids found in our cheeses. This reaction is considered to be a form of

detoxification, because methyl ketones are less toxic than acids (Kinderlerer, 1993; Spinnler,

2011). Interestingly, this pathway appeared to be more active in the cheese *P. roqueforti*

populations, as methyl ketone levels were lower in cheeses made with lumber (four-fold

difference) and silage (10-fold lower) strains than in cheeses made with cheese population

strains. Methyl ketone concentrations were not directly associated with the concentrations of

their precursors (acids), the highest concentrations being found in the lumber and Roquefort

populations. The biosynthesis pathway producing methyl ketones must, therefore, be more

efficient in cheese populations, particularly the non-Roquefort population. The cheese *P.*

roqueforti populations were probably selected for their higher acid detoxification capacity, as

this produces aromatic compounds with a very positive impact on flavour (Spinnler, 2011).

The concentrations of secondary alcohols (resulting from the reduction of methyl ketones)

were also higher in cheeses produced by cheese *P. roqueforti* strains, particularly those of the

Roquefort population, for which they were seven times higher than for the non-Roquefort

cheese population and 20 times higher than for the silage/lumber populations; 2-heptanol was

the major alcoholic compound produced. The reduction of 2-heptanone to 2-heptanol occurs

specifically in anaerobic conditions and is much stronger in the Roquefort population; aerobic

conditions were similar for all the populations. The Roquefort *P. roqueforti* population may

also have been selected for this feature, as secondary alcohols provide "fruity notes", which

are associated with better aromatic quality (Spinnler, 2011). Methyl ketones may be reduced

to alcohols by an alcohol dehydrogenase, as occurs when aldehyde is reduced to alcohol via

the Ehrlich pathway. Alcohol dehydrogenase genes may thus have been targets of selection in

the Roquefort *P. roqueforti* population, although they were not detected as evolving under

positive selection in a previous study (Dumas *et al.*, 2020).

We also found higher levels of esters in cheeses made with cheese *P. roqueforti* populations.

Esters are produced principally by the esterification of ethanol with acids generated by beta-

oxidation. *Leuconostoc* starters can produce ethanol, and ester synthesis has also been

described as a detoxification mechanism (Mason *et al.*, 2000). These results further indicate

that cheese *P. roqueforti* populations, particularly the Roquefort population, have been

selected for acid detoxification capacity, leading to a large large variety of less toxic aromatic

compounds with strong aromas and flavors.

Overall, the aromas of cheeses made with cheese *P. roqueforti* strains had more appealing

aromas, and this was particularly true for cheeses made with Roquefort strains. These aroma

properties probably reflect selection by humans. The cheeses made with silage and lumber

populations had a mild unpleasant smell, whereas those made with cheese strains smelled like

typical blue cheeses, with cheeses made with Roquefort strains having the strongest smell.

This may reflect previously reported horizontal gene transfers in cheese populations,

involving genes with predicted functions in lipolysis or amino-acid catabolism, and the

positive selection of genes involved in aroma production (Dumas *et al.*, 2020). We compared

P. roqueforti populations between cheeses made following commercial modes of production,

which represents a major advance relative to previous studies based on experimental models

or synthetic media (Gillot *et al.*, 2017; Dumas *et al.*, 2020). We used unpasteurized ewe's

milk, in accordance with the requirements for Roquefort PDO production, which also affects

cheese aromas. In future studies, it would be interesting to determine whether the use of

pasteurized or unpasteurized ewe's milk or cow's milk leads to similar specific features of the

Roquefort versus non-Roquefort cheese *P. roqueforti* populations, as there may have been

selection during domestication, leading to an adaptation of the Roquefort population for the

catabolism of unpasteurized ewe's milk.

Conclusion. We showed that the *P. roqueforti* population had a strong impact on cheese

quality, appearance and aroma. The populations used for cheesemaking led to bluer cheeses,

with better aromas, probably due to domestication involving the selection of multiple fungal

traits by humans seeking to make the best possible cheeses. French cheese producers have

been inoculating cheeses with *P. roqueforti* spores from moldy rye bread since the end of the

19[th] century (Labbe *et al.,* 2004, 2009; Vabre, 2015). This process made it possible for them

to re-inoculate with the strains producing the best cheeses, thereby applying a strong selection

pressure. The two cheese populations displayed a number of specific features, with the

Roquefort population notably producing more intense and specific aromas and flavors. The

selection of different fungal varieties for different usages has also been reported in the

fermenting yeast *Saccharomyces cerevisiae* (Gallone *et al.,* 2016; Legras *et al.,* 2018).

Previous studies on *P. roqueforti* detected recurrent changes in amino acids and horizontal

gene transfers in cheese populations, both of which facilitated rapid adaptation (Dumas *et al.,*

2020; Ropars *et al.,* 2015). Our findings provide greater insight into *P. roqueforti*

domestication and pave the way for strain improvement through the targeting of relevant

traits. A protocol inducing sexual selection has been developed in *P. roqueforti* (Ropars *et al.,*

2014), making it possible to perform crosses between strains from the two cheese

populations, each of which harbors very little genetic diversity (Dumas *et al.,* 2020), to

generate variability and to identify strains with high levels of performance; the results of this

study will facilitate the choice of the parental strains for crossing and of the most important

phenotypes to be measured in the offspring. Parental strains with strongly contrasting

phenotypes for the traits important for cheesemaking that we found to be differentiated

between populations (such as volatile compound production, lipolysis and proteolysis) should

be used, to maximize variability in the progeny.

References

Almeida, M., Hébert, A., Abraham, A.-L., Rasmussen, S., Monnet, C., Pons, N., Delbès-Paus, C., Loux, V., Batto, J.-M., Leonard, P., Kennedy, S., Ehrlich, S., Pop, M., Montel, M.-C., Irlinger, F. & Renault, P. (2014). Construction of a dairy microbial genome catalog opens new perspectives for the metagenomic analysis of dairy fermented products. *BMC Genomics*, *15*, 1101.

Alonso, L., Juarez, M., Ramos, M., & Martin Alvarez, P. J. (1987). Effects of changes during ripening and frozen storage on the physicochemical and sensory characteristics of Cabrales cheeses. *International Journal of Food Science & Technology*, *22*(5), 525–534.

Andersen, L. T., Ardö, Y., & Bredie, W. L. P. (2010). Study of taste-active compounds in the water-soluble extract of mature Cheddar cheese. *International Dairy Journal*, *20*(8), 528–536.

Andersson, A. F., Lindberg, M., Jakobsson, H., Bäckhed, F., Nyrén, P., & Engstrand, L. (2008). Comparative analysis of human gut microbiota by barcoded pyrosequencing. *PLoS ONE*, *3*(7), e2836.

Anderson, D. F., & Day, E. A. (1966). Quantitation, evaluation, and effect of certain microorganisms on flavor components of blue cheese. *Journal of Agricultural and Food Chemistry*, *14*(3), 241–245.

Ardö, Y., McSweeney, P. L. H., Magboul, A. A. A., Upadhyay, V. K., & Fox, P. F. (2017). Biochemistry of cheese ripening: proteolysis. In P. L. H. McSweeney, P. F. Fox, P. D. Cotter, & D. W. Everett (Eds.), *Cheese: Chemistry, Physics and Microbiology* (Fourth Edi, pp. 445–482). Elsevier Ltd.

Ardö, Y. (2006). Flavour formation by amino acid catabolism. *Biotechnology Advances*, *24*(2), 238–242.

Auguie, B. (2017). gridExtra: miscellaneous functions for "Grid" graphics. Retrieved from https://cran.r-project.org/web/packages/gridExtra/index.html

Baker, E. P., Wang, B., Bellora, N., Peris, D., Hulfachor, A. B., Koshalek, J. A., Libkind, D. & Hittinger, C. T.(2015). The genome sequence of *Saccharomyces eubayanus* and the domestication of Lager-brewing yeasts. *Molecular Biology and Evolution*, *32*(11), 2818–2831.

Bates, D., Mächloer, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1), 1–48.

Bigelis, R. (2001). Fungal fermentation: industrial. *Encyclopedia of Life Sciences*, 1–8.

Bjørn-Helge, M., Wehrens, R., & Hovde Liland, K. (2018). *pls: partial least squares and principal component regression* (R package version 2.7-0). Retrieved from https://cran.r-project.org/package=pls

Cerning, J., Gripon, J.-C., Lamberet, G., & Lenoir, J. (1987). Les activités biochimiques des *Penicillium* utilisés en fromagerie. *Le Lait*, *67*(1), 3–39.

Cheese and processed cheese: determination of total solids content. International standards
4a. (1982). *International Dairy Federation, Brussels, Belgium*. Brussels, Belgium.
http://doi.org/ISO 5534:2004 [IDF 4:2004]

Cheese and processed cheese: determination of chloride content potentiometric titration
method. International standards 4a. International Dairy Federation, Brussels, Belgium. 1988.
https://www.iso.org/standard/43922.html. ISO 5943:2006 [IDF 88:2006]

Cheese and processed cheese: determination of chloride content potentiometric titration
method. International standards 4a. *International Dairy Federation, Brussels, Belgium*.
Brussels, Belgium. http://doi.org/ISO 5943:2006 [IDF 88:2006]

Cheeseman, K., Ropars, J., Renault, P., Dupont, J., Gouzy, J., Branca, A., *et al.*. (2014).
Multiple recent horizontal transfers of a large genomic region in cheese making fungi. *Nature*
Communications, 5, 2876.

Collins, Y. F., McSweeney, P. L. H., & Wilkinson, M. G. (2003). Lipolysis and free fatty
acid catabolism in cheese: a review of current knowledge. *International Dairy Journal*,
13(11), 841–866.

De Man, J. C., Rogosa, M., & Sharpe, M. E. (1960). A medium used for the cultivation of
Lactobacilli. *Journal of Applied Bacteriology*, *23*, 130–135.

Denis, C., Gueguen, M., Henry, E., & Levert, D. (2001). New media for the numeration of
cheese surface bacteria. *Le Lait, 81*(3), 365–379.

Devoyod, J. J., & Muller, M. (1969). La flore microbienne du fromage de Roquefort III. Les
streptocoques lactiques et les *Leuconostoc* ; Influence de différents micro-organismes de
contamination. *Le Lait, 487,* 369–399. Retrieved from https://doi.org/10.1051/lait:196948715

Diezhandino, I., Fernández, D., González, L., McSweeney, P. L. H., & Fresno, J. M. (2015).
Microbiological, physico-chemical and proteolytic changes in a spanish blue cheese during
ripening (Valdeón cheese). *Food Chemistry, 168*, 134–141.

Dumas, É., Feurtey, A., Rodríguez de la Vega, R. C., Le Prieur, S., Snirc, A., Coton, M.,
Thierry, A., Coton, E., Le Piver, M., Roueyre, D., Ropars, J., Branca, A. & Giraud, T. (2020).
Independent domestication events in the blue-cheese fungus *Penicillium roqueforti*.
Molecular Ecology, 33913(January), 451773.

Dupont, J., Dequin, S., Giraud, T., Le Tacon, F., Masit, S., Ropars, J., Richard, F. & Selosse,
M.-A. (2017). Fungi as a source of food. *Microbiology Spectrum, 5*(3), 1–22.

Duval, P., Chatelard-Chauvin, C., Gayard, C., Rifa, E., Bouchard, P., Hulin, S., Picque, D. &
Montel, M.-C. (2016). Microbial dynamics in industrial blue veined cheeses in different
packaging. *International Dairy Journal, 56,* 198–207.

Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., & Knight, R. (2011). UCHIME
improves sensitivity and speed of chimera detection. *Bioinformatics, 27*(16), 2194–2200.

Escudié, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., Maman, S.,
Hernandez-Raquet, G., Combes, S. & Pascal, G. (2018). FROGS: Find, Rapidly, OTUs with
Galaxy Solution. *Bioinformatics, 34*(8), 1287–1294.

Fleming, A. (1929). On the antibacterial action of cultures of a *Penicillium,* with special
reference to their use in the isolation of *B. influenzæ. British Journal of Experimental*
Pathology, 10(3), 226–236.

Fox, J., & Weisberg, S. (2019). *An {R} Companion to applied Regression.* (T. Oaks, Ed.)
(Third). Sage. Retrieved from https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Gallone, B., Steensels, J., Prahl, T., Soriaga, L., Saels, V., Herrera-Malaver, B., Merlevede A,
Roncoroni M, Voordeckers K, Miraglia L., Teiling, C., Steffy, B., Taylor, M., Schwartz, A.,
Richardson, T., White, C., Baele, G., Maere, S., & Verstrepen, K. J. (2016). Domestication
and divergence of *Saccharomyces cerevisiae* beer yeasts. *Cell, 166*(6), 1397–1410.

Ganesan, B., & Weimer, B. C. (2017). Amino acid catabolism and its relationship to cheese
flavor outcomes. In P. L. H. McSweeney, P. F. Fox, P. D. Cotter, & D. W. Everett (Eds.),
Cheese: Chemistry, Physics and Microbiology: Fourth Edition (Fourth Edi, Vol. 1, pp. 483–
516). Elsevier Ltd.

Gatto, L., & Lilley, K. S. (2012). Msnbase-an R/Bioconductor package for isobaric tagged
mass spectrometry data visualization, processing and quantitation. *Bioinformatics, 28*(2),
288–289.

Gibbons, J. G., Salichos, L., Slot, J. C., Rinker, D. C., McGary, K. L., King, J. G., Klich
M.A., Tabb D.L., McDonald W.H. & Rokas, A. (2012). The evolutionary imprint of
domestication on genome variation and function of the filamentous fungus *Aspergillus*
oryzae. Current Biology, 22(15), 1403–1409.

Gillot, G., Jany, J.-L., Coton, M., Le Floch, G., Debaets, S., Ropars, J., López-Villavicencio
M., Dupont J., Branca A., Giraud T. & Coton, E. (2015). (Fifty shades of blue :) Insights into
Penicillium roqueforti morphological and genetic diversity. *PLoS ONE, 10*(6), e0129849.

Gillot, G., Jany, J.-L., Poirier, E., Maillard, M., Debaets, S., Thierry, A., Coton E. & Coton,
M. (2017). Functional diversity within the *Penicillium roqueforti* species. *International*
Journal of Food Microbiology, 241, 141–150.

Giraud, T., Koskella, B., & Laine, A.-L. (2017). Introduction: microbial local adaptation:
insights from natural populations, genomics and experimental evolution. *Molecular Ecology,*
26(7), 1703–1710.

Gladieux, P., Ropars, J., Badouin, H., Branca, A., Aguileta, G., De Vienne, D. M., Rodríguez
De La Vega R.C., Branco S. & Giraud, T. (2014). Fungal evolutionary genomics provides
insight into the mechanisms of adaptive divergence in eukaryotes. *Molecular Ecology, 23*(4),
753–773.

Gomri, G. (1946). Buffers in the range of pH 6.5 to 9.6. *Proceedings of Society for*
Experimental Biology and Medicine, 6(1), 33–34.

Gonçalves, M., Pontes, A., Almeida, P., Barbosa, R., Serra, M., Libkind, D., Hutzler M.,
Gonçalves P. & Sampaio, J. P. (2016). Distinct domestication trajectories in top-fermenting
beer yeasts and wine yeasts. *Current Biology, 26*(20), 2750–2761.

González de Llano, D., Ramos, M., Polo, C., Sanz, J., & Martinez-Castro, I. (1990).
Evolution of the volatile components of an artisanal blue cheese during ripening. *Journal of*
Dairy Science, 73(7), 1676–1683.

González de Llano, D., Ramos, M., Rodriguez, A., Montilla, A., & Juarez, M. (1992).
Microbiological and physicochemical characteristics of Gamonedo blue cheese during
ripening. *International Dairy Journal*, *2*(2), 121–135.

Hébert, A., Forquin-Gomez, M. P., Roux, A., Aubert, J., Junot, C., Heilier, J. F., Landaud S.,
Bonnarmeb P. & Beckerich, J. M. (2013). New insights into sulfur metabolism in yeasts as
revealed by studies of *Yarrowia lipolytica*. *Applied and Environmental Microbiology, 79*(4),
1200–1211.

ISO. (2004). Determination of water activity NF ISO 21807. *International Standardisation*
Organisation. http://doi.org/ISO 21807:2004

Kinderlerer, J. L. (1993). Fungal strategies for detoxification of medium chain fatty acids.
International Biodeterioration and Biodegradation, 32(1–3), 213–224.

Kinsella, J. E., & Hwang, D. H. (1976). Enzymes of *Penicillium roqueforti* involved in the
biosynthesis of cheese flavor. *C R C Critical Reviews in Food Science and Nutrition*, *8*(2),
191–228.

Kopp, B., & Rehm, H. J. (1979). Antimicrobial action of roquefortine. *European Journal of*
Applied Microbiology and Biotechnology, *6*, 397–401 (1979).

Kuhl, C., Tautenhahn, R., Böttcher, C., Larson, T. R., & Neumann, S. (2012). CAMERA: An
integrated strategy for compound spectra extraction and annotation of liquid
chromatography/mass spectrometry data sets. *Analytical Chemistry*, *84*(1), 283–289.

Labbe, M., & Serres, J. P. (2004). *Chroniques du Roquefort - De la préhistoire à l'aube*
industrielle. Grand Imprimeur, La Primaube, France.

Labbe, M., & Serres, J. P. (2009). *Chroniques du Roquefort: des Hommes, des entreprises,*
des marques, période moderne. Grand Imprimeur, La Primaube, France.

Larsen, M. D., & Jensen, K. (1999). The effects of environmental conditions on the lipolytic
activity of strains of *Penicillium roqueforti*. *International Journal of Food Microbiology*,
46(2), 159–166.

Larsen, M. D., Kristiansen, K. R., & Hansen, T. K. (1998). Characterization of the proteolytic
activity of starter cultures of *Penicillium roqueforti* for production of blue veined cheeses.
International Journal of Food Microbiology, *43*(3), 215–221.

Larson, G., & Fuller, D. Q. (2014). The evolution of animal domestication. *Annual Review of*
Ecology, Evolution, and Systematics, *45*, 115–136.

Lazuka, A., Auer, L., Bozonnet, S., Morgavi, D. P., O'Donohue, M., & Hernandez-Raquet,
G. (2015). Efficient anaerobic transformation of raw wheat straw by a robust cow rumen-
derived microbial consortium. *Bioresource Technology*, *196*, 241–249.

Le Boucher, C., Courant, F., Jeanson, S., Chereau, S., Maillard, M., Royer, A. L., Thierry A.,
Dervilly-Pinel G., Le Bizec B. & Lortal, S. (2013). First mass spectrometry metabolic
fingerprinting of bacterial metabolism in a model cheese. *Food Chemistry*, *141*(2), 1032–
1040.

Legras, J.-L., Galeote, V., Bigey, F., Camarasa, C., Marsit, S., Nidelet, T., Couloux A., Guy
J., Franco-Duarte R. Marcet-Houben, M. Gabaldon, T., Schuller, D., Sampaio, J. P. &
Dequin, S. (2018). Adaptation of *S. cerevisiae* to fermented food environments reveals
remarkable genome plasticity and the footprints of domestication. *Molecular Biology and*
Evolution, *35*(7), 1712–1727.

Libkind, D., Hittinger, C. T., Valeŕio, E., Gonçalves, C., Dover, J., Johnston, M., Gonçalves,
P. & Sampaio, J. P. (2011). Microbe domestication and the identification of the wild genetic
stock of lager-brewing yeast. *Proceedings of the National Academy of Sciences of the United*
States of America, *108*(35), 14539–14544.

Liu, Z., Lozupone, C., Hamady, M., Bushman, F. D., & Knight, R. (2007). Short
pyrosequencing reads suffice for accurate microbial community analysis. *Nucleic Acids*
Research, *35*(18), e120.

Ludemann, V., Greco, M., Rodríguez, M. P., Basílico, J. C., & Pardo, A. G. (2010). Conidial
production by *Penicillium nalgiovense* for use as starter cultures in dry fermented sausages
by solid state fermentation. *LWT - Food Science and Technology*, *43*(2), 315–318.

Mahé, F., Rognes, T., Quince, C., de Vargas, C., & Dunthorn, M. (2014). Swarm: robust and
fast clustering method for amplicon-based studies. *PeerJ*, *2*, e593.

Martín, J.-F., & Coton, M. (2016). Blue cheese: microbiota and fungal metabolites. In J.
Frias, C. Martinez-Villaluenga, & E. Peñas (Eds.), *Fermented Foods in Health and Disease*
Prevention (pp. 275–303). Elsevier Inc.

Mason, A. B., & Dufour, J. P. (2000). Alcohol acetyltransferases and the significance of ester
synthesis in yeast. *Yeast*, *16*(14), 1287–1298.

Mayeux, J. V, Sandine, W. E., & Elliker, P. R. (1962). A selective medium for detecting
Leuconostoc organisms in mixed-strain starter cultures. *Journal of Dairy Science*, *45*, 655–
656.

McSweeney, P. L. H., & Sousa, M. J. (2000). Biochemical pathways for the production of
flavour compounds in cheeses during ripening: A review. *Le Lait*, *80*(3), 293–324.

McSweeney, P. L. H., Fox, P. F., & Ciocia, F. (2017). Metabolism of residual lactose and of
lactate and citrate. In P. L. H. McSweeney, P. F. Fox, P. D. Cotter, & D. W. Everett (Eds.),
Cheese: Chemistry, Physics and Microbiology: Fourth Edition (Fourth Edi, pp. 411–421).
Elsevier Ltd.

McSweeney, P. L. H. (1997). The flavour of milk and dairy products: III. Cheese: Taste.
International Journal of Dairy Technology, *50*(4), 123–128.

Moio, L., Piombino, P., & Addeo, F. (2000). Odour-impact compounds of Gorgonzola
cheese. *Journal of Dairy Research*, *67*(2), 273–285.

Moreau, C. (1980). Le *Penicillium roqueforti,* morphologie, physiologie, intérêt en industrie
fromagère, mycotoxines. *Le Lait, 60*(595–596), 254–271.

Mossel, D. A. A., Eelderink, I., Koopmans, M., & Van Rossem, F. (1978). Optimisation of a
MacConkey-type medium for the enumeration of *Enterobacteriaceae*. *Laboratory Practices*,
27, 1049–1050.

Mossel, D. A. A., Kleynen   Semmeling, A. M. C., Vincentie, H. M., Beerens, H., &
Catsaras, M. (1970). Oxytetracycline   Glucose   Yeast extract agar for selective enumeration
of moulds and yeasts in foods and clinical material. *Journal of Applied Bacteriology*, *33*(3),
454–457.

Nelson, F. E. (1940). The effect of the new standard milk agar on the plate count of dairy
products. *Journal of Bacteriology*, *39*(3), 263–272. Retrieved from
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC374570/

Oksanen, J., Blanchet, G. F., Firendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin,
P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M. H. H., Szoecs, E. & Wagner, H.
(2019). vegan: community ecology package. Retrieved from
https://cran.r-project.org/package=vegan

Perrone, G., Samson, R. A., Frisvad, J. C., Susca, A., Gunde-Cimerman, N., Epifani, F., &
Houbraken, J. A. M. P. (2015). *Penicillium salamii,* a new species occurring during seasoning
of dry-cured meat. *International Journal of Food Microbiology*, *193*, 91–98.

Pham, V. T., Lacroix, C., Braegger, C. P., & Chassard, C. (2016). Early colonization of
functional groups of microbes in the infant gut. *Environmental Microbiology*, *18*(7), 2246–
2258.

Pitt, J. I., & Hocking, A. D. (2009). *Fungi and food spoilage* (Springer). Springer, Boston,
MA.

Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile
open source tool for metagenomics. *PeerJ, 4,* e2584.

Ropars, J., Cruaud, C., Lacoste, S., & Dupont, J. (2012). A taxonomic and ecological
overview of cheese fungi. *International Journal of Food Microbiology*, *155*(3), 199–210.

Ropars, J., Lo, Y.-C., Dumas, É., Snirc, A., Begerow, D., Rollnik, T., Lacoste, S., Dupont, J.,
Giraud, T. & López-Villavicencio, M. (2016). Fertility depression among cheese-making
Penicillium roqueforti strains suggests degeneration during domestication. *Evolution*, *70*(9),
2099–2109.

Ropars, J., López-Villavicencio, M., Dupont, J., Snirc, A., Gillot, G., Coton, M., Jany, J.-L.,
Coton, E. & Giraud, T. (2014). Induction of sexual reproduction and genetic diversity in the
cheese fungus *Penicillium roqueforti*. *Evolutionary Applications*, *7*(4), 433–441.

Ropars, J., López-Villavicencio, M., Snirc, A., Lacoste, S., & Giraud, T. (2017). Blue cheese-
making has shaped the population genetic structure of the mould *Penicillium roqueforti*.
PLoS ONE, 12(3), e0171387.

Ropars, J., Rodríguez de la Vega, R. C., Gouzy, J., Dupont, J., Swennen, D., Dumas, É.,
Giraud, T. & Branca, A. (2016). Diversity and mechanisms of genomic adaptation in
Penicillium. In R. P. de Vries, I. Benoit Gelber, & A. Rordam (Eds.), *Aspergillus and*
Penicillium in the post-genomic era (HAL). Caister Academic Press. http://doi.org/⟨hal–
01302706⟩

Ropars, J., Rodríguez de la Vega, R. C., López-Villavicencio, M., Gouzy, J., Sallet, E.,
Dumas, É., Lacoste, S., Debuchy, R., Dupont, J., Branca, A. & Giraud, T. (2015). Adaptive
horizontal gene transfers between multiple cheese-associated fungi. *Current Biology, 25*(19),
2562–2569.

Rothe, M., Engst, W., & Erhardt, V. (1982). Studies on characterization of blue cheese
flavour. *Die Nahrung, 26*(7/8), 591–602.

Roudot-Algaron, F. (1996). Le goût des acides aminés, des peptides et des protéines:
Exemple de peptides sapides dans les hydrolysats de caséines. *Lait, 76*(4), 313–348.

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years
of image analysis. *Nature Methods, 9*(7), 671–675.

Schnurer, J. (1993). Comparison of methods for estimating the biomass of three food-borne
fungi with different growth patterns. *Applied and Environmental Microbiology, 59*(2), 552–
555.

Searle, S. R., Speed, F. M., & Milliken, G. A. (1980). Population marginal means in the
linear model: An alternative to least squares means. *American Statistician, 34*(4), 216–221.

Sicard, D., & Legras, J.-L. (2011). Bread, beer and wine: Yeast domestication in the
Saccharomyces sensu stricto complex. *Comptes Rendus - Biologies, 334*(3), 229–236.

Smith, C. A., Want, E. J., O'Maille, G., Abagyan, R., & Siuzdak, G. (2006). XCMS:
Processing mass spectrometry data for metabolite profiling using nonlinear peak alignment,
matching, and identification. *Analytical Chemistry, 78*(3), 779–787.

Snow, G. (2016). TeachingDemos: Demonstrations for teaching and learning. Retrieved from
https://cran.r-project.org/web/packages/TeachingDemos/index.html

Spinnler, H.-E. (2011). Rôle des lipides dans la perception olfactive des produits laitiers.
Sciences Des Aliments, 30, 103–120.

Terzaghi, B. E., & Sandine, W. E. (1975). Improved medium for lactic streptococci and their
bacteriophages. *Applied Microbiology, 29*(6), 807–813.

Thierry, A., Collins, Y. F., Abeijón Mukdsi, M. C., McSweeney, P. L. H., Wilkinson, M. G.,
& Spinnler, H.-E. (2017). Lipolysis and metabolism of fatty acids in cheese. In P. L. H.
McSweeney, P. F. Fox, P. D. Cotter, & D. W. Everett (Eds.), *Cheese: Chemistry, Physics and*
Microbiology: Fourth Edition (Fourth Edi, pp. 423–444). Elsevier Ltd.

Urbanek, S., & Horner, J. (2019). Cairo: R graphics device using Cairo graphics library for
creating high-quality bitmap (PNG, JPEG, TIFF), vector (PDF, SVG, PostScript) and display

(X11 and Win32) output. Retrieved from
https://cran.r-project.org/web/packages/Cairo/index.html

Vabre, S. (2015). *Le sacre du Roquefort*. Presses universitaires François Rabelais, Tours,
France.

Vallone, L., Giardini, A., & Soncini, G. (2014). Secondary metabolites from *Penicillium*
roqueforti, a starter for the production of Gorgonzola cheese. *Italian Journal of Food Safety*,
3(3), 173–177 (2118).

Van Den Boogaart, K. G., Tolosana-Delgado, R., & Bren, M. (2018). compositions: A
unified R package to analyze compositional data. *Computers & Geosciences*, *34*(4), 320–338.

Villanueva, R. A. M., Chen, Z. J., & Wickham, H. (2016). *ggplot2: Elegant graphics for data*
analysis using the grammar of graphics (2nd Editio). Springer-Verlag New York.

Wickham, H. (2011). The split-apply-combine strategy for data analysis. *Journal of*
Statistical Software, *40*(1), 1–29.

Wilke, C. O. (2019). cowplot: Streamlined plot theme and plot annotations for "ggplot2".
Retrieved from https://cran.r-project.org/web/packages/cowplot/index.html

Williams, A. G., Beattie, S. H., & Banks, J. M. (2004). Enzymes involved in flavour
formation by bacteria isolated from the smear population of surface-ripened cheese.
International Journal of Dairy Technology, *57*(1), 7–13.

Woo, A. H., & Lindsay, R. C. (1984). Concentrations of major free fatty acids and flavor
development in italian cheese varieties. *Journal of Dairy Science*, *67*(5), 960–968.

Figure legends
893

Figure 1: Experimental cheesemaking. (A) Experimental design for cheesemaking, using one

strain per cheese, and three different strains from each of the four *Penicillium roqueforti*

populations (non-Roquefort cheese in blue, Roquefort cheese in purple, silage/food spoiler in

orange, lumber/food spoiler in green, the lineages of which are shown on the left). Each assay

(February, March, April) corresponded to a single strain from each of the four populations,

with three production replicates at different times, different batches of unpasteurized milk

and with two cheeses produced per strain in each replicate. The identities of the strains used

are indicated on the left of each assay, for each of the four *P. roqueforti* populations. (B)

Picture of the experimental cheeses at 20 days of maturation.

Figure 2: Mean percentage blue area per cheese slice at 20, 90 and 180 days of maturation,

for cheeses made with strains from the four *Penicillium roqueforti* populations (non-

Roquefort cheese in blue, Roquefort cheese in purple, silage/food spoiler in orange and

lumber/food spoiler in green). Error bars indicate 95% confidence intervals.

Figure 3: Illustration of the differences in the mean percentage blue area per cheese slice at

180 days of maturation between the four *Penicillium roqueforti* populations (non-Roquefort

cheese in blue, Roquefort cheese in purple, silage/food spoiler in orange and lumber/food

spoiler in green). Contrast and brightness have been standardized and the edges cropped.

Figure 4: Sums of 3,864 non-targeted aqueous signal peak areas, weighted by their mass-to-

charge ratios ("m/z"), obtained in positive ionization mode in 90-day cheeses made with

strains from the four *Penicillium roqueforti* populations (lumber/food spoiler in green, non-

Roquefort cheese in blue, Roquefort cheese in purple and silage/food spoiler in orange).

Figure 5: Sums of 3,094 non-targeted organic signal peak areas, weighted by their mass-to-

charge ratios ("m/z"), obtained in negative ionization mode in 90-day cheeses made with

strains from the four *Penicillium roqueforti* populations (lumber/food spoiler in green, non-

Roquefort cheese in blue, Roquefort cheese in purple and silage/food spoiler in orange).

Figure 6: Volatile compound production (integrated peak areas from chromatograms in

arbitrary units) in 90-day cheeses inoculated with strains from the four *Penicillium roqueforti*

populations (non-Roquefort cheese in blue, Roquefort cheese in purple, silage/food spoiler in

orange and lumber/food spoiler in green). The areas for each family of compounds are the

sum of the integrated areas of the compounds belonging to the family concerned. Alcohols I

and II are derived from proteolysis and lipolysis, respectively. Acids I, II and III are derived

from proteolysis, glycolysis and lipolysis, respectively (Supplementary Table 3). The color of

the titles indicates the affiliation of the compounds to their families, as in Figure S7.

Figure 7: Integrated surface area (from chromatograms in arbitrary units) of methyl ketones

(A) and secondary alcohols (B) for each assay (February, March, April) for the three strains

of each *Penicillium roqueforti* population (lumber/food spoiler in green, non-Roquefort

cheese in blue, Roquefort cheese in purple, silage/food spoiler in orange). Error bars

represent standard deviations across cheese replicates.

Figure legends for the Supplementary Material

Figure S1: Abundance of microorganisms in experimental cheeses. A. Abundance (in log

colony-forming units/g) of the eight types of microorganisms monitored at various stages of

cheese maturation (i.e. unpasteurized milk, 9, 20, 90 and 180 days), for each of the four

Penicillium roqueforti populations used to inoculate the cheeses (non-Roquefort cheese in

blue, Roquefort cheese in purple, silage/food spoiler in orange and lumber/food spoiler in

green). Error bars represent standard deviations across assays. B. Relative abundance of the

six main bacterial operational taxonomic units in cheeses made with strains from the four

Penicillium roqueforti populations (non-Roquefort cheese in blue, Roquefort cheese in

purple, silage/food spoiler in orange and lumber/food spoiler in green) in each assay

(February in light gray, March in mid-gray and April in dark gray) in cheeses at nine (45°

hatching) and 20 (135° hatching) days of maturation.

Figure S2: Mean bacterial genus diversity. A: Shannon index, B: Inverse of Simpson index=

1 - Simpson index) for the operational taxonomic units detected by metabarcoding in 9-day

cheeses (left) and 20-day cheeses (right) made with strains from the four *Penicillium*

roqueforti populations (lumber/food spoiler in green, non-Roquefort cheese in blue,

Roquefort cheese in purple and silage/food spoiler in orange).

Figure S3: Illustration of image processing for estimation of the percentage blue area on

cheese slices: (a) example of an unprocessed image of a cheese slice; (b) image after

brightness and contrast standardization; (c) image after cropping; (d) corresponding image

binarization with a grayscale of 102 on the red channel. White and black correspond to pixel

classification: in white, the inner part of the cheese and empty cavities; in black, cavities

filled with the fungus.

Figure S4: Differences in amino acid content between cheeses according to the population-

of-origin of the *Penicillium roqueforti* strains. A. Discrimination between 90-day cheeses

made with cheese (blue) and non-cheese (green) *P. roqueforti* populations (left), or Roquefort

cheese (purple) and non-Roquefort cheese (blue) *P. roqueforti* populations (right), based on

the amounts of the 23 identified amino acids present, according to an orthogonal signal-

corrected partial least squares (PLS) discriminant analysis. Vertical and horizontal axes

represent PLS1 and PLS 2 scores and gray arrows represent the relative contribution of

loadings of signals significantly discriminating the group considered in a *t*-test with jackknife

resampling. B. Amounts of molecules from particular classes detected in cheeses: mean

integrated peak area from chromatograms in arbitrary units (bars, left axis) and cumulative

percentage (line with dots, right axis) of aqueous extracts across all 90-day cheeses.

Figure S5: Sums of 8,472 non-targeted organic signal peak areas, weighted by their mass-to-

charge ratios ("m/z"), obtained in positive ionization mode in 90-day cheeses made with

strains from the four *Penicillium roqueforti* populations (lumber/food spoiler in green, non-

Roquefort cheese in blue, Roquefort cheese in purple and silage/food spoiler in orange).

Figure S6: Non-protein nitrogen levels at 20, 90 and 180 days of maturation, and water

activity at 90 and 180 days of maturation. Comparison of cheeses made with strains from

different *Penicillium roqueforti* populations (non-Roquefort cheese in blue, Roquefort cheese

in purple, silage/food spoiler in orange and lumber/food spoiler in green). Error bars indicate

95% confidence intervals.

Figure S7: Discrimination between 90-day cheeses inoculated with strains from the four

Penicillium roqueforti populations (non-Roquefort cheese in blue, Roquefort cheese in

purple, silage/food spoiler in yellow and lumber/food spoiler in green), based on the amounts

of 41 volatile compounds in an orthogonal signal-corrected partial least squares (PLS)

discriminant analysis. Vertical and horizontal axes represent the PLS1 and PLS2 variances,

and arrows represent the relative contributions of compound odor loadings significantly

discriminating the group considered (according to www.thegoodscentscompany.com) in a *t*-

test with jackknife resampling. The odor colors indicate the families in Figure 6 to which the

associated compounds belong.

Featured image: Roquefort cheese slice with symbols for two methyl ketones (2-heptanone

and 2-pentanone).

Figure 1: Experimental cheesemaking. (A) Experimental design for cheesemaking, using one strain per cheese, and three different strains from each of the four *Penicillium roqueforti* populations (non-Roquefort cheese in blue, Roquefort cheese in purple, silage/food spoiler in orange, lumber/food spoiler in green, the lineages of which are shown on the left). Each assay (February, March, April) corresponded to a single strain from each of the four populations, with three production replicates at different times, different batches of unpasteurized milk and with two cheeses produced per strain in each replicate. The identities of the strains used are indicated on the left of each assay, for each of the four *P. roqueforti* populations. (B) Picture of the experimental cheeses at 20 days of maturation.

Figure 2: Mean percentage blue area per cheese slice at 20, 90 and 180 days of maturation, for cheeses made with strains from the four *Penicillium roqueforti* populations (non-Roquefort cheese in blue, Roquefort cheese in purple, silage/food spoiler in orange and lumber/food spoiler in green). Error bars indicate 95% confidence intervals.

Non-cheese *P. roqueforti* populations	Cheese *P. roqueforti* populations
lumber/food spoiler (5.2 %)

 | non-Roquefort (10.2 %)

 |
| silage/food spoiler (5.5%)

 | Roquefort (11.8 %)

 |

Figure 3: Illustration of the differences in the mean percentage blue area per cheese slice at 180 days of maturation between the four *Penicillium roqueforti* populations (non-Roquefort cheese in blue, Roquefort cheese in purple, silage/food spoiler in orange and lumber/food spoiler in green). Contrast and brightness have been standardized and the edges cropped.

Figure 4: Sums of 3,864 non-targeted aqueous signal peak areas, weighted by their mass-to-charge ratios ("m/z"), obtained in positive ionization mode in 90-day cheeses made with strains from the four *Penicillium roqueforti* populations (lumber/food spoiler in green, non-Roquefort cheese in blue, Roquefort cheese in purple and silage/food spoiler in orange).

Figure 5: Sums of 3,094 non-targeted organic signal peak areas, weighted by their mass-to-charge ratios ("m/z"), obtained in negative ionization mode in 90-day cheeses made with strains from the four *Penicillium roqueforti* populations (lumber/food spoiler in green, non-Roquefort cheese in blue, Roquefort cheese in purple and silage/food spoiler in orange).

Figure 6: Volatile compound production (integrated peak areas from chromatograms in arbitrary units) in 90-day cheeses inoculated with strains from the four *Penicillium roqueforti* populations (non-Roquefort cheese in blue, Roquefort cheese in purple, silage/food spoiler in orange and lumber/food spoiler in green). The areas for each family of compounds are the sum of the integrated areas of the compounds belonging to the family concerned. Alcohols I and II are derived from proteolysis and lipolysis, respectively. Acids I, II and III are derived from proteolysis, glycolysis and lipolysis, respectively (Supplementary Table 3). The color of the titles indicates the affiliation of the compounds to their families, as in Figure S7.