Full Title: Boysenberry and apple juice reduces acute lung inflammation

through increased alternatively activated macrophage activity in an acute

mouse model of allergic airways disease

Odette M. Shaw[1*], Janine Cooney[2], Gregory M. Sawyer[1], Hannah Dinnan[1], Sheridan

Martell[1]

[1]Nutrition & Health Group, Food Innovation Portfolio, The New Zealand Institute for

Plant and Food Research Limited, Palmerston North, New Zealand

[2]Biological Chemistry & Bioactives Group, Food Innovation Portfolio, The New

Zealand Institute for Plant and Food Research Limited, Hamilton, New Zealand

***Correspondence:**

Odette Shaw

odette.shaw@plantandfood.co.nz

Short title: M2 macrophage reduce acute allergic airways inflammation

Abstract:

Bioactive compounds such as anthocyanins, proanthocyanins and other polyphenols

are found in a wide variety of fruits and vegetables, and consumption of these have

been associated with reduced lung inflammation and improved lung function in

asthma and other lung diseases. We investigated whether a combination of

Boysenberry and apple juice, found in BerriQi® Boysenberry and apple juice

concentrate, could reduce the allergic airways inflammation associated with asthma.

We characterised the polyphenolic components in BerriQi® Boysenberry and apple

juice concentrate and identified the main compounds as cyanidin glycosides,

ellagitannins, and chlorogenic acid. We found that consumption of 2.5 mg/kg total

anthocyanins from the BerriQi® Boysenberry and apple juice concentrate

significantly reduced eosinophil infiltration following acute ovalbumin (OVA) exposure

in a mouse model of allergic airways inflammation. We found that BerriQi®

Boysenberry and apple juice concentrate consumption increased M2 (CD206+)

macrophages and production of the M2-associated cytokines CXCL10 and CCL4

within the lung. These results suggest that consumption of BerriQi® Boysenberry

and apple juice concentrate could promote a shift towards an anti-inflammatory

environment within the lung leading to reduced immune cell infiltration and tissue

damage.

Keywords: allergic airways inflammation, Boysenberry, apple, anthocyanins,

ellagitannins, alternatively activated macrophages.

1. Introduction

Asthma is a heterogeneous, chronic, inflammatory lung disease characterized by reversible airways obstruction, bronchospasm and infiltration of immune cells (1-3). It is estimated that 150 million people are affected by asthma worldwide, with a 5–15% prevalence in children (4) and there is evidence that early life exposure to air pollution caused by vehicle exhaust, environmental dust and industrial processes, increases the severity of asthma in children (5-7). The respiratory symptoms such as cough, and wheeze are worsened by exposure to pollution (8). Proinflammatory cytokine production in response to allergens by immune cells are further increased with concomitant pollution exposure (9-13). Eosinophils, in particular, produce reactive oxygen species, and cytokines, leading to epithelial damage and contribute to mucosal inflammation and the recruitment of other proinflammatory immune cells (14-17). These repeated acute inflammatory responses leads to tissue damage and remodelling, contributing to airway hyperresponsiveness, mucus cell hyperplasia, fixed airway flow obstruction, and loss of lung function over time (18-21).

Large-scale epidemiological studies have found that increased fruit and vegetable consumption correlates with reduced asthma symptoms (22-25). These dietary-related improvements in lung function benefits are also seen in people living in polluted environments (26-28). Fruits and vegetables contain numerous bioactive compounds, including anthocyanins and procyanidins, which have been shown to attenuate lung inflammation in cell and animal models of allergy and asthma (29-34). Human population studies have identified that dietary intake of foods high in polyphenols (28) such as apples, pears (35), carrots, tomatoes (25) and citrus is inversely correlated with the frequency and severity of reported asthma symptoms, especially wheezing and coughing (22, 25, 35, 36). Previously, we have identified

that Boysenberry consumption led to decreased chronic lung inflammation, and improved lung tissue repair in an animal model of chronic allergic lung inflammation (33). Boysenberries contain high concentrations of anthocyanins and ellagitannins (37, 38). We have also found that procyanidin-rich extracts from apple suppressed IL-4 mediated cytokine production in cell culture models of lung epithelial allergic inflammation (32, 39).

There is increasing interest in understanding the mechanisms of action that specific plant bioactives have in the human body. This is partially to better understand the benefits of consuming specific fruits and vegetables, and partially to add value to specific foods through validated health claims. There is also interest in determining if combining specific plants containing different polyphenols can augment the health benefits above those seen with the individual plant. Use of animal models, where dietary intake can be tightly controlled are useful for both demonstrating/revealing the efficacy for identified compounds, and determining the biological mechanisms of action. The aim of this study was to determine whether the combination of Boysenberries and apple, as found in BerriQi® Boysenberry and apple juice concentrate with 2.5 mg/kg total anthocyanins (TAC) could reduce allergic airways inflammation in response to acute ovalbumin exposure in a mouse model system. We also sought to determine the mechanisms involved in any ameliorating effect.

2. Methods

2.1 Mice and Materials

C57BL/6J male mice were group housed on 12h light/dark cycle in a conventional animal facility at The New Zealand Institute for Plant and Food Research Limited (Palmerston North, New Zealand). Mice were fed Prodiet RMH1800 standard chow

for rodents (Lab Diet, St Louis, MO, USA) and filtered water ad libitum throughout

the study, all attempts to minimise suffering were made. All experimental procedures

were approved by the AgResearch Grasslands Animal Ethics Committee (AE

approvals #14839, #14731 and #14016) and carried out in accordance with the

Animal Welfare Act (1999). A proprietary Boysenberry and apple juice concentrate

(BerriQi®) was supplied by Anagenix Ltd (Auckland, New Zealand). Legendplex™

13-plex Th cytokine, proinflammatory cytokine and proinflammatory chemokine

panels, Zombie NIR™ fixable viability dye, and anti-mouse CD3 (clone 17A2), CD4

(clone GK1.5), CD8a (clone 53-6.7), CD80 (clone 16-10A1), CD86 (clone GL-1),

CD11c (clone N418), CD45 (clone 30-F11), CD206 (clone C068C2), CD14 (clone

Sa14-2), Ly6C (clone HK1.4), Gr-1 (clone RB6-8C5), I-A/I-E (MHC class II; clone

M5/114.15.2), and F4/80 (clone BM8) were purchased from Biolegend (San Diego,

CA, USA). Anti-mouse SiglecF (clone E50-2440) and CD11b (clone M1/70) were

from BD Biosciences (San Jose, CA, USA). Ovalbumin (OVA), and Alum were

purchased from Sigma (Auckland, New Zealand). iScript Advanced cDNA kit was

from Bio-Rad Laboratories (Hercules, CA, USA). Unless otherwise stated, all cell

culture media, supplements, Taqman probes and buffers were purchased from Life

Technologies NZ (Auckland, NZ).

2.2 BerriQi® Formulation and Chemical Composition Analysis

Boysenberries (*Rubus ursinus* var loganbaccus cv Boysenberry), harvested between

January-March 2017 from the Nelson region of NZ, were processed to a juice

concentrate by Boysenberries NZ (Nelson, NZ). Apples (*Malus domestica* –mixed

varieties) were harvested between April-June 2017 from the Hawkes Bay region of

New Zealand, and juice concentrates prepared by Profruit Ltd. (Hastings, NZ). The

freshly harvested Boysenberries and apples were graded, milled and pressed into

juice, de-seeded by centrifugation, pasteurised, depectinised by enzymatic treatment, clarified by centrifugation, and concentrated by evaporation by their respective manufacturers to yield fruit juice concentrates, each with an 8-fold concentration factor.

Shott NZ Ltd. (Wellington, NZ) then blended the juice concentrates (proprietary proportions), and added potassium sorbate preservative to a final concentration of 0.12% (w/w), yielding a dark red, non-cloudy syrup. The final BerriQi® Boysenberry and apple juice concentrate product measured pH 3.39 (by pH meter) and was 68.3° Brix (by refractometry), with a specific gravity of 1.35 mg/mL comprising 68% solids measured by both gravimetric and drying methods.

The polyphenol content of the BerriQi® Boysenberry and apple juice concentrate was determined by liquid chromatography-mass spectrometry (LC-MS) using an LTQ linear ion trap mass spectrometer fitted with an ESI interface (ThermoFisher Scientific, San Jose, CA, USA) coupled to an Ultimate 3000 UHPLC and PDA detector (Dionex, Sunnyvale, CA, USA). Anthocyanins were separated on a Poroshell 120 SB-C18, 2.1x150 mm, 2.7 µm, analytical LC column (Agilent, Torrance, CA, USA), maintained at 70°C. The solvents were (A) 5:3:92 acetonitrile:formic acid:water v/v/v and (B) acetonitrile + 0.1% formic acid (flow rate, 200 µL/min). The initial mobile phase, 100% A, was held for 2 min before being ramped linearly to 88% A at 14 min, returning to 5% A at 15 min and held for 4 min before resetting to the original conditions. The sample injection volume was 10 µL. The MS data were acquired in the positive mode. Anthocyanin concentrations are reported as cyanidin-3-O-glucoside equivalents.

Other phenolic compound separation was achieved using a Hypersil GOLD aQ 1.9μ

C18 175Å (Thermo Scientific, Waltham, MA, USA), 150 × 2.1 mm column

maintained at 45°C. The solvents were (A) water + 0.1% formic acid and (B)

acetonitrile + 0.1% formic acid (flow rate, 200 μL/min). The initial mobile phase, 95%

A/5% B, was ramped linearly to 85% A at 10 min, held for 3.75 min, then ramped

linearly to 75% A at 18 min, 67.2% A at 25 min, 50% A at 28 min, 3% A at 29 min and

held for 4 min before resetting to the original conditions. The sample injection volume

was 4 μL. The MS data were acquired in the negative mode.

2.3 OVA-Induced Airway Inflammation Model

Allergic airway disease was induced as previously described (34, 40). For the

Boysenberry and apple interventions, mice were randomized into receiving either

water (vehicle control) or 2.5 mg/kg TAC in the BerriQi® Boysenberry and apple

juice concentrate as previously described (34). Briefly, mice were fasted for 4 h

before being orally gavaged with water (control) or 2.5 mg/kg TAC in the BerriQi®

Boysenberry and apple juice concentrate made up to a total volume of 200μL in

water 1h before OVA challenge and again 2 days post-challenge. Mice were

sacrificed 4 days following intranasal ovalbumin challenge and immune parameters

were analyzed.

2.4 Immune Parameter Analysis

Bronchoalveolar lavage fluid (BALF) and lung tissues were collected as previously

described and immune cells were phenotyped by flow cytometry (40). Lung tissue

supernatant for cytokine analysis was prepared as previously described (34). Briefly,

the left lung lobe was minced into 500 μL phenol red-free complete RPMI media and

incubated at 37˚C for 30 min before being filtered through a 40 μm mesh to and

centrifuged to remove cellular material. The resulting supernatant was analysed for cytokine concentration using Legendplex bead-based multiplex immunoassays as per the manufacturer's instruction. Both cell phenotyping and the cytokine multiplex assays were analyzed using a BD FACSverse (BD Biosciences, San Jose, CA, USA). H&E and AB-PAS histological staining were performed by Massey IVABS histology unit.

2.5 Real-time qPCR analysis

Mouse lung tissue was collected four days following OVA challenge and snap frozen in liquid nitrogen. The lung samples were crushed into powder using a mortar and pestle with liquid nitrogen to keep the samples frozen. The RNA was extracted from the powder using a TRIZOL total RNA extraction protocol. RNA was quantified using an LVis plate in a POLARstar Omega plate reader (BMG) and the quality of the ribosomal RNA bands confirmed by agarose gel electrophoresis (data not shown). 5 µg RNA from each sample was used as the template for cDNA synthesis using the iScript™ cDNA Synthesis Kit. Taqman® Gene Expression Assays were purchased for each gene of interest. Two housekeeping genes, GAPDH and β-actin, were used as controls to determine the amount of relative gene expression. Taqman® Gene Expression Master Mix was used to PCR amplify the genes in a Bio-Rad™ CFX384™ Real-Time PCR Dectection System. Three lung samples per treatment group were prepared and amplified in quadruplicate with the housekeeping genes amplified on the sample 384-well plate.

2.6 Statistical Analysis

Data were analyzed using one-way analysis of variance (ANOVA) with a Tukey's

post hoc test and graphed in SigmaPlot 12.5 (Systat Software Inc., San Jose, CA,

USA).

3. Results

3.1 Chemical Composition of the Boysenberry and Apple Juice Concentrate

The results of the LC-MS analysis showed that cyanidin glycosides, ellagitannins,

and chlorogenic acid were the major components in BerriQi® Boysenberry and apple

juice concentrate (Table 1, supplementary figures 1, 2). Minor components included

phloretin 2-O-glucoside, and a mix of phenolic acids, flavonol glycosides, flavanol

monomers and procyanidins. The major classes of phenolic compounds were

anthocyanins (1969 µg/mL) and hydrolysable tannins (946 µg/mL), accounting for

56% and 27%, respectively, of the total phenolics quantified. The most abundant

tannins were ellagic acid (449 µg/mL) and sanguiin H6 (213 µg/mL).

3.2 Effect of Boysenberry and Apple juice concentrate Intervention on

Ovalbumin-Induced Allergic Airways Inflammation

Acute intranasal OVA exposure resulted in an infiltration of immune cells into the

lung (Figure 1A) and increased mucous production (Figure 1B). Consumption of 2.5

mg/kg TAC BerriQi® Boysenberry and apple juice concentrate reduced the

infiltration of immune cells and decreased OVA-induced mucous production (Figure

1A-B). We quantified the type and number of immune cells infiltrating into the lung,

and found that acute OVA exposure significantly increased ($P<0.001$) infiltrating

eosinophils (CD45+/CD11b+/SiglecF+), neutrophils (CD45+Ly6C+Gr-1+) and T-cells

(CD45+/CD3+/CD4+ or CD45+/CD3+/CD8a+) in the lung, compared with the lung of

naïve animals (Figure 1C-F). Compared with animals only exposed to OVA, those

that also consumed 2.5 mg/kg TAC BerriQi® Boysenberry and apple juice

concentrate showed a significant decrease (P<0.001) in the number of infiltrating

eosinophils, neutrophils and T cells in the lung (Figure 1C-F). We saw no change in

the number of CD4+ or CD8+ T cells in the mediastinal (lung-draining) lymph node

for any of the treatment groups (Figure 2A-B).

There was a trend towards an increased percentage of CD206+/CD14-

macrophages in the lungs of mice that consumed 2.5 mg/kg TAC BerriQi®

Boysenberry and apple juice concentrate (Figure 3A). We measured the gene

expression of Arg1, Ym-1 and Fizz1 in lung tissue and found that 2.5 mg/kg TAC

BerriQi® Boysenberry and apple juice concentrate consumption led to a significant

fold- increase in Arg1 (P<0.05) and Fizz1 (P<0.01) gene expression compared to

naïve mice (Table 2). We found no significant fold-change in Nos2 or Ym-1 gene

expression between any of the treatment groups (Table 2).

Consumption of 2.5 mg/kg TAC BerriQi® Boysenberry and apple juice concentrate

led to increased levels of the cytokines IL-17A, CXCL10, and CCL4 (Figure 3B-D)

persisting at four days following OVA challenge, but did not affect the IL-17F

concentration (Figure 3E). We saw no effect on the concentrations of the classical

Th1/2 cytokines IFNγ, IL-5, IL-9 or IL-10 in either the BerriQi® Boysenberry and

apple juice concentrate treated or the OVA alone mice compared to naïve controls

(Figure 2C-F).

4. Discussion

We evaluated the effects of dietary supplementation with BerriQi® Boysenberry and

apple juice concentrate containing 2.5 mg/kg TAC, on the immune responses in a

mouse model of acute allergic airways inflammation. Our results show that

consumption of 2.5 mg/kg TAC BerriQi® Boysenberry and apple juice concentrate

reduced granulocyte and local T cell infiltration into the lung after OVA challenge, but

did not alter T cell activation within the lung draining lymph node or the levels of

classical Th-2 and Th-1 cytokines in the lung at four days following OVA challenge.

Our current results indicated that BerriQi® Boysenberry and apple juice concentrate

had little impact on the Th-2/Th-1 mediated allergic response of mice, but rather

targeted innate proinflammatory immune pathways. This is consistent with our

previously reported finding in a mouse model of chronic allergic airways inflammation

using 10mg/kg TAC Boysenberry juice concentrate (33). Chemical composition

analysis showed that the BerriQi® Boysenberry and apple juice concentrate

formulation contained high concentrations of cyanidin glycosides, ellagitannins, and

chlorogenic acid. These compounds have been previously shown to reduce

inflammatory signalling in vitro (38, 41, 42), and in vivo animal models of

inflammation (34, 43-46). Our current results suggest that consumption of 2.5 mg/kg

TAC BerriQi® Boysenberry and apple juice concentrate, which also contains high

levels of ellagitannins and chlorogenic acid, could have broader lung health benefits

beyond allergic asthma disease by promoting the resolution of inflammation caused

by innate immune cell overactivation. Current asthma therapies also suppress both

the adaptive and innate immune responses without affecting the aberrant sensitivity

to the allergen and this can lead to adverse events.

Consumption of BerriQi® Boysenberry and apple juice concentrate had less of an

effect on monocyte/macrophage infiltration than on granulocyte infiltration, and there

was increased percentage of CD206+ monocytes. This could represent a shift to an

M2 anti-inflammatory phenotype. We then measured the changes in gene

expression for Arg1, Ym-1 and Fizz1, the classic genes for identifying alternatively

activated macrophages (47, 48). Arg1 and Fizz1 gene expression was significantly

increased with   BerriQi® Boysenberry and apple juice concentrate consumption.

Consistent with a shift to a more anti-inflammatory macrophage phenotype, we

detected increased levels of CXCL10 and CCL4 cytokines, which are produced by

M2 macrophages, in the lungs of mice that consumed BerriQi® Boysenberry and

apple juice concentrate. These results suggest that the consumption of Boysenberry

and apple juice concentrate led to a switch to the M2 phenotype in OVA-challenged

mice. This could be one of the mechanisms by which BerriQi® Boysenberry and

apple juice concentrate consumption contributed to the resolution of inflammation.

Previously, we reported that 10mg/kg TAC Boysenberry juice concentrate can

increase the abundance of alternatively activated (M2) macrophages, which promote

tissue repair in a chronic model of airways inflammation (33). Our current results

suggest that BerriQi® Boysenberry and apple juice concentrate, had a similar effect

in this mouse model of acute inflammation, in particular the increase in Arg1 gene

expression is similar to our previously reported study showing increased arginase

protein expression by alternatively activated macrophages (33)Further, research

looking at an animal model Th2-mediated inflammation has identified M2

macrophage derived Fizz1as a key limiting factor for Th2-mediated pulmonary

inflammation (49).

The mice that consumed 2.5 mg/kg TAC BerriQi® Boysenberry and apple juice

concentrate showed increased levels of the cytokines IL-17A, CXCL10, and CCL4,

but the levels of IL-17F were not affected. High IL-17 and IL-17F levels have been

implicated in asthma pathogenesis, however, there is also evidence that IL-17A (50)

can increase the abundance of MMP-9, an important tissue remodelling protein in

asthma (33) as well as inducing apoptosis of neutrophils and eosinophils (50).

CXCL10 and CCL4 are chemokines that attract monocytes/macrophages, and

CXCL10 may also inhibit the infiltration of eosinophils in response to allergic airways

inflammation (51). The combination of increased IL-17A-mediated granulocyte

apoptosis and CXCL10-mediated inhibition of granulocyte infiltration could explain

how the consumption of the BerriQi® Boysenberry and apple juice concentrate

resulted in decreased allergic airways inflammation, in particular the reduced number

of eosinophils and neutrophils.

This switch to a M2 macrophage phenotype may be through the Boysenberry and

apple polyphenols identified in the BerriQi® Boysenberry and apple juice concentrate

directly inhibiting proinflammatory pathways, or through an indirect shortening of the

proinflammatory phase. Other studies have shown that increased dietary fibre

shortened the duration of the proinflammatory phase, leading to reduced tissue

damage in a chronic mouse model of allergic asthma (52). Based on the estimated

fibre content in BerriQi® Boysenberry and apple juice concentrate (Supplementary

table 1), it is unlikely that the dietary fibre component played a significant role in the

immune modulation we observed. However, the combination of anthocyanins with

other polyphenols identified in the BerriQi® Boysenberry and apple juice concentrate

could have a similar effect on inflammation either by shortening the proinflammatory

time course, or promoting the production of anti-inflammatory proteins. Ellagitannins

have been shown in cell culture and animal models of chronic inflammatory diseases

to reduce proinflammatory prostaglandins (53), cytokines (45, 54), and other proteins

(42, 55, 56). Anthocyanins have also been shown to inhibit proinflammatory proteins

(57, 58), and activate anti-inflammatory pathways in models of inflammation (59-63).

It is possible that the combination of the different polyphenols in the BerriQi®

Boysenberry and apple juice concentrate act on a number of different immune

pathways to regulate the immune responses to OVA.

We found that mice that consumed BerriQi® Boysenberry and apple juice

concentrate had reduced immune cell infiltration in response to acute OVA challenge

and this could be as a result of a shift towards an anti-inflammatory environment

within the lung. These results highlight the potential of anthocyanin-rich Boysenberry

and apple dietary supplementation to modulate innate immune pathways during

acute allergic lung inflammation. Further work is needed to determine if these

pathways are also altered in other lung inflammatory conditions, such as air pollution

exposure. Clinical studies are needed to show if these findings are translatable to

human health.

Conflict of Interest

All Authors were employed by The New Zealand Institute of Plant & Food Research Limited, a New Zealand Crown Research Institute wholly owned by the New Zealand Government for the purposes of research into sustainable production, elite breeding, food and health science of horticultural, arable, and seafood products.

OMS report that they are named on patents related to the formulation of BerriQi® Boysenberry and apple juice concentrate, but have not received any financial compensation, nor will receive any personal royalty payments as a result of this. None of the other authors declare any other conflicts of interest. Under the terms of the Innovation Cell™ Collaboration Agreement Plant & Food Research and Anagenix have a royalty sharing agreement for any royalties that result from the sale of BerriQi® Boysenberry and apple juice concentrate product.

Author Contributions

OMS designed, performed, analyzed and interpreted the in vivo studies, and wrote and helped to edit the manuscript; JC performed, analyzed and interpreted the chemcial composition experiment and GMS performed, analyzed and interpreted in vivo studies, and both contributed to the writing and editing of the manuscript; HD and SM performed the in vivo studies and helped edit the manuscript.

Funding

This work was funded by The New Zealand Institute of Plant & Food Research Limited as a contribution to the Innovation Cell™ Collaboration Agreement executed between Anagenix Limited and The New Zealand Institute for Plant and Food Research Limited on 5 September 2016 (PFR reference #33609).

The authors declare that this study received funding from The New Zealand Institute

of Plant & Food Research Limited. The funder was not involved in the study design,

collection, analysis, and interpretation of data, the writing of this article or the

decision to submit it for publication.

Acknowledgements

The authors acknowledge the efforts of Roger Hurst, TC Chadderton, and Brendan

Vercoe in securing internal funding for this work. The authors would like to thank

Andrew Carroll for assisting with data analysis and Jenny Smith for her invaluable

help preparing the manuscript.

5. References

1.   Agache I, Akdis CA. Endotypes of allergic diseases and asthma: An important

step in building blocks for the future of precision medicine. Allergol Int.

2016;65(3):243-52.

2.   Agrawal DK, Bharadwaj A. Allergic airway inflammation. Curr Allergy Asthma

Rep. 2005;5(2):142-8.

3.   Barnes PJ. Pathophysiology of asthma. Br J Clin Pharmacol. 1996;42(1):3-10.

4.   WHO. Prevention of allergy and allergic asthma: based on the WHO/WAO

Meeting on the Prevention of Allergy and Allergic Asthma, Geneva, 8-9 January

2002. Management DoND, Arthritis CRDa, Organization WA, Organization. WH,

editors. Geneva: World Health Organization; 2003.

5.   Hsu HH, Chiu YH, Coull BA, Kloog I, Schwartz J, Lee A, et al. Prenatal

Particulate Air Pollution and Asthma Onset in Urban Children. Identifying Sensitive

Windows and Sex Differences. Am J Respir Crit Care Med. 2015;192(9):1052-9.

6.   Miller RL, Peden DB. Environmental effects on immune responses in patients

with atopy and asthma. J Allergy Clin Immunol. 2014;134(5):1001-8.

7.   Jung DY, Leem JH, Kim HC, Kim JH, Hwang SS, Lee JY, et al. Effect of

Traffic-Related Air Pollution on Allergic Disease: Results of the Children's Health and

Environmental Research. Allergy Asthma Immunol Res. 2015;7(4):359-66.

8.   Hoek G, Pattenden S, Willers S, Antova T, Fabianova E, Braun-Fahrlander C,

et al. PM10, and children's respiratory symptoms and lung function in the PATY

study. Eur Respir J. 2012;40(3):538-47.

9.   Acciani TH, Brandt EB, Khurana Hershey GK, Le Cras TD. Diesel exhaust

particle exposure increases severity of allergic asthma in young mice. Clin Exp

Allergy. 2013;43(12):1406-18.

10.   Brandt EB, Biagini Myers JM, Acciani TH, Ryan PH, Sivaprasad U, Ruff B, et

al. Exposure to allergen and diesel exhaust particles potentiates secondary allergen-

specific memory responses, promoting asthma susceptibility. J Allergy Clin Immunol.

2015;136(2):295-303 e7.

11.   Carlsten C, Blomberg A, Pui M, Sandstrom T, Wong SW, Alexis N, et al.

Diesel exhaust augments allergen-induced lower airway inflammation in allergic

individuals: a controlled human exposure study. Thorax. 2016;71(1):35-44.

12.   Kim J, Natarajan S, Vaickus LJ, Bouchard JC, Beal D, Cruikshank WW, et al.

Diesel exhaust particulates exacerbate asthma-like inflammation by increasing CXC

chemokines. Am J Pathol. 2011;179(6):2730-9.

13.   van Voorhis M, Knopp S, Julliard W, Fechner JH, Zhang X, Schauer JJ, et al.

Exposure to atmospheric particulate matter enhances Th17 polarization through the

aryl hydrocarbon receptor. PLoS One. 2013;8(12):e82545.

14.   Trivedi SG, Lloyd CM. Eosinophils in the pathogenesis of allergic airways

disease. Cell Mol Life Sci. 2007;64(10):1269-89.

15.   Amin K, Janson C, Bystrom J. Role of Eosinophil Granulocytes in Allergic

Airway Inflammation Endotypes. Scand J Immunol. 2016;84(2):75-85.

16.   Bossley CJ, Fleming L, Gupta A, Regamey N, Frith J, Oates T, et al. Pediatric

severe asthma is characterized by eosinophilia and remodeling without T(H)2

cytokines. J Allergy Clin Immunol. 2012;129(4):974-82 e13.

17.   Brown JR, Kleimberg J, Marini M, Sun G, Bellini A, Mattoli S. Kinetics of

eotaxin expression and its relationship to eosinophil accumulation and activation in

bronchial biopsies and bronchoalveolar lavage (BAL) of asthmatic patients after

allergen inhalation. Clin Exp Immunol. 1998;114(2):137-46.

18.   Bergeron C, Al-Ramli W, Hamid Q. Remodeling in asthma. Proc Am Thorac

Soc. 2009;6(3):301-5.

19.   Ahdieh M, Vandenbos T, Youakim A. Lung epithelial barrier function and

wound healing are decreased by IL-4 and IL-13 and enhanced by IFN-gamma. Am J

Physiol Cell Physiol. 2001;281(6):C2029-38.

20.   Al-Muhsen S, Johnson JR, Hamid Q. Remodeling in asthma. J Allergy Clin

Immunol. 2011;128(3):451-62.

21.   Brightling CE, Gupta S, Gonem S, Siddiqui S. Lung damage and airway

remodelling in severe asthma. Clin Exp Allergy. 2012;42(5):638-49.

22.   McKeever TM, Britton J. Diet and asthma. Am J Respir Crit Care Med.

2004;170(7):725-9.

23.   Nurmatov U, Devereux G, Sheikh A. Nutrients and foods for the primary

prevention of asthma and allergy: systematic review and meta-analysis. J Allergy

Clin Immunol. 2011;127(3):724-33 e1-30.

24.   Okoko BJ, Burney PG, Newson RB, Potts JF, Shaheen SO. Childhood asthma and fruit consumption. Eur Respir J. 2007;29(6):1161-8.

25.   Rosenlund H, Kull I, Pershagen G, Wolk A, Wickman M, Bergstrom A. Fruit and vegetable consumption in relation to allergy: disease-related modification of consumption? J Allergy Clin Immunol. 2011;127(5):1219-25.

26.   Stevens EL, Rosser F, Forno E, Peden D, Celedon JC. Can the effects of outdoor air pollution on asthma be mitigated? J Allergy Clin Immunol. 2019;143(6):2016-8 e1.

27.   Burbank AJ, Duran CG, Pan Y, Burns P, Jones S, Jiang Q, et al. Gamma tocopherol-enriched supplement reduces sputum eosinophilia and endotoxin-induced sputum neutrophilia in volunteers with asthma. J Allergy Clin Immunol. 2018;141(4):1231-8 e1.

28.   Pounis G, Arcari A, Costanzo S, Di Castelnuovo A, Bonaccio M, Persichillo M, et al. Favorable association of polyphenol-rich diets with lung function: Cross-sectional findings from the Moli-sani study. Respir Med. 2018;136:48-57.

29.   Park SJ, Shin WH, Seo JW, Kim EJ. Anthocyanins inhibit airway inflammation and hyperresponsiveness in a murine asthma model. Food Chem Toxicol. 2007;45(8):1459-67.

30.   Coleman SL, Shaw OM. Progress in the understanding of the pathology of allergic asthma and the potential of fruit proanthocyanidins as modulators of airway inflammation. Food Funct. 2017;8(12):4315-24.

31.   Coleman S, Kruger M, Sawyer G, Hurst R. Procyanidin A2 Modulates IL-4-Induced CCL26 Production in Human Alveolar Epithelial Cells. Int J Mol Sci. 2016;17(11):1888.

32.   Sawyer GM, Stevenson DE, McGhie TK, Hurst RD. Suppression of CCL26

and CCL11 generation in human alveolar epithelial cells by apple extracts containing

procyanidins. J Funct Foods. 2017;31:141-51.

33.   Shaw OM, Hurst RD, Harper JL. Boysenberry ingestion supports fibrolytic

macrophages with the capacity to ameliorate chronic lung remodeling. Am J Physiol

Lung Cell Mol Physiol. 2016;311(3):L628-38.

34.   Shaw OM, Nyanhanda T, McGhie TK, Harper JL, Hurst RD. Blackcurrant

anthocyanins modulate CCL11 secretion and suppress allergic airway inflammation.

Mol Nutr Food Res. 2017;61(9):1600868.

35.   Garcia-Larsen V, Thawer N, Charles D, Cassidy A, van Zele T, Thilsing T, et

al. Dietary Intake of Flavonoids and Ventilatory Function in European Adults: A

GA(2)LEN Study. Nutrients. 2018;10(1).

36.   Garcia V, Arts IC, Sterne JA, Thompson RL, Shaheen SO. Dietary intake of

flavonoids and asthma in adults. Eur Respir J. 2005;26(3):449-52.

37.   Cooney JM, Jensen DJ, McGhie TK. LC-MS identification of anthocyanins in

boysenberry extract and anthocyanin metabolites in human urine following dosing. J

Sci Food Agr. 2004;84(3):237-45.

38.   McGhie TK, Martin H, Lunken RC. The combination of analytical-scale HPLC

separation with a TR-FRET assay to investigate JAK2 inhibitory compounds in a

Boysenberry drink. Food Funct. 2012;3(11):1170-5.

39.   Coleman SL, Hurst RD, Sawyer GM, Kruger MC. The in vitro evaluation of

isolated procyanidins as modulators of cytokine-induced eotaxin production in

human alveolar epithelial cells. J Berry Res. 2016;6(2):115-24.

40.   Shaw OM, Harper JL. An efficient single prime protocol for the induction of

antigen-induced airways inflammation. J Immunol Methods. 2013;395(1-2):79-82.

41. Cassidy A, Rogers G, Peterson JJ, Dwyer JT, Lin H, Jacques PF. Higher

dietary anthocyanin and flavonol intakes are associated with anti-inflammatory

effects in a population of US adults. Am J Clin Nutr. 2015;102(1):172-81.

42. El-Shitany NA, El-Bastawissy EA, El-desoky K. Ellagic acid protects against

carrageenan-induced acute inflammation through inhibition of nuclear factor kappa

B, inducible cyclooxygenase and proinflammatory cytokines and enhancement of

interleukin-10 via an antioxidant mechanism. Int Immunopharmacol. 2014;19(2):290-

9.

43. Auclair S, Silberberg M, Gueux E, Morand C, Mazur A, Milenkovic D, et al.

Apple polyphenols and fibers attenuate atherosclerosis in apolipoprotein E-deficient

mice. J Agric Food Chem. 2008;56(14):5558-63.

44. Denis MC, Roy D, Yeganeh PR, Desjardins Y, Varin T, Haddad N, et al. Apple

peel polyphenols: a key player in the prevention and treatment of experimental

inflammatory bowel disease. Clin Sci (Lond). 2016;130(23):2217-37.

45. Guan S, Zheng YY, Yu X, Li WL, Han B, Lu J. Ellagic acid protects against

LPS-induced acute lung injury through inhibition of nuclear factor kappa B,

proinflammatory cytokines and enhancement of interleukin-10. Food Agr Immunol.

2017;28(6):1347-61.

46. Impellizzeri D, Talero E, Siracusa R, Alcaide A, Cordaro M, Maria Zubelia J,

et al. Protective effect of polyphenols in an inflammatory process associated with

experimental pulmonary fibrosis in mice. Br J Nutr. 2015;114(6):853-65.

47. Chung S, Lee TJ, Reader BF, Kim JY, Lee YG, Park GY, et al. FoxO1

regulates allergic asthmatic inflammation through regulating polarization of the

macrophage inflammatory phenotype. Oncotarget. 2016;7(14):17532-46.

48.   Kurowska-Stolarska M, Stolarski B, Kewin P, Murphy G, Corrigan CJ, Ying S,

et al. IL-33 amplifies the polarization of alternatively activated macrophages that

contribute to airway inflammation. J Immunol. 2009;183(10):6469-77.

49.   Nair MG, Du Y, Perrigoue JG, Zaph C, Taylor JJ, Goldschmidt M, et al.

Alternatively activated macrophage-derived RELM-{alpha} is a negative regulator of

type 2 inflammation in the lung. J Exp Med. 2009;206(4):937-52.

50.   Linden A, Dahlen B. Interleukin-17 cytokine signalling in patients with asthma.

Eur Respir J. 2014;44(5):1319-31.

51.   Su YC, Rolph MS, Hansbro NG, Mackay CR, Sewell WA. Granulocyte-

macrophage colony-stimulating factor is required for bronchial eosinophilia in a

murine model of allergic airway inflammation. J Immunol. 2008;180(4):2600-7.

52.   Trompette A, Gollwitzer ES, Yadava K, Sichelstiel AK, Sprenger N, Ngom-Bru

C, et al. Gut microbiota metabolism of dietary fiber influences allergic airway disease

and hematopoiesis. Nat Med. 2014;20(2):159-66.

53.   Karlsson S, Nanberg E, Fjaeraa C, Wijkander J. Ellagic acid inhibits

lipopolysaccharide-induced expression of enzymes involved in the synthesis of

prostaglandin E2 in human monocytes. Br J Nutr. 2010;103(8):1102-9.

54.   Cornelio Favarin D, Martins Teixeira M, Lemos de Andrade E, de Freitas

Alves C, Lazo Chica JE, Arterio Sorgi C, et al. Anti-inflammatory effects of ellagic

acid on acute lung injury induced by acid in mice. Mediators Inflamm.

2013;2013:164202.

55.   Marin M, Maria Giner R, Rios JL, Recio MC. Intestinal anti-inflammatory

activity of ellagic acid in the acute and chronic dextrane sulfate sodium models of

mice colitis. J Ethnopharmacol. 2013;150(3):925-34.

56. Saba, Khan S, Parvez S, Chaudhari B, Ahmad F, Anjum S, et al. Ellagic acid

attenuates bleomycin and cyclophosphamide-induced pulmonary toxicity in Wistar

rats. Food Chem Toxicol. 2013;58:210-9.

57. Esposito D, Chen A, Grace MH, Komarnytsky S, Lila MA. Inhibitory effects of

wild blueberry anthocyanins and other flavonoids on biomarkers of acute and chronic

inflammation in vitro. J Agric Food Chem. 2014;62(29):7022-8.

58. Fu Y, Zhou E, Wei Z, Wang W, Wang T, Yang Z, et al. Cyanidin-3-O-beta-

glucoside ameliorates lipopolysaccharide-induced acute lung injury by reducing

TLR4 recruitment into lipid rafts. Biochem Pharmacol. 2014;90(2):126-34.

59. Chen YF, Shibu MA, Fan MJ, Chen MC, Viswanadha VP, Lin YL, et al. Purple

rice anthocyanin extract protects cardiac function in STZ-induced diabetes rat hearts

by inhibiting cardiac hypertrophy and fibrosis. J Nutr Biochem. 2016;31:98-105.

60. Edirisinghe I, Banaszewski K, Cappozzo J, McCarthy D, Burton-Freeman BM.

Effect of black currant anthocyanins on the activation of endothelial nitric oxide

synthase (eNOS) in vitro in human endothelial cells. J Agric Food Chem.

2011;59(16):8616-24.

61. Khanna S, Roy S, Bagchi D, Bagchi M, Sen CK. Upregulation of oxidant-

induced VEGF expression in cultured keratinocytes by a grape seed

proanthocyanidin extract. Free Radic Biol Med. 2001;31(1):38-42.

62. Koh ES, Lim JH, Kim MY, Chung S, Shin SJ, Choi BS, et al. Anthocyanin-rich

Seoritae extract ameliorates renal lipotoxicity via activation of AMP-activated protein

kinase in diabetic mice. J Transl Med. 2015;13(1):203.

63. Liu Y, Tan D, Tong C, Zhang Y, Xu Y, Liu X, et al. Blueberry anthocyanins

ameliorate radiation-induced lung injury through the protein kinase RNA-activated

pathway. Chem Biol Interact. 2015;242:363-71.

 Table 1. Phenolic compounds detected in BerriQi® (µg/mL). M+ and (M-H)- ions are the pseudomolecular ions used for identification of compounds by liquid chromatography-mass spectrometry (LC-MS). All identifications confirmed by MS/MSn experiments. Peak numbers refer to chromatograms shown in supplementary data. # detected as [M+formate]- adduct

Peak	M+	(M-H)⁻	Compound	µg/mL
			Anthocyanins	
1	611		Cyanidin 3-*O*-sophoroside	883
2	449		Cyanidin 3-*O*-glucoside	571
3	481		Cyanidin 3-*O*-sambubioside	24
4	757		Cyanidin 3-*O*-(2-glucosylrutinoside)	411
5	595		Cyanidin 3-*O*-rutinoside	62
6	727		Cyanidin 3-*O*-xylosylrutinoside	18
			Phenolic acids	
7		169	Gallic acid	140
8		153	Protocatechuric acid	35
10		353	Chlorogenic acid	69
11		179	Caffeic acid	7
15		337	4-*p*-Coumaroylquinic acid	26
17		337	5-*p*-Coumaroylquinic acid	3
			Flavan-3-ols and procyanidins	
9		335#	Catechin	3
12		577	Procyanidin B2	6
13		335#	*Epi*-catechin	21
25		579	unknown procyanidin isomer	3
			Hydrolysable tannins	
14		1567	Sanguiin H10 isomer 1	9
16		469	Sanguisorbic acid dilactone	120
19		2036.5	Galloyl-SH6	66
20		1567	Sanguiin H10 isomer 2	61
21		2501	Lambertian C (minus ellagic acid)	11
22		2803	Lambertian C	17
24		1869	Sanguiin H6	213
26		301	Ellagic acid	449
			Flavonols	
27		609	Quercetin 3-*O*-rutinoside	6
28		463	Quercetin 3-*O*-galactoside	17

29	477	Quercetin 3-O-glucuronide	36
30	463	Quercetin 3-O-glucoside	13
31	433	Quercetin 3-O-pentoside 1	9
32	433	Quercetin 3-O-pentoside 3	6
33	433	Quercetin 3-O-pentoside 2	13
35	447	Quercetin 3-O-rhamnoside	15
37	301	Quercetin	19
		Chalcones	
34	567	Phloretin 2-O-xylo-glucoside	9
36	481[#]	Phloretin 2-O-glucoside	62
		Unknowns	
18	563[#]	unknown	71
23	639	unknown	22
		Total	**1557**

Table 2: BerriQi® Boysenberry and apple juice concentrate increases alternatively activated macrophage gene expression in the lung. Mice were primed with ovalbumin (OVA)/Alum i.p. and then challenged 7 days later with OVA i.n. (Day 0). Mice were orally gavaged with 2.5 mg/kg total anthocyanins (TAC) in the BerriQi® Boysenberry and apple juice concentrate (BerriQi) 1h before OVA challenge and again 2 days post-challenge. Mean fold-change (SEM) in gene expression was measured by real-time qPCR in lung tissue4 days post-OVA challenge. $*P<0.05$, $**P<0.01$ compared with naïve (one-way ANOVA with Tukey's Post Hoc test) for four experimental replicates with n=3 per treatment groups.

Target Gene	Naïve	OVA	OVA + BerriQi
Arg1	1 (0.5)	9.4 (5.3)	26.0 (7.9) *
Ym-1	1 (0.4)	4.4 (1.9)	5.5 (1.0)
Fizz1	1 (0.5)	15.4 (0.9) **	19.6 (3.3) **
Nos2	1 (0.3)	1.5 (0.8)	1.7 (0.5)

Figure 1: BerriQi® Boysenberry and apple juice concentrate suppresses

ovalbumin-induced airway inflammation, and immune cell infiltration. Mice

were primed with ovalbumin (OVA)/Alum i.p. and then challenged 7 days later with

OVA i.n. (Day 0). Mice were orally gavaged with 2.5 mg/kg total anthocyanins (TAC)

in the BerriQi® Boysenberry and apple juice concentrate (BerriQi) 1h before OVA

challenge and again 2 days post-challenge. (A) Haematoxylin and eosin stained lung

tissue from naïve, OVA-challenged and OVA-challenged mice treated with BerriQi®

Boysenberry and apple juice concentrate. Magnification 10x (top) and 20x (bottom)

Asterisk =cell infiltration. (B) Alcian-blue Periodic acid-Schiff stained lung tissue from

naïve, OVA-challenged and OVA-challenged mice treated with BerriQi®

Boysenberry and apple juice concentrate. Magnification 10x (top) and 20x (bottom).

Arrow=mucous producing goblet cells. (C) Total eosinophil, (D) Total neutrophil, (E)

CD4+ T cells and CD8+ T cells in bronchioalveolar lavage fluid (BALF) were

determined 4 days post-OVA challenge. Data presented as mean ± SEM $P < 0.001$

compared with naïve and OVA challenge + BerriQi® Boysenberry and apple juice

concentrate (one-way ANOVA with Tukey's Post Hoc test) for two experimental

replicates with n=10 per treatment groups.

Figure 2: BerriQi® Boysenberry and apple juice concentrate does not alter

classical Th1/Th2 cells and cytokines. Mice were primed with ovalbumin

(OVA)/Alum i.p. and then challenged 7 days later with OVA i.n. (Day 0). Mice were

orally gavaged with 2.5 mg/kg total anthocyanins (TAC) in the BerriQi® Boysenberry

and apple juice concentrate (BerriQi) 1h before OVA challenge and again 2 days

post-challenge. Mediastinal lymph node (MSLN) (A) CD4+ and (B) CD8+ T cells

number; and lung tissue production of (C) IL-5, (D) IL-9, (E) IL-10 and (F) IFNγ were

determined 4 days post-OVA challenge by Legendplex. Data presented as mean ±

SEM for two experimental replicates with n=10 per treatment groups.

Figure 3: BerriQi® Boysenberry and apple juice concentrate suppresses

ovalbumin-induced airway inflammation through increased IL-17A, CXCL10

and CCL4 concentration. Mice were primed with ovalbumin (OVA)/Alum i.p. and

then challenged 7 days later with OVA i.n. (Day 0). Mice were orally gavaged with

2.5 mg/kg total anthocyanins (TAC) in the BerriQi® Boysenberry and apple juice

concentrate (BerriQi) 1h before OVA challenge and again 2 days post-challenge. (A)

Percentage of CD206+ macrophages in bronchioalveolar lavage fluid (BALF) and

lung tissue production of (B) CXCL10, (C) IL-17A, (D) CCL4 and (E) IL-17F was

determined 4 days post-OVA challenge by Legendplex. Data presented as mean ±

SEM, P<0.05 compared with OVA challenge + BerriQi® Boysenberry and apple juice

concentrate, P<0.01 compared with naïve and OVA challenge (one-way ANOVA with

Tukey's Post Hoc test) for two experimental replicates with n=10 per treatment

groups.

SUPPLEMENTARY DATA

Supplementary Figure 1. Liquid Chromatography-Mass Spectrometry (LC-MS)

chromatogram for BerriQiTM concentrate, showing the anthocyanin UV/VIS profile

measured at 520 nm. Peak numbers refer to compounds listed in Table 1.

Supplementary Figure 2. Liquid Chromatography-Mass Spectrometry (LC-MS)

chromatograms for BerriQiTM concentrate, showing PDA (photodiode array)

phenolic profiles measured at 200–600 nm, 280 nm and 360 nm. Peak numbers

refer to compounds listed in Table 1. X, denotes sorbic acid, a preservative added to

the BerriQi concentrate during formulation. This was present at a concentration of

~2000 µg/mL.

A

B

C

D

E

F